

G-Computation

Malcolm Barrett

2021-09-01

Normal regression estimates associations. But we want *causal* estimates: what would happen if *everyone* in the study were exposed to x vs if *no one* was exposed.

G-Computation/G-Formula

- 1 Fit a model for $y \sim x + z$ where z is all covariates
- 2 Create a duplicate of your data set for each level of x
- 3 Set the value of x to a single value for each cloned data set (e.g $x = 1$ for one, $x = 0$ for the other)

G-Computation / G-Formula

Advantages of the parametric G-formula

Often more statistically precise than propensity-based methods

Incredibly flexible

Basis of other important causal models, e.g. causal survival analysis and TMLE

Greek Pantheon data (greek_data)

The name of a Greek god	A prognostic factor	The treatment, a heart transplant	The outcome, death
Rheia	0	0	0
Kronos	0	0	1
Demeter	0	0	0
Hades	0	0	0
Hestia	0	1	0
Poseidon	0	1	0
Hera	0	1	0
Zeus	0	1	1
Artemis	1	0	1
Apollo	1	0	1

+ 10 more rows

1. Fit a model for $y \sim a + 1$

```
1 greek_model <- lm(y ~ a + 1, data = greek_data)
```

2. Create a duplicate of your data set for each level of a

The name of a Greek god	A prognostic factor	The treatment, a heart transplant	The outcome, death
Rheia	0	0	0
Kronos	0	0	1
Demeter	0	0	0
Hades	0	0	0
Hestia	0	1	0
Poseidon	0	1	0
Hera	0	1	0
Zeus	0	1	1
Artemis	1	0	1
Apollo	1	0	1

2. Create a duplicate of your data set for each level of a

The name of a Greek god	A prognostic factor	The treatment, a heart transplant	The outcome, death
Rheia	0	0	0
Kronos	0	0	1
Demeter	0	0	0
Hades	0	0	0
Hestia	0	1	0
Poseidon	0	1	0
Hera	0	1	0
Zeus	0	1	1
Artemis	1	0	1
Apollo	1	0	1

The name of a Greek god	A prognostic factor	The treatment, a heart transplant	The outcome, death
Rheia	0	0	0
Kronos	0	0	1
Demeter	0	0	0
Hades	0	0	0
Hestia	0	1	0
Poseidon	0	1	0
Hera	0	1	0
Zeus	0	1	1
Artemis	1	0	1
Apollo	1	0	1

3. Set the value of **a** to a single value for each cloned data set

The name of a Greek god	A prognostic factor	a	The outcome, death
Rheia	0	0	0
Kronos	0	0	1
Demeter	0	0	0
Hades	0	0	0
Hestia	0	0	0
Poseidon	0	0	0
Hera	0	0	0
Zeus	0	0	1
Artemis	1	0	1
Apollo	1	0	1

The name of a Greek god	A prognostic factor	a	The outcome, death
Rheia	0	1	0
Kronos	0	1	1
Demeter	0	1	0
Hades	0	1	0
Hestia	0	1	0
Poseidon	0	1	0
Hera	0	1	0
Zeus	0	1	1
Artemis	1	1	1
Apollo	1	1	1

3. Set the value of **a** to a single value for each cloned data set

```
1 # set all participants to have a = 0
2 untreated_data <- greek_data |>
3   mutate(a = 0) #<<
4
5 # set all participants to have a = 1
6 treated_data <- greek_data |>
7   mutate(a = 1) #<<
```

4. Make predictions using the model on the cloned data sets

```
1 # predict under the data where everyone is untreated
2 predicted_untreated <- greek_model |>#<<
3   augment(newdata = untreated_data) |>#<<
4   select(untreated = .fitted)
5
6 # predict under the data where everyone is treated
7 predicted_treated <- greek_model |>#<<
8   augment(newdata = treated_data) |>#<<
9   select(treated = .fitted)
10
11 predictions <- bind_cols(
12   predicted_untreated,
13   predicted_treated
14 )
```

5. Calculate the estimate you want

```
1 predictions |>
2   summarise(
3     mean_treated = mean(treated),
4     mean_untreated = mean(untreated),
5     difference = mean_treated - mean_untreated #<<
6   )
```

A tibble: 1 × 3


	mean_treated	mean_untreated	difference
	<dbl>	<dbl>	<dbl>
1	0.5	0.5	0

Your Turn


Work through Your Turns 1-3 in `07-g-computation-exercises.qmd`

Detour: Colliders, selection bias, and loss to follow-up


Confounders and chains



Colliders



Colliders



Let's prove it!

```
1 set.seed(1234)
2 collider_data <- collider_triangle() |>
3   simulate_data(-.6)
```


Let's prove it!

```
1 collider_data
```

```
# A tibble: 500 × 3
```


	m	x	y
	<dbl>	<dbl>	<dbl>
1	-0.457	-0.410	1.28
2	0.281	1.79	-0.550
3	0.0835	1.31	-0.169
4	0.640	1.06	-1.40
5	-1.30	0.435	1.16
6	-0.569	0.630	-0.000667
7	-0.793	1.50	-1.10
8	-0.482	0.748	-0.411
9	-0.706	1.03	-0.381
10	1.42	-0.841	-0.420
"			

Let's prove it!



correct effect size: 0

Loss to follow-up



Adjusting for selection bias

- 1 Fit a probability of censoring model,
e.g. *glm(censoring ~ predictors, family
= binomial())*
- 2 Create weights using inverse
probability strategy
- 3 Use weights in your causal model

We won't do it here, but you can include many types of weights in a given model. Just take their product, e.g. *multiply inverse propensity of treatment weights by inverse propensity of censoring weights.*

Your Turn

Work through Your Turns 4-6 in 07-g-computation-exercises.qmd

