

When Standard Methods Succeed

Lucy D'Agostino McGowan

Wake Forest University

2022-07-05 (updated: 2022-10-16)

when correlation **is** causation

When you have no confounders and there is a linear relationship between the exposure and the outcome, that **correlation is a causal relationship**



When you have no confounders and there is a linear relationship between the **exposure** and the outcome, that correlation is a causal relationship



When you have no confounders and there is a linear relationship between the exposure and the **outcome**, that correlation is a causal relationship



When you have no **confounders** and there is a linear relationship between the exposure and the outcome, that correlation is a causal relationship



randomized controlled trials

A/B testing

**Even in these cases, using the
methods you will learn here can
help!**

- 1 Adjusting for baseline covariates can make an estimate **more efficient**
- 2 Propensity score weighting is **more efficient** than direct adjustment
- 3 Sometimes we are **more comfortable with the functional form of the propensity score (predicting exposure) than the outcome model**

Example

simulated data (100 observations)

Example

simulated data (100 observations)

Treatment is **randomly** assigned

Example

simulated data (100 observations)

Treatment is **randomly** assigned

There are **two baseline covariates**:
age and weight

Unadjusted model

```
lm(y ~ treatment, data = data)
```

Characteristic	Beta	SE ¹	95% CI ¹	p-value
treatment	1.6	0.803	-0.04, 3.1	0.056

¹ SE = Standard Error, CI = Confidence Interval

Adjusted model

```
lm(y ~ treatment + weight + age,
```

```
### Warning in modify_column_unhide(.  
### std.error): partial argument match  
### 'columns'
```

```
### Warning in match.call(definition,  
### envir): partial argument match of  
### 'columns'
```

Characteristic	Beta	SE ¹	95% CI ¹	p-value
treatment	1.5	0.204	1.1, 1.9	<0.001
weight	0.18	0.103	-0.03, 0.38	0.087
age	0.20	0.005	0.19, 0.21	<0.001

¹ SE = Standard Error, CI = Confidence Interval

Propensity score adjusted model

Characteristic	Beta	SE	95% CI	p-value
----------------	------	----	--------	---------

Example

simulated data (10,000 observations)

Treatment is **randomly** assigned

There are **two baseline covariates**:
age and weight

Unadjusted model

```
lm(y ~ treatment, data = data)
```

Characteristic	Beta	SE ¹	95% CI ¹	p-value
treatment	0.89	0.082	0.73, 1.1	<0.001

¹ SE = Standard Error, CI = Confidence Interval

Adjusted model

```
lm(y ~ treatment + weight + age,
```

```
### Warning in modify_column_unhide(.  
### std.error): partial argument match  
### 'columns'
```

```
### Warning in match.call(definition,  
### envir): partial argument match of  
### 'columns'
```

Characteristic	Beta	SE ¹	95% CI ¹	p-value
treatment	1.0	0.020	1.0, 1.0	<0.001
weight	0.19	0.010	0.17, 0.21	<0.001
age	0.20	0.001	0.20, 0.20	<0.001

¹ SE = Standard Error, CI = Confidence Interval

Propensity score adjusted model

Characteristic	Beta	SE	95% CI	p-value
----------------	------	----	--------	---------

time-varying confounding