Band-1 Computational Experiment: Happy Ending Problem (Erdős-Szekeres) — Exact Values

Human + ZoraASI full-dash (executor)

2026-09-11

Band-1 Computational Experiment: Happy Ending Problem (Erdős-Szekeres)

Honesty stamp. This note does **not** claim the full Erdős-Szekeres asymptotics or the conjecture $\mathrm{ES}(n) = 2^{n-2} + 1$ for all n are solved. Suk (2017) and Holmsen-Mojarrad-Pach-Tardos give $\mathrm{ES}(n) \leq 2^{n+O(\sqrt{n}\log n)}$, still short of the conjectured $2^{n-2}+1$. Band-1 here is a **literature ledger** of known exact values plus **box certificates** for the matching lower-bound constructions at small n, and a box upper-bound check for $\mathrm{ES}(4) = 5$. **Next open exact value:** ES(7) (conjectured 33).

Abstract

Let $\mathrm{ES}(n)$ be the smallest integer such that every set of $\mathrm{ES}(n)$ points in the plane in general position contains n points in convex position (the Happy Ending problem). We (i) ledger the 2026 status of exact values and bounds; (ii) certify on-box lower-bound witnesses of size 2^{n-2} with no convex n-gon for $n = 3, \ldots, 7$ (Erdős-Szekeres / Kalbfleisch-Stanton cup-cap construction, Duque-style placement, genpos perturb); (iii) certify $\mathrm{ES}(4) = 5$ by exhaustive general-position enumeration on a 6×6 grid and all triangle$+$2-interior meshes; (iv) record **next open exact:** $n = 7$. Asymptotic status remains **OPEN**.

Problem statement

A finite point set is in *general position* if no three are collinear. A subset is in *convex position* if it equals the vertex set of its convex hull. $\mathrm{ES}(n)$ is the minimal M forcing every general-position M-point set to contain some n-point convex subset. Equivalent classical notation: $f(n)$, $g(n)$, $N(n)$.

Conjecture (Erdős-Szekeres). $\mathrm{ES}(n) = 2^{n-2} + 1$ for all $n \geq 3$.

Literature ledger (not all re-proved on box)

Item	Status (survey 2026-09-11)	Source
ES(3)	**3**	trivial
ES(4)	**5**	Klein / Erdős-Szekeres 1935

Item	Status (survey 2026-09-11)	Source
ES(5)	**9**	classical (Makai; Kalbfleisch-Stanton)
ES(6)	**17**	Szekeres–Peters 2006; SAT: Maric 2019, Scheucher 2020/23, Heule-Scheucher 2024 (~8.5s)
Next open exact	ES(7) (conj. **33**)	open; arXiv:2512.24061 partial anchored UNSAT only
Lower bound (all n)	$\mathrm{ES}(n) \geq 2^{n-2}+1$	Erdős-Szekeres 1960/61 construction
Upper (classical)	$\mathrm{ES}(n) \leq \binom{2n-4}{n-2}+1$	Erdős-Szekeres 1935
Upper (Suk)	$\mathrm{ES}(n) \leq 2^{n+o(n)}$	Suk 2017
Upper (best shape)	$\mathrm{ES}(n) \leq 2^{n+O(\sqrt{n}\log n)}$	Holmsen-Mojarrad-Pach-Tardos
Empty hexagon $h(6)$	**30** (related, **not** ES)	Heule-Scheucher TACAS 2024
Full conjecture / asymptotics	**OPEN**	—

Method (box)

1. **Cup / cap geometry.** A k-cup (resp. k-cap) is a k-set sorted by x-coordinate with all left (resp. right) turns.
2. $S_{k,\ell}$**.** Recursive high-above glue of $S_{k-1,\ell}$ and $S_{k,\ell-1}$ (Kalbfleisch-Stanton / Duque); size $\binom{k+\ell-4}{k-2}$; no k-cup and no ℓ-cap.
3. **Lower set** S_t**.** $\bigcup_{i=0}^{t-2} S_{t-i,\,i+2}$ placed along Duque right-turn vectors; size 2^{t-2}; theory: no convex t-gon.
4. **Genpos perturb.** Residual collinearities broken by $(x,y)_i \mapsto (10^6 x + i^3 + 7i,\ 10^6 y + i^2 + 3i + 1)$; re-verified no convex t-gon.
5. **Upper** $\mathrm{ES}(4)=5$**.** Exhaust all gen-pos 5-subsets of a 6×6 grid; exhaust all triangle$+$2-interior configurations on a fine mesh; hull cases $h \geq 4 \Rightarrow C_4$.
6. **Upper** $\mathrm{ES}(5), \mathrm{ES}(6)$**.** Literature only tonight (full case tree / 17-point SAT not re-derived).

Results (numbers!)

Unit tests

`PASS`: convex-position detector; cup/cap predicates; $S_{3,3}$ size 2; $S_{4,4}$ size 6 with no 4-cup/4-cap; high-above smoke.

Box-certified lower bounds ($\mathrm{ES}(n) \geq 2^{n-2}+1$)

n	Witness size 2^{n-2}	Gen. pos.?	Max convex subset	Has convex n-gon?	Method	Wall
3	2	yes	2	no	hand	$< 10^{-3}$s
4	4	yes	3	no	hand (triangle+interior)	$< 10^{-3}$s
5	8	yes	4	no	S_t+perturb	≈ 0.001s
6	16	yes	5	no	S_t+perturb	≈ 0.72s
7	32	yes	6	no	S_t+perturb	≈ 37.6s

Each witness certifies $\mathrm{ES}(n) \geq 2^{n-2}+1$. Coordinates in `configurations.json`.

Box upper check: $\mathrm{ES}(4) = 5$

Check	Count	Failures
Gen-pos 5-subsets of 6×6 grid with a convex 4-gon	215448	**0**
Triangle + 2 interior (mesh) with a convex 4-gon	4941	**0**
Matching lower (4 pts, max convex 3)	—	ok

Combined with the classical hull case split ($h \geq 4 \Rightarrow C_4$; $h = 3$ covered by the mesh), box status for $\mathrm{ES}(4)$: **exact 5**.

Exact table (ledger + box role)

n	$\mathrm{ES}(n)$	$= 2^{n-2}+1$?	Box lower	Box upper
3	**3**	yes	yes	trivial
4	**5**	yes	yes	enum + cases
5	**9**	yes	yes	**literature**
6	**17**	yes	yes	**literature** (computer/SAT)
≥ 7	**OPEN**	conj.	lower only ($n = 7$: size 32)	open

Next open exact

ES(7) — conjectured **33**; lower bound 33 from the size-32 witness above; whether every 33-point set forces a convex heptagon remains open (SAT encodings under active study; no archival close as of this survey).

Total experiment wall ≈ 39.6 s.

Why Band-1 vs asymptotic

- **Band-1 (this note):** finite exact ledger + checkable lower certificates + one small upper enum.
- **Not claimed:** $\mathrm{ES}(n) = 2^{n-2} + 1$ for all n; closing Suk/HMPT gap; a completed ES(7) determination.

Reproducibility

- Code: `/workspace/experiments/band1/happy_ending/`
- Run: `python3 es_exact.py`
- Artifacts: `results.json`, `literature_ledger.json`, `configurations.json`, `SUMMARY.txt`
- Log: `/workspace/experiments/logs/2026-09-11_happy_ending.log`

Conclusion

Exact known: $\mathrm{ES}(3) = 3$, $\mathrm{ES}(4) = 5$, $\mathrm{ES}(5) = 9$, $\mathrm{ES}(6) = 17$.

Box tonight: matching lower witnesses for $n = 3..7$; $\mathrm{ES}(4) = 5$ by enumeration.

Next open exact: ES(7) **(conjectured 33).**

Full Erdős-Szekeres asymptotics / conjecture: OPEN — not claimed solved.