Process Cookbook

 $^{2}$   $\dots^{1}$ 

3 ....

Author Note

5 ....

1

6 Correspondence concerning this article should be addressed to ..., .... E-mail: ...

7 Abstract

8 Begin here...

9 Keywords: ....

Word count: 95

11

12

19

31

# Process Cookbook

# Hook thoughts

# 3 Hook from last meeting

- A "it is confusing and we would like to organize" argument.
- New researchers are increasingly asked to collect longitudinal data. We care about longitudinal data.
- The longitudinal literature appears more confusing than it is because people propose different hypotheses, use different models, and make different inferences.
  - We organize and discuss the similarities.

# 20 Bliese paper hook

- A "this other technique can augment what you are doing" argument.
- We increasingly have access to longitudinal data that contain discontinuities.
- Examples.
- These studies use a variety of techniques to understand discontinuous patterns, but
  we believe that discontinuous growth models can augment existing techniques...
- Practical advice is spread across numerous sources and is basic we review.

# 27 Aguinis paper hook

- A "this literature is technical and mathy, not accessible to org
- researchers" argument.
  - We care about A or B.
- There is a large technical literature on A or B.

- Much of this work is not easily accessible to researchers with the usual methodological and statistical background obtained from doctoral level training in management.
- We provide recommendations.

## 35 Thoughts from last meeting

36

38

# A no argument but describing paper.

- We care about longitudinal data
  - Our literature now contains a lot of cool examples and inferences
- We are going to describe them
- This paper is for newcomers who want some examples and for longitudinal modelers
  who want other options to augment what they are doing
- Organizational researchers are increasingly interested in longitudinal data structures. 42 These data align with the processes we study, which "are not static but instead develop, 43 change, and evolve over time" (Pitariu & Ployhart, p. 405) or are "sequences of events that play out within each person's stream of experience" (Beal, 2015, p. 5). If researchers can collect longitudinal data, they are then in a better position to understand patterns over time. Bell and Kozlowski (2003), for example, state that "longitudinal designs... will be far more revealing of the team phenomeon under investigation" (p. 59). Similarly – in their review of emotional labor – Grandey and Gabriel (2015) suggest that future researchers may be much better equipped to demonstrate emotion regulation patters "with longitudinal methods" (p. 329). Not only do longitudinal data collections help researchers observe relationships over time, many argue that doing so helps improve our theories. For instance, Pitariu and Ployhart (2010) suggest that collecting longitudinal data, observing change, and testing dynamic models "stimulates greater refinement in our existing theories" (p. 411), and Grandey and Gabriel (2015) urge for greater theoretical development of 55

emotions by studying their "reciprocal and unfolding...processes...through momentary

assessments or lagged effects" (p. 322). These quotes reveal our field's emphasis on
 collecting longitudinal data to sufficiently observe a process and build better theory.

A greater emphasis on longitudinal data is also apparent in our empirical literature,
where collecting repeated measures is now common. For instance, Jones et al. (2016)
observed the work attitudes of pregnant women in their second trimester every week until
they gave birth. (???) assessed job satisfaction across three time points that were each
separated by six weeks. Matthews, Wayne, and Ford (2014) studied work-family conflict
across several months. Meier and Spector (2013) examined counterproductive work
behavior over five waves. Hardy, Day, and Steele (2018) investigated self-regulation over 20
lab trials. Gabriel, Diefendorff, Chandler, Moran, and Greguras (2014) measured fit and
affect five times per day for 10 consecutive work days. Finally, Johnson, Lanaj, and Barnes
(2014) observed justice behavior and resource depletion across 10 consecutive workdays.

Armed with repeated observations, these studies then explore a variety of interesting inferences. Hardy et al. (2018) propose and find support for growth trends in self-regulation, metacognition, and performance. Jones et al. (2016) conclude that concealing behaviors among pregnant women lead to greater subsequent physical health. Johnson et al. (2014) establish that "justice is dynamic: The frequency of actors' justice behaviors varies day to day" (p. 10), and these daily fluctuations predict daily changes in depletion. Finally, Meier and Spector (2013) suggest that the effects of work stressors on counterproductive work behaviors are not substantially different across different time lags.

Moreover, we find different types of sophisticated modeling strategies throughout.

Meier and Spector (2013) present a sequence of path models that test increasingly longer

time lags. Hardy et al. (2018) and Jones et al. (2016) employ bivariate cross-lagged latent

growth curves, an approach similar to the latent change model used by (???). We also find

complex hierarchical linear models in various event-sampling studies (e.g., ???; Rosen,

Koopman, Gabriel, & Johnson, 2016).

In summary, our increasing theoretical emphasis on longitudinal data and its growing prevalence in our empirical literature has led to many interesting inferences and model applications. Researchers are beginning to propose hypotheses with lags and examine the reciprocal nature between two or more variables. Studies now commonly make inferences about change, growth, and dynamics. And sophisticated models, such as multi-level and latent growth curves, make their appearance in almost every longitudinal study.

We want to bring attention to this new and exciting literature. Specifically, in this 89 paper we describe some of the common inferences and models researchers employ with 90 longitudinal data, provide examples from our literature, point out the strengths and 91 weaknesses of each approach, and discuss recommendations. Many of these studies appear 92 unrelated due to the different language used among various content or statistical modeling 93 areas – but we believe our discussion will reveal some of their covert similarities. Our paper is meant for two audiences. First, we believe this summary will help any newcomers who want to make inferences with their longitudinal data but are unaware of what types of inferences and models are available. Second, this paper is meant to show seasoned researchers other possibilities – other inferences and models that may augment how they currently approach longitudinal data.

Below, we discuss four common inferences researchers make with longitudinal data:
relationships over time, growth, change, and dynamics. These do not represent every
possible domain we can explore with longitudinal data, but they do cover a large portion of
the techniques currently employed in our literature. We begin by explaining what we mean
by longitudinal research. We then unpack a framework proposed by Xu and DeShon that
relates the four different research streams. In the core of the paper, we discuss each stream
in depth, provide examples from the literature, point researchers to potential models, and
acknowledge the pitfalls and limitations of each approach.

108 Intro

# 109 Define longitudinal

This paper is exclusively devoted to the inferences we make with repeated 110 observations, so we begin by identifying a few labels and definitions. Authors typically 111 identify a "longitudinal" study by making a contrast with respect to either a) research 112 designs or b) data structures. Longitudinal research is different from cross-sectional 113 research because longitudinal designs entail three or more repeated observations (Ployhart 114 & Bliese, Singer & Willett). We therefore emphasize differences on the number of 115 observations when we distinguish longitudinal from other types of research. Longitudinal 116 data are repeated observations on several units (i.e., N or i > 1), whereas panel data are 117 observations of one unit over time – a distinction that focuses on the amount of people in 118 our study (given repeated measures). Most organizational studies collect data on more 119 than one unit, therefore our discussion below focuses on longitudinal research with 120 longitudinal data, or designs with N > 1, t >= 3, and the same construct(s) measured on 121 (potentially) each i at (potentially) each t.

#### 123 Introduce framework


124

125

126

Presenting the entire inference and modeling literature that uses longitudinal data would be impossible. Instead, we focus on four related streams that we feel can be organized nicely using a framework proposed by Xu and DeShon. Figure one shows each inference we will discuss in this paper: relationships, growth, change, and dynamics.

In each panel in figure one, time is on the x-axis to portray that we are investigating these inferences over time. Each slice contains an observation of y, such that at time t we observe  $y_t$  and at t+1 we observe  $y_{t+1}$ . What differentiates the panels – the inferences – is the pattern of relationships we investigate – and we add complexity as we move from A to D. For example, researchers do not include lag effects when they are interested in relationships over time (panel A), but they do include a lag effect when they study change or dynamics (panels C and D). We will devote a section to each of these inferences below, but we first describe some preliminary pieces about code and data that we refer to repeatedly in this paper.



# 8 Introduce data sets and generice variables

137

We will use two generic variables – affect (x) and performance (y) – throughout this paper. These variables will hopefully provide continuity across the inferences and also provide an illuminating backdrop after we refer to x or y generically. Moreover, we use

code examples in each section that refer to data sets that contain measures of affect and performance on 50 simulated subjects across five time points.

These data are contained in two data sets: long\_df and wide\_df. The raw data are
the same, but their formatting differs because different estimation techniques require
different data structures. Structural equations modeling (SEM) requires wide data, whereas
HLM requires long data. The first data set, long\_df contain the data in long form,

affect	performance	id	time
10.269606	17.23310	1	1
9.370015	22.81949	2	1
10.868660	17.64258	3	1
11.727196	20.56741	4	1
10.024188	12.08548	5	1
10.368025	13.17514	6	1
8.690796	12.76671	7	1
10.738622	13.62823	8	1

148

151

where id refers to a person identifier, time refers to the observation period, and scores on affect and performance are listed in their respective columns.

The second data set, wide\_df contain the data in wide form,

152

id	affect.1	performance.1	affect.2	performance.2	affect.3	performance.3
1	10.269606	17.23310	9.945866	13.653924	9.059233	20.944278
2	9.370015	22.81949	10.461612	10.886965	10.093910	16.578068
3	10.868660	17.64258	9.403230	9.386411	10.066253	10.834513
4	11.727196	20.56741	11.263258	20.491540	9.009344	8.547953
5	10.024188	12.08548	8.854671	16.052982	9.927885	16.759775
6	10.368025	13.17514	11.084624	13.839404	9.776743	26.595386
7	8.690796	12.76671	8.471005	13.229533	9.857223	10.831559
8	10.738622	13.62823	8.426258	16.364850	10.535407	8.794605

where id is defined above and now affect and performance are given new columns at
each measurement period – such that, as an example, person one reports 10.3 for affect at
time one and 9.9 for affect at time two. Again, we will refer to these variables and data sets
as we unpack each section, but remember that the values within each data set are the same

- the only difference is their format.

### 58 Introduce Code and Its Use

Finally, we will also present code snippets that estimate models related to each inference below. Code is typically published in methods literature to demonstrate how to estimate models, but it can also be used as a tool – a language – to build a greater understanding of the phenomenon (nature of code cite). We would like to use it for both reasons. Our goal is to provide readers with code for estimating models, but also allow them to see the inferences represented in various ways – words and code – to help clarify what they represent. The code snippets will take consistent forms throughout. HLM models will be expressed as follows:

```
hlm_model <- lme(

code
more code

data = long_df
)</pre>
```

where hlm\_model is an object that stores the results of our model, lme is a function call to
a linear mixed-effects regression (HLM), and within that we specify our effects and
reference our data. In the examples below, code and more code will be replaced by the
effects we specify, and data will always reference long\_df because HLM requires long
formatted data. Finally, after storing our results in the object hlm\_model we can actually
view them with:

```
summary(hlm_model)
```

Similarly, SEM models will be expressed as:

where sem\_model is an object that stores the results of our SEM model and we refer to the
wide\_df data set because SEM requires wide data. Notice that in the SEM code snippet we
first create a string object, sem\_string to specify our effects. Again, the code and more
code will be replaced by actual effects when we get to our inferences. Just like the HLM
code, we then view our SEM results with:

### summary(sem model)

In summary, the style of code snippets just presented will be used throughout each inference section. What will change are the code and more code pieces, and in those areas we will impute effects specific to each inference. If you wish to run these models on your own computer, you will also need to load their packages in your r script with load(nlme) for HLM and load(lavaan) for SEM.

We now turn to the inferences...

## Relationships

# 186 Explanation

184

185

A common inference in our longitudinal literature is the relationship between an 187 outcome (y) and one or more predictors  $(x_p)$  over time. As shown in figure one, researchers 188 observe y and one or more predictors at each time point, and are then interested in the 189 immediate effect of the predictors on y. Although there are multiple slices in our figure, 190 typically the effect of x on y is treated as stable over time and therefore the analysis 191 returns an estimate of a single parameter (e.g., a single beta weight). This parameter is 192 essentially a summary statement of the immediate effect of x on y at any point in time. In 193 other words, researchers observe x and y at every t from time t to t+5, as an example, and 194 then report a statement of the effect of  $x_t$  on  $y_t$ , or the expected immediate effect of x on y195 at any possible moment. 196

Although the effect (i.e., the parameter relating x to y) is treated as stable over time, the values on x and y – the raw data – are typically allowed to vary. When the values of x(potentially) change at each observation, the analysis is referred to as a time-varying covariates analysis – whereas a time-invariant analysis would be one where the raw data on x does not change at each observation (e.g., gender).

To clarify, consider this inference with respect to our generic variables: affect x and 202 performance y. Researchers would collect performance and affect data on multiple people 203 at each observation over several time points. Affect may vary at each observation – it may 204 be 5 at time t and 9 at time t+1 – so this analysis is referred to as a time-varying 205 covariates analysis. Researchers are then interested in the stable, immediate influence of 206 affect on performance. That is, they are interested in the effect of affect, on performance, 207 at any potential t, where, on average, they expect the effect of affect on performance to be 208 close to their parameter value. 209

There is a distinction with how we use the term "stable" that merits more 210 explanation. In a relationships inference, "stable" means that we expect the parameter 211 value relating affect to performance to be the same at each moment. In other words, the 212 relationship between affect and performance will be the same at each t – high values of 213 affect will result in high values of performance (if the parameter is positive) and low values 214 of affect will result in low values of performance. "Stable" in this inference context does not mean that affect has a lasting impression on performance, or that affect at time t influences 216 performance at some later time. This distinction is a difficult one, but it represents a major 217 difference between making a relationships inference versus some of the others we have yet 218 to explore. A relationships inference, therefore, is concerned with the average influence over 219 time, or what we expect the immediate effect of x on y to be at any given moment. 220

### 1 Code

We gain even more clarity on this inference when we consider the code used to
estimate models. A typical SEM model would be estimated with something similar to the
following code.

```
performance.1 ~ b1*affect.1

performance.2 ~ b1*affect.2

performance.3 ~ b1*affect.3

performance.4 ~ b1*affect.4

performance.5 ~ b1*affect.5
```

where this code snippet is identical to our introductory SEM code but we have replaced code and more code with actual effects we wish to estimate. In this case, we regress performance at time t on affect at time t and estimate b1 – the parameter relating affect to performance over time. The analysis will return one number for b1 because, again, we are treating this as a stable estimate over time.

The HLM equivalent would be:

230

```
lme(

fixed = performance ~ affect,

random = ~1|id,
```

```
data = long_df
)
```

where we again replace code and more code from the introductory HLM snippet with a
regression of performance on affect. Notice that we do not need to type the performance on
affect regressions at each time point as we did with the SEM code. Typically in SEM
software you type out the entire model, whereas HLM packages use condensed code. There
are two other new pieces of code: fixed and random. These commands specify how we
want HLM to estimate the parameters we are interested in, but at this point we do not
need to worry about them.

We will devote an entire section to the statistical properties of HLM and SEM below,
what is important here is our regression of performance on affect. In both HLM and SEM
we estimate a stable parameter relating affect at time t to performance at time t – in SEM
software we explicitly identify a parameter (b1) whereas HLM software typically does so
automatically.

### Hypotheses

249

Barnes, Schaubroeck, Huth, and Ghumman (2011) and Chi, Chang, and Huang
(2015) present hypotheses that are consistent with a relationships inference. Barnes et al.
(2011) predict a negative relationship between poor sleep and cognitive self control.
Similarly, Chi et al. (2015) hypothesize that daily negative mood negatively relates to daily
task performance.

### $Longitudinal\ Data\ Structure.$

Researchers that infer relationships over time either a) measure their variables at the same time points or b) treat the data as if they were measured as so by the constraints of

their analysis. Barnes et al. (2011) give an example of the former; they measure sleep and cognitive self control every morning for five consecutive days (study 4). Chi et al. (2015), as an example of the latter, measure negative mood in the morning and task performance in the afternoon – but these variables are treated as if they were taken at the same time (e.g., day 4) when entered into their analysis (described later).

There is no limit to the number of possible time points, but studies of this type 257 typically collect between three and ten repeated measurements. Our examples above 258 measure their variables once per day, but other frequencies could also be used. Methods 259 literature typically recommends that researchers maintain the same interval between each 260 measurement across their study, but this can be difficult when studies last longer than a 261 week. For example, Chi et al. (2015) take their measurements for ten work days. In the 262 middle of those assessments, however, their sample employees go home for the weekend and 263 are not measured. The space, therefore, between t and t+1 is not the same across their analysis. Finally, our organizational literature almost always takes measures on more than 265 one unit across time.

#### Models.

267

Both Barnes et al. (2011) and Chi et al. (2015) use hiearchical linear modeling 268 (HLM) – also known as multi-level or random coefficient models – to estimate the 269 parameters that generate their observed data. Random coefficient models assume that 270 their random effects are distributed  $N(\gamma, \sigma^2)$  – but this statement is easier to understand 271 with a contrast. Imagine that everyone in the sample receives their own  $X_t$  on  $Y_t$  effect and we then summarize those effects by taking their average. HLM, conversely, estimates one  $X_t$  on  $Y_t$  effect across units (people) but does so by imposing a normality constraint. 274 Finally, Barnes et al. (2011) and Chi et al. (2015) use models that fall into the broader 275 time-varying analysis class because their measurements of X are allowed to vary over time. 276 In time-invariant analyses, conversely, values of the covariate remain stable (e.g., gender). 277

Pitfalls and Recommendations. What can go wrong. Recommendation. Why.

Growth Growth

#### 80 Inference 1

278

281

282

283

288

293

What is the level of a construct at a given point in time?

# Examples.

# Hypotheses.

There are few (if any) studies that make direct predictions about construct levels at certain time points. Almost every longitudinal study, however, estimates an intercept term to approximate the answer. These terms are then discussed in the context of broader growth inferences that we discuss below.

# Longitudinal Data Structure.

This inference requires several repeated measures on one variable. For example, Zhu,
Wanberg, Harrison, and Diehn (2016) measure work adjustment once a month for nine
months (among expatriates) and Jones et al. (2016) measure concealing behaviors (among
pregnant women) every week – but it is not clear for how long they do so.

#### Models.

Zhu et al. (2016) use HLM to estimate their parmaters, whereas Jones et al. (2016)
do so in a structural equation modeling framework. As stated, researchers are interested in
an intercept estimate for this inference, and although they can place it on any time point,
researchers typically use their first observation. In SEM, the intercept is a latent variable
regressed on either an observed indicator (again, usually the first observation) or its latent
equivalent. Jones et al. (2016), for example, regress their latent concealing behaviors
intercept term on the latent variable for concealing behaviors at time one. In HLM, the
investigator first creates a variable that represents time and then regresses the outcome on

this time indicator to estimate an intercept term. For example, Zhu et al. (2016) regress work adjustment on time using HLM to estimate its initial level.

Pitfalls and Recommendations. What can go wrong. Recommendations. Why.

### OF Inference 2

304

308

309

313

317

What are the patterns of a trajectory or trajectories across time? How does the construct change over time?

# Examples.

## Hypotheses.

Zhu et al. (2016) predict that expatriate work adjustment follows a positive trajectory, increasing over the time of the assignment. Similarly, Jones et al. (2016) hypothesize that concealing behaviors will have negative slopes over time.

# Longitudinal Data Structure.

As with our previous inference, the only requirement here is a repeated assessment on one variable over several time points – both Zhu et al. (2016) and Jones et al. (2016) meet these requirements.

# Models.

Researchers use slope terms in their models for making inferences about trajectory patterns. In HLM, the same procedure we described above – regressing the outcome on time – also provides a slope estimate. Typically researchers do not allow their slope estimates to vary over units, but the models can easily support random slope terms. Zhu et al. (2016), for example, estimate a random slope effect by regressing work adjustment on time. In SEM, a latent slope term is regressed onto all time points except for the time point that was used by the latent intercept term. Jones et al. (2016) regress a latent concealing behaviors slope term on latent concealing behaviors at all time points beyond  $t_1$ .

Researchers may also be interested in curvilinear slopes. The only additional requirement for doing so is to create other time indicators with respect to the higher order polynomial. Zhu et al. (2016), for example, included linear, quadratic, and cubic time variables to estimate different slope curves. Note that these are still "linear" processes because they are linear in parameters.

**Pitfalls and Recommendations.** What can go wrong. Recommendations. Why.

#### Inference 3

331

333

334

335

339

Are there between person differences in trajectories or levels?

### Examples.

# Hypotheses.

Similar to our first growth inference, researchers rarely make a direct prediction
about between person differences in trajectory or level, but it is almost always included as
part of a larger growth analysis.

#### $Longitudinal\ Data\ Structure.$

We now require repeated observations on multiple units to examine between person (unit) differences. For example, Zhu et al. (2016) collect data from 179 expatriates across nine consecutive months.

#### Models.

Models that estimate intercept or slope terms also estimate a variance on those
terms, and these are then subjected to significance tests. When the variance on the
intercept is significant, that is taken to mean that there are between unit differences in
intercept (levels), whereas significant variance estimates on the slope terms indicates
between unit differences in slope (trajectory). Zhu et al. (2016) use HLM to estimate the

variance on their intercept and slope terms and find evidence of between person differences in work adjustment intercept (level) and slope (trajectory).

**Pitfalls and Recommendations.** What can go wrong. Recommendations. Why.

### Inference 4

351

355

356

364

372

How is the level of a construct related to the trajectory of the same or different constructs? What is the correlation between intercept and slope terms?

## Examples.

# Hypotheses.

Schaubroeck, Lam, and Peng (2016) predict that initial levels of peer leader's transformational leadership will positively relate to the slope of employee beliefs. Zhu et al. (2016) hypothesize that initial level of expatriate work adjustment is negatively related to the speed of change in work adjustment. Notice that the first prediction is about how the level of one construct is related to the slope of another, whereas the second focuses only on a single variable. Although it is uncommon in our literature, developmental researchers will also predict associations between slope and the final – rather than initial – observation.

#### $Longitudinal\ Data\ Structure.$

Data structures that researchers use to examine this inference are consistent with what we have already reviewed. Schaubroeck et al. (2016) measured transformational leadership and employee beliefs at three time points. Their time two assessment was 10 weeks after their first observation, whereas their final measurement was taken 10 months after their second observation. That is, the data structure for their respective time points were t, t + 10, and t + 40. As stated above, Zhu et al. (2016) collected data once a month for nine months.

#### Models.

Both HLM and SEM provide estimates of the covariance between the intercept and 373 slope terms discussed above, and this covariance indicates the association between level 374 and trajectory. For example, Zhu et al. (2016) report a negative covariance estimate 375 between the initial level of expatriate work adjustment (intercept) and its linear trajectory 376 over time (slope). As stated, researchers can also examine this association but among the 377 level of one variable and the trajectory of a different variable. Schaubroeck et al. (2016) 378 report an estimate of the relationship between transformational leadership intercept and 379 employee beliefs slope. It is often helpful to generate predicted values using the analysis 380 estimates and then plot the levels and trajectories to interpret them appropriately. 381

**Pitfalls and Recommendations.** What can go wrong. Recommendations. Why.

### 83 Inference 5

382

386

387

395

What inter-individual characteristics relate to intra-individual differences in level or slope?

# Examples.

#### Hypotheses.

Inferences of this type take the form of stable unit effects and their relationship with trajectory or level. Li, Burch, and Lee (2017) predict a positive relationship between job strain and job complexity trajectory. Zhu et al. (2016) hypothesize that the trajectory of expatriate work adjustment is positively related to perceived career instrumentality. Finally, Jones et al. (2016) make two inferences of this type: contextual support is a) negatively related to average levels of concealing but b) positively related to the slope of concealing.

### Longitudinal Data Structures.

We discussed the data structures for Zhu et al. (2016) and Jones et al. (2016) above.

Li et al. (2017) use a similar design but their data are collected once per year. Job

complexity is measured once every year for three years, and job strain is measured once at the final time point (year 3).

#### Models.

400

409

412

413

414

Both Li et al. (2017) and Zhu et al. (2016) use HLM to estimate their parameters, 401 whereas Jones et al. (2016) do so with SEM. In these models, the intercept or slope terms 402 discussed above are regressed onto the inter-individual characteristic. For example, Li et al. 403 (2017) report the estimate of job strain predicting job complexity trajectory and Zhu et al. 404 (2016) do the same but for career instrumentality and job complexity trajectory. In the 405 SEM framework, Jones et al. (2016) report estimates of contextual support predicting 406 concealing slope and intercept. Note that these models fit into the broader time-invariant 407 analysis class that contrast with the time-variant covariates analyses discussed earlier. 408

**Pitfalls and Recommendations.** What can go wrong. Recommendations. Why.

410 Change

### Inference 1

How are the changes of one variable associated with changes in another over time?

# Examples.

### Hypotheses.

Change inferences are similar to relationship inferences in that they relate a predictor to a response at the same time point, but the researcher now wants to know whether the predictor is associated with an increase or decrease of the outcome. For example, Johnson et al. (2014) predict that, within individuals, exhibiting daily procedural justice behavior is associated with an increase in resource depletion. Similarly, Lanaj, Johnson, and Lee (2016) hypothesize that, within individuals, transformational leadership is associated with a decrease in negative affect.

422

431

444

446

## Longitudinal Data Structures.

Johnson et al. (2014) measured justice behavior and resource depletion in the
afternoon of 10 consecutive workdays. Lanaj et al. (2016) measured their variables at
different frequencies. They measured transformational leadership and negative affect in the
afternoon of 15 consecutive workdays, but they also took an additional measurement of
negative affect every morning – creating a data structure of three variables: negative affect
from t to t + 15, transformational leadership from t to t + 15 and morning negative affect
from t to t + 15. That is, they essentially doubled the time points of negative affect – from t to t + 30 by measuring it twice as frequently.

#### Models.

Johnson et al. (2014) and Lanaj et al. (2016) employ HLM change models that are 432 consistent with our literature's preference for using partialed values of the outcome variable 433 rather than difference scores. In models of this type, researchers regress their outcome on 434 the predictor at the same time point and the prior value of the outcome variable. For 435 example, Johnson et al. (2014) regressed resource depletion at time t on 1) procedural 436 justice behavior at time t and 2) resource depletion at time t-1; they then report the 437 relationship between justice behavior and resource depletion change. Similarly, Lanaj et al. 438 (2016) regress negative affect at time t on 1) transformational leadership at time t and 2) 439 negative affect at time t-1 (i.e., morning negative affect) and report the association between transformational leadership and affect change. Note that, due to the different data 441 sampling strategies, Johnson et al. (2014) reports change from the prior day, whereas Lanaj et al. (2016) report change from the morning to afternoon.

**Pitfalls and Recommendations.** What can go wrong. Recommendations. Why.

### 445 Dynamics

The core concept is: how are prior states related to future states?

### Inference 1

- How is one variable related to its future self?
- Examples.
- Hypotheses.
- ${\it Longitudinal\ Data\ Structures.}$
- Models.
- Pitfalls and Recommendations.

### 454 Inference 2

- How is one variable related to a different variable in the future?
- Examples.
- Hypotheses.
- Gabriel et al. (2014) hypothesize that perceived P-O fit is positively related to subsequent positive affect.
- ${\it Longitudinal\ Data\ Structure}.$
- Gabriel et al. (2014) measured perceived P-O fit and positive affect five times per day for 10 consecutive work days.
- Models.
- Researchers that use models related to this inference regress their outcome at time t on a prior value of the independent variable. Gabriel et al. (2014) do so using HLM and regress perceive P-O fit at time t on 1) positive affect at time t-1 and 2) perceived P-O fit at time t-1.
- Pitfalls and Recommendations. What can go wrong. Recommendations. Why.

### Inference 3

471

472

479

485

Directionality – what is the relationship direction among a set of variables?

# Examples.

# Hypotheses.

A directionality inference is typically explored using a pair of dynamic hypotheses.

For example, Hardy et al. (2018) predict that 1) prior self efficacy negatively relates to

subsequent meta cognition and 2) prior meta cognition positively relates to subsequent self

efficacy. Similarly, Matthews et al. (2014) contrast two theories, one that predicts that

work-family conflict is negatively related to subsequent well-being, whereas the other

predicts that well-being is negatively related to work-family conflict.

## $Longitudinal\ Data\ Structures.$

Hardy et al. (2018) measured self efficacy and meta cognition after five evenly spaced computer game trails – although they also measured baseline self efficacy. Matthews et al. (2014) assessed work-family conflict and well-being concurrently at three points in time: t, and t+7 where the intervals are months – meaning that the final survey was assessed six months after the survey before it.

# Models.

Both Hardy et al. (2018) and Matthews et al. (2014) use an SEM approach to
estimate their parameters, but Matthews et al. (2014) restrict their model to a path
analysis. Their modeling strategies are the same as those just presented, but they examine
them as a set to explore the relationship direction among their variables. Matthews et al.
(2014) find significant effects in both directions – WFC predicting subsequent well-being
and well-being predicting subsequent WFC – and therefore suggest that the relationship
works in both directions. Hardy et al. (2018) also report the estimates of self efficacy
predicting subsequent meta cognition and vice versa, but only find significant effects for the

latter. They conclude that "self-regulatory processes (i.e., exploration and meta cognition)
positively influenced subsequent self-efficacy... In contrast, we only found limited support
for the notion that self-efficacy significantly influences subsequent self-regulated learning
processes" (p. 24).

**Pitfalls and Recommendations.** What can go wrong. Recommendations. Why.

#### Inference 4

502

503

511

512

513

Reciprocal – Is the pattern of relationships reciprocal, such that one variable influences another, and this latter variable then goes back to influence the first?

# Examples.

## Hypotheses.

Kaltiainen, Lipponen, and Holtz (2017) present a set of hypotheses to explore
reciprocal dynamics among justice perceptions and trust. They predict that "process
justice perceptions and cognitive trust have positive reciprocal relations over time.

Specifically, planning stage (time 1) process justice (cognitive trust) perceptions will have a
positive relationship with subsequent post-merger (time 2) cognitive trust (process justice)
perceptions, which in turn will have a positive relationship with later post-merger (time 3)
process justice (cognitive trust) perceptions" (p. idk).

#### Longitudinal Data Structure.

They assess cognitive trust and justice perceptions once a year for three years.

# Models.

Kaltiainen et al. (2017) evaluate a number of models but find best fit for a model that includes reciprocal relationships among justice and trust. In that model, trust at time t is regressed on trust at time t-1 and justice at time t-1, and justice at time t is regressed on trust at time t-1 and justice at time t-1. They find significant cross-lag effects for trust on justice at all times but only one significant cross-lag effect of justice on trust, and ultimately conclude that "the justice  $\Rightarrow$  trust  $\Rightarrow$  justice reciprocal relationship was supported but the trust  $\Rightarrow$  justice  $\Rightarrow$  trust relationship was not supported as the relationship between justice at time 2 and cognitive trust at time 3 was not statistically significant" (p. 14).

Pitfalls and Recommendations. What can go wrong. Recommendations. Why.

### Inference 5

523

525

526

527

528

529

530

531

532

534

535

537

Mediation – one variable influences an outcome through an intermediate variable

## Things to notice and questions

- 1) Not going example by example, I am combining them.
  - Originally I thought it would be more clear this way, but I could be wrong. I do
    not have enough examples to pair every inference with its own unique empirical
    example.
- 2) Pitfalls and recommendations after each inference (as I did here...) or after each entire inference section (i.e., one for relationships, one for growth, etc.)?
- 3) "We" or "they" phrasing?
  - I.e., "we as a field typically do this..." or "researchers tend to do this..."
  - Not for stylistic reasons but for authoritative reasons.
  - 4) Several issues that came up while writing the data structures sections
  - 1) They are repetitive

538

539

540

541

542

543

544

545

546

547

548

549

551

556

• 2) Each example has their own issue (e.g., spacing between t and t + 1 isn't consistent, or the variables were measured at different frequencies). If I bring them up out of the blue it is really easy to lose the reader.

# • Solutions

- I could write one data structure piece per inference section rather than one per inference
- I could skip data structures in the inference examples just provide
   hypotheses and models and then give them their own section later
- I could give pictures like figure 1 in Xu and DeShon in each inference example section and then give data structures and issues their own section later. That is, in the examples I simply speak about hypotheses and models as if every study measured their variables from t to t+10 on the same interval. Then, at the end I come back with data collection problems/discrepancies I noticed in the articles. I like this option.

552 Extra

This emphasis has translated into a growing number of longitudinal studies in our empirical literature. This interest is apparent not only in reviews and theoretical discussions, but also in the increasing

Longitudinal studies are also increasingly common in our empirical literature.

References

- Barnes, C. M., Schaubroeck, J., Huth, M., & Ghumman, S. (2011). Lack of sleep and
   unethical conduct. Organizational Behavior and Human Decision Processes, 115(2),
   169–180.
- Chi, N.-W., Chang, H.-T., & Huang, H.-L. (2015). Can personality traits and daily
  positive mood buffer the harmful effects of daily negative mood on task performance
  and service sabotage? A self-control perspective. Organizational Behavior and
  Human Decision Processes, 131, 1–15.
- Gabriel, A. S., Diefendorff, J. M., Chandler, M. M., Moran, C. M., & Greguras, G. J.

  (2014). The dynamic relationships of work affect and job satisfaction with

  perceptions of fit. *Personnel Psychology*, 67(2), 389–420.
- Hardy, J. H., Day, E. A., & Steele, L. M. (2018). Interrelationships among self-regulated
   learning processes: Toward a dynamic process-based model of self-regulated
   learning. Journal of Management, 0149206318780440.
- Johnson, R. E., Lanaj, K., & Barnes, C. M. (2014). The good and bad of being fair:

  Effects of procedural and interpersonal justice behaviors on regulatory resources.

  Journal of Applied Psychology, 99(4), 635.
- Jones, K. P., King, E. B., Gilrane, V. L., McCausland, T. C., Cortina, J. M., & Grimm, K. J. (2016). The baby bump: Managing a dynamic stigma over time. *Journal of Management*, 42(6), 1530–1556.
- Kaltiainen, J., Lipponen, J., & Holtz, B. C. (2017). Dynamic interplay between merger process justice and cognitive trust in top management: A longitudinal study.

  Journal of Applied Psychology, 102(4), 636.
- Lanaj, K., Johnson, R. E., & Lee, S. M. (2016). Benefits of transformational behaviors for leaders: A daily investigation of leader behaviors and need fulfillment. *Journal of*

- Applied Psychology, 101(2), 237.
- Li, J., Burch, T. C., & Lee, T. W. (2017). Intra-individual variability in job complexity
  over time: Examining the effect of job complexity trajectory on employee job strain.

  Journal of Organizational Behavior, 38(5), 671–691.
- Matthews, R. A., Wayne, J. H., & Ford, M. T. (2014). A work–family conflict/subjective
   well-being process model: A test of competing theories of longitudinal effects.
   Journal of Applied Psychology, 99(6), 1173.
- Meier, L. L., & Spector, P. E. (2013). Reciprocal effects of work stressors and
   counterproductive work behavior: A five-wave longitudinal study. Journal of
   Applied Psychology, 98(3), 529.
- Rosen, C. C., Koopman, J., Gabriel, A. S., & Johnson, R. E. (2016). Who strikes back? A daily investigation of when and why incivility begets incivility. *Journal of Applied*Psychology, 101(11), 1620.
- Schaubroeck, J., Lam, S. S., & Peng, A. C. (2016). Can peers' ethical and transformational
   leadership improve coworkers' service quality? A latent growth analysis.
   Organizational Behavior and Human Decision Processes, 133, 45–58.
- Zhu, J., Wanberg, C. R., Harrison, D. A., & Diehn, E. W. (2016). Ups and downs of the
  expatriate experience? Understanding work adjustment trajectories and career
  outcomes. *Journal of Applied Psychology*, 101(4), 549.