Asymptotic Methods

Part II

Recorded from the lecturer in 2026-2027.

Pre-requisites

- Complex methods / analysis.
- Refer to Section II in Stuart.

1. Intro & Motivation

- Most real problems don't have closed form solutions, or if they do, they are unworkable or difficult to interpret.
- Numerical approximations can be useful, but they can be expensive, they may have bad numerical properties, or solutions may be inscrutable.
- Taylor series, etc. can be limited in applicability.
- Asymptotic methods emphasise usefulness, interpretability as well as accuracy.

Examples

e.g. 1)

$$\Gamma(z) = \int_0^\infty t^{z-1} e^{-t}\, dt, \qquad \mathrm{Re}(z) > 0.$$

For n an integer,

$$\Gamma(n+1) = n! \sim \sqrt{2\pi}\, n^{n+\frac{1}{2}} e^{-n} \left(1 + \frac{1}{12n} + \frac{1}{288n^2} + \cdots\right) \qquad \text{as } n \to \infty.$$

e.g. 2)

$$x^2 y'' + xy' - x^2 y = 0.$$

Exact series solution:

$$I_0(x) = \sum_{n=0}^{\infty} \frac{x^{2n}}{2^{2n}(n!)^2}.$$

Liouville–Green:

$$\implies I_0(x) \sim (2\pi x)^{-1/2} e^x \qquad \text{as } x \to \infty.$$

e.g. 3)

$$y'' + k^2 e^{-x^2/2} y = 0,$$

with

$$y \to y_0 \qquad \text{as } x \to \infty,$$

$$y \to \pm y_0 \qquad \text{as } x \to -\infty.$$

$$k = ?, \qquad k = k_0, k_1, k_2, \ldots$$

$$\implies k_n \sim \frac{\sqrt{\pi}}{2}\left(n + \frac{1}{2}\right) \qquad \text{as } n \to \infty.$$

WKBJ.

- Convergent methods typically give arbitrarily accurate approximations with more & more terms/iterations.
- Asymptotic methods give arbitrarily accurate approximations as some variable approaches a limit. Thus they always come with a qualifier, e.g. $z \to 0$ or $n \to \infty$.

2. Asymptotic Expansions

2.1 Definitions & Notation

(a) Asymptotic Equality/Equivalence

For 2 functions $f(x)$ and $g(x)$, we have

$$f(x) \sim g(x) \qquad \text{about } x = x_0$$

if

$$\boxed{\lim_{x \to x_0} \frac{f(x)}{g(x)} = 1.}$$

Remarks:

- Usually we want to approximate f with g.
- We can write
$$f(x) = \underbrace{g(x)}_{\text{dominant}} + \underbrace{[f(x) - g(x)]}_{\text{subdominant}}.$$
- Note $\sim$ is an equivalence relation (that is reflexive, transitive, symmetric).
- It respects products:
$$f_1 \sim g_1, \qquad f_2 \sim g_2$$
$$\implies f_1 f_2 \sim g_1 g_2.$$
- It is linear:
$$\alpha f_1 + \beta f_2 \sim \alpha g_1 + \beta g_2,$$
where α, β are constants.
- Definition has problems when dealing with oscillatory functions that go through 0 as $x \to x_0$.

(b) "Little o"

Suppose f & g are real functions.

We write

$$\boxed{f(x) = o(g(x)) \qquad \text{as } x \to x_0}$$

if

$$\boxed{\lim_{x \to x_0} \frac{f(x)}{g(x)} = 0.}$$

Remark:

$$f(x) = o(g(x))$$

can also be written as

$$f(x) \ll g(x) \qquad \text{as } x \to x_0.$$

- Can be framed in inequalities, e.g. $x_0 = \infty$ and let $g > 0$.
  Then $f = o(g)$ as $x \to \infty$ is equivalent to

$$\forall \varepsilon > 0, \quad \exists a \quad \text{s.t. when } x > a,$$

$$\boxed{|f| \leq \varepsilon g.}$$

(c) "Big O"

We write

$$f(x) = O(g(x)) \qquad \text{as } x \to x_0$$

for $g > 0$, if

$$\exists C > 0 \quad \text{s.t.}$$

$$\boxed{|f(x)| \leq C g(x)}$$

for x sufficiently close to x_0.

Warning:

In some areas of physics people use big-O and $\sim$ interchangeably.

Little Exercises: Show That

(a)

$$f(x) \sim g(x) \iff f - g = o(g) \qquad \text{as } x \to x_0.$$

(b)

$$\sin x \ll x^2 \qquad \text{as } x \to \infty.$$

(c)

$$12x^5 + 6x^3 + x = O(x^5) \qquad \text{as } x \to \infty.$$

(d)

$$e^{ax} \ll e^{bx} \qquad \text{as } x \to \infty,$$

if $b > a > 0$.

(e)

$$e^x \gg x^N \qquad \text{as } x \to \infty,$$

$$\forall N \in \mathbb{Z}^+.$$

2.2 Asymptotic Expansions

- Have unworkable/unknown $f(x)$.
- Want to know its behaviour in a limit.
- Want to find a simpler/known $g(x)$ so that

$$f \sim g.$$

- Also want to break up g into a hierarchy of approximations, s.t.

$$g(x) = \sum_{k=0}^{N} a_k \phi_k(x),$$

where a_k are coefficients, ϕ_k are known & workable, and where adding more terms gives a better approximation that can be estimated.

(a) Asymptotic Sequence

A sequence of functions

$$\{\phi_0(x), \phi_1(x), \ldots\}$$

is an "asymptotic sequence" as $x \to x_0$ if

$$\boxed{\phi_{k+1}(x) \ll \phi_k(x) \qquad \text{as } x \to x_0, \quad \forall k.}$$

e.g. 1)

$$\phi_k = (x-a)^k \qquad \text{around } x = a.$$

Check:

$$\frac{\phi_{k+1}}{\phi_k} = x - a \to 0 \qquad \text{as } x \to a. \quad \checkmark$$

e.g. 2)

$$\phi_k = e^{-kx} x^k \qquad \text{as } x \to \infty \text{ or } x \to -\infty.$$

Check:

$$\frac{\phi_{k+1}}{\phi_k} = e^{-x} x.$$

$$e^{-x} x \to 0 \quad \text{as } x \to \infty \qquad \checkmark$$
$$e^{-x} x \to \infty \quad \text{as } x \to -\infty \qquad \times$$

e.g. 3)

$$\phi_k = \cos(nx) \qquad \text{as } x \to \infty.$$

$\times$

e.g. 4)

$$\{|x|, \log|x|, \log|\log|x||, \ldots\} \qquad \text{as } x \to \infty.$$

?

(b) Asymptotic Expansion

- Suppose $f(x)$ is a function we want to understand near $x = x_0$.
- Let $\{\phi_k\}_{k=0}^{\infty}$ be an asymptotic sequence.

Then, the finite sum

$$\sum_{k=0}^{N} a_k \phi_k(x)$$

for some coefficients, is an (N-term) asymptotic approximation of $f(x)$ as $x \to x_0$ if:

$$\boxed{f(x) = \sum_{k=0}^{N} a_k \phi_k(x) + o(\phi_N(x))} \qquad \text{as } x \to x_0.$$

Alternatively, we can write the above as

$$\boxed{\frac{f(x) - \sum_{k=0}^{N} a_k \phi_k(x)}{\phi_N(x)} \longrightarrow 0} \qquad \text{as } x \to x_0.$$

End of Lecture 1