Tower Breakers - The Final Battle



Our unsung tower-breaking heroes (players P_1 and P_2) only have one tower left, and they've decided to break it for a special game commemorating the end of 5 days of Game Theory! The rules are as follows:

- ullet P_1 always moves first, and both players always move optimally.
- ullet Initially there is 1 tower of height N.
- The players move in alternating turns. The moves performed by each player are different:
 - 1. At each turn, P_1 divides the current tower into some number of smaller towers. If the turn starts with a tower of height H and P_1 breaks it into $x \geq 2$ smaller towers, the following condition must apply: $H = h_1 + h_2 + \ldots + h_x$, where h_i denotes the height of the i^{th} new tower.
 - 2. At each turn, P_2 chooses some tower k of the x new towers made by P_1 (where $1 \le k \le x$). Then P_1 must pay k^2 coins to P_2 . After that, P_1 gets another turn with tower h_k and the game continues.
- ullet The game is over when no valid move can be made by P_1 , meaning that H=1.
- ullet P_1 's goal is to pay as few coins as possible, and P_2 's goal is to earn as many coins as possible.

Can you predict the number of coins that P_2 will earn?

Input Format

The first line contains a single integer, T, denoting the number of test cases. Each of the T subsequent lines contains a single integer, N, defining the initial tower height for a test case.


Constraints

- $1 \le T \le 100$
- $2 < N < 10^{18}$


Output Format

For each test case, print a single integer denoting the number of coins earned by $P_{\mathbf{2}}$ on a new line.

Sample Input



Sample Output



Explanation

Our players make the following moves:

1.
$$H = N = 4$$

- 1. P_1 splits the initial tower into ${f 2}$ smaller towers of sizes ${f 3}$ and ${f 1}.$
- 2. P_2 chooses the first tower and earns $\mathbf{1}^2=\mathbf{1}$ coin.

2. H = 3

- 1. P_1 splits the tower into ${f 2}$ smaller towers of sizes ${f 2}$ and ${f 1}$.
- 2. P_2 chooses the first tower and earns $\mathbf{1^2}=\mathbf{1}$ coin.

3. H = 2

- 1. P_1 splits the tower into ${\bf 2}$ smaller towers of size ${\bf 1}$.
- 2. P_2 chooses the second tower and earns $\mathbf{2^2}=\mathbf{4}$ coins.

The total number of coins earned by P_2 is 1+1+4=6 , so we print 6 on a new line.