House Location



Adam and Martha are planning to leave the city after their retirement and build a house in a huge land belonging to their family. To keep everyone happy, they want to build the house at a location having distance $_a*d1_$ from aunt Kimberly's house, where a is some ratio and d1 is the distance of that location to uncle Bob's house. Also, the house should be at a distance $_b*d2_$ from uncle Jack's house where b is some ratio and d2 is the distance of the location to aunt Janet's house.

You need to help them find the location of their house.

Input Format

The first line of input contains two integers *a* and *b* (the ratios above). In the next four lines, there are 4 pairs of integers that indicate the coordinates of Kimberly's, Bob's, Jack's, and Janet's houses, respectively.

Output Format

You must output the coordinate of house with exactly two points after decimal point (rounded to closest one hundredth). If there is no location satisfying the above constraints, output **Impossible!** If there are more than one possible locations, output a location with minimum x-coordinate and among the ones having the minimum x-coordinate, a location with minimum y-coordinate.

Constraints

```
1 < a, b <= 1000
```

-1000 <= all input coordinates <= 1000

Sample Input

```
3 4
4 0
0 0
-2 -4
-2 -1
```

Sample Output

-2.00 0.00

Explanation

As required, the point (-2.00, 0.00) has distance 2 from Bob's house and distance 3*2=6 from Kimberly's house. It also has distance 1 from Janet's house and distance 4*1=4 from Jack's house.