

Thesis

Thomas Wiskow Thorbjørnsen

February 14, 2021

1 Introduction

This is an introduction, welcome!

Introduce notation which will be used in text. A list of notation and description would be nice, so that the reader might scroll back up if something is unclear.

2 Triangulated Categories

Probably introduce this section, what is happening and what will be done etc. I can maybe say something about algebraic triangulated categories and topological triangulated categories, and explaining the name cone, fiber and cofiber.

2.1 Definition and First Properties

In this section \mathcal{T} denotes an additive category and $T : \mathcal{T} \rightarrow \mathcal{T}$ is an additive autoequivalence of \mathcal{T} , which is often called translation or suspension functor.

Definition 2.1. A sextuple is a collection (A, B, C, a, b, c) of objects $A, B, C \in \mathcal{T}$ and morphisms $a : A \rightarrow B$, $b : B \rightarrow C$, $c : C \rightarrow TA$. These sextuples can be drawn as diagrams in the following way:

$$A \xrightarrow{a} B \xrightarrow{b} C \xrightarrow{c} TA$$

A morphism between sextuples is a triple of morphism (α, β, γ) , where $\alpha : A \rightarrow A'$, $\beta : B \rightarrow B'$ and $\gamma : C \rightarrow C'$ such that the following diagram commutes:

$$\begin{array}{ccccccc} A & \xrightarrow{a} & B & \xrightarrow{b} & C & \xrightarrow{c} & TA \\ \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma & & \downarrow T\alpha \\ A' & \xrightarrow{a'} & B' & \xrightarrow{b'} & C' & \xrightarrow{c'} & TA' \end{array}$$

The naming convention of the sextuples isn't standardized, some literatures calls the sextuples for triangles instead [literature here, learn bibtex you lazy fuck]. This name arises from an alternate description of the diagrams given above. To remove confusion about the domain or codomain of the arrows, one arrow of the triangle is decorated with " $_T$ ". This decorator means that the functor T has to be applied to the corresponding edge of the arrow. Thus the c arrow points to TA , not A .

$$\begin{array}{ccc} A & \xrightarrow{a} & B \\ T \uparrow & & \uparrow T \\ C & \xleftarrow{b} & A \end{array} \quad \begin{array}{ccccccc} A & \xrightarrow{\phi_a} & A' & & & & \\ T \uparrow & \searrow a & & \swarrow a' & & & T \uparrow \\ & B & \xrightarrow{\phi_b} & B' & & & \\ & \swarrow b & & \searrow b' & & & \\ C & \xleftarrow{\phi_c} & C' & & & & \end{array}$$

A triangulated category is an additive category together with a translation functor T and a triangulation Δ consisting of sextuples. When a sextuple is an element of Δ it is usually called a distinguished triangle, an exact triangles or just a triangle. Note that if sextuples are referred to as triangles it is common to either call the elements of Δ for distinguished triangles or exact triangles. As this is not the case for this thesis these objects will be referred to as triangles.

Definition 2.2. A triangulation of an additive category \mathcal{T} with translation T is a collection Δ of triangles consisting of sextuples in \mathcal{T} satisfying the following axioms:

1. (TR1) Formation axiom

(a) A sextuple isomorphic to a triangle is a triangle.

(b) Every morphism $a : A \rightarrow B$ can be embedded into a triangle (A, B, C, a, b, c) :

$$A \xrightarrow{a} B \xrightarrow{b} C \xrightarrow{c} TA$$

(c) For every object A there is a triangle $(A, A, 0, id_A, 0, 0)$:

$$A \xrightarrow{id_A} A \xrightarrow{0} 0 \xrightarrow{0} TA$$

2. (TR2) Rotation axiom

For every triangle (A, B, C, a, b, c) there is a triangle (B, C, TA, b, c, Ta)

$$A \xrightarrow{a} B \xrightarrow{b} C \xrightarrow{c} TA \implies B \xrightarrow{b} C \xrightarrow{c} TA \xrightarrow{-Ta} TB$$

3. (TR3) Morphism axiom

Given the two triangles (A, B, C, a, b, c) (1) and (A', B', C', a', b', c') (2)

$$(1) \quad A \xrightarrow{a} B \xrightarrow{b} C \xrightarrow{c} TA \quad (2) \quad A' \xrightarrow{a'} B' \xrightarrow{b'} C' \xrightarrow{c'} TA'$$

and morphisms $\phi_A : A \rightarrow A'$ and $\phi_B : B \rightarrow B'$ such that the square (1) commutes, then there is a morphism $\phi_C : C \rightarrow C'$ (not necessarily unique) such that (ϕ_A, ϕ_B, ϕ_C) is a morphism of triangles (2).

$$(1) \quad \begin{array}{ccc} A & \xrightarrow{a} & B \\ \downarrow \phi_A & & \downarrow \phi_B \\ A' & \xrightarrow{a'} & B' \end{array} \quad (2) \quad \begin{array}{ccccccc} A & \xrightarrow{a} & B & \xrightarrow{b} & C & \xrightarrow{c} & TA \\ \downarrow \phi_A & & \downarrow \phi_B & & \downarrow \phi_C & & \downarrow T\phi_A \\ A' & \xrightarrow{a'} & B' & \xrightarrow{b'} & C' & \xrightarrow{c'} & TA' \end{array}$$

4. (TR4) Octahedron axiom

Given the triangles (A, B, C', a, x, x') (1), (B, C, A', b, y, y') (2) and $(A, C, B', b \circ a, z, z')$ (3)

$$(1) \quad A \xrightarrow{a} B \xrightarrow{x} C' \xrightarrow{x'} TA$$

$$(2) \quad B \xrightarrow{b} C \xrightarrow{y} A' \xrightarrow{y'} TB$$

$$(3) \quad A \xrightarrow{b \circ a} C \xrightarrow{z} B' \xrightarrow{z'} TA$$

then there exist morphisms $f : C' \rightarrow B'$ and $g : B' \rightarrow A'$, the following diagram commutes and the third row is a triangle:

$$\begin{array}{ccccccc}
T^{-1}B' & \xrightarrow{T^{-1}z'} & A & \xrightarrow{id_A} & A & & \\
\downarrow T^{-1}g & & \downarrow a & & \downarrow b \circ a & & \\
T^{-1}A' & \xrightarrow{T^{-1}y'} & B & \xrightarrow{b} & C & \xrightarrow{y} & A' \xrightarrow{y'} TB \\
& & \downarrow x & & \downarrow z & & \parallel id_{A'} \downarrow Tx' \\
& & C' & \xrightarrow{f} & B' & \xrightarrow{g} & A' \xrightarrow{Ti \circ y'} TC' \\
& & \downarrow x' & & \downarrow z' & & \\
& & TA & \xrightarrow{id_{TA}} & TA & &
\end{array}$$


A triangulated category is denoted as (\mathcal{T}, T, Δ) , where \mathcal{T} is the additive category, T is the translation and Δ is the triangulation. When \mathcal{T} is called a triangulated category, it should be understood as the triple given above.

Remark. The rotation axiom has a dual, and it can be thought of as a rotation in the opposite direction. This dual can be proved by the other axioms, so it is here omitted as an axiom. The dual rotation axiom goes as:


Given a triangle $A \xrightarrow{a} B \xrightarrow{b} C \xrightarrow{c} TA$, there is a triangle $T^{-1}C \xrightarrow{T^{-1}c} A \xrightarrow{a} B \xrightarrow{b} C$. To be able to prove this, some more lemmata are needed.

Remark. The final axiom is referred to as the octahedron axiom. By using the alternative description of the triangle diagram, it is possible to rewrite the diagram as an octahedron. The axiom can be restated as the following:

Given the triangles (A, B, C', a, x, x') (1), (B, C, A', b, y, y') (2) and $(A, C, B', b \circ a, z, z')$ (3)



then there exists morphisms $f : C' \rightarrow B'$ and $g : B' \rightarrow A'$, the following diagram commutes and the teal back face is a triangle.



Lemma 2.1. *Let (A, B, C, a, b, c) be a triangle, then $b \circ a = 0$*

Proof. By TR2 the triangle (A, B, C, a, b, c) can be rotated to (B, C, TA, b, c, Ta) .

$$\begin{array}{ccc} A & \xrightarrow{a} & B \\ \uparrow T & & \uparrow T \\ C & \xleftarrow{b} & TA \end{array} \quad \begin{array}{c} \downarrow c \\ \downarrow -Ta \end{array} \quad \Rightarrow \quad \begin{array}{ccc} B & \xrightarrow{b} & C \\ \uparrow T & & \uparrow T \\ TA & \xleftarrow{c} & TB \end{array}$$

The triangle exists $(C, C, 0, id_C, 0, 0)$ by TR1 and TR3 says there exists a morphism from TA to 0 making the diagram below commute.

$$\begin{array}{ccccccc} B & \xrightarrow{b} & C & \xrightarrow{c} & TA & \xrightarrow{-Ta} & TB \\ \downarrow b & & \downarrow id_C & & \downarrow 0 & & \downarrow Tb \\ C & \xrightarrow{id_C} & C & \xrightarrow{0} & 0 & \xrightarrow{0} & TC \end{array}$$

Thus $0 = Tb \circ -Ta = T(-ba) \Rightarrow b \circ a = 0$ as T is a translation. \square

Definition 2.3. An additive functor between triangulated categories $F : (\mathcal{T}, T, \Delta) \rightarrow (\mathcal{R}, R, \Gamma)$ is called exact or triangulated if there exist a natural isomorphisms $\alpha : FT \rightarrow RF$ such that $F(\Delta) \subseteq \Gamma$.

A functor $F : \mathcal{T} \rightarrow \mathcal{R}$ is called a triangle-equivalence if it is triangulated and an equivalence of categories. In this case \mathcal{T} and \mathcal{R} are called triangle-equivalent.

Definition 2.4. Let \mathcal{T} be a triangulated category and \mathcal{A} be an abelian category. A covariant functor $H : \mathcal{T} \rightarrow \mathcal{A}$ is called a homological functor if $\forall (A, B, C, a, b, c) : \Delta$ there is a long exact sequence in \mathcal{A} .

$$\begin{array}{ccc} A & \xrightarrow{a} & B \\ \uparrow T & & \uparrow T \\ C & \xleftarrow{b} & TA \end{array} \quad \begin{array}{c} \downarrow c \\ \downarrow -Ta \end{array} \quad \Rightarrow \quad \begin{array}{c} \dots \longrightarrow H(T^i A) \xrightarrow{H(T^i a)} H(T^i B) \xrightarrow{H(T^i b)} H(T^i C) \\ \hspace{10em} \downarrow H(T^i c) \\ \hspace{10em} \longrightarrow H(T^{i+1} A) \xrightarrow{H(T^{i+1} a)} H(T^{i+1} B) \xrightarrow{H(T^{i+1} b)} H(T^{i+1} C) \longrightarrow \dots \end{array}$$

Dually, a contravariant functor $H : \mathcal{T} \rightarrow \mathcal{A}$ is called cohomological if $\forall (A, B, C, a, b, c) : \Delta$ there is a long exact sequence in \mathcal{A} .

$$\begin{array}{ccc} A & \xrightarrow{a} & B \\ \uparrow T & & \uparrow T \\ C & \xleftarrow{b} & TA \end{array} \quad \begin{array}{c} \downarrow c \\ \downarrow -Ta \end{array} \quad \Rightarrow \quad \begin{array}{c} \dots \longleftarrow H(T^{i-1} A) \xleftarrow{H(T^{i-1} a)} H(T^{i-1} B) \xleftarrow{H(T^{i-1} b)} H(T^{i-1} C) \\ \hspace{10em} \downarrow H(T^{i-1} c) \\ \hspace{10em} \longleftarrow H(T^i A) \xleftarrow{H(T^i a)} H(T^i B) \xleftarrow{H(T^i b)} H(T^i C) \longleftarrow \dots \end{array}$$

Lemma 2.2. *Let $M : \mathcal{T}$ be any object of \mathcal{T} , then the represented functors $\mathcal{T}(M, -)$ is homological and $\mathcal{T}(-, M)$ is cohomological.*

Proof. Only the covariant case needs to be proved, as the contravariant case is dual. For $\mathcal{T}(M, -)$ to be homological, it has to create long exact sequences for every triangle in Δ . Let $(A, B, C, a, b, c) : \Delta$ be a triangle, then there can be extracted sequences in Ab for any $i : \mathbb{N}$.

$$\begin{array}{c}
A \\
\uparrow c \\
C
\end{array}
\begin{array}{c}
\searrow a \\
\swarrow b
\end{array}
B \implies \mathcal{T}(M, T^i A) \xrightarrow{T^i a_*} \mathcal{T}(M, T^i B) \xrightarrow{T^i b_*} \mathcal{T}(M, T^i C)$$

Observe that it is enough to prove that these types of diagrams are exact, as the other diagrams can be obtained by the rotation axiom, thus reducing it to same case. The goal is then to prove that $\text{Im}(T^i a_*) = \text{Ker}(T^i b_*)$.

Since $ba = 0$ it follows that $\text{Im}(T^i a_*) \subseteq \text{Ker}(T^i b_*)$. Assume that $f : \text{Ker}(T^i b_*)$, that is $f : M \rightarrow T^i B$ such that $b_*(f) = 0$. The current goal is to show that f factors through $T^i A$, as this means that $\text{Ker}(T^i b_*) \subseteq \text{Im}(T^i a_*)$. Note that since T is a translation, it is necessarily a right adjoint to the inverse translation; thus $\mathcal{T}(M, T^i B) \simeq \mathcal{T}(T^{-i} M, B)$, and by this assertion it suffices to assume that $f : T^{-i} M \rightarrow B$ such that $b \circ f = 0$. By TR1 and TR2 there exists triangles $(T^{-i} M, 0, T^{-i+1} M, 0, 0, -T^{-i+1} \text{id})$ and $(B, C, TA, b, c, -Ta)$.

$$\begin{array}{ccccccc}
T^{-i} M & \xrightarrow{0} & 0 & \xrightarrow{0} & T^{-i+1} M & \xrightarrow{-T^{-i+1} \text{id}} & T^{-i+1} M \\
\downarrow f & & \downarrow 0 & & \downarrow g & & \downarrow Tf \\
B & \xrightarrow{b} & C & \xrightarrow{c} & TA & \xrightarrow{-Ta} & TB
\end{array}$$

The left square commutes by the assumption, thus the morphism g exist by TR3, such that $-Ta \circ h = -Tf \circ T^{-i+1} \text{id} = -Tf \implies Ta \circ h = Tf$, thus $f = a \circ T^{-1} h$ asserting that f factors through A . \square

Lemma 2.3. *Let $(\phi_A, \phi_B, \phi_C) : (A, B, C, a, b, c) \rightarrow (A', B', C', a', b', c')$ be a morphism of triangles. If 2 of the maps are isomorphisms, then the last one is an isomorphism as well.*

$$\begin{array}{ccccccc}
A & \xrightarrow{a} & B & \xrightarrow{b} & C & \xrightarrow{c} & TA \\
\wr \downarrow \phi_A & & \wr \downarrow \phi_B & & \wr \downarrow \phi_C & & \wr \downarrow T\phi_A \\
A' & \xrightarrow{a'} & B' & \xrightarrow{b'} & C' & \xrightarrow{c'} & TA'
\end{array}$$

Proof. Without loss of generality, assume that ϕ_A and ϕ_B are the isomorphisms. This can be done as the rotation axiom reduce the other cases to this case. Then we have the following diagram:

$$\begin{array}{ccccccc}
A & \xrightarrow{a} & B & \xrightarrow{b} & C & \xrightarrow{c} & TA \\
\wr \downarrow \phi_A & & \wr \downarrow \phi_B & & \downarrow \phi_C & & \wr \downarrow T\phi_A \\
A' & \xrightarrow{a'} & B' & \xrightarrow{b'} & C' & \xrightarrow{c'} & TA'
\end{array}$$

By applying the functor $\mathcal{T}(C', -)$ we get the following diagram in Ab:

$$\begin{array}{ccccccccc}
\mathcal{T}(C', A) & \xrightarrow{a_*} & \mathcal{T}(C', B) & \xrightarrow{b_*} & \mathcal{T}(C', C) & \xrightarrow{c_*} & \mathcal{T}(C', TA) & \xrightarrow{Ta_*} & \mathcal{T}(C', TB) \\
\wr \downarrow (\phi_A)_* & & \wr \downarrow (\phi_B)_* & & \downarrow (\phi_C)_* & & \wr \downarrow (T\phi_A)_* & & \wr \downarrow (T\phi_B)_* \\
\mathcal{T}(C', A') & \xrightarrow{a'_*} & \mathcal{T}(C', B') & \xrightarrow{b'_*} & \mathcal{T}(C', C') & \xrightarrow{c'_*} & \mathcal{T}(C', TA') & \xrightarrow{Ta'_*} & \mathcal{T}(C', TB)
\end{array}$$

By the 5-lemma, we get that $(\phi_C)_*$ is an isomorphisms, i.e. $(\phi_C)_*$ is both mono and epi. Thus for some unique s in $\mathcal{T}(C', C)$, $\phi_{C*}(s) = id_{C'}$.

By applying the functor $\mathcal{T}(-, C)$ we get the diagram:

$$\begin{array}{ccccccccc}
\mathcal{T}(A, C) & \xleftarrow{a^*} & \mathcal{T}(B, C) & \xleftarrow{b^*} & \mathcal{T}(C, C) & \xleftarrow{c^*} & \mathcal{T}(TA, C) & \xleftarrow{Ta^*} & \mathcal{T}(TB, C) \\
(\phi_A)^* \uparrow \wr | & & (\phi_B)^* \uparrow \wr | & & (\phi_C)^* \uparrow & & (\phi_{TA})^* \uparrow \wr | & & (\phi_{TB})^* \uparrow \wr | \\
\mathcal{T}(A', C) & \xleftarrow{a'^*} & \mathcal{T}(B, C) & \xleftarrow{b'^*} & \mathcal{T}(C', C) & \xleftarrow{c'^*} & \mathcal{T}(TA', C) & \xleftarrow{Ta'^*} & \mathcal{T}(TB', C)
\end{array}$$

By the 5-lemma, we get that $(\phi_C)^*$ is an isomorphism. By the same argument $id_C = s' \circ \phi_C$ for some unique s' . ϕ_C is both split mono and split epi, which means it is an isomorphism. \square

Corollary 2.3.1. $(A, B, 0, a, 0, 0)$ is a triangle if and only if a is an isomorphism.

Proof. Assume that a is an isomorphism. Then it is seen that $(a, id_B, 0)$ is an isomorphism of triangles.

$$\begin{array}{ccccccc} A & \xrightarrow{a} & B & \xrightarrow{0} & 0 & \xrightarrow{0} & TA \\ \wr \downarrow a & & \wr \downarrow id_B & & \wr \downarrow 0 & & \wr \downarrow Ta \\ B & \xrightarrow{id_B} & B & \xrightarrow{0} & 0 & \xrightarrow{0} & TB \end{array}$$

Conversely, assume that $(A, B, 0, a, 0, 0)$ is a triangle. Then the same diagram as above can be constructed, and by the 2 out of 3 property, a has to be an isomorphism. \square

Lemma 2.4. *For a triangle (A, B, C, a, b, c) the following are equivalent:*

- a is split mono
- b is split epi
- $c = 0$

Proof. The proof has two parts. First assume that a is split mono, and prove that b is split epi, and $c = 0$. By duality, it is then known that b being split epi implies that a is split mono and $c = 0$. The final part is to assume that $c = 0$, and prove either a is split mono or b is split epi.

Assume that a is split mono, then there exist an a^{-1} such that $id_A = a^{-1}a$. Let $M : \mathcal{T}$ be any object, then we can make a long exact sequence:

$$\mathcal{T}(M, T^{-1}C) \xrightarrow{T^{-1}c_*} \mathcal{T}(M, A) \begin{array}{c} \xrightarrow{a_*} \\ \xleftarrow{a_*^{-1}} \end{array} \mathcal{T}(M, B) \xrightarrow{b_*} \mathcal{T}(M, C) \xrightarrow{c_*} \mathcal{T}(M, TA)$$

By assumption a_* is split mono, thus $T^{-1}c_* = 0$ and in particular $c = 0$. This implies that b_* is epi, making a split short exact sequence.

$$0 \xrightarrow{0} \mathcal{T}(M, A) \xrightarrow{a_*} \mathcal{T}(M, B) \xrightarrow{b_*} \mathcal{T}(M, C) \xrightarrow{0} 0$$

$\xleftarrow{a_*^{-1}}$ $\xleftarrow{b_*^{-1}}$

This gives that b is split epi, completing the first part of the proof.

For the next part, assume that $c = 0$; then we can construct the following triangles.

$$(1) \begin{array}{ccc} A & \xrightarrow{a} & B \\ T \uparrow & & \downarrow 0 \\ C & \xleftarrow{b} & \end{array} \implies \begin{array}{ccc} C & \xrightarrow{0} & TA \\ T \uparrow & & \downarrow -Tb \\ TB & \xleftarrow{-Ta} & \end{array}$$

$$(2) \begin{array}{ccc} A & \xrightarrow{id_A} & A \\ T \uparrow & & \downarrow 0 \\ 0 & \xleftarrow{0} & \end{array} \implies \begin{array}{ccc} 0 & \xrightarrow{0} & TA \\ T \uparrow & & \downarrow 0 \\ TA & \xleftarrow{-id_{TA}} & \end{array}$$

(1) is constructed by applying TR2 twice, while (2) is constructed with TR1 and TR2 twice. Observe that there is a commutative square between the triangles, allowing for TR3 to make a morphism of triangles.

$$\begin{array}{ccccccc} C & \xrightarrow{0} & TA & \xrightarrow{-Ta} & TB & \xrightarrow{-Tb} & TC \\ \downarrow 0 & & \parallel id_{TA} & & \downarrow Ta^{-1} & & \downarrow 0 \\ 0 & \xrightarrow{0} & TA & \xrightarrow{-id_{TA}} & TA & \xrightarrow{0} & 0 \end{array}$$

Thus $T(a^{-1}a) = id_{TA} = T(id_A) \implies id_A = a^{-1}a$, making a split mono. \square

Lemma 2.5. *Given two triangles (A, B, C, a, b, c) and (A', B', C', a', b', c') the following are equivalent:*

$$\begin{array}{ccccccc} A & \xrightarrow{a} & B & \xrightarrow{b} & C & \xrightarrow{c} & TA \\ \downarrow f & & \downarrow g & & \downarrow h & & \downarrow Tf \\ A' & \xrightarrow{a'} & B' & \xrightarrow{b'} & C' & \xrightarrow{c'} & TA' \end{array} \quad \begin{array}{l} 1. (f, g, h) \text{ is a morphism of triangles} \\ 2. \exists g : B \rightarrow B' \text{ such that } b'ga = 0 \end{array}$$

Moreover, if $\mathcal{T}(A, T^{-1}C') \simeq 0$, then f and h are unique.

Proof. 1. \implies 2. as the composition $ba = 0 = b'a'$, so assume 2. The existence of f and h is evident from the long exact sequence of the bottom triangle at the functor represented by A .

$$\mathcal{T}(A, T^{-1}C') \xrightarrow{T^{-1}c'_*} \mathcal{T}(A, A') \xrightarrow{a'_*} \mathcal{T}(A, B') \xrightarrow{b'_*} \mathcal{T}(A, C')$$

The morphism $ga : \mathcal{T}(A, B')$ such that $b'ga = b'_*(ga) = 0$, thus $ga : Ker(b'_*)$. By exactness $\exists f : \mathcal{T}(A, A')$ such that $a'_*f = ga$, and by TR3 $\exists h : C \rightarrow C'$ such that (f, g, h) is a morphism of triangles. Now assume that $\mathcal{T}(A, T^{-1}C') \simeq 0$. Exactness determines that a'_* is a monomorphism, and f is then unique. Since T is a translation, we have that $\mathcal{T}(A, T^{-1}C') \simeq \mathcal{T}(TA, C')$. By using the functor $\mathcal{T}(-, C')$ at the top triangle, we get that b^* is a monomorphism, and thus h is chosen uniquely. \square

Lemma 2.6. *If (A, B, C, a, b, c) is a triangle, then $(T^{-1}C, A, B, T^{-1}c, a, b)$ is a triangle.*

Proof. By TR2 we can construct a triangle.

$$\begin{array}{ccc}
A & \xrightarrow{a} & B \\
\uparrow T & & \downarrow c \\
C & \xleftarrow{b} &
\end{array}
\implies
\begin{array}{ccc}
C & \xrightarrow{c} & TA \\
\uparrow T & & \downarrow Tb \\
TB & \xleftarrow{Ta} &
\end{array}$$

By TR1 we can create a triangle $(T^{-1}C, A, B', T^{-1}c, a', b')$, and then use TR3 to find a morphism.

$$\begin{array}{ccccccc}
C & \xrightarrow{c} & TA & \xrightarrow{Ta'} & TB' & \xrightarrow{Tb'} & TC \\
\parallel id_C & & \parallel id_{TA} & & \downarrow h & & \parallel id_{TC} \\
C & \xrightarrow{c} & TA & \xrightarrow{Ta} & TB & \xrightarrow{Tb} & TC
\end{array}$$

By the 2 out of 3 property it is seen that h is an isomorphism, so the triple $(id_{T^{-1}C}, id_A, T^{-1}h)$ is an isomorphism of sextuples, and by TR1, is an isomorphism of triangles, asserting that $(T^{-1}C, A, B, T^{-1}c, a, b)$ is in fact a triangle. \square

Remark. The Octahedron axiom have not been used once in this section. This motivates the definition of a pre-triangulated category, that is a triangulation satisfying all axioms except TR4. Note that these properties also holds for pre-triangulated categories. However, the octahedron axiom is still important, since TR3 can be proved from the other 3 axioms. [Need sources here]

2.2 $K[x]/(x^n)$

In this section quotients of the module categories of the K-algebras $K[x]/(x^n)$ will be studied and proved to be triangulated. As a first example it would be more instructive with a more canonical triangulated category, such as the homotopy category of chain complexes. In this type of categories one would be able to see the underlying triangulated structure much more clearly, and how it interplays with long exact sequences. Never the less, this class of categories will be more instructive in showing how flexible the notion of a triangulated category may be.

The more skilled reader may notice that the algebra $K[x]/(x^n)$ is self-injective and is of finite length. Therefore the category of modules over $K[x]/(x^n)$ is a Frobenius category, and the stabilization admits a triangulation. What all of this means will be clarified in the next section, for now we will focus on the structure of this very specific category, and prove that it's stabilized module category has a triangulation.

Definition 2.5. The injective stable $\text{mod-}K[x]/(x^n)$ category is the quotient category of morphisms factoring over injective objects. This means that if I_1, I_2 is injective and $f, g : A \rightarrow B$ are morphisms factoring over I_1 and I_2 respectively, then f and g are related in the quotient.

The projective stable categories are defined in the same way, but with projective objects instead.

Remark. Cosyzygies are in fact a functor for injective stable categories, and syzygies are functors for the projective stable categories.

Write about the indecomposable objects in this category. Find the projective and injective modules, i.e. show that $K[x]/(x^n)$ is injective.

2.3 Localizations of Triangulated Categories

Verdier quotient goes here. And more stuff about these localizations, probably state Gabriel-Zisman and how this is often a good trick too prove that the localization actually exists.

2.4 Discussion of Triangulations

Maybe do some Yoneda-embedding into sheaves and use that to deduce how the triangulation are in fact a shadow of some other abelian category

3 Exact Categories

3.1 Definitions and First Properties

3.2 The Frobenis Category

3.3 The Homotopy Category

4 The Derived Category

4.1 Admissable Morphisms

4.2 Homology and Long Exact Sequences

4.3 The Derived Category

4.4 If time, derived functors as well

5 Auslander-Reiten Triangles

5.1 Krull-Schmidt Categories

5.2 Definition and First Properties

5.3 Description of Derived Categories

Appendix A: Category theory

5.4 Quotient Categories

5.5 Additive and Abelian Categories

5.6 Freyd-Mitchell Embedding???