

# On the Derived Category of Strongly Homotopy Associative Algebras

Thomas Wiskow Thorbjørnsen

August 31, 2022

## **Abstract**

*Fill inn abstract*

## **Sammendrag**

*Fyll inn sammendraget*

## **Acknowledgements**

*Thank the people in your life who has made this journey easier :D*

# Contents

<b>Contents</b>	<b>iii</b>
<b>1 Bar and Cobar Construction</b>	<b>1</b>
1.1 Preliminaries	2
1.1.1 Algebras	2
1.1.2 Coalgebras	6
1.1.3 Electric Circuits	10
1.1.4 Derivations and DG-Algebras	13
1.2 Cobar-Bar Adjunction	21
1.2.1 Convolution Algebras	21
1.2.2 Twisting Morphisms	24
1.2.3 Bar and Cobar Construction	25
1.3 Strongly Homotopy Associative Algebras and Coalgebras	32
1.3.1 SHA-Algebras	32
1.3.2 $A_\infty$ -Coalgebras	37
<b>2 Homotopy Theory of Algebras</b>	<b>39</b>
2.1 Model categories	40
2.1.1 Model categories	41
2.1.2 Homotopy category	44

2.1.3	Quillen adjoints . . . . .	54
2.2	Model structures on Algebraic Categories . . . . .	58
2.2.1	DG-Algebras as a Model Category . . . . .	58
2.2.2	A Model Structure on DG-Coalgebras . . . . .	63
2.2.3	Homotopy theory of $A_\infty$ -algebras . . . . .	70
2.3	The Homotopy Category of $Alg_\infty$ . . . . .	74
<b>3</b>	<b>Derived Categories of Strongly Homotopy Associative Algebras . . . . .</b>	<b>77</b>
3.1	Twisting Morphisms . . . . .	77
3.1.1	Twisted Tensor Products . . . . .	78
3.1.2	Model Structure on Module Categories . . . . .	81
3.1.3	Model Structure on Comodule Categories . . . . .	81
3.1.4	Triangulation of Homotopy Categories . . . . .	83
3.1.5	The Fundamental Theorem of Twisting Morphisms . . . . .	92
3.2	Polydules . . . . .	93
3.2.1	The Bar Construction . . . . .	93
3.2.2	Polydules of SHA-algebras . . . . .	95
3.2.3	Universal Enveloping Algebra . . . . .	96
3.2.4	The Derived Category $D_\infty A$ of Augmented $A_\infty$ Algebras . . . . .	97
3.3	The Derived Category $D_\infty A$ . . . . .	100
	<b>Bibliography . . . . .</b>	<b>101</b>

# Chapter 1

## Bar and Cobar Construction

In Stasheff's papers [1] and [2], a strongly homotopy associative algebra, or  $A_\infty$ -algebra, over a field is a graded vector space together with homogenous linear maps  $m_n : A^{\otimes n} \rightarrow A$  of degree  $2-n$  satisfying some homotopical relations. This will be made precise later. We may regard  $m_2$  to be a multiplication of  $A$ , it is however not a priori associative. The associator of  $m_2$  is taken to be the homotopical relation of  $m_3$ . Thus, we know that the homotopy of  $A$  is an associative algebra. The maps  $m_n$  corresponds uniquely to a map  $m^c : BA \rightarrow \bar{A}[1]$ , which extends to a coderivation  $m^c : BA \rightarrow BA$  of the bar construction of  $A$ . So we could instead define an  $A_\infty$ -algebra to be a coalgebra on the form  $BA$ .

In order to understand the bar construction we will first study it on associative algebras. Given a differential graded coassociative coalgebra  $C$  and a differential graded associative algebra  $A$ , we say that a homogenous linear transformation  $\alpha : C \rightarrow A$  is twisting if it satisfies the Maurer-Cartan equation:

$$\partial\alpha + \alpha \star \alpha = 0.$$

Let  $Tw(C, A)$  be the set of twisting morphisms, then considering it as a functor  $Tw : \mathbf{CoAlg}_{\mathbb{K}}^{op} \times \mathbf{Alg}_{\mathbb{K}} \rightarrow \mathbf{Ab}$  we want to show that it is represented in both arguments. Moreover, these representations give rise to an adjoint pair of functors, called the bar and cobar construction.

$$\begin{array}{ccc} & B & \\ \text{Alg}_{\mathbb{K},+}^{\bullet} & \xrightarrow{\quad} & \text{Coalg}_{\mathbb{K},\text{conil}}^{\bullet} \\ & \Omega & \end{array} \quad \top$$

The bar and cobar construction will be the basis for our discussion of  $A_\infty$ -algebras. As the bar construction can be used to define  $A_\infty$ -algebras, we may easily dualize this to define  $A_\infty$ -coalgebras in terms of the cobar construction. This chapter will follow the notations and progression presented in Loday and Vallette [3] to develop the theory for the bar-cobar adjunction.

## 1.1 Preliminaries

### 1.1.1 Algebras

This section is a review of associative algebras over a field  $\mathbb{K}$ . The main goal for this section is to establish some conventions and see how they may be dualized to coalgebras. This collection of algebras together with ring homomorphisms form the category  $\text{Alg}_{\mathbb{K}}$  of associative algebras over  $\mathbb{K}$ .

**Definition 1.1.1** ( $\mathbb{K}$ -Algebra). Let  $\mathbb{K}$  be a field with unit 1. A  $\mathbb{K}$ -algebra  $A$ , or an algebra  $A$  over  $\mathbb{K}$  is a vector space with structure morphisms called multiplication and unit,

$$\begin{aligned} (\cdot_A) : A \otimes_{\mathbb{K}} A &\rightarrow A \\ 1_A : \mathbb{K} &\rightarrow A, \end{aligned}$$

satisfying the associativity and identity laws.

$$\begin{aligned} \text{(associativity)} \quad (a \cdot_A b) \cdot_A c &= a \cdot_A (b \cdot_A c) \\ \text{(unitality)} \quad 1_A(1) \cdot_A a &= a = a \cdot_A 1_A(1) \end{aligned}$$

Whenever  $A$  does not possess a unit morphism, we will call  $A$  a non-unital algebra. In this case only the associativity law must hold.

By abuse of notation we will confuse the unit of  $\mathbb{K}$  by the unit of  $A$ . Since  $1_A$  is a ring homomorphism, this should not be a cause of confusion. However, when we want to use the unit as a morphism, we will stick to the  $1_A$  notation. When there are no confusion, we will also exchange  $(\cdot_A)$  concatenation of variables as strings.

**Definition 1.1.2** (Algebra homomorphisms). Let  $A$  and  $B$  be algebras. Then  $f : A \rightarrow B$  is an algebra homomorphism if

1.  $f$  is  $\mathbb{K}$ -linear
2.  $f(ab) = f(a)f(b)$
3.  $f \circ 1_A = 1_B$

Whenever  $A$  and  $B$  are non-unital, we must drop the condition for  $f$  to preserve units.

**Definition 1.1.3** (Category of algebras). We let  $\text{Alg}_{\mathbb{K}}$  denote the category of algebras. Its objects consists of every algebra  $A$ , and the morphisms are algebra homomorphisms. The sets of morphisms between  $A$  and  $B$  are denoted as  $\text{Alg}_{\mathbb{K}}(A, B)$ .

Let  $\widehat{\text{Alg}}_{\mathbb{K}}$  denote the category of non-unital algebras. Its objects consists of every non-unital algebra  $A$ , and the morphisms are non-unital algebra homomorphisms. The sets of morphisms between  $A$  and  $B$  are denoted as  $\widehat{\text{Alg}}_{\mathbb{K}}(A, B)$ .

Observe that for an algebra  $A$ , the triple  $(A, (\cdot_A), 1_A)$  is a monoid in  $Mod_{\mathbb{K}}$ . Thus the category of algebras corresponds to the category of monoids in  $Mod_{\mathbb{K}}$ . We reformulate the algebra axioms as the following commutative diagrams:

$$\begin{array}{ccc}
 A \otimes_{\mathbb{K}} A \otimes_{\mathbb{K}} A & \xrightarrow{(\cdot_A) \otimes id_{\mathbb{K}}} & A \otimes_{\mathbb{K}} A \\
 \downarrow id_{\mathbb{K}} \otimes (\cdot_A) & & \downarrow (\cdot_A) \\
 A \otimes_{\mathbb{K}} A & \xrightarrow{(\cdot_A)} & A
 \end{array}
 \qquad
 \begin{array}{ccccc}
 A \otimes_{\mathbb{K}} \mathbb{K} & \xrightarrow{id_A \otimes 1_A} & A \otimes_{\mathbb{K}} A & \xleftarrow{1_A \otimes id_A} & \mathbb{K} \otimes_{\mathbb{K}} A \\
 & \searrow \cong & \downarrow (\cdot_A) & & \swarrow \cong \\
 & & A & & 
 \end{array}$$

We will soon give a third and equivalent of defining a diagram. By using electric circuits [3], we get a diagram calculus to manipulate morphisms of diagrams.

We supply some examples of algebras one may encounter in nature.

*Example 1.1.4.* Let  $\mathbb{K}$  be any field. The field is trivially an algebra over itself.

*Example 1.1.5.* The complex numbers  $\mathbb{C}$  is an algebra over  $\mathbb{R}$ . This is true since it is a vector space over  $\mathbb{R}$ , and complex multiplication respects scalar multiplication.

*Example 1.1.6.* Let  $\mathbb{K}$  be any field. The ring of  $n$ -dimensional matrices  $M_n(\mathbb{K})$  is an algebra over  $\mathbb{K}$ . The multiplication is taken to be the matrix multiplication and the unit is the  $n$ -dimensional identity matrix.

Augmented algebras will be central to our discussion. An algebra  $A$  is augmented if it is equipped with an algebra homomorphism which splits the algebra into an ideal component and a unit component. We make this precise with the following definition

**Definition 1.1.7** (Augmented algebras). Let  $A$  be an algebra. It is called augmented if there is an algebra homomorphism  $\varepsilon_A : A \rightarrow \mathbb{K}$ .

Given this algebra homomorphism we know that it has to preserve the unit. Thus the kernel  $\text{Ker} \varepsilon_A \subseteq A$  is almost  $A$ , but without its unit. In the module category  $Mod_{\mathbb{K}}$  the morphism  $\varepsilon_A$  is automatically a split-epimorphism, where the splitting is given by the unit  $1_A$ . Thus as a module we have  $A \simeq \bar{A} \oplus \mathbb{K}$ , where  $\bar{A} = \text{Ker} \varepsilon_A$ .  $\bar{A}$  is called the augmentation ideal, or the reduced algebra of  $A$ .

A morphism  $f : A \rightarrow B$  of augmented algebras is an algebra homomorphism, but with the added condition that it must preserve the augmentation; i.e.  $\varepsilon_B \circ f = \varepsilon_A$ . The collection of all augmented algebras over  $\mathbb{K}$  together with the morphisms describes constitutes the category of augmented algebras over  $\mathbb{K}$ :  $\text{Alg}_{\mathbb{K},+}$ .

Given an augmented algebra  $A$ , taking kernels of  $\varepsilon_A$  gives a functor  $\bar{\phantom{x}} : \text{Alg}_{\mathbb{K},+} \rightarrow \widehat{\text{Alg}}_{\mathbb{K}}$ . This functor is well defined on morphisms of augmented algebras, as each morphism is required to preserve the splitting. This functor employs a quasi-inverse, given by the free augmentation  $\bar{\phantom{x}}^+ : \widehat{\text{Alg}}_{\mathbb{K}} \rightarrow \text{Alg}_{\mathbb{K},+}$ . Given a non-unital algebra  $A$ , the free augmentation is defined as  $A^+ = A \oplus \mathbb{K}$ ,

where the multiplication is given by:

$$(a, k)(a', k') = (aa' + ak' + a'k, kk').$$

The unit is given by the element  $(0, 1)$ . We summarize this statement.

**Proposition 1.1.8.** *The functors  $\_ -$  and  $\_ +$  are quasi-inverse to each other.*

*Proof.* Supply proof?

□

There are many augmented algebras to encounter in nature. We will note some examples.

*Example 1.1.9* (Group algebra). Pick any group  $G$  and any field  $\mathbb{K}$ . The group ring  $K[G]$  is an augmented algebra where the augmentation is given by the sum

$$\sum_{g \in G} k_g g = \sum_{g \in G} k_g$$

Among our most important example of algebras is the tensor algebra. This is also the free algebra over  $\mathbb{K}$ .

*Example 1.1.10* (Tensor algebra). Let  $V$  be a  $\mathbb{K}$ -module. We define the tensor algebra  $T(V)$  of  $V$  as the module

$$T(V) = \mathbb{K} \oplus V \oplus V^{\otimes 2} \oplus V^{\otimes 3} \oplus \dots$$

Given two strings  $v^1 \dots v^i$  and  $w^1 \dots w^j$  in  $T(V)$  we define the multiplication by the concatenation operation.

$$\begin{aligned} \nabla_{T(V)} : T(V) \otimes_{\mathbb{K}} T(V) &\rightarrow T(V) \\ (v^1 \dots v^i) \otimes (w^1 \dots w^j) &\mapsto v^1 \dots v^i w^1 \dots w^j \end{aligned}$$

The unit is given by including  $\mathbb{K}$  into  $T(V)$ .

$$\begin{aligned} v_{T(V)} : \mathbb{K} &\rightarrow T(V) \\ 1 &\mapsto 1 \end{aligned}$$

Observe that the tensor algebra is augmented. The projection from  $T(V)$  into  $\mathbb{K}$  is an algebra homomorphism, so we may split the tensor algebra into its unit and its augmentation ideal  $T(V) \simeq \mathbb{K} \oplus \bar{T}(V)$ . We call  $\bar{T}(V)$  the reduced tensor algebra.

**Proposition 1.1.11** (Tensor algebras are free). *The tensor algebras are the free algebras over the category of  $\mathbb{K}$ -modules, i.e. for any  $\mathbb{K}$ -module  $V$  there is a natural isomorphism  $\text{Hom}_{\mathbb{K}}(V, A) \simeq \text{Alg}_{\mathbb{K}}(T(V), A)$ .*

*The reduced tensor algebra is the free non-unital algebra over the category of  $\mathbb{K}$ -modules, i.e. for any  $\mathbb{K}$ -module  $V$  there is a natural isomorphism  $\text{Hom}_{\mathbb{K}}(V, A) \simeq \widehat{\text{Alg}}_{\mathbb{K}}(\bar{T}(V), A)$ .*



*Proof.* This proposition should be evident from the description of an algebra homomorphism from a tensor algebra. If  $f : T(V) \rightarrow A$  is an algebra homomorphism, then  $f$  must satisfy the following conditions:

- Unitality:  $f(1) = 1$
- Homomorphism property: Given  $v, w \in V$ , then  $f(vw) = f(v) \cdot_A f(w)$

By induction, we see that  $f$  is completely determined by where it sends the elements of  $V$ . Thus restriction by the inclusion of  $V$  into  $T(V)$  induces a bijection.  $\square$

Just as for rings, every algebra  $A$  has a module category of its own. The definition is very much like that for rings.

**Definition 1.1.12** (Modules). Let  $A$  be an algebra over  $\mathbb{K}$ . A  $\mathbb{K}$ -module  $M$  is said to be a left (right)  $A$ -module if there exists a structure morphism  $\mu_M : A \otimes_{\mathbb{K}} M \rightarrow M$  ( $\mu_M : M \otimes_{\mathbb{K}} A \rightarrow M$ ) called multiplication. We require that  $\mu_M$  is associative and preserves the unit of  $A$ . I.e. we have the commutative diagrams in  $Mod_{\mathbb{K}}$ .

$$\begin{array}{ccccc}
 A \otimes_{\mathbb{K}} A \otimes_{\mathbb{K}} M & \xrightarrow{A \otimes \mu_M} & A \otimes_{\mathbb{K}} M & & \mathbb{K} \otimes_{\mathbb{K}} M \xrightarrow{1_A \otimes M} A \otimes_{\mathbb{K}} M \\
 \downarrow \cdot_A \otimes M & & \downarrow \mu_M & & \searrow \simeq \downarrow \mu_M \\
 A \otimes_{\mathbb{K}} M & \xrightarrow{\mu_M} & M & & M
 \end{array}$$

**Definition 1.1.13** ( $A$ -linear homomorphisms). Let  $M, N$  be two left  $A$ -modules. A morphism  $f : M \rightarrow N$  is called  $A$ -linear if it is  $\mathbb{K}$ -linear and for any  $a$  in  $A$ , i.e.  $f(am) = af(m)$ .

The category of left  $A$ -modules is denoted as  $Mod_A$ , where the morphisms  $\text{Hom}_A(-, -)$  are  $A$ -linear. Likewise, the category of right  $A$ -modules is denoted as  $Mod^A$ . There is a free functor from the category of  $\mathbb{K}$ -modules to left  $A$ -modules.

**Proposition 1.1.14.** Let  $M$  be a  $\mathbb{K}$ -module. The module  $A \otimes_{\mathbb{K}} M$  is a left  $A$ -module. Moreover, it is the free left module over  $\mathbb{K}$ -modules, i.e. there is a natural isomorphism  $\text{Hom}_{\mathbb{K}}(M, N) \simeq \text{Hom}_A(A \otimes_{\mathbb{K}} M, N)$ .

*Proof.* We use the tensor-hom adjunction to arrive at the conclusion.

$$\text{Hom}_A(A \otimes_{\mathbb{K}} M, N) \simeq \text{Hom}_A(A, \text{Hom}_{\mathbb{K}}(M, N)) \simeq \text{Hom}_{\mathbb{K}}(M, N)$$

$\square$

### 1.1.2 Coalgebras

In this section we want to dualize the definitions given in the last section. As we will see, some strange artifacts from the structure of algebras appears in the dualization process. We would like to define a coalgebra as an algebra, but all of the diagrams are reversed. For many purposes this dualization is good, but as we will encounter some finiteness conditions are necessary. The category of "ill-behaved" coalgebras will be denoted  $\text{CoAlg}_{\mathbb{K}}$ , and we will look at this first.

**Definition 1.1.15** ( $\mathbb{K}$ -Coalgebra). Let  $\mathbb{K}$  be a field. A coalgebra  $C$  over  $\mathbb{K}$  is a  $\mathbb{K}$ -module with structure morphisms called comultiplication and counit,

$$\begin{aligned} (\Delta_C) : C &\rightarrow C \otimes_{\mathbb{K}} C \\ \varepsilon_C : C &\rightarrow \mathbb{K}, \end{aligned}$$

satisfying the coassociativity and coidentity laws.

$$\begin{aligned} \text{(coassociativity)} \quad & (\Delta_C \otimes id_C) \circ \Delta_C(c) = (id_C \otimes \Delta_C) \circ \Delta_C(c) \\ \text{(counitality)} \quad & (id_C \otimes \varepsilon_C) \circ \Delta_C(c) = c = (\varepsilon_C \otimes id_C) \circ \Delta_C(c) \end{aligned}$$

In the same way as for algebras, we say that a coalgebra is non-counital if it is without a counit.

Unlike algebras, coalgebras does not admit a single intuitive method for writing repeated application of the comultiplication. For instance, given an element  $c \in C$  we may apply the comultiplication twice on  $c$  in two different ways:

$$\begin{aligned} \Delta_{C,(1)}^2(c) &= (\Delta_C \otimes C) \Delta_C(c), \\ \Delta_{C,(2)}^2(c) &= (C \otimes \Delta_C) \Delta_C(c). \end{aligned}$$

One should immediately note that  $\Delta_{C,(1)}^2(c) = \Delta_{C,(2)}^2(c)$  is coassociativity. So there is in fact a unique way to do repeated application of  $\Delta_C$  on  $c$ . We denote the  $n$ -fold repeated application of  $\Delta_C$  by  $\Delta_C^n$ . Since the element  $\Delta_C^n(c)$  represents a finite sum in  $C^{\otimes n}$ , one may use Sweedlers notation [3] as it is well-defined.

$$\Delta_C^n(c) = \sum c_{(1)} \otimes \dots \otimes c_{(n)}$$

**Definition 1.1.16** (Coalgebra homomorphism). Let  $C$  and  $D$  be coalgebras. Then  $f : C \rightarrow D$  is a coalgebra morphism if

1.  $f$  is  $\mathbb{K}$ -linear
2.  $(f \otimes f) \circ \Delta_C(c) = \Delta_D(f(c))$
3.  $\varepsilon_D \circ f = \varepsilon_C$

Whenever  $C$  and  $D$  are non-counital, we only require 1 and 2 for a homomorphism of non-counital coalgebras.

**Definition 1.1.17** (Category of Coalgebras). Let  $\text{CoAlg}_{\mathbb{K}}$  denote the category of coalgebras. Its objects consists of every coalgebra  $C$ , and the morphisms are coalgebra homomorphisms. The sets of morphisms between  $C$  and  $D$  are denoted as  $\text{CoAlg}_{\mathbb{K}}(C, D)$ .

Let  $\widehat{\text{CoAlg}}_{\mathbb{K}}$  denote the category of non-unital algebras. Its objects consists of every non-unital algebra  $C$ , and the morphisms are non-unital algebra homomorphisms. The sets of morphisms between  $C$  and  $D$  are denoted as  $\widehat{\text{CoAlg}}_{\mathbb{K}}(C, D)$ .

At a first glance, coalgebras may seem weird and unnatural, but they appear many places in nature.

*Example 1.1.18* ( $\mathbb{K}$  as a coalgebra). The field  $\mathbb{K}$  can be given a coalgebra structure over itself. Since  $\{1\}$  is a basis for  $\mathbb{K}$  we define the structure morphisms as

$$\begin{aligned}\Delta_{\mathbb{K}}(1) &= 1 \otimes 1 \\ \varepsilon(1) &= 1.\end{aligned}$$

One may check that these morphisms are indeed coassociative and counital. Thus we may regard our field as either an algebra or coalgebra over itself.

*Example 1.1.19* ( $\mathbb{K}[G]$  as a coalgebra). The group algebra has a natural coalgebra structure. We may take duplication of group elements as the comultiplication, i.e.

$$\Delta_{\mathbb{K}[G]}(kg) = kg \otimes g.$$

Coincidentally we have already defined the counit, this is the augmentation  $\varepsilon_C$  for the group algebra  $\mathbb{K}[G]$ . Recall that this was

$$\varepsilon_C(\sum k_g g) = \sum k_g.$$

One may see that these morphisms satisfy coassociativity and counitality.

Before we state our main example, we will first introduce part of its essential structure.

**Definition 1.1.20** (Coaugmented coalgebras). Let  $C$  be a coalgebra.  $C$  is coaugmented if there is a coalgebra homomorphism  $\eta_C : \mathbb{K} \rightarrow C$ .

Similar to augmented algebras, each coaugmented coalgebra splits in the category  $\text{Mod}_{\mathbb{K}}$ . We first notice that given a coalgebra homomorphism  $f$ , the cokernel  $\text{Cok} f$  is also a coalgebra. This is some induced quotient structure. Given a coaugmentation  $\eta_C : \mathbb{K} \rightarrow C$ , we call  $\text{Cok} \eta_C = \bar{C}$  for the coaugmentation quotient or reduced coalgebra of  $C$ . Then we obtain the splitting  $C \simeq \bar{C} \oplus \mathbb{K}$ . The reduced coproduct, denoted  $\bar{\Delta}_C$  may explicitly be given as

$$\bar{\Delta}_C(c) = \Delta_C(c) - 1 \otimes c - c \otimes 1.$$

*Example 1.1.21* (Tensor Coalgebras). Let  $V$  be a  $\mathbb{K}$ -module. We define the tensor coalgebra  $T^c(V)$  of  $V$  as the module

$$T^c(V) = \mathbb{K} \oplus V \oplus V^{\otimes 2} \oplus V^{\otimes 3} \oplus \dots$$

Given a string  $v^1 \dots v^i$  in  $T(V)$  we define the comultiplication by the deconcatenation operation.

$$\begin{aligned} \Delta_{T^c(V)} : T^c(V) &\rightarrow T^c(V) \otimes_{\mathbb{K}} T^c(V) \\ v^1 \dots v^i &\mapsto 1 \otimes (v^1 \dots v^i) + \left( \sum_{j=1}^{n-1} (v^1 \dots v^j) \otimes (v^{j+1} \dots v^i) \right) + (v^1 \dots v^i) \otimes 1 \end{aligned}$$

The counit is given by projecting  $T^c(V)$  onto  $\mathbb{K}$ .

$$\begin{aligned} \varepsilon_{T^c(V)} : T^c(V) &\rightarrow \mathbb{K} \\ 1 &\mapsto 1 \\ v^1 \dots v^i &\mapsto 0 \end{aligned}$$

We observe that the tensor coalgebra is coaugmented. Its coaugmentation is given by the inclusion of  $\mathbb{K}$  into  $T^c(V)$ . We can split  $T^c(V) \simeq \mathbb{K} \oplus \bar{T}^c(V)$ , where  $\bar{T}^c(V)$  denotes the reduced tensor coalgebra.

Cofreeness does not come for free for the tensor coalgebra. Our problem is a mismatch in the behavior of algebras and coalgebras. The problem arises when we try to do evaluation. Suppose that  $A$  is an algebra and that we have  $n$  elements of  $A$ , i.e. an element of  $A^{\otimes n}$ . On this element we may apply the multiplication of  $A$  a maximum of  $n$ -times, we cannot multiply an element with nothing non-trivially. However, given an element in a coalgebra  $C$ , we may use the comultiplication on this element  $n$  times,  $n+1$  times and so on ad infinitum. In the coalgebra we may comultiply an element possibly an infinite amount of times. This is a property which is sometimes ill-behaved with our dualization of algebras to coalgebras.

However, when we dualized the tensor algebra to the tensor coalgebra we did not lose this property, that an element may only be comultiplied a finite number of times. This is due to the composition of  $T^c(V)$  as a direct sum of  $V^{\otimes n}$ ; i.e. any element is a finite sum of finite tensors.

This extra assumption we need for coalgebras will be called conilpotent. Let  $C \simeq \mathbb{K} \oplus \bar{C}$  be a coaugmented coalgebra, we define the coradical filtration of  $C$  as a filtration  $Fr_0 C \subseteq Fr_1 C \subseteq \dots \subseteq Fr_r C \subseteq \dots$  by the submodules:

$$\begin{aligned} Fr_0 C &= \mathbb{K} \\ Fr_r C &= \mathbb{K} \oplus \{c \in \bar{C} \mid \forall n \geq r \bar{\Delta}_C(c) = 0\}. \end{aligned}$$

**Definition 1.1.22** (Conilpotent coalgebras). Let  $C$  be a coaugmented coalgebra. We say that  $C$  is conilpotent if its coradical filtration is exhaustive, i.e.

$$\varinjlim_r Fr_r C \simeq C.$$

The subcategory of conilpotent coalgebras will be denoted as  $\text{CoAlg}_{\mathbb{K}, \text{conil}}$ .

**Proposition 1.1.23** (Conilpotent tensor coalgebra). Let  $V$  be a  $\mathbb{K}$ -module. The tensor coalgebra  $T^c(V)$  is conilpotent.

*Proof.* Let  $v \in V$ , then  $\Delta_{T^c(V)}(v) = 1 \otimes v + v \otimes 1$  and  $\bar{\Delta}_{T^c(V)}(v) = 0$ . We then observe the following:

$$\begin{aligned} Fr_0 T^c(V) &= \mathbb{K} \\ Fr_1 T^c(V) &= \mathbb{K} \oplus V \\ Fr_r T^c(V) &= \bigoplus_{i \leq r} V^{\otimes i} \end{aligned}$$

This shows that the coradical filtration is exhaustive.  $\square$

**Proposition 1.1.24** (Cofree tensor coalgebra). *The tensor coalgebra is the cofree conilpotent coalgebra over the category of  $\mathbb{K}$ -modules, i.e. for any  $\mathbb{K}$ -module  $V$  and any conilpotent coalgebra  $C$  there is a natural isomorphism  $\text{Hom}_{\mathbb{K}}(C, V) \simeq \text{CoAlg}_{\mathbb{K}, \text{conil}}(C, T^c(V))$ .*

Fiks dette?

*Proof.* This proposition should be evident from the description of a coalgebra homomorphism into the a tensor coalgebra. If  $g : C \rightarrow T^c(V)$  is a coalgebra homomorphism, then  $g$  must satisfy the following conditions:

1. (Coaugmentation)  $g(1) = 1$
2. (Counitality) Given  $c \in \bar{C}$  then  $\varepsilon_{T^c(V)} \circ g(c) = 0$
3. (Homomorphism property) Given  $c \in C$  then  $\Delta_{T^c(V)}(g(c)) = (g \otimes g) \circ \Delta_C(c)$

We will construct the maps for the isomorphism explicitly. If  $g : C \rightarrow T^c(V)$  is a coalgebra homomorphism, then composing with projection gives a map  $\pi \circ g : C \rightarrow V$ . Note that  $\pi \circ g(1) = 0$ , so this is essentially a map  $\pi \circ g : \bar{C} \rightarrow V$ . For the other direction, let  $\bar{g} : \bar{C} \rightarrow V$ . We will then define  $g$  as

$$g = id_{\mathbb{K}} \oplus \sum_{i=1}^{\infty} (\otimes^i \bar{g}) \bar{\Delta}_C^{i-1}.$$

Observe that  $g$  is well defined, since convergence of the sum follows from conilpotency of  $C$ . One may then check that  $g$  is a coalgebra homomorphism, which yields the result.  $\square$

Integral to our dualization are comodules. We quickly give a definition of them.

**Definition 1.1.25** (Comodules). Let  $C$  be a coalgebra. A  $\mathbb{K}$ -module  $M$  is said to be left (right)  $C$ -comodule if there exist a structure morphism  $\omega_M : M \rightarrow C \otimes_{\mathbb{K}} M$  ( $\omega_M : M \rightarrow M \otimes_{\mathbb{K}} C$ ) called comultiplication. We require that  $\omega_M$  is coassociative with respect to the comultiplication of  $C$  and preserves the counit of  $C$ . I.e. we have the following commutative diagrams in  $\text{Mod}_{\mathbb{K}}$ .

$$\begin{array}{ccc} C \otimes_{\mathbb{K}} C \otimes_{\mathbb{K}} M & \xleftarrow{C \otimes \omega_M} & C \otimes_{\mathbb{K}} M \\ \Delta_C \otimes M \uparrow & & \omega_M \uparrow \\ C \otimes_{\mathbb{K}} M & \xleftarrow{\omega_M} & M \end{array} \quad \begin{array}{ccc} \mathbb{K} \otimes_{\mathbb{K}} M & \xleftarrow{\varepsilon_C \otimes M} & C \otimes_{\mathbb{K}} M \\ & \searrow \simeq & \omega_M \uparrow \\ & & M \end{array}$$

**Definition 1.1.26** (C-colinear homomorphism). Let  $M, N$  be two left  $C$ -comodules. A morphism  $g : M \rightarrow N$  is called  $C$ -colinear if it is  $\mathbb{K}$ -linear and for any  $m$  in  $M$ ,  $\omega_N(g(m)) = (id_C \otimes g)\omega_M(m)$ . In Sweedlers notation this looks like:

$$\sum g(c)_{(1)} \otimes g(c)_{(2)} = \sum c_{(1)} \otimes g(c_{(2)}).$$

The category of left  $C$ -comodules is denoted as  $\text{CoMod}_C$ , where the morphisms  $\text{Hom}_C(\_, \_)$  are  $C$ -colinear. We would also like to restrict our attention to those  $C$ -comodules which are conilpotent, i.e. those comodules which have an exhaustive coradical filtration. The coradical filtration is defined analogously, as we only care for the  $\mathbb{K}$ -module structure. Notice that for conilpotent coalgebras this requirement is automatic. Likewise, the category of right  $C$ -comodules is denoted as  $\text{CoMod}^C$ .

**Proposition 1.1.27.** *Let  $M$  be a  $\mathbb{K}$ -module. The module  $C \otimes_{\mathbb{K}} M$  is a left  $C$ -comodule. Moreover, it is the cofree left comodule over  $\mathbb{K}$ -modules, i.e. there is an isomorphism  $\text{Hom}_{\mathbb{K}}(N, M) \simeq \text{Hom}_C(N, C \otimes_{\mathbb{K}} M)$ .*

*Proof.* This is dual to 1.1.14. □

### 1.1.3 Electric Circuits

Calculations involving both algebras and coalgebras tends to become convoluted and unmanagable. Since we want to study the interplay between algebras and coalgebras using other tools to write down equations can be handy. We will develop the graphical calculus briefly mentioned in [3]. This graphical calculus will consists of string diagrams, referred to as electric circuits, which describes function compositions of tensors. Since we only care about the interplay of tensors, we may develop this graphical calculus in any closed symmetric monoidal category. Why do we want to introduce this abstraction? A closed symmetric monoidal category is a good category to model functions, or morphisms, which may take several variables in its argument. Moreover, in the next section we are going to switch categories. Thus the same notions may be applied and the differences are only seen in the implementations.

appendix?

The definition of a closed symmetric monoidal category may be found in the appendix. We recall that it is a category  $\mathcal{C}$  together with a bifunctor  $\_ \otimes \_ : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$  usually called tensor, and a unit object  $Z \in \mathcal{C}$ . We are also given four natural isomorphisms relating the functors and the unit to what they are meant to be:

$$\begin{aligned} \text{Associator} \quad \alpha &: (A \otimes B) \otimes C \rightarrow A \otimes (B \otimes C). \\ \text{Right unit} \quad \rho &: A \otimes Z \rightarrow A. \\ \text{Left unit} \quad \lambda &: Z \otimes A \rightarrow A. \\ \text{Braiding/Symmetry} \quad \beta &: A \otimes B \rightarrow B \otimes A. \end{aligned}$$


These natural isomorphisms are supposed to satisfy some laws as well. See the appendix for the full definition.

In an electric circuit we want to rewrite equations possibly involving tensors into a string diagram. To illustrate with some simple examples, let  $f : A \rightarrow B$ ,  $g : B \rightarrow C$  and  $h : D \rightarrow E$ . We may consider the composition

$$(g \otimes E) \circ (f \otimes h) : A \otimes D \rightarrow C \otimes E.$$

An electric circuit is written from top to bottom and is composed of levels. This means that the first morphisms which we apply will be at the top, descending downwards with each function composition. We write each argument in the composition as its own string. Thus this example above will look like the circuit below. Notice how  $f$  and  $h$  are at the same level, indicating that they should be thought of as  $f \otimes h$ . Thus an  $\otimes$  indicates a change of string, while a  $\circ$  indicates a change of level.


Beware that when many tensors are used, we should really remember how each string is tensored. We may call adding tensors for horizontal composition and composition of morphism for vertical composition. Both of these have a choice in how we associate them, but both have unique choices up to isomorphism given by the associator.



The true power of electric circuit comes to light when we consider morphisms that in some sense "creates" or "destroys" strings. E.g. a morphism of 2 variables "destroys" a string by in some sense applying them to each other. Consider now a morphism  $f : A \otimes B \rightarrow C$ , we represent this morphism in an electric circuit by using a converging fork. Likewise, "creation" of strings is seen as a diverging fork.



We may write the unit object  $Z$  without any strings in a circuit. By the right and left unit, an object is isomorphic to itself tensored with the unit. In this manner whenever we have a morphism entering or exiting the unit  $Z$  we start a new string by using a sink or a source. E.g. consider  $f$  as before and a morphism  $g : Z \rightarrow A$ , then we may write  $f \circ (g \otimes B)$  as the circuit below. Notice that again this is only well-defined up to isomorphism by the right and left unit.



The final operation we have is braiding. Whenever we apply the braiding morphism on the tensors, we may simply denote this as interchanging the strings. E.g.  $\beta_{A,B} : A \otimes B \rightarrow B \otimes A$  is the circuit below. Notice that by naturality of  $\beta$  we may move a braiding along the circuit. In this manner, if we have two braids, they may undo each other. A braid may also be moved to an end, in this manner they may be ignored during calculations.



With the language of electric circuits, we may now write down the axioms of an algebra or coalgebra by circuits. The axioms states the existence of morphisms, as well as commutativity of some diagrams. We may do this electrically. The structure maps of algebras and coalgebras will be given special notation, since we will use these often.

For convenience we will let  $\mathcal{C} = \text{Mod}_{\mathbb{K}}$ . This is a closed symmetric monoidal category, with  $\otimes_{\mathbb{K}}$  as  $\otimes$ . Recall that an algebra is a  $\mathbb{K}$ -module  $A$  together with maps  $(\cdot_A) : A \otimes A \rightarrow A$  and  $1_A : \mathbb{K} \rightarrow A$ . These morphisms will be denoted electrically as the diagrams below.

$$(\cdot_A) = \begin{array}{c} \diagup \quad \diagdown \\ | \\ \diagdown \quad \diagup \end{array} \quad 1_A = \begin{array}{c} \circ \\ | \end{array}$$

We write the electric laws for an algebra as how one would write equations. Associativity and unitality then becomes as follows.

$$\text{Associativity} \quad \begin{array}{c} \diagup \quad \diagdown \\ | \quad | \\ \diagdown \quad \diagup \end{array} = \begin{array}{c} \diagup \quad \diagdown \\ | \quad | \\ \diagdown \quad \diagup \end{array}$$


$$\text{Unitality} \quad \begin{array}{c} \diagup \quad \diagdown \\ | \quad | \\ \diagdown \quad \diagup \end{array} = \begin{array}{c} | \end{array} = \begin{array}{c} \circ \\ | \end{array}$$


Dually, given a coalgebra  $C$  we make the same notation, but everything will be turned upside down. The maps  $\Delta_C : C \rightarrow C \otimes C$  and  $\varepsilon_C : C \rightarrow \mathbb{K}$  will be denoted as the following circuits.

$$\Delta_C = \begin{array}{c} | \\ \diagdown \quad \diagup \end{array} \quad \varepsilon_C = \begin{array}{c} | \\ \circ \end{array}$$

The electric laws for  $C$  becomes the following diagrams.




Coassociativity 

Counitality 

This notation will be adopted for our algebras and coalgebras when convenient. This will compactify the notation for comultiplication and make tensors more readable. Moreover, the intuition for algebras are easier to dualize to coalgebras with this notation. One simply has to work out an algebra statement and then turn the diagram upside down to make it into a coalgebra statement.

Previously we talked about braiding and how that relates to interchanging strings. In the same manner that we have a horizontal and vertical associator, we also have a vertical and horizontal braiding. The braiding first encountered is the horizontal braiding. The vertical braiding refers then to function composition of tensors. This manifests in electric circuits as sliding a morphism along a string. Whenever the given braiding of  $\mathcal{C}$  is nice enough we can get away by ignoring it whenever we move a morphism along a string. For instance, look at the category of  $\mathbb{K}$ -modules where we may define the braiding on elementary tensors as  $\beta(a \otimes b) = b \otimes a$ . In this case the braiding is agnostic to how we move our morphisms along a string. This means that we have the following equality of circuits.



However, in nature we may encounter a braiding which is not nice enough. Here we should take a step back to figure out how we can move morphisms along strings, before we continue to use this graphical calculus of function composition. We will meet such a braiding soon.

### 1.1.4 Derivations and DG-Algebras

This section aims to define differential graded algebras and their modules. Given an algebra  $A$ , we may define a derivation as a map satisfying the Leibniz rule. In the dual case for a coalgebra we may define a coderivation as a map satisfying the Zinbiel rule, but for brevity we will refer to these maps as derivations. Once we get a grasp on how to make derivations we introduce graded algebras and modules, to equip these with derivations. This will give us the categories of differential graded algebras and chain complexes. Throughout this section we will also develop electric circuits for these notions.

**Definition 1.1.28** (Derivations and Coderivations). Let  $M$  be an  $A$ -bimodule. A  $\mathbb{K}$ -linear morphism  $d : A \rightarrow M$  is called a derivation if  $d(ab) = d(a)b + ad(b)$ , i.e. electrically:

Let  $N$  be a  $C$ -bicomodule. A  $\mathbb{K}$ -linear morphism  $d : N \rightarrow C$  is called a coderivation if  $\Delta_C \circ d = (d \otimes id_C) \circ \omega_N^r + (id_C \otimes d) \circ \omega_N^l$ , i.e. electrically:

We remark that this translation between equations and electric circuits is not at the same level of generalization. Due to this, the electric circuit description has more advantages as it allows us to think with elements when we are really only dealing with morphisms. This will later be abused, to derive results which is e.g. independent of the given braiding on the category.

A useful fact about derivations is that they will always map the identity to 0. We obtain this from Leibniz rule as one would get  $d(1) = 2d(1)$ , and thus  $d(1) = 0$ .

**Proposition 1.1.29.** *Let  $V$  be a  $\mathbb{K}$ -module and  $M$  be a  $T(V)$ -bimodule. A  $\mathbb{K}$ -linear morphism  $f : V \rightarrow M$  uniquely determines a derivation  $d_f : T(V) \rightarrow M$ , i.e. there is an isomorphism  $\text{Hom}_{\mathbb{K}}(V, M) \simeq \text{Der}(T(V), M)$ .*

*Let  $N$  be a  $T^c(V)$ -cobimodule. A  $\mathbb{K}$ -linear morphism  $g : M \rightarrow V$  uniquely determines a coderivation  $d_g^c : N \rightarrow T^c(V)$ , i.e. there is an isomorphism  $\text{Hom}_{\mathbb{K}}(N, V) \simeq \text{Coder}(N, T^c(V))$ .*

**Proof.** Let  $a_1 \otimes \dots \otimes a_n$  be an elementary tensor of  $T(V)$ . We define a map  $d_f : T(V) \rightarrow M$  as

$$d_f(a_1 \otimes \dots \otimes a_n) = \sum_{i=1}^n a_1 \dots f(a_i) \dots a_n$$

$$d_f(1) = 0.$$

$d_f$  is a derivation by definition.

Restriction to  $V$  gives the natural isomorphism. Let  $i : V \rightarrow T(V)$ , then  $i^* d_f = f$ . Let  $d : T(V) \rightarrow M$  be a derivation, then  $d_{i^* d} = d$ . Suppose that  $g : M \rightarrow N$  is a morphism between  $T(V)$ -bimodules, then naturality follows from bilinearity.

In the dual case,  $d_g^c : N \rightarrow T^c(V)$  is a bit tricky to define. Let  $\omega_N^l : N \rightarrow N \otimes T^c(V)$  and  $\omega_N^r : N \rightarrow T^c(V) \otimes N$  denote the coactions on  $N$ . Since  $T^c(V)$  is conilpotent we get the same kind of finiteness restrictions on  $N$ . Define the reduced coactions as  $\bar{\omega}_N^l = \omega_N^l - \_ \otimes 1$  and  $\bar{\omega}_N^r = \omega_N^r - 1 \otimes \_$ , this is well-defined by coassociativity. Observe that for any  $n \in N$  there are  $k$  and  $k' > 0$  such that  $\bar{\omega}_N^{l^k}(n) = 0$  and  $\bar{\omega}_N^{r^{k'}}(n) = 0$ .

Let  $n_{(k)}^{(i)}$  denote the extension of  $n$  by  $k$  coactions at position  $i$ , i.e.

$$n_{(k)}^{(i)} = \bar{\omega}_N^{r^i} \bar{\omega}_N^{l^{k-i}}(n).$$

The extension of  $n$  by  $k$  coactions is then the sum over every position  $i$ ,

$$n_{(k)} = \sum_{i=0}^k n_{(k)}^{(i)}.$$

Observe that  $n_{(0)} = n$ . The grade of  $n$  may be thought of as the smallest  $k$  such that  $n_{(k)}$  is zero. This grading gives us the coradical filtration of  $N$ , and it is exhaustive by the finiteness restrictions given above. So every element of  $N$  may be given a finite grade.

If  $g : N \rightarrow V$  is a linear map, we may think of it as a map sending every element of  $N$  to an element of  $T^c(V)$  of grade 1. To get a map which sends element of grade  $k$  to grade  $k$ , we must extend the morphism. Let  $\pi : T^c(V) \rightarrow V$  be the linear projection and define  $g_{(k)}^{(i)} = \pi \otimes \dots \otimes g \otimes \pi$  as a morphism which of  $k$  tensors which is  $g$  at the  $i$ -th argument, but the projection otherwise.  $d_g^c$  is then defined as the sum over each coaction and coordinate.


$$d_g^c(n) = \sum_{k=0}^{\infty} \sum_{i=0}^k g_{(k)}^{(i)}(n_{(k)}^{(i)})$$

Upon closer inspection we may observe that this is the dual construction of the derivation morphism. It is well-defined as the sum is finite by the finiteness restrictions. The map is a coderivation by duality, and the natural isomorphism is given by composition with the projection map  $\pi$ .  $\square$


We now wish to get into the territory of derivations as we know them from homological algebra. We get this by choosing an algebra and equipping it with a derivation. This will give us a universal derivation to work with when we want to define a differential on a graded module.

**Definition 1.1.30** (Differential algebra). Let  $A$  be an algebra. We say that  $A$  is a differential algebra if it is equipped with a derivation  $d : A \rightarrow A$ . Dually, a coalgebra  $C$  is called a differential coalgebra if it is equipped with at least one coderivation  $d : C \rightarrow C$ .


**Definition 1.1.31** ( $A$ -derivation). Let  $(A, d_A)$  be a differential algebra and  $M$  a left  $A$ -module. A  $\mathbb{K}$ -linear morphism  $d_M : M \rightarrow M$  is called an  $A$ -derivation if  $d_M(am) = d_A(a)m + ad_M(m)$ , or electrically:



Dually, given a differential coalgebra  $(C, d_C)$  and  $N$  a left  $C$ -comodule, a  $\mathbb{K}$ -linear morphism  $d_N : N \rightarrow N$  is a coderivation if  $\omega_N \circ d_N = (d_C \otimes id_N + id_C \otimes d_N) \circ \omega_N$ , or electrically:



When there is no ambiguity, we will start to adopt writing the differential in electric circuits as a triangle.



**Proposition 1.1.32.** *Let  $A$  be a differential algebra and  $M$  a  $\mathbb{K}$ -module. A  $\mathbb{K}$ -linear morphism  $f : M \rightarrow A \otimes_{\mathbb{K}} M$  uniquely determines a derivation  $d_f : A \otimes_{\mathbb{K}} M \rightarrow A \otimes_{\mathbb{K}} M$ , i.e. there is an isomorphism  $\text{Hom}_{\mathbb{K}}(M, A \otimes_{\mathbb{K}} M) \simeq \text{Der}(A \otimes_{\mathbb{K}} M)$ . Moreover,  $d_f$  is given as  $(\nabla_A \otimes id_M) \circ (id_A \otimes f) + d_A \otimes id_M$ .*

*Dually, if  $C$  is a differential coalgebra and  $N$  is a  $\mathbb{K}$ -module, then a  $\mathbb{K}$ -linear morphism  $g : C \otimes_{\mathbb{K}} N \rightarrow N$  uniquely determines a coderivation  $d_g : C \otimes_{\mathbb{K}} N \rightarrow C \otimes_{\mathbb{K}} N$ . There is an isomorphism  $\text{Hom}_{\mathbb{K}}(C \otimes_{\mathbb{K}} N, N) \simeq \text{Coder}(C \otimes_{\mathbb{K}} N)$ , and  $d_g$  is given as  $(id_C \otimes g) \circ (\Delta_C \otimes id_N) + d_C \otimes id_N$ .*

*Proof.* This is only proved in the case of an algebra. The case of a coalgebra is dual.

We first got to prove that the morphism  $d_- : \text{Hom}_{\mathbb{K}}(M, A \otimes_{\mathbb{K}} M) \rightarrow \text{Der}(A \otimes_{\mathbb{K}} M)$  is well-defined. This means to check that for any morphism  $f : M \rightarrow A \otimes_{\mathbb{K}} M$  the morphism  $d_f$  satisfies the Leibniz rule.

Assume that we have elements  $a, b \in A$  and  $m \in M$ . Then  $d_f(ab \otimes m) = d_f(a(b \otimes m))$ . We now make an abuse of notation to write an equality between an element and a circuit. Recall that this really means that we have to think of  $a, b$  and  $m$  as generalized elements.

$$\begin{aligned}
 d_f(ab \otimes m) &= \text{circuit with } f \text{ and } \nabla_A \text{ and } d_A \text{ components} = \text{circuit with } f \text{ and } \nabla_A \text{ and } d_A \text{ components} \\
 &= d_A(a)b \otimes m + ad_f(b \otimes m).
 \end{aligned}$$

Next we show that  $d_-$  has an inverse. This is given by "restriction to  $M$ ", also known as  $(1_A \otimes M)^* : \text{Hom}_{\mathbb{K}}(A \otimes_{\mathbb{K}} M, N) \rightarrow \text{Hom}_{\mathbb{K}}(M, N)$ . Let  $f : M \rightarrow A \otimes_{\mathbb{K}} M$  be a linear map and  $D : A \otimes_{\mathbb{K}} M \rightarrow A \otimes_{\mathbb{K}} M$  be a derivation.

$$d_f \circ (1_A \otimes M) = \text{diagram} + \text{diagram} = f$$

$$d_{D \circ (1_A \otimes M)} = \text{diagram} + \text{diagram} = D$$

Notice that in the last equation we use the Leibniz rule to get the equality to  $D$ .

□

We are now going to slowly work our way towards the context of homological algebra. We say that a  $\mathbb{K}$ -module  $M^*$  admits a  $\mathbb{Z}$ -grading if it decomposes into either summands or factors

$$M^* = \bigoplus_{z \in \mathbb{Z}} M^z \text{ or } M^* = \prod_{z \in \mathbb{Z}} M^z.$$

Gir dette m  
definere?

An element of  $m \in M$  is said to be homogenous if it is properly contained in a single summand,  $m$  is then given the degree of the summand. We say that a morphism of graded modules  $f : M^* \rightarrow N^*$  is homogenous of degree  $n$  if it preserves the grading, i.e.  $f(M^i) \subseteq N^{n+i}$ . The degree of a homogenous element  $m$  or morphism  $f$  is denoted as  $|m|$  or  $|f|$ .

There is a distinction between the ordinary and self-enriched category of graded modules. We are going to work with this category as it is both the ordinary and self-enriched category. Its hom objects is the graded module of homogenous morphism. We denote a factor in the grading as  $\text{Hom}_{\mathbb{K}}^w(M^*, N^*) = \{f : M^* \rightarrow N^* \mid f \text{ is homogenous and } |f| = w\}$ , so the graded hom is

Kan jeg skr  
dette?

$$\text{Hom}_{\mathbb{K}}^* = \prod_{w \in \mathbb{Z}} \text{Hom}_{\mathbb{K}}^w.$$

The ordinary hom objects are the hom objects of the underlying modules,  $\text{Hom}_{\mathbb{K}}(M, N)$ . This category is denoted as  $\text{Mod}_{\mathbb{K}}^*$ . In general and whenever it makes sense, we denote  $\mathcal{C}^*$  as the category of  $\mathbb{Z}$ -graded objects from  $\mathcal{C}$ .

The category  $\text{Mod}_{\mathbb{K}}^*$  is a closed symmetric monoidal category. The tensor is given by the following formula, using the underlying tensor of  $\text{Mod}_{\mathbb{K}}$

$$M^* \otimes N^* = \bigoplus_{n \in \mathbb{Z}} \bigoplus_{p \in \mathbb{Z}} M^p \otimes_{\mathbb{K}} N^q, \text{ where } q = n - p.$$

The associator of  $\text{Mod}_{\mathbb{K}}$  may be lifted to this tensor. The unit is the module  $\mathbb{K}^* \simeq \mathbb{K}$  concentrated in degree 0. The right and left unit transformation may also be lifted from  $\mathbb{K}$ .

The category  $Mod_{\mathbb{K}}^*$  is closed. This means that our graded tensor fixed in one variable is left adjoint to the graded hom. We may obtain the graded hom as a right adjoint for the other variable as well by using the braiding which we will define later. Showing closedness is a routine calculation by using the tensor-hom adjunction from  $Mod_{\mathbb{K}}$ .

$$\begin{aligned}
Hom_{\mathbb{K}}^*(A^* \otimes B^*, C^*) &= \prod_{w \in \mathbb{Z}} \prod_{n \in \mathbb{Z}} \prod_{p \in \mathbb{Z}} Hom_{\mathbb{K}}^w(\bigoplus A^p \otimes_{\mathbb{K}} B^{n-p}, C^n) \\
&= \prod_{w \in \mathbb{Z}} \prod_{n \in \mathbb{Z}} \prod_{p \in \mathbb{Z}} Hom_{\mathbb{K}}(A^p \otimes B^{n-(p+w)}, C^n) \simeq \prod_{w \in \mathbb{Z}} \prod_{n \in \mathbb{Z}} \prod_{p \in \mathbb{Z}} Hom_{\mathbb{K}}(A^p, Hom_{\mathbb{K}}(B^{n-(p+w)}, C^n)) \\
&\simeq \prod_{w \in \mathbb{Z}} \prod_{p \in \mathbb{Z}} Hom_{\mathbb{K}}(A^p, \prod_{n \in \mathbb{Z}} Hom_{\mathbb{K}}(B^{n-(p+w)}, C^n)) = \prod_{w \in \mathbb{Z}} \prod_{p \in \mathbb{Z}} Hom_{\mathbb{K}}(A^p, Hom_{\mathbb{K}}^{p+w}(B^*, C^*)) \\
&= \prod_{w \in \mathbb{Z}} Hom_{\mathbb{K}}^w(A^*, Hom_{\mathbb{K}}^*(B^*, C^*)) = Hom_{\mathbb{K}}^*(A^*, Hom_{\mathbb{K}}^*(B^*, C^*))
\end{aligned}$$

We give a braiding on homogenous elementary tensors as

$$\beta(a \otimes b) = (-1)^{|a||b|} b \otimes a.$$

It is immediate that  $\beta_{A,b}$  is inverse to  $\beta_{B,A}$ . Observe that this category also admits a braiding where we don't introduce a sign. This does however not work when we want to add differentials to our graded modules, so this is why we stick with this sign. This braiding is also commonly known as Koszul sign convention.

Since  $Mod_{\mathbb{K}}^*$  is a closed symmetric monoidal category it admits electric circuits. Thus the previous results which we have proved by electric circuits also apply to this category, as the proof is identical in this language. One should note that the specific implementation may be different as vertical braiding may work different. We will now study the vertical braiding in more detail. By definition [4] the application of two homogenous morphisms  $f : A \rightarrow A'$  and  $g : B \rightarrow B'$  on elements  $a \in A$  and  $b \in B$  on tensors is

$$(f \otimes g)(a \otimes b) = (-1)^{|g||a|} f(a) \otimes g(b).$$

Viewing  $a$  and  $b$  as generalized elements again, we get Koszul sign rule on morphisms, i.e. given homogenous composable morphisms  $f, f', g, g'$  we get that

$$(f' \otimes g') \circ (f \otimes g) = (-1)^{|g'||f|} (f' \circ f) \otimes (g' \circ g).$$

Electrically we may represent this as a 2 string circuit where a morphism on the left wants to downwards pass a morphism on its right,

$$\begin{array}{c} \text{---} \circ \text{---} \\ \text{---} \end{array} = (-1)^{|g||f|} \begin{array}{c} \text{---} \circ \text{---} \\ \text{---} \end{array}$$

Unlike the other electric equations in which we may substitute parts of an electric circuit with other equal parts this does not however work a priori in this context. This is because how we defined levels. If we are given a 3 string circuit, then the formula changes, this is because we want to manipulate every element on a level simultaneously. In the case of moving a left-most component downwards pass many components, we may regard them as a component on a single string. We will use this to prove a interchange of components on an  $n$ -string circuit formula.

**Proposition 1.1.33.** *Let  $n \geq 1$  and suppose that we have  $a_i \in A_i \rightarrow B_i$  and  $b_i : B_i \rightarrow C_i$  for any  $0 < i \leq n$ . Then we get that*

$$(b_i \circ a_i) \otimes \dots \otimes (b_n \circ a_n) = (-1)^s (b_1 \otimes \dots \otimes b_n) \circ (a_1 \otimes \dots \otimes a_n),$$

$$\text{where } s = \sum_{i=1}^n |b_i| \left( \sum_{1 \leq j < i} |a_j| \right)$$

*Proof.* We prove this by induction. If  $n = 1$  this is true.  $s = 0$  since the sum is empty, so  $b_1 \circ a_1 = (-1)^s b_1 \circ a_1$ .

Assume that the conclusion holds for  $n - 1$ . Assume that we have  $a_i$  and  $b_i$  as in the hypothesis.

Let  $s' = \sum_{i=1}^{n-1} |b_i| \left( \sum_{1 \leq j < i} |a_j| \right)$ , then

$$s = s' + |b_n| \left( \sum_{i=1}^{n-1} |a_i| \right).$$

The conclusion follows from this calculation.

$$\begin{aligned} (b_1 \circ a_1) \otimes \dots \otimes (b_n \circ a_n) &= (-1)^{s'} ((b_1 \otimes \dots \otimes b_{n-1}) \circ (a_1 \otimes \dots \otimes a_{n-1})) \otimes (b_n \circ a_n) \\ &= (-1)^{s' + |b_n| (\sum_{i=1}^{n-1} |a_i|)} (b_1 \otimes \dots \otimes b_n) \circ (a_1 \otimes \dots \otimes a_n) \end{aligned}$$

□

Now that we have developed a graphical calculus for  $Mod_{\mathbb{K}}^*$  we are finally ready to scout for some algebras. We are only concerned with algebras which comes equipped with a derivation.

**Definition 1.1.34** (Differential graded algebra).  $(A^\bullet, d_A)$  is a differential graded algebra if:

- $A^\bullet$  is a differential algebra in  $Mod_{\mathbb{K}}^*$
- The structure morphisms  $(\cdot_A)$  and  $1_A$  are homogenous of degree 0
- The differential  $d_A$  is homogenous and of degree 1
- $d_A^2 = 0$

*Example 1.1.35* (The unit).  $\mathbb{K}^\bullet = (\mathbb{K}, 0)$  is a differential graded algebra in the trivial way. It is concentrated in degree 0 and the differential is the trivial derivation.

**Example 1.1.36** (De Rham complex). Given a manifold  $M$ , the exterior algebra  $\Lambda M$  is a differential graded algebra. See Tu [5] for a deeper explanation.

$M^\bullet$  is called a cochain complex if it comes equipped with a homogenous morphism of degree 1, like  $d_M^\bullet : M^\bullet \rightarrow M^\bullet$ , such that  $d_M^{\bullet 2} = 0$ . This morphism is called differential. A chain morphism  $f : M^\bullet \rightarrow N^\bullet$  is a homogenous  $\mathbb{K}$ -linear morphism of degree 0, such that  $f \circ d_M^\bullet = d_N^\bullet \circ f$ . The category of chain complexes will be denoted as  $ChMod_{\mathbb{K}}$  or  $Mod_{\mathbb{K}}^\bullet$ . Generally  $\mathcal{C}^\bullet$  is the category of chain complexes whenever it makes sense, and the  $\mathbb{K}$ -module of morphisms between two chain complexes is denoted as  $Hom_{\mathbb{K}}^\bullet(M^\bullet, N^\bullet)$ .

The functor  $_{-}[n] : Mod_{\mathbb{K}}^\bullet \rightarrow Mod_{\mathbb{K}}^\bullet$  shifts the degree on each object by adding  $n$  to each grade, it is called the shift functor. Let  $\otimes$  denote the total tensor product in  $Mod_{\mathbb{K}}^\bullet$ . There is an isomorphism between the identity shift functor and total tensor of the stalk of  $\mathbb{K}$ ,  $_{-}[0] \simeq \mathbb{K} \otimes _-$ . In the same manner, shifting  $n$ -fold becomes isomorphic to tensoring with the shifted stalk of  $\mathbb{K}$ ,  $_{-}[n] \simeq \bar{\mathbb{K}}[n] \otimes _-$ . For our purposes we will let  $(A^\bullet, d_A^\bullet)[n] = (A^{\bullet+n}, -d_A^{\bullet+n})$ . The koszul sign rule gives us a switching map for the tensor product. Thus, if  $f^* : A^\bullet \rightarrow B^\bullet$  is a morphism of degree  $k$ , then  $f^*[n] = (-1)^{k \cdot n} f^{*+n}$ .

**Proposition 1.1.37.** *Let  $M^\bullet$  and  $N^\bullet$  be two chain complexes. The graded module of morphisms  $Hom_{\mathbb{K}}^*(M^\bullet, N^\bullet)$  is a chain complex, given by the differential  $\partial(f) = d_N^\bullet \circ f - (-1)^{|f|} f \circ d_M^\bullet$ .*

*Proof.* We observe that  $\partial : Hom_{\mathbb{K}}^*(M^\bullet, N^\bullet) \rightarrow Hom_{\mathbb{K}}^*(M^\bullet, N^\bullet)$  is a morphism of degree 1. It remains to check that  $\partial^2 = 0$ . Pick any homogenous morphism  $f : M^\bullet \rightarrow N^\bullet$ .

$$\begin{aligned} \partial^2(f) &= \partial(d_N^\bullet \circ f - (-1)^{|f|} f \circ d_M^\bullet) = \partial(d_N^\bullet \circ f) - (-1)^{|f|} \partial(f \circ d_M^\bullet) \\ &= -(-1)^{|d_N^\bullet \circ f|} d_N^\bullet \circ f \circ d_M^\bullet - (-1)^{|f|} d_N^\bullet \circ f \circ d_M^\bullet = 0 \end{aligned}$$

□

In an electric diagram we write  $\partial f$  as a sum of circuits.

$$\partial f = \begin{array}{c} \circlearrowleft \\ | \\ \nabla \end{array} + (-1)^{|f|} \begin{array}{c} \nabla \\ | \\ \circlearrowleft \end{array}$$

Observe that  $f : M^\bullet \rightarrow N^\bullet$  of degree 0 is a chain morphism if and only if  $\partial(f) = 0$ . We then observe that  $Hom_{\mathbb{K}}^\bullet(M^\bullet, N^\bullet) \simeq Z^0 Hom_{\mathbb{K}}^*(M^\bullet)$ . Moreover, if  $f$  is a boundary, i.e. there is some  $h$  such that  $f = \partial h$ , then  $f$  null-homotopic. Thus the 0'th homology  $H^0 Hom_{\mathbb{K}}^*(M^\bullet, N^\bullet)$  is the set of every chain map which is not null-homotopic up to null homotopy. We obtain this lemma immediate from this discussion.

**Lemma 1.1.38.** *Suppose that  $N^\bullet$  is a contractible chain complex. Then for any chain complex  $M^\bullet$ , the 0'th homology of  $Hom_{\mathbb{K}}^*(M^\bullet, N^\bullet)$  is  $H^0 Hom_{\mathbb{K}}^*(M^\bullet, N^\bullet) \simeq 0$*



To complete the definitions of graded modules and chain complexes to algebras we would like the structure morphisms to respect the given structure. E.g. if  $a$  and  $b$  are homogenous elements, we would like that the degree of  $ab$  is the sum of its parts, i.e.  $|ab| = |a| + |b|$ . Since multiplication by identity doesn't do anything, we want that the identity lives in the 0'th degree, and so forth.

**Definition 1.1.39** (Graded algebra). Let  $A^*$  be a graded  $\mathbb{K}$ -module. We say that  $A^*$  is a graded algebra if  $A^*$  is an algebra such that  $\nabla_A$  and  $v_A$  are homogenous and of degree 0. Dually,  $C^*$  is a graded coalgebra if  $\Delta_C$  and  $\varepsilon_C$  are homogenous and of degree 0.

**Definition 1.1.40** (Differential graded algebra). Let  $A^\bullet$  be a chain complex over  $\mathbb{K}$ . We say that  $A^\bullet$  is a differential graded algebra, or dg algebra, if it is a graded algebra and the differential is a graded derivation, i.e.  $d_A(ab) = d_A(a)b + (-1)^{|a|}ad_A(b)$ .

Dually,  $C^\bullet$  is a differential graded coalgebra if  $C^\bullet$  is a graded coalgebra and the differential is a graded coderivation.

## 1.2 Cobar-Bar Adjunction

### 1.2.1 Convolution Algebras

Let  $C$  be a coalgebra and  $A$  an algebra, then if  $f, g : C \rightarrow A$  are  $\mathbb{K}$ -linear morphism we may define  $f \star g = \nabla_A(f \otimes g)\Delta_C$ . We call the operation  $\star$  for convolution.

$$f \star g = \begin{array}{c} \diagup \quad \diagdown \\ \textcircled{f} \quad \textcircled{g} \\ \diagdown \quad \diagup \end{array}$$

**Proposition 1.2.1** (Convolution algebra). *The  $\mathbb{K}$ -module  $\text{Hom}_{\mathbb{K}}(C, A)$  is an associative algebra when equipped with convolution  $\star : \text{Hom}_{\mathbb{K}}(C, A) \rightarrow \text{Hom}_{\mathbb{K}}(C, A)$ . The unit is given by  $1 \mapsto v_A \circ \varepsilon_C$ .*

*Proof.* This proposition follows from (co)associativity and (co)unitality of  $(C, A)$ .

$$\begin{aligned} (f \star g) \star h &= \begin{array}{c} \diagup \quad \diagdown \\ \textcircled{f} \quad \textcircled{g} \quad \textcircled{h} \\ \diagdown \quad \diagup \end{array} = \begin{array}{c} \diagup \quad \diagdown \\ \textcircled{f} \quad \textcircled{g} \quad \textcircled{h} \\ \diagdown \quad \diagup \end{array} = \begin{array}{c} \diagup \quad \diagdown \\ \textcircled{f} \quad \textcircled{g} \quad \textcircled{h} \\ \diagdown \quad \diagup \end{array} = f \star (g \star h) \\ (v_A \circ \varepsilon_C) \star f &= \begin{array}{c} \diagup \quad \diagdown \\ \circ \quad \textcircled{f} \\ \diagdown \quad \diagup \end{array} = \begin{array}{c} \diagup \quad \diagdown \\ \textcircled{f} \\ \diagdown \quad \diagup \end{array} = \begin{array}{c} \diagup \quad \diagdown \\ \textcircled{f} \quad \circ \\ \diagdown \quad \diagup \end{array} = f \star (v_A \circ \varepsilon_C) \end{aligned}$$

□

If  $A$  is an algebra and  $C$  is a coalgebra, then they may be given the structure of a differential algebra by attaching the 0 morphism to each algebra as the (co)derivation. In this case proposition 1.1.32 says that a morphism  $f : M \rightarrow A \otimes_{\mathbb{K}} M$  determines the derivation given as  $d_f = (\nabla_A \otimes id_M) \circ (id_A \otimes f)$ . Dually, a morphism  $g : C \otimes_{\mathbb{K}} M \rightarrow M$  determines the coderivation  $d_g = (id_C \otimes g) \circ (\Delta_C \otimes id_N)$ .

If  $\alpha : C \rightarrow A$  is a  $\mathbb{K}$ -linear morphism, then there are two ways to extend  $\alpha$  to obtain a (co)derivation. Precomposing with  $C$ 's comultiplication gives us a morphism from  $C$  to the free  $A$ -module  $A \otimes_{\mathbb{K}} C$ .

$$(\alpha \otimes id_C) \circ \Delta_C : C \rightarrow A \otimes_{\mathbb{K}} C$$

Postcomposing with the multiplication of  $A$  gives us a morphism from to the cofree  $C$ -comodule  $C \otimes_{\mathbb{K}} A$  to  $A$ .

$$\nabla_A \circ (\alpha \otimes id_A) : C \otimes_{\mathbb{K}} A \rightarrow A$$

Notice that when applying proposition 1.1.32 to both morphisms yields the same map, and it is thus both a derivation and a coderivation.

$$d_\alpha = (\nabla_A \otimes id_C) \circ (id_A \otimes \alpha \otimes id_C) \circ (id_A \otimes \Delta_C)$$

$$d_\alpha = \text{[Diagram: A vertical line on the left and a vertical line on the right. A horizontal line connects them at the top. A circle labeled \alpha is on the horizontal line, with a vertical line passing through it. The line enters the circle from the left and exits to the right.]}$$

**Proposition 1.2.2.**  $d_{(\_)} : Hom_{\mathbb{K}}(C, A) \rightarrow End(C \otimes_{\mathbb{K}} A)$  is a morphism of algebras. Moreover, if  $\alpha \star \alpha = 0$ , then  $d_\alpha^2 = 0$ .

*Proof.* The proof quickly follows from (co)associativity and (co)unitality.

$$d_{\alpha \star \beta} = \text{[Diagram: A vertical line on the left and a vertical line on the right. A horizontal line connects them at the top. Two circles labeled \alpha and \beta are on the horizontal line, with a vertical line passing through them. The line enters the first circle from the left and exits to the right, then enters the second circle from the left and exits to the right.]} = \text{[Diagram: A vertical line on the left and a vertical line on the right. A horizontal line connects them at the top. Two circles labeled \alpha and \beta are on the horizontal line, with a vertical line passing through them. The line enters the first circle from the left and exits to the right, then enters the second circle from the left and exits to the right.]} = d_\alpha \circ d_\beta$$

$$d_{v_A \circ \varepsilon_C} = \begin{array}{c} \text{---} \diagup \text{---} \text{---} \text{---} \diagdown \text{---} \\ | \quad | \quad | \quad | \quad | \\ \text{---} \diagdown \text{---} \text{---} \text{---} \diagup \text{---} \end{array} = \begin{array}{c} | \quad | \\ | \quad | \end{array} = id_{C \otimes_{\mathbb{K}} A}$$

□

Suppose that  $C$  and  $A$  are differential graded (co)algebras. We want to expect that the differential  $\partial$  makes  $\text{Hom}_{\mathbb{K}}^*(C, A)$  into a dg-algebra.

**Proposition 1.2.3.** *The convolution algebra  $(\text{Hom}_{\mathbb{K}}^*(C, A), \star)$  is a dg-algebra with differential  $\partial$ .*

*Proof.* We know that  $(\text{Hom}_{\mathbb{K}}^*(C, A), \star)$  is a convolution algebra and that  $(\text{Hom}_{\mathbb{K}}^*(C, A), \partial)$  is a chain complex. It remains to verify that the differential is compatible with the multiplication, i.e.  $\partial(f \star g) = \partial f \star g + (-1)^{|f|} f \star \partial g$ .

Let  $f, g \in \text{Hom}_{\mathbb{K}}^*(C, A)$  be two homogenous morphisms. The key property to arrive at the result is that the differential in a dg-(co)algebra is a (co)derivation. We denote the degree of  $f \star g$  as  $|f \star g| = |f| + |g| = d$

$$\begin{aligned} \partial(f \star g) &= \partial \left( \begin{array}{c} \text{---} \diagup \text{---} \text{---} \text{---} \diagdown \text{---} \\ | \quad | \quad | \quad | \quad | \\ \text{---} \diagdown \text{---} \text{---} \text{---} \diagup \text{---} \end{array} \right) = \begin{array}{c} \text{---} \diagup \text{---} \text{---} \text{---} \diagdown \text{---} \\ | \quad | \quad | \quad | \quad | \\ \text{---} \diagdown \text{---} \text{---} \text{---} \diagup \text{---} \end{array} - (-1)^d \begin{array}{c} \text{---} \diagup \text{---} \text{---} \text{---} \diagdown \text{---} \\ | \quad | \quad | \quad | \quad | \\ \text{---} \diagdown \text{---} \text{---} \text{---} \diagup \text{---} \end{array} \\ &= \begin{array}{c} \text{---} \diagup \text{---} \text{---} \text{---} \diagdown \text{---} \\ | \quad | \quad | \quad | \quad | \\ \text{---} \diagdown \text{---} \text{---} \text{---} \diagup \text{---} \end{array} + (-1)^{|f|} \begin{array}{c} \text{---} \diagup \text{---} \text{---} \text{---} \diagdown \text{---} \\ | \quad | \quad | \quad | \quad | \\ \text{---} \diagdown \text{---} \text{---} \text{---} \diagup \text{---} \end{array} - (-1)^d ((-1)^{|g|} \begin{array}{c} \text{---} \diagup \text{---} \text{---} \text{---} \diagdown \text{---} \\ | \quad | \quad | \quad | \quad | \\ \text{---} \diagdown \text{---} \text{---} \text{---} \diagup \text{---} \end{array} + \begin{array}{c} \text{---} \diagup \text{---} \text{---} \text{---} \diagdown \text{---} \\ | \quad | \quad | \quad | \quad | \\ \text{---} \diagdown \text{---} \text{---} \text{---} \diagup \text{---} \end{array}) \\ &= \begin{array}{c} \text{---} \diagup \text{---} \text{---} \text{---} \diagdown \text{---} \\ | \quad | \quad | \quad | \quad | \\ \text{---} \diagdown \text{---} \text{---} \text{---} \diagup \text{---} \end{array} - (-1)^{|f|} \begin{array}{c} \text{---} \diagup \text{---} \text{---} \text{---} \diagdown \text{---} \\ | \quad | \quad | \quad | \quad | \\ \text{---} \diagdown \text{---} \text{---} \text{---} \diagup \text{---} \end{array} + (-1)^{|f|} \begin{array}{c} \text{---} \diagup \text{---} \text{---} \text{---} \diagdown \text{---} \\ | \quad | \quad | \quad | \quad | \\ \text{---} \diagdown \text{---} \text{---} \text{---} \diagup \text{---} \end{array} - (-1)^{|g|} \begin{array}{c} \text{---} \diagup \text{---} \text{---} \text{---} \diagdown \text{---} \\ | \quad | \quad | \quad | \quad | \\ \text{---} \diagdown \text{---} \text{---} \text{---} \diagup \text{---} \end{array} \\ &= \begin{array}{c} \text{---} \diagup \text{---} \text{---} \text{---} \diagdown \text{---} \\ | \quad | \quad | \quad | \quad | \\ \text{---} \diagdown \text{---} \text{---} \text{---} \diagup \text{---} \end{array} + (-1)^{|f|} \begin{array}{c} \text{---} \diagup \text{---} \text{---} \text{---} \diagdown \text{---} \\ | \quad | \quad | \quad | \quad | \\ \text{---} \diagdown \text{---} \text{---} \text{---} \diagup \text{---} \end{array} = \partial(f) \star g + (-1)^{|f|} f \star \partial(g) \end{aligned}$$

□

### 1.2.2 Twisting Morphisms

In this section we will define twisting morphisms from coalgebras to algebras. They are of importance as the bifunctor  $Tw(C, A)$  is represented in both arguments. To understand the elements of  $Tw$  we start this section by reviewing the Maurer-Cartan equation.

Suppose that  $C$  is a dg-coalgebra and  $A$  is a dg-algebra. We say that a morphism  $\alpha \in \text{Hom}_{\mathbb{K}}^*(C, A)$  is twisting if it is of degree 1 and satisfies the Maurer-Cartan equation:

$$\partial\alpha + \alpha \star \alpha = 0.$$

We say that  $\alpha$  is an element of  $Tw(C, A) \subset \text{Hom}_{\mathbb{K}}^1(C, A) \subset \text{Hom}_{\mathbb{K}}^*(C, A)$ . In light of proposition 1.2.2, every morphism between coalgebras and algebras extend to a unique (co)derivation on the tensor product  $C \otimes_{\mathbb{K}} A$ . Let  $d_{\alpha}^r$  denote this unique morphism. In the case of dg-coalgebras and dg-algebras we perturbate the total differential on the tensor with  $d_{\alpha}^r$ , as in proposition 1.1.32. We call this derivation for the perturbed derivative.


$$d_{\alpha}^{\bullet} = d_{C \otimes_{\mathbb{K}} A}^{\bullet} + d_{\alpha}^r = d_C^{\bullet} \otimes id_A + id_C \otimes d_A^{\bullet} + d_{\alpha}^r$$

**Proposition 1.2.4.** *Suppose that  $C$  is a dg-coalgebra and  $A$  is a dg-algebra, and  $\alpha \in \text{Hom}_{\mathbb{K}}^{-1}(C, A)$ . The perturbed derivation satisfies the following relation.*

$$d_{\alpha}^{\bullet 2} = d_{\partial\alpha + \alpha \star \alpha}^r$$

Moreover, a morphism is twisting if and only if the perturbed derivative is a differential.

*Proof.*  $d_{\alpha}^{\bullet 2} = d_{C \otimes_{\mathbb{K}} A}^{\bullet} \circ d_{\alpha}^r + d_{\alpha}^r \circ d_{C \otimes_{\mathbb{K}} A}^{\bullet} + d_{\alpha}^{r 2}$ . By proposition 1.2.2  $d_{\alpha}^r$  is an algebra homomorphism from the convolution algebra to the endomorphism algebra, thus  $d_{\alpha}^{r 2} = d_{\alpha \star \alpha}^r$ .



By summing the above terms we get

$$d_{C \otimes_{\mathbb{K}} A}^\bullet \circ d_\alpha^r + d_\alpha^r \circ d_{C \otimes_{\mathbb{K}} A}^\bullet = d_{d_A^\bullet \circ \alpha + \alpha \circ d_C^\bullet}^r = d_{\partial \alpha}^r,$$

to obtain the result.

$$d_\alpha^{\bullet 2} = d_{C \otimes_{\mathbb{K}} A}^\bullet \circ d_\alpha^r + d_\alpha^r \circ d_{C \otimes_{\mathbb{K}} A}^\bullet + d\alpha^{r2} = d_{\partial \alpha}^r + d_{\alpha \star \alpha}^r = d_{\partial \alpha + \alpha \star \alpha}$$

□

**Corollary 1.2.4.1.** *If  $\alpha : C \rightarrow A$  is a twisting morphism, then  $(C \otimes_{\mathbb{K}} A, d_\alpha^\bullet)$  is a chain complex. It is called the right twisted tensor product and is denoted as  $C \otimes_\alpha A$ .*

Normally  $A \otimes C$  and  $C \otimes A$  are isomorphic as modules. In general, it is not true that  $C \otimes_\alpha A$  and  $A \otimes_\alpha C$  are isomorphic, since we choose a particular side to perform the twisting. However, if  $A$  is commutative and  $C$  is cocommutative then they are isomorphic. To illustrate we realize the unique derivation above as a right derivative. The left derivative  $d_\alpha^l$  is then defined analogously.

$$d_\alpha^l =$$

*Remark 1.2.5.* Functoriality of  $\otimes_\alpha$  is obtained from the category of elements. I propose that there is an equivalence of categories, that is:

$$\int_{(C,A)} Tw(C, A) \simeq \text{right twisted tensors.}$$

### 1.2.3 Bar and Cobar Construction

The bar construction was first formalized for augmented skew-commutative dg-rings by Eilenberg and Mac Lane [6]. The bar construction then served as a tool to calculate the homology of the Eilenberg-Mac Lane spaces. This construction was later dualized by Adams [7] to obtain the cobar construction. It's first purpose was to serve as a method for constructing an injective resolution in order to calculate the cotensor resolution [8]. With time, the bar-cobar construction have been subjected to much generalization, such as a fattened tensor product on simplicially enriched, tensored and cotensored categories [9]. We will mainly follow the work of [3] to obtain the bar and cobar construction. The approach which we are going to take is also slightly inspired by MacLanes[10] canonical resolutions of comonads.

For our purposes, the bar construction of an augmented algebra is a simplicial resolution with the cofree coalgebra structure. For a dg-algebra, we will realize this resolution as the total complex of its resolution. Dually, the cobar construction of a conilpotent coalgebra is a cosimplicial resolution with the free algebra structure. We will see that these constructions defines an adjoint pair of functors.

**Definition 1.2.6.** The simplex category  $\Delta$  consists of ordered sets  $[n] = \{0, \dots, n\}$  for any  $n \in \mathbb{N}$ . A morphism in  $\Delta$  is a monotone function between the sets.

$\Delta_+$  is the augmented simplex category, where we add the object  $[-1] = \emptyset$ .  $\Delta^+$  is the non-full subcategory of  $\Delta$ , where all morphisms are injective functions.

The simplex category comes equipped with coface and codegeneracy morphisms. The coface maps are the injective morphisms  $\delta_i : [n] \rightarrow [n+1]$ , and the codegeneracy maps are the surjective morphisms  $\sigma_i : [n] \rightarrow [n-1]$ .

$$\delta_i(k) = \begin{cases} k, & \text{if } k < i \\ k+1, & \text{otherwise} \end{cases} \quad \sigma_i(k) = \begin{cases} k, & \text{if } k \leq i \\ k-1, & \text{otherwise} \end{cases}$$

Every morphism in  $\Delta$  may be realized as a composition of coface and codegeneracy maps, see [10]. Furthermore, these maps are characterized by some identities, called the cosimplicial identities.

1.  $\delta_j \delta_i = \delta_i \delta_{j-1}$ , if  $i < j$
2.  $\sigma_j \delta_i = \delta_i \sigma_{j-1}$ , if  $i < j$
3.  $\sigma_j \delta_i = id$ , if  $i = j$  or  $i = j + 1$
4.  $\sigma_j \delta_i = \delta_{i-1} \sigma_j$ , if  $i > j + 1$
5.  $\sigma_j \sigma_i = \sigma_i \sigma_{j+1}$ , if  $i \leq j$

We may arrange the arrows of the augmented simplex category in the following way:

$$\begin{array}{ccccccc} [-1] & \longrightarrow & [0] & \xrightarrow{\delta_i} & [1] & \xrightarrow{\delta_i} & [2] & \xrightarrow{\delta_i} & \dots \\ & & & & & & & & \\ [-1] & & [0] & \xleftarrow{\sigma_1} & [1] & \xleftarrow{\sigma_i} & [2] & \xleftarrow{\sigma_i} & \dots \end{array}$$

Let  $\mathcal{C}$  be a category. A simplicial object in  $\mathcal{C}$  is a functor  $S : \Delta^{op} \rightarrow \mathcal{C}$ . It may be viewed as a collection of objects  $\{S_n\}_{n \in \mathbb{N}}$  together with face maps  $d^i : S_n \rightarrow S_{n-1}$  and degeneracy maps  $s^i : S_n \rightarrow S_{n+1}$  satisfying the simplicial identities. An augmented simplicial object is a functor  $S : \Delta_+^{op} \rightarrow \mathcal{C}$ . The restricted functor  $\bar{S} : \Delta^{op} \rightarrow \mathcal{C}$  is the augmentation ideal of  $S$ . An augmented semi-simplicial object is a functor  $S : (\Delta_+^+)^{op} \rightarrow \mathcal{C}$ . Dually, a cosimplicial object is a functor  $S : \Delta \rightarrow \mathcal{C}$ , it may be regarded as a sequence of objects with coface and codegeneracy maps satisfying the cosimplicial identities.

Let  $\mathcal{A}$  be an abelian category. To each semi-simplicial object  $M : (\Delta^+)^{op} \rightarrow \mathcal{A}$  there is an associated chain complex  $M^\bullet$ . Let  $M^{-i} = M_i$  with differential  $d_M^{-n} = \sum_{i=1}^{n+1} (-1)^{i-1} d^i$ . This differential is well-defined by simplicial identity 1.

$$\dots \longrightarrow M_2 \xrightarrow{d^1-d^2+d^3} M_1 \xrightarrow{d^1-d^2} M_0 \xrightarrow{0} 0 \longrightarrow \dots$$

As face maps and degeneracy maps have the same identities, but flipped around, we could also have defined a chain complex by using the degeneracies instead.

The augmented simplex category has a universal monoid. Let  $+$  :  $\Delta_+ \times \Delta_+ \rightarrow \Delta_+$  be a functor acting on objects and morphisms as:

$$[m] + [n] = [m + n + 1]$$

$$(f + g)(k) = \begin{cases} f(k), & \text{if } k \leq m \\ g(k) + m, & \text{otherwise} \end{cases}$$

Notice that  $[-1] + \_ \simeq Id_{\Delta}$ , so  $(\Delta, +, [-1])$  is a monoidal category. Since  $[0]$  is terminal in  $\Delta$  it becomes a monoid with  $\delta_{-1} : [-1] \rightarrow [0]$  as unit and  $\sigma_0 : [1] \rightarrow [0]$  as multiplication. Associativity and unitality is satisfied by uniqueness of morphisms  $f : [n] \rightarrow [0]$ .

**Proposition 1.2.7.** *Let  $(\mathcal{C}, \otimes, Z)$  be a monoidal category. If  $(C, \eta, \mu)$  is a monoid in  $\mathcal{C}$ , then there is a strong monoidal functor  $F : \Delta_+ \rightarrow \mathcal{C}$ , such that  $F[0] \simeq C$ ,  $F\delta_{-1} \simeq \eta$  and  $F\sigma_0 \simeq \mu$ .*

*Proof.* This is proved in Mac Lanes book [10]. □

An algebra  $A$  is a monoid in the monoidal category  $(Mod_{\mathbb{K}}, \otimes_{\mathbb{K}}, \mathbb{K})$ . By proposition 1.2.7 we may think of  $A$  as an augmented cosimplicial object  $A : \Delta_+ \rightarrow Mod_{\mathbb{K}}$ . Notice that all of the cosimplicial identities follow from associativity and unitality. If  $A$  is an augmented algebra, we may instead give it the structure of an augmented simplicial set. Let  $d_0^0 = \varepsilon_A$  be the augmentation. We define  $d_n^n = A^{\otimes n-1} \otimes \varepsilon_A$  and set  $d_n^i = A^{i-1} \otimes \nabla_A \otimes A^{\otimes n-i-1}$ . All the degeneracies are set to be the units, i.e.  $s_n^i = A^{\otimes i} \otimes v_A \otimes A^{\otimes n-i-1}$ . One may check that this structure defines an augmented simplicial object  $A : \Delta_+^{op} \rightarrow Mod_{\mathbb{K}}$ . Observe that the associated chain complex  $A^\bullet$  is exactly the Hochschild complex of  $A$ . We depict the simplicial object as the following diagram:

$$\begin{array}{ccccccc} \mathbb{K} & \xleftarrow{\varepsilon_A} & A & \xleftarrow[A \otimes \varepsilon_A]{\nabla_A} & A^{\otimes 2} & \xleftarrow[A^{\otimes 2} \otimes \varepsilon_A]{\nabla_A} & A^{\otimes 3} & \xleftarrow[A^{\otimes 4} \otimes \varepsilon_A]{\nabla_A} & \dots \\ & & & & & & & & \\ \mathbb{K} & & A & \xrightarrow{s^1} & A^{\otimes 2} & \xrightarrow{s^i} & A^{\otimes 3} & \xrightarrow{s^i} & \dots \end{array}$$

The augmentation ideal  $\bar{A}$  carries a natural semi-simplicial structure induced by  $A$ . By restricting each of the face maps  $\bar{d}^i = d^i|_{\bar{A}} : \bar{A}^{\otimes n} \rightarrow \bar{A}^{\otimes n-1}$  we obtain the maps together with the simplicial identity 1. This is the non-unital Hochschild complex of  $A$ . We may depict the semi-simplicial object as the following diagram:

$$\mathbb{K} \xleftarrow{0} \bar{A} \xleftarrow[0]{\nabla_A} \bar{A}^{\otimes 2} \xleftarrow[0]{\nabla_A} \bar{A}^{\otimes 3} \xleftarrow[0]{\nabla_A} \dots$$

As graded modules, the chain complex  $\bar{A}^\bullet$  is isomorphic to  $T^c(\bar{A})$ . Here we think of the grading  $T^c(\bar{A})$  as starting at 0 and going down to negative degrees. Consider instead the looped, or shifted, non-unital algebra  $\bar{A}[1]$ . Every algebra may be considered as a graded algebra concentrated in degree 0, the shift functor then recontextualize the degree the algebra is concentrated in. Since the tensor product respects shifting, we see that shifting may be regarded as a tensor.

$$M[n] \simeq \mathbb{K} \otimes M[n] \simeq \mathbb{K}[n] \otimes M \simeq \mathbb{K}\{\sigma\} \otimes M$$

Here  $\sigma$  is a formal generator of  $\mathbb{K}$  such that  $|\sigma| = -n$ . If we let  $\omega$  be a formal generator s.t.  $|\omega| = -1$ , then  $\bar{A}[1] \simeq \mathbb{K}\{\omega\} \otimes \bar{A}$ . Every element of  $\bar{A}[1]$  may then be written on the form  $\omega a$ , where  $a \in \bar{A}$ . Using MacLanes and Eilenbergs notation [6], we would write  $[a]$  instead.

The looped algebra  $\bar{A}[1]$  is no longer associative. For  $\mathbb{K}\{\omega\}$  we define a map  $(\cdot)$ .

$$\begin{aligned} (\cdot) : \mathbb{K}\{\omega\}^{\otimes 2} &\rightarrow \mathbb{K}\{\omega\} \\ \omega \cdot \omega &= \omega \end{aligned}$$

The looped multiplication may then be defined as the composite  $\nabla_{\bar{A}[1]} = ((\cdot) \otimes \nabla_{\bar{A}}) \circ (\mathbb{K}\{\omega\} \otimes \beta \otimes \bar{A})$ . Multiplying elements like  $\omega a_1$  and  $\omega a_2$  would then look like:

$$\nabla_{\bar{A}[1]}(\omega a_1 \otimes \omega a_2) = (-1)^{|a_1||\omega|}((\cdot) \otimes \nabla_{\bar{A}})(\omega^{\otimes 2} \otimes a_1 \otimes a_2) = (-1)^{|a_1|} \omega a_1 a_2$$

It is now also evident that the new map  $\nabla_{\bar{A}[1]}$  is no longer a morphism of degree 0, but of degree 1.

**Proposition 1.2.8.** *The differential  $d_{\bar{A}[1]}^\bullet$  is a coderivation for the cofree coalgebra  $T^c(\bar{A}[1])$ . Thus  $(\bar{A}[1]^\bullet, d_{\bar{A}[1]}^\bullet)$  is a dg-coalgebra.*

*Proof.* By injecting  $\bar{A}[1]$  into  $T^c(\bar{A}[1])$  we may think of  $\nabla_{\bar{A}[1]} : \bar{A}[1]^{\otimes 2} \rightarrow T^c(\bar{A}[1])$  as a morphism into the tensor coalgebra. By using proposition 1.1.29,  $\nabla_{\bar{A}[1]}$  extends uniquely into a coderivation:

$$d_{\bar{A}[1]}^c = \sum_{n=0}^{\infty} \sum_{i=0}^n (\nabla_{\bar{A}[1]})_{(i)}^{(n)} = d_{\bar{A}[1]}^\bullet.$$

□

If  $(A, d_A^\bullet)$  is an augmented dg-algebra, then  $A$  is a simplicial object of  $Mod_{\mathbb{K}}^\bullet$ . It has an associated chain complex. Taking the alternate sum of face maps gives us a double complex as below. We define the double complex  $A^\bullet$  as the associated chain complex to  $A$ .



$$\begin{array}{ccccccc}
\cdots & \xrightarrow{0} & 0 & \xrightarrow{0} & \mathbb{K} & \xrightarrow{0} & 0 \xrightarrow{0} \cdots & A[1]^{\otimes 0} \\
& & \varepsilon_A \uparrow & & \varepsilon_A \uparrow & & \varepsilon_A \uparrow & \uparrow \\
\cdots & \xrightarrow{d_{A[1]}^\bullet} & A^{-1} & \xrightarrow{d_{A[1]}^\bullet} & A^0 & \xrightarrow{d_{A[1]}^\bullet} & A^1 \xrightarrow{d_{A[1]}^\bullet} \cdots & A[1]^{\otimes 1} \\
& & \nabla_A \uparrow \uparrow A \otimes \varepsilon_A & & \nabla_A \uparrow \uparrow A \otimes \varepsilon_A & & \nabla_A \uparrow \uparrow A \otimes \varepsilon_A & = \uparrow \\
\cdots & \xrightarrow{d_{A[1]^{\otimes 2}}^\bullet} & (A^{\otimes 2})^{-1} & \xrightarrow{d_{A[1]^{\otimes 2}}^\bullet} & (A^{\otimes 2})^0 & \xrightarrow{d_{A[1]^{\otimes 2}}^\bullet} & (A^{\otimes 2})^1 \xrightarrow{d_{A[1]^{\otimes 2}}^\bullet} \cdots & A[1]^{\otimes 2} \\
& & \nabla_A \uparrow \uparrow \uparrow A^{\otimes 2} \otimes \varepsilon_A & & \nabla_A \uparrow \uparrow \uparrow A^{\otimes 2} \otimes \varepsilon_A & & \nabla_A \uparrow \uparrow \uparrow A^{\otimes 2} \otimes \varepsilon_A & \uparrow \\
& & \vdots & & \vdots & & \vdots & \vdots
\end{array}$$

For simplicity we write  $d_1$  for the horizontal differential and  $d_2$  for the vertical differential. The total associated chain complex is the total complex for  $Tot(A^\bullet)$ , denoted  $A^\bullet$  if there are no confusion. Whenever we go down to the total complex we should switch out the multiplication with the suspended multiplication. The following proposition shows that this is well-defined, among other things. This total complex is then isomorphic, as graded modules, to  $T^c(\bar{A}[1])$

**Proposition 1.2.9.** *Let  $A$  an augmented dg-algebra. The bar complex  $BA$  is the total associated chain complex of the augmentation ideal  $\bar{A}$ .  $(BA, d_{BA}^\bullet)$  is the cofree conilpotent coalgebra equipped with  $d_{BA}^\bullet = d_1 + d_2$  as coderivation.*

*Proof.* It is apparent that  $d_1$  and  $d_2$  are coderivations with respect to deconcatenation. Since the multiplication  $\nabla_A$  is a chain map, we should have  $d_{BA}^{\bullet 2} = d_1 \circ d_2 + d_2 \circ d_1 = 0$ . This will be shown for each element in  $A^{\otimes 2}$ , the result may then be extended to all of  $BA$ . In the following calculation we will not decorate each  $a$  with an  $\omega$ , and it should be understood that these elements are looped.

$$\begin{aligned}
d_1 \circ d_2(a_1 \otimes a_2) &= (-1)^{|a_1|} d_1(a_1 a_2) = (-1)^{|a_1|} d_{A[1]}^\bullet(a_1 a_2) \\
&= (-1)^{|a_1|+1} d_A^\bullet(a_1 a_2) = (-1)^{|a_1|+1} (d_A^\bullet(a_1) a_2 + (-1)^{|a_1|} a_1 d_A^\bullet(a_2)) \\
&= (-1)^{|a_1|+1} d_A^\bullet(a_1) a_2 - a_1 d_A^\bullet(a_2)
\end{aligned}$$

$$\begin{aligned}
d_2 \circ d_1(a_1 \otimes a_2) &= d_2 \circ (d_{A[1]}^\bullet \otimes id_{A[1]} + id_{A[1]} \otimes d_{A[1]}^\bullet[1])(a_1 \otimes a_2) \\
&= -d_2 \circ (d_A^\bullet(a_1) \otimes a_2 + (-1)^{|a_1|+1} a_1 \otimes d_A^\bullet(a_2)) \\
&= (-1)^{|d_A^\bullet(a_1)|+1} d_A^\bullet(a_1) a_2 + (-1)^{2|a_1|+2} a_1 d_A^\bullet d_A^\bullet(a_2) \\
&= (-1)^{|a_1|} d_A^\bullet(a_1) a_2 + a_1 d_A^\bullet(a_2) = -d_1 \circ d_2(a_1 \otimes a_2)
\end{aligned}$$

□

**Remark 1.2.10.** For now we don't need to show that  $BA$  is a functor. This property follows from  $BA$  being the representing object of  $Tw(\_, A)$ .

On the other hand, a coalgebra  $C$  is a comonoid in  $Mod_{\mathbb{K}}$ . By the dual of proposition 1.2.7 we may think of it as an augmented simplicial object  $C : (\Delta_+)^{op} \rightarrow Mod_{\mathbb{K}}$ . Dually, all of the simplicial identities follows from coassociativity and counitality. A coaugmented coalgebra  $C$  may be given an augmented cosimplicial structure in the opposite way of algebras. We then get that the coaugmentation quotient  $\bar{C}$  is a semi-cosimplicial object of  $Mod_{\mathbb{K}}$ . Observe that  $\bar{C}$  has an associated chain complex like  $\bar{A}$ , but every arrow goes in the opposite direction.

$$\begin{array}{ccccccc} \mathbb{K} & \xrightarrow{v_C} & C & \xrightleftharpoons[A \otimes v_C]{\Delta_C} & C^{\otimes 2} & \xrightleftharpoons[C^{\otimes 2} \otimes v_C]{\Delta_C} & C^{\otimes 3} \xrightleftharpoons[C^{\otimes 3} \otimes v_C]{\Delta_C} \dots \\ & & & & & & \\ \mathbb{K} & & C & \xleftarrow{s_1} & C^{\otimes 2} & \xleftarrow{s_i} & C^{\otimes 3} \xleftarrow{s_i} \dots \end{array}$$

The cobar construction is made from the inverse shifted, or suspended coalgebra  $C[-1]$ . We may also denote suspension by tensoring with the formal generator  $s$ , where  $|s| = 1$ . That is we have the isomorphism  $C[-1] \simeq \mathbb{K}\{s\} \otimes C$ . The cobar construction is realized as the free tensor algebra  $T(\bar{C}[-1])$ , where the comultiplication  $\Delta_{\bar{C}[-1]}$  induces a derivation  $d_{\bar{C}[-1]}$  by proposition 1.1.29.

**Remark 1.2.11.** As we have chosen to define  $\nabla_{A[1]}(a_1 \otimes a_2) = (-1)^{|a_1|} a_1 a_2$ , we are forced by the linear dual to define  $\Delta_{C[-1]}(c) = -(-1)^{|c(1)|} c_{(1)} \otimes c_{(2)}$ . Here we use the Sweedlers notation without sums to denote the comultiplication. Note that this really should be a sum of many different elementary tensors. Lastly, observe that this definition also agrees with Koszuls sign rule.

**Proposition 1.2.12.** *Let  $C$  be a coaugmented dg-coalgebra. The cobar complex  $\Omega C$  is the total associated chain complex of the coaugmentation quotient  $\bar{C}$ .  $(\Omega C, D_{\Omega C}^\bullet)$  is the free algebra equipped with  $d_{\Omega C}^\bullet = d_1 + d_2$  as derivation.*

$$\begin{array}{ccccccc} & \vdots & & \vdots & & \vdots & \\ & \Delta_C \otimes \bar{C} \uparrow \uparrow & & \Delta_C \otimes \bar{C} \uparrow \uparrow & & \Delta_C \otimes \bar{C} \uparrow \uparrow & \\ \dots & \xrightarrow{d_{\bar{C}[-1]}^{\otimes 2}} & (\bar{C}^{\otimes 2})^{-1} & \xrightarrow{d_{\bar{C}[-1]}^{\otimes 2}} & (\bar{C}^{\otimes 2})^0 & \xrightarrow{d_{\bar{C}[-1]}^{\otimes 2}} & (\bar{C}^{\otimes 2})^1 \xrightarrow{d_{\bar{C}[-1]}^{\otimes 2}} \dots \\ & \Delta_C \uparrow & & \Delta_C \uparrow & & \Delta_C \uparrow & \\ \dots & \xrightarrow{d_{\bar{C}[-1]}} & \bar{C}^{-1} & \xrightarrow{d_{\bar{C}[-1]}} & \bar{C}^0 & \xrightarrow{d_{\bar{C}[-1]}} & \bar{C}^1 \xrightarrow{d_{\bar{C}[-1]}} \dots \\ & \uparrow & & 0 \uparrow & & \uparrow & \\ \dots & \longrightarrow & 0 & \longrightarrow & \mathbb{K} & \longrightarrow & 0 \longrightarrow \dots \end{array}$$

We will now see that the bar and cobar construction defines an adjoint pair of functors. Note that since for any conilpotent dg-coalgebra  $C$ , the object  $\Omega C$  represents the functor in the category

of augmented algebras. By Yoneda's lemma, the data of morphisms are then defined, so  $\Omega$  does truly define a functor.

**Theorem 1.2.13.** *Let  $C$  be a conilpotent dg-coalgebra and  $A$  an augmented dg-algebra. The functor  $Tw(C, A)$  is represented in both arguments, i.e.*

$$Alg_{\mathbb{K},+}^{\bullet}(\Omega C, A) \simeq Tw(C, A) \simeq CoAlg_{\mathbb{K},Conil}^{\bullet}(C, BA).$$

*Proof.* We will show that  $\Omega C$  represents the set of twisting morphisms in the first argument. Showing that  $BA$  represents the second argument uses every dual proposition. Thus, it is necessary that  $C$  is conilpotent, in order to dualize the arguments.

Suppose that  $f : \Omega C \rightarrow A$  is an augmented dg-algebra homomorphism.  $f$  is then a morphism of degree 0. By freeness,  $f$  is uniquely determined by a morphism  $f|_{\bar{C}[-1]} : \bar{C}[-1] \rightarrow \bar{A}$  of degree 0, which corresponds to a morphism  $f' : C \rightarrow A$  of degree 1.

Since  $f$  is a morphism of chain complexes it commutes with the differential, i.e.

$$\begin{aligned} f \circ d_{\Omega C}^{\bullet} &= d_A^{\bullet} \circ f \\ f \circ (d_1 + d_2) &= d_A^{\bullet} \circ f \end{aligned}$$

By free-forget, this is equivalent to say that  $-f' \circ d_C^{\bullet} - f' \star f' = d_A^{\bullet} \circ f'$ . Thus  $f'$  is a twisting morphism as desired.  $\square$

For our convenience, we will give these isomorphisms some names. Whenever  $\tau : C \rightarrow A$  is a twisting morphism, the induced morphism of algebras is denoted  $f_{\tau} : \Omega C \rightarrow A$  and the induced morphism of coalgebras is denoted  $g_{\tau} : C \rightarrow BA$ .

*Remark 1.2.14.* These functors consists of a composition with the augmentation ideal (quotient) and then the (co)free tensor (co)algebra. By reversing these operations we obtain another adjunction which is more or less the same adjunction. By abuse of notation we will call these functors for the bar and cobar construction as well, and they induce an adjoint pair between dg-algebras and reduced conilpotent dg-coalgebras. In other words, given a dg-algebra  $A$  and a reduced conilpotent dg-coalgebra  $C$ ,  $BA = \bar{T}^c(A[1])$  and  $\Omega C = \bar{T}(C[-1])$ .

$$\begin{array}{ccc} & \Omega & \\ \swarrow & & \searrow \\ Alg_{\mathbb{K}}^{\bullet} & \perp & CoAlg_{\mathbb{K},conil}^{\bullet} \\ \searrow & & \swarrow \\ & B & \end{array}$$

Associated to this adjunction, we obtain universal elements, together with universal properties. Let  $A$  be an augmented dg-algebra, then the identity of the coalgebras  $id_{BA} : BA \rightarrow BA$ , the counit  $\varepsilon_A : \Omega BA \rightarrow A$  and a twisting morphism  $\pi_A : BA \rightarrow A$  are equivalent by the adjunction and representation. Dually, the identity of algebras  $id_{\Omega C} : \Omega C \rightarrow \Omega C$ , the unit  $\eta_C : C \rightarrow B\Omega C$  and the twisting morphism  $\iota_C : C \rightarrow \Omega C$  are equivalent. The morphisms  $\pi_A$  and  $\iota_C$  are called the universal elements. We summarize their universal property in the following corollary.

**Corollary 1.2.14.1.** *Let  $A$  be an augmented dg-algebra, and  $C$  a conilpotent dg-coalgebra. Any twisting morphism  $\alpha : C \rightarrow A$  factors uniquely through either  $\pi_A$  or  $\iota_C$ .*

$$\begin{array}{ccc}
 & \Omega C & \\
 \iota_C \nearrow & & \searrow g_\alpha \\
 C & \xrightarrow{\alpha} & A \\
 \searrow f_\alpha & & \nearrow \pi_A \\
 & BA &
 \end{array}$$

Moreover, the morphism  $f_\alpha$  is a morphism of dg-coalgebras, and  $g_\alpha$  is a morphism of dg-algebras.

**Definition 1.2.15** (Augmented Bar-Cobar construction). Let  $A$  be an augmented dg-algebra. The (right) augmented bar construction is the right twisted tensor product  $BA \otimes_{\pi_A} A$ , where  $\pi_A$  is the universal twisting morphism.

Let  $C$  be a conilpotent dg-coalgebra. The (right) augmented cobar construction is the right twisted tensor product  $C \otimes_{\iota_C} \Omega C$ , where  $\iota_C$  is the universal twisting morphism.

*Remark 1.2.16.* We could have defined the augmented bar-cobar construction as the left twisted tensor product. There is really no preference of handedness. Whenever we wish to be precise which handedness we will use it will be specified, e.g. the left augmented bar construction of  $A$ .

**Proposition 1.2.17.** *The augmentation ideal of the augmented bar (cobar) construction is acyclic, i.e.  $BA \bar{\otimes}_{\pi_A} A$  ( $A \bar{\otimes}_{\pi_A} BA$ ) and  $C \bar{\otimes}_{\iota_C} \Omega C$  ( $\Omega C \bar{\otimes}_{\iota_C} C$ ) are acyclic.*

*Proof.* We will postpone this proof until chapter 3. This is a part of the fundamental theorem of twisting morphisms, and will not be relevant until then.  $\square$

## 1.3 Strongly Homotopy Associative Algebras and Coalgebras

### 1.3.1 SHA-Algebras

We have seen from corollary 1.2.8 that any dg-algebra  $A$  defines a dg-coalgebra  $T^c(A[1])$ , the bar construction, with a coderivation  $m^c$  of degree  $-1$ . Does this however work in reverse? I.e. if  $A$  is a vector space such that the coalgebra  $T^c(A[1])$  together with a coderivation  $m^c$  is a dg-coalgebra, is then  $A$  an algebra? The answer to this is no, but it leads to the definition of a strongly homotopy associative algebra.

**Definition 1.3.1.** An  $A_\infty$ -algebra is a graded vector space  $A$  together with a differential  $m : \bar{T}^c(A[1]) \rightarrow \bar{T}^c(A[1])$  that is a coderivation of degree 1.

The differential  $m$  induces structure morphisms on  $A[1]$ . By proposition 1.1.29 there is a natural bijection  $\text{Hom}_{\mathbb{K}}(\bar{T}^c(A[1]), A[1]) \simeq \text{Coder}(\bar{T}^c(A[1]), \bar{T}^c(A[1]))$  given by the projection onto  $A[1]$ . Thus  $m : \bar{T}^c(A[1]) \rightarrow \bar{T}^c(A[1])$  corresponds to maps  $\tilde{m}_n : A[1]^{\otimes n} \rightarrow A[1]$  of degree 1 for any  $n \geq 1$ . We define maps  $m_n : A^{\otimes n} \rightarrow A$  by the composite  $s\tilde{m}_n\omega^{\otimes n}$ . Since  $\omega^{\otimes n}$  is of degree  $-n$ ,  $\tilde{m}_n$  and  $s$  is of degree 1, we get that  $m_n$  is of degree  $2 - n$ .

$$\begin{array}{ccc} A^{\otimes n} & \xrightarrow{m_n} & A \\ \omega^{\otimes n} \downarrow \simeq & & \uparrow \simeq s \\ A[1]^{\otimes n} & \xrightarrow{\tilde{m}_n} & A[1] \end{array}$$

**Proposition 1.3.2.** *An  $A_{\infty}$ -algebra is equivalent to a graded vector space  $A$  together with homogenous morphisms  $m_n : A^{\otimes n} \rightarrow A$  of degree  $2 - n$ . Moreover, the morphism must satisfy the following relations for any  $n \geq 1$ :*

$$(\text{rel}_n) \quad \sum_{p+q+r=n} (-1)^{pq+r} m_{p+1+r} \circ (id^{\otimes p} \otimes m_q \otimes id^{\otimes r}) = 0$$

**Remark 1.3.3.** We make a more convenient notation for  $(\text{rel}_n)$ , called partial composition  $\circ_i$ .

$$\begin{aligned} m_{p+1+r} \circ_{p+1} m_q &= m_k \circ (id^{\otimes p} \otimes m_q \otimes id^{\otimes r}) \\ (\text{rel}_n) \quad \sum_{p+q+r=n} (-1)^{pq+r} m_{p+1+r} \circ_{p+1} m_q &= 0 \end{aligned}$$

Before starting with the proof we will use a lemma for checking whether a coderivation  $m : T^c(A) \rightarrow T^c(A)$  is a differential.

**Lemma 1.3.4.** *Let  $m : T^c(A) \rightarrow T^c(A)$  be a coderivation, and denote  $m_n = m|_{A^{\otimes n}}$ .  $m$  is a differential if and only if the following relations are satisfied:*

$$\sum_{p+q+r=n} m_{p+1+r} \circ_{p+1} m_q = 0$$

*Proof.* By proposition 1.1.29 we may write  $m = \sum_{n=0}^{\infty} \sum_{i=0}^n m_{(n)}^{(i)}$ . By using partial composition, we rewrite its  $n$ 'th component as:

$$m_n = \sum_{q=1}^n \sum_{p=1}^n id^{\otimes(n-q)} \circ_p m_q = \sum_{p+q+r=n} id^{\otimes(p+1+r)} \circ_{p+1} m_q$$

For  $m^2$  we denote its  $n$ 'th component as  $m_n^2$ . Observe the following:

$$\begin{aligned} m_n^2 &= m \circ m_n = m \circ \sum_{p+q+r=n} id^{\otimes(p+1+r)} \circ_{p+1} m_q = \sum_{p+q+r=n} m \circ_{p+1} m_q \\ \pi m_n^2 &= \pi \sum_{p+q+r=n} m \circ_{p+1} m_q = \sum_{p+q+r=n} m_{p+1+r} \circ_{p+1} m_q \end{aligned}$$

Since every coderivation are uniquely determined by  $\pi$ , its projection onto  $A$  we get that  $m^2 = 0$  if and only if

$$\sum_{p+q+r=n} m_{p+1+r} \circ_{p+1} m_q = 0.$$

□

*Proof of proposition 1.3.2.* Let  $(A, m)$  be an  $A_\infty$ -algebra. We denote the  $n$ 'th component of  $m$  as  $\tilde{m}_n$ . The  $n$ 'th components thus define maps  $m_n : A^{\otimes n} \rightarrow A$  as  $m_n = s\tilde{m}_n\omega^{\otimes n}$ .

By the above lemma we know that the  $n$ 'th component of  $m^2$  is:

$$\begin{aligned} \sum_{p+q+r=n} \tilde{m}_{p+1+r} \circ_{p+1} \tilde{m}_q \\ = \sum_{p+q+r=n} \omega m_{p+1+r} s^{\otimes(p+1+r)} \circ_{p+1} \omega m_q s^{\otimes q} = \sum_{p+q+r=n} (-1)^{pq+r} \omega m_{p+1+r} \circ_{p+1} m_q s^{\otimes n} \end{aligned}$$

Since suspension and desuspension are isomorphism we get that  $m^2 = 0$  if and only if  $(\text{rel}_n)$  are 0 for every  $n \geq 1$ , i.e.

$$\sum_{p+q+r=n} (-1)^{pq+r} m_{p+1+r} \circ_{p+1} m_q = 0$$

□

Given an  $A_\infty$  algebra  $A$  we may either think of it as a differential tensor coalgebra  $\bar{T}^c(A[1])$  with differential  $m : \bar{T}^c(A[1]) \rightarrow \bar{T}^c(A[1])$  or as a graded vector space with morphisms  $m_n : A^{\otimes n} \rightarrow A$  satisfying  $(\text{rel}_n)$ . We will calculate  $(\text{rel}_n)$  for 1, 2, 3:

$$\begin{aligned} (\text{rel}_1) \quad m_1 \circ m_1 &= 0 \\ (\text{rel}_2) \quad m_1 \circ m_2 - m_2 \circ_1 m_1 - m_2 \circ_2 m_1 &= 0 \\ (\text{rel}_3) \quad m_1 \circ m_3 - m_2 \circ_1 m_2 + m_2 \circ_2 m_2 + m_3 \circ_1 m_1 + m_3 \circ_2 m_1 + m_3 \circ_3 m_1 &= 0 \end{aligned}$$

We see that  $(\text{rel}_1)$  states that  $m_1$  should be a differential, we may thus think of  $(A, m_1)$  as a chain complex. Furthermore,  $(\text{rel}_2)$  says that  $m_2 : (A^{\otimes 2}, m_1 \otimes id_A + id_A \otimes m_1) \rightarrow (A, m_1)$  is a morphism of chain complexes. Lastly,  $(\text{rel}_3)$  gives us a homotopy for the associator of  $m_2$ , namely  $m_3$ . Thus we may regard  $(A, m_1, m_2)$  as an algebra which is associative up to homotopy. Regarding  $A$  as a chain complex instead we obtain our final definition of an  $A_\infty$ -algebra.

**Proposition 1.3.5.** Suppose that  $(A, d)$  is a chain complex, and that there exists morphisms  $m_n : A^{\otimes n} \rightarrow A$  of degree  $2 - n$  for any  $n \geq 2$ .  $A$  is an  $A_\infty$ -algebra if and only it satisfies the following relations:

$$(\text{rel}'_n) \quad \partial(m_n) = - \sum_{\substack{n=p+q+r \\ k=p+1+r \\ k>1, q>1}} (-1)^{pq+r} m_k \circ_{p+1} m_q$$

We define the homotopy of an  $A_\infty$ -algebra to be the homology of the chain complex  $(A, m_1)$ . Since  $\partial(m_3) = m_2 \circ_1 m_2 - m_2 \circ_2 m_2$ , we get that  $m_2$  is associative in homology. Thus for any  $A_\infty$ -algebra  $A$ , the homotopy  $HA$  is an associative algebra. The operadic homology of  $A$  is defined as the homology of  $(T^c(A[1]), m)$ , which is the non-unital augmented Hochschild homology of  $A$ .

**Example 1.3.6.** Suppose that  $V$  is a cochain complex with differential  $d$ . Then  $V$  is an  $A_\infty$ -algebra with trivial multiplication. In other words  $m^1 = d$  and  $m^i = 0$  for any  $i > 1$ .

**Example 1.3.7.** Suppose that  $A$  is a dg-algebra. Then  $A$  is an  $A_\infty$ -algebra where  $m^1 = d$ ,  $m^2 = \cdot$  and  $m^i = 0$  for any  $i > 2$ .

**Example 1.3.8.** Let  $A$  be a connected weight-graded algebra over  $\mathbb{K}$ . We may then think of  $\mathbb{K}$  as an  $A$ -module with trivial action, i.e. as the quotient  $A/A^{(i)}_{(i \geq 1)}$ , then  $Ext_A^*(\mathbb{K}, \mathbb{K})$  is an  $A_\infty$ -algebra. This was shown by Lu, Palmari, Wu and Zhang [11].

Next we want to understand the category of  $A_\infty$ -algebras. A morphism between  $A_\infty$ -algebras is called an  $\infty$ -morphism. We define such an  $\infty$ -morphism  $f : A \rightsquigarrow B$  between  $A_\infty$ -algebras as associated dg-coalgebra homomorphism  $Bf : (\bar{T}^c(A[1]), m^A) \rightarrow (\bar{T}^c(B[1]), m^B)$ . Here  $Bf$  is purely formal, we will make sense of this soon.

**Proposition 1.3.9.** Let  $A, B$  be two  $A_\infty$ -algebras. A collection of morphisms  $f_n : A^{\otimes n} \rightarrow B$  of degree  $1 - n$  for any  $n \geq 1$  defines an  $\infty$ -morphism  $f : A \rightsquigarrow B$  if and only if  $f_1$  is a morphism of chain complexes and for any  $n \geq 2$  the following relations are satisfied:

$$(rel_n) \quad \partial(f_n) = \sum_{\substack{p+1+r=k \\ p+q+r=n}} (-1)^{pq+r} f_k \circ_{p+1} m_q^A - \sum_{\substack{k \geq 2 \\ i_1 + \dots + i_k = n}} (-1)^e m_k^B \circ (f_{i_1} \otimes f_{i_2} \otimes \dots \otimes f_{i_k})$$

where  $e$  is given as:  $e = (k-1)(1-i_1) + (k-2)(1-i_2) + \dots + 2(1-i_{k-2}) + (1-i_{k-1})$

*Proof.* This is immediate by the universal property of cofree coalgebras. To see how we get the exponent  $e$  we explain the first term. The factor  $(1-i_1)$  is precisely the degree of  $f_{i_1}$ , and  $k-1$  is exactly how many tensors which appear after  $f_{i_1}$ . We get the first term as we have to move  $k-1$   $\omega$ s over  $f_{i_1}$ . The same applies to every other tensor after  $f_{i_1}$ , but for  $f_{i_2}$  there are only  $k-2$  tensors which comes after, and so on.  $\square$

Since the composition of two dg-coalgebra homomorphism is again a dg-algebra homomorphism, we get that the composition of two  $\infty$ -morphisms is again an  $\infty$ -morphism. More explicitly if  $f : A \rightsquigarrow B$  and  $g : B \rightsquigarrow C$  are two  $\infty$ -morphisms, then their composition is defined as:

$$(fg)_n = \sum_r \sum_{i_1 + \dots + i_r = n} (-1)^e g_r(f_{i_1} \otimes \dots \otimes f_{i_r}).$$

**Definition 1.3.10.** An  $\infty$ -morphism  $f : A \rightsquigarrow B$  is called strict if  $f_n = 0$  for any  $n \geq 2$ .

**Definition 1.3.11.**  $\text{Alg}_\infty$  denotes the category of  $A_\infty$ -algebras, and the morphisms in this category are the  $\infty$ -morphisms.

Observe that we may extend the bar construction to  $B : Alg_\infty \rightarrow CoAlg_{\mathbb{K}, conil}^\bullet$  to a fully faithful functor. This may be done explicitly by using proposition 1.1.29. The subcategory of the essential image is the full subcategory of every dg-coalgebra that is isomorphic, as a graded coalgebra, to a cofree tensor coalgebra. Notice that the bar construction on the category of dg-algebras is a non-full injection into the category of  $A_\infty$ -algebras. Observe that this inclusion is the recontextualization of a dg-algebra into an  $A_\infty$ -algebra.

We call a quasi-isomorphism between  $A_\infty$ -algebras for an  $\infty$ -quasi-isomorphism. Given an  $\infty$ -morphism  $f : A \rightsquigarrow B$ , then we say that it is an  $\infty$ -quasi-isomorphism if  $f_1$  is a quasi-isomorphism. If we wanted to be more stringent with this definition we would want an  $\infty$ -quasi-isomorphism to be an  $\infty$ -morphism which is a quasi-isomorphism of dg-coalgebras. We will see that these definitions are equivalent later.

A homotopy between two  $A_\infty$ -algebras is a homotopy between the dg-coalgebras they define. We may however trace this definition back along the quasi-inverse of the bar construction to get a new definition in terms of many morphisms. Let  $f, g : A \rightsquigarrow B$  be two  $\infty$ -morphisms, we say that  $f - g$  is null-homotopic if there is a collection of morphisms  $h_n : A^{\otimes n} \rightarrow B$  of degree  $n$  such that the following relations are satisfied for any  $n \geq 1$ :

$$f_n - g_n = \sum (-1)^s m_{r+1+t}^B \circ (f_{i_1} \otimes \dots \otimes f_{i_r} \otimes h_k \otimes g_{j_1} \otimes \dots \otimes g_{j_t}) + \sum (-1)^{j+kl} h_i \circ_{j+1} m_k^A$$

Where  $s$  is some constant depending on  $t, r$  and  $k$ . More details may be found in [12].

As in the same case for algebras, there is also a notion of unital  $A_\infty$ -algebras and augmented  $A_\infty$ -algebras. For this discussion it is important to note that the field  $\mathbb{K}$  is also an  $A_\infty$ -algebra. This algebra will serve as the initial algebra, in the same way as it did for ordinary algebras.

**Definition 1.3.12.** A strictly unital  $A_\infty$ -algebra is an  $A_\infty$ -algebra  $A$  together with a unit morphism  $v_A : \mathbb{K} \rightarrow A$  of degree 0 such that the following are satisfied:

- $m_1 \circ v_A = 0$ .
- $m_2(id_A \otimes v_A) = id_A = m_2(v_A \otimes id_A)$ .
- $m_i \circ_k v_A = 0$  for any  $i \geq 3$  and  $1 \leq k < i$ .

A strictly unital  $\infty$ -morphism  $f : A \rightsquigarrow B$  between strictly unital  $A_\infty$ -algebras is a morphism which preserves the unit. This means that  $f_1 v_A = v_B$  and  $f_i \circ_k v_A = 0$  for any  $i \geq 2$  and  $1 \leq k < i$ . The collection of strictly unital  $A_\infty$ -algebras and strictly unital  $\infty$ -morphisms form a non-full subcategory of  $A_\infty$ -algebras. A strict  $\infty$ -morphism which is unital at the level of chain complexes is automatically strictly unital. Strict unital will then mean strict and strictly unital. Note that  $\mathbb{K}$  is strictly unital where the unit is  $id_{\mathbb{K}}$ .

**Definition 1.3.13.** An augmented  $A_\infty$ -algebra is a strictly unital  $A_\infty$ -algebra  $A$  together with a strict unital morphism  $\varepsilon_A : A \rightarrow \mathbb{K}$ . The  $\infty$ -morphism  $\varepsilon_A$  is called the augmentation of  $A$ .

The collection of augmented  $A_\infty$ -algebras and strictly unital morphism is the category of augmented  $A_\infty$ -algebras, denoted as  $Alg_{\infty,+}$ . As in the same way for algebras, there is an equivalence



of categories  $\text{Alg}_\infty \simeq \text{Alg}_{\infty,+}$ . The augmentation ideal, or the reduced  $A_\infty$ -algebra is the kernel of the augmentation  $\varepsilon_A$ . A priori, it does not make sense to talk about this limit as we do not know if it exists. However, we will see in section 2.3.3 that such morphisms does in fact have a kernel. This defines a functor,  $\bar{\_} : \text{Alg}_{\infty,+} \rightarrow \text{Alg}_\infty$ , where  $\text{Ker} \varepsilon_A = \bar{A}$ . The quasi-inverse to this functor is free augmentations. Given an  $A_\infty$ -algebra  $A$ , we may construct the  $A_\infty$ -algebra  $A \oplus \mathbb{K}$ . The structure morphisms are given by  $m_i^A$ , but there is now a unit  $v_{A \oplus \mathbb{K}}$ . Thus we get that  $m_1(1) = 0$ ,  $m_2(a \otimes 1) = a$  and  $m_i \circ_k 1 = 0$  in the same manner. We obtain a functor  $\bar{\_}^+ : \text{Alg}_\infty \rightarrow \text{Alg}_{\infty,+}$ , where  $A \oplus \mathbb{K} = \bar{A}^+$ .

### 1.3.2 $A_\infty$ -Coalgebras

Dual to  $A_\infty$ -algebras we got conilpotent  $A_\infty$ -coalgebras. Here we instead ask ourselves if the cobar construction has some converse. I.e. if  $C$  is a graded vector space such that  $T(C[-1])$  together with a derivation  $m$  is a dg-algebra, is then  $C$  a coalgebra? Again, the answer to this is no, but we do obtain a definition for conilpotent  $A_\infty$ -coalgebras.

**Definition 1.3.14.** A graded vector space  $C$  is called a conilpotent  $A_\infty$ -coalgebra if it is a dg-algebra of the form  $(\bar{T}(C[-1]), d)$  where  $d$  is a derivation of degree 11.

*Remark 1.3.15.* For the rest of this thesis, an  $A_\infty$ -coalgebra should be understood as a conilpotent  $A_\infty$ -coalgebra unless otherwise specified.

**Corollary 1.3.15.1.**  $C$  is an  $A_\infty$ -coalgebra with differential  $d$  then there is a chain complex  $(C, d^1)$ , where  $d^1$  is of degree 1, and together with morphisms  $d^n : C \rightarrow C^{\otimes n}$  such that  $d$  uniquely determines each  $d^i$  for any  $i > 0$ . Conversely, if the morphisms  $d^i$  satisfy  $(\text{rel})_n$ , then they uniquely determine a  $d$  such that  $C$  is an  $A_\infty$ -coalgebra.

$$(\text{rel}_n) \quad \sum_{p+q+r=n} (-1)^{pq+r} d^{p+1+q} \circ_{p+1}^{op} d^q = 0$$

A morphism of  $A_\infty$ -coalgebras would be defined in the same manner, but opposite. So an  $\infty$ -comorphism  $f : C \rightsquigarrow D$  is either a morphism  $\tilde{f} : (T(C[-1]), m^C) \rightarrow (T(D[-1]), m^D)$  of dg-algebras. Equivalently such an  $\infty$ -comorphism is a collection of morphisms  $f_n : C \rightarrow D^{\otimes n}$  of degree  $1 - n$  such that  $f_1$  is a morphism of chain complexes and for any  $n \geq 2$  the following relations are satisfied:

$$(\text{rel}_n) \quad \partial(f_n) = \sum_{\substack{p+1+r=k \\ p+q+r=n}} (-1)^{pq+r} f_k \circ_{p+1}^{op} m_q^D - \sum_{\substack{k \geq 2 \\ i_1 + \dots + i_k = n}} (-1)^e m_k^C \circ^{op} (f_{i_1} \otimes f_{i_2} \otimes \dots \otimes f_{i_k})$$

where  $e$  is given as:  $e = (k-1)(1-i_1) + (k-2)(1-i_2) + \dots + 2(1-i_{k-2}) + (1-i_{k-1})$

We denote  $\text{Coalg}_\infty$  as the category of  $A_\infty$ -coalgebras. In the same manner, the cobar construction extends to this category and gives us an identification of  $A_\infty$ -coalgebras and a subcategory of dg-algebras. This subcategory consists of every dg-algebra that is isomorphic, as an algebra, to a free tensor algebra. Lastly, every dg-coalgebra is an  $A_\infty$ -coalgebra by letting every morphism  $m^i = 0$  where  $i > 2$ . This gives a non full inclusion.



## Chapter 2

# Homotopy Theory of Algebras

Quillen envisioned a more general approach to homotopy theory, which he dubbed homotopical algebra. A homotopy theory was first enclosed by the structure of a model category, and now we mostly consider closed model category. Many of the results from classical homotopy theory was recovered in this new setting of model categories. The theorem which we are most concerned about is Whiteheads theorem:

**Theorem 2.0.1** (Whiteheads Theorem). *Let  $X$  and  $Y$  be two CW-complexes. If  $f : X \rightarrow Y$  is a weak equivalence, then it is also a homotopy equivalence. I.e. there exists a morphism  $g : Y \rightarrow X$  such that  $gf \sim id_X$  and  $fg \sim id_Y$ .*

If we employ Quillens model category onto the category  $\mathbf{Top}$ , we get that a space  $X$  is bifibrant if and only if it is a CW-complex. The natural generalization is then to not ask of  $X$  to be a CW-complex, but a bifibrant object.

**Theorem 2.0.2** (Generalized Whiteheads Theorem). *Let  $\mathcal{C}$  be a model category. Suppose that  $X$  and  $Y$  are bifibrant objects of  $\mathcal{C}$ , and that there is a weak-equivalence  $f : X \rightarrow Y$ . Then  $f$  is also a homotopy equivalence, i.e. there exists a morphism  $g : Y \rightarrow X$  such that  $gf \sim id_X$  and  $fg \sim id_Y$ .*

The category of differential graded algebras employs such a model category, and here we let the weak-equivalences be quasi-isomorphisms. On the other hand the category of differential graded coalgebras has a model structure where the weak equivalences are the maps which are sent to quasi-isomorphism by the cobar construction. Moreover, in this case the bar and cobar construction defines a Quillen equivalence between these model structures. As we will see, a dg-coalgebra will be bifibrant exactly when it is an  $A_\infty$ -algebra. Thus, by Whiteheads theorem, quasi-isomorphisms lift to homotopy equivalences. In this case the derived category of  $A_\infty$ -algebras is equivalent to the homotopy category of  $A_\infty$ -algebras.

We will conclude this section by looking at the category of algebras as a subcategory of  $A_\infty$ -algebras. The derived category may then be expressed as the homotopy category  $A_\infty$ -algebras,

restricted to algebras.

## 2.1 Model categories

In this section we will define Quillens model category. As one may see is that in practice there are a plethora of semantically different definitions of model categories, however they are all made to be equivalent. The difference comes down to preference. In this thesis we will use the definitions as they are developed in Mark Hoveys book [13]. We will then go on to prove the basic results known about model categories, its associated homotopy category and Quillen functors between model categories.

Before we state the definition of a model category we need some preliminary definitions. For this section, let  $\mathcal{C}$  be a category.

**Definition 2.1.1** (Retract). A morphism  $f : A \rightarrow B$  in  $\mathcal{C}$  is a retract of a morphism  $g : c \rightarrow D$  if it fits in a commutative diagram:

$$\begin{array}{ccccc}
 & & id_A & & \\
 & \curvearrowright & & \curvearrowleft & \\
 A & \longrightarrow & C & \longrightarrow & A \\
 \downarrow f & & \downarrow g & & \downarrow f \\
 B & \longrightarrow & D & \longrightarrow & B \\
 & \curvearrowleft & & \curvearrowright & \\
 & & id_B & & 
 \end{array}$$

**Definition 2.1.2** (Functorial factorization). A pair of functors  $\alpha, \beta : \mathcal{C}^{\rightarrow} \rightarrow \mathcal{C}^{\rightarrow}$  is called a functorial factorization if for any morphism  $f = \beta(f) \circ \alpha(f)$ . We will denote the morphisms in the factorization as  $f_{\alpha}$  and  $f_{\beta}$ . The functorial factorization may be depicted by the following commutative diagram:

$$\begin{array}{ccc}
 A & \xrightarrow{f} & B \\
 & \searrow f_{\alpha} & \nearrow f_{\beta} \\
 & C & 
 \end{array}$$

**Definition 2.1.3** (Lifting properties). Suppose that the morphisms  $i : A \rightarrow B$  and  $p : C \rightarrow D$  fits inside a commutative square.  $i$  is said to have the left lifting property with respect to  $p$ , or  $p$  has the right lifting property with respect to  $i$ , if there is an  $h : B \rightarrow C$  such that the two triangles commute.

$$\begin{array}{ccc}
 A & \longrightarrow & C \\
 \downarrow i & \nearrow h & \downarrow p \\
 B & \longrightarrow & D
 \end{array}$$

*Remark 2.1.4.* We will call the left lifting property for LLP and the right lifting property for RLP.

### 2.1.1 Model categories

**Definition 2.1.5** (Model category). Let  $\mathcal{C}$  be a category with all finite limits and colimits.  $\mathcal{C}$  admits a model structure if there are three wide subcategories each defining a class of morphisms:

- $Ac \subset Mor(\mathcal{C})$  are called weak equivalences
- $Cof \subset Mor(\mathcal{C})$  are called cofibrations
- $Fib \subset Mor(\mathcal{C})$  are called fibrations

In addition we call morphisms in  $Cof \cap Ac$  for acyclic cofibrations and  $Fib \cap Ac$  for acyclic fibrations. Moreover,  $\mathcal{C}$  has two functorial factorizations  $(\alpha, \beta)$  and  $(\gamma, \delta)$ . The following axioms should be satisfied:

- MC1** The class of weak equivalences satisfy the 2-out-of-3 property, i.e. if  $f$  and  $g$  are composable morphisms such that 2 out of  $f$ ,  $g$  and  $gf$  are weak equivalences, then so is the third.
- MC2** The three classes  $Ac$ ,  $Cof$  and  $Fib$  are retraction closed, i.e. if  $f$  is a retraction of  $g$ , and  $g$  is either a weak-equivalence, cofibration or fibration, then so is  $f$ .
- MC3** The class of cofibrations have the left lifting property with respect to acyclic fibrations, and fibrations have the right lifting property with respect to acyclic cofibrations.
- MC4** Given any morphism  $f$ ,  $f_\alpha$  is a cofibration,  $f_\beta$  is an acyclic fibration,  $f_\gamma$  is an acyclic cofibration and  $f_\delta$  is a fibration.

*Remark 2.1.6.* The class  $Ac$  has every isomorphism. This is because every isomorphism is a retract of some identity morphism.

*Remark 2.1.7.* The type of category which has been introduced above was first called a closed model category by Quillen [14]. In his sense, a model category does not require to have either finite limits or finite colimits. In our case, we will explicitly state whenever a model category is non-closed, i.e. does not have every limit or colimit.

A model category  $\mathcal{C}$  is now defined to be a category equipped with a particular model structure. Notice that a category may admit several model structures. We will postpone examples until sufficient theory have been developed. For more topological examples, we refer to Dwyer-Spalinski [15] and Hovey [13].

An interesting and a not so non-trivial property of model categories is that giving all three classes  $Ac$ ,  $Cof$  and  $Fib$  is redundant. Given the class of weak equivalences and either cofibrations or fibrations, the model structure is determined. Thus the classes of fibrations are determined by acyclic cofibrations and cofibrations are determined by fibrations. The next two results will show this.

**Lemma 2.1.8** (The retract argument). *Let  $\mathcal{C}$  be a category. Suppose there is a factorization  $f = pi$  and that  $f$  has LLP with respect to  $p$ , then  $f$  is a retract of  $i$ . Dually, if  $f$  has RLP to  $i$ , then it is a retract of  $p$ .*

*Proof.* We assume that  $f : A \rightarrow C$  has LLP with respect to  $p : B \rightarrow C$ . Then we may find a lift  $r : C \rightarrow B$ , which realize  $f$  as a retract of  $i$ .

$$\begin{array}{ccc} A & \xrightarrow{i} & B \\ \downarrow f & \nearrow r & \downarrow p \\ C & \xlongequal{\quad} & C \end{array} \implies \begin{array}{ccccc} A & \xlongequal{\quad} & A & \xlongequal{\quad} & A \\ \downarrow f & & \downarrow i & & \downarrow f \\ C & \xrightarrow{r} & B & \xrightarrow{p} & C \end{array}$$

□

**Proposition 2.1.9.** *Let  $\mathcal{C}$  be a model category. A morphism  $f$  is a cofibration (acyclic cofibration) if and only if  $f$  has LLP with respect to acyclic fibrations (fibrations). Dually,  $f$  is a fibration (acyclic fibration) if and only if it has RLP with respect to acyclic cofibrations (cofibrations).*

*Proof.* Assume that  $f$  is a cofibration. By MC3, we know that  $f$  has LLP with respect to acyclic fibrations. Assume instead that  $f$  has LLP with respect to every acyclic fibration. By MC4 we factor  $f = f_\alpha \circ f_\beta$ , where  $f_\alpha$  is a cofibration and  $f_\beta$  is an acyclic fibration. Since we assume  $f$  to have LLP with respect to  $f_\beta$ , by lemma 2.1.8 we know that  $f$  is a retract of  $f_\alpha$ . Thus by MC2, we know that  $f$  is a cofibration. □

**Corollary 2.1.9.1.** *Let  $\mathcal{C}$  be a model category. (Acyclic) Cofibrations are stable under pushouts, i.e. if  $f$  is an (acyclic) cofibration, then  $f'$  is an (acyclic) cofibration.*

$$\begin{array}{ccc} A & \longrightarrow & C \\ \downarrow f & \lrcorner & \downarrow f' \\ B & \longrightarrow & D \end{array}$$

*Dually, fibrations are stable under pullbacks.*

*Proof.* This is clear by the universal property of pushouts. □

Since we assume that every model category  $\mathcal{C}$  admits finite limits and colimits, we know that it has both an initial and a terminal object. We let  $\emptyset$  denote the initial object and  $*$  denote the terminal object.

**Definition 2.1.10** (Cofibrant, fibrant and bifibrant objects). *Let  $\mathcal{C}$  be a model category. An object  $X$  is called cofibrant if the unique morphism  $\emptyset \rightarrow X$  is a cofibration. Dually,  $X$  is called fibrant if the unique morphism  $X \rightarrow *$  is fibrant. If  $X$  is both cofibrant and fibrant, we call it bifibrant.*

There is no reason for every object to be either cofibrant or fibrant. However, we may see that every object is weakly equivalent to an object which is either fibrant or cofibrant. We will make it precise what it means for two objects to be weakly equivalent later.

**Construction 2.1.11.** Let  $X$  be an object of a model category  $\mathcal{C}$ . The morphism  $i : \emptyset \rightarrow X$  has a functorial factorization  $i = i_\beta \circ i_\alpha$ , where  $i_\alpha : \emptyset \rightarrow QX$  is a cofibration and  $i_\beta : QX \rightarrow X$  is an acyclic fibration. By definition  $QX$  is cofibrant and weakly equivalent to  $X$ .

$Q : \mathcal{C} \rightarrow \mathcal{C}$  defines a functor called the cofibrant replacement. To see this we first look at the slice category  $\emptyset/\mathcal{C}$ . The objects are morphisms  $f : \emptyset \rightarrow X$  for any object  $X$  in  $\mathcal{C}$ , while morphisms are commutative triangles. We first observe that  $\emptyset/\mathcal{C} \subset \mathcal{C}^\rightarrow$  is a subcategory of the arrow category. Thus  $(\alpha, \beta)$  may be interpreted as functors on the slice category to the arrow category. Moreover, since every arrow  $f : \emptyset \rightarrow X$  is unique, we observe that this category is equivalent to  $\mathcal{C}$ . Thus  $(\alpha, \beta)$  may be interpreted as functors on  $\mathcal{C}$  into arrows. We define  $Q$  as the composition  $Q = \text{tar} \circ \alpha$ .

Dually, we get a fibrant replacement  $R$  by dualizing the above argument.


We collect the following properties

**Lemma 2.1.12.** *The cofibrant replacement  $Q$  and fibrant replacement  $R$  preserves weak equivalences.*

*Proof.* Clear from the 2-out-of-3 property. □

**Lemma 2.1.13** (Ken Brown's lemma). *Let  $\mathcal{C}$  be a model category and  $\mathcal{D}$  be a category with weak equivalences satisfying the 2-out-of-3 property. If  $F : \mathcal{C} \rightarrow \mathcal{D}$  is a functor sending acyclic cofibrations between cofibrant objects to weak equivalences, then  $F$  takes all weak equivalences between cofibrant objects to weak equivalences. Dually, if  $F$  takes all acyclic fibrations between fibrant objects to weak equivalences, then  $F$  takes all weak equivalences between fibrant objects to weak equivalences.*

*Proof.* Suppose that  $A$  and  $B$  are cofibrant objects and that  $f : A \rightarrow B$  is a weak equivalence. Using the universal property of the coproduct we define the map  $(f, id_B) = p : A \coprod B \rightarrow B$ .  $p$  has a functorial factorization into a cofibration and acyclic fibration,  $p = p_\beta \circ p_\alpha$ . We recollect the maps in the following pushout diagram:




By lemma 2.1.9.1 both  $i_1$  and  $i_2$  are cofibrations. Since  $f$ ,  $id_B$  and  $p_\beta$  are weak equivalences, so are  $p_\alpha \circ i_1$  and  $p_\alpha \circ i_2$  by MC2. Moreover, they are acyclic cofibrations.

Assume that  $F : \mathcal{C} \rightarrow \mathcal{D}$  is a functor as described above. Then by assumption,  $F(p_\alpha \circ i_1)$  and  $F(p_\alpha \circ i_2)$  are weak equivalences. Since a functor sends identity to identity, we also know that  $F(id_B)$  is a weak equivalence. Thus by the 2-out-of-3 property  $F(p_\beta)$  is a weak equivalence, as  $F(p_\beta) \circ F(p_\alpha \circ i_2) = id_{F(B)}$ . Again, by 2-out-of-3 property  $F(f)$  is a weak equivalence, as  $F(f) = F(p_\beta) \circ F(p_\alpha \circ i_1)$ .  $\square$

## 2.1.2 Homotopy category

Homotopy theory at its most abstract is the study of categories and functors up to weak equivalences. Here, a weak equivalence may be anything, but most commonly it is a weak equivalence in topological homotopy or a quasi-isomorphism in homological algebra. The biggest concern when dealing with such concepts is to make a functor well-defined up to these chosen weak equivalences. To this end, there is a construction to amend these problems, known as derived functors. We define a homotopical category in the sense of Riehl [16].

**Definition 2.1.14** (Homotopical Category). Let  $\mathcal{C}$  be a category.  $\mathcal{C}$  is Homotopical if there is a wide subcategory constituting a class of morphisms known as weak equivalences,  $Ac \subset Mor\mathcal{C}$ . The weak equivalences should satisfy the **2-out-of-6 property**, i.e. given three composable morphisms  $f, g$  and  $g, h$  if  $gf$  and  $hg$  are weak equivalences, then so are  $f, g, h$  and  $hgf$ .



*Remark 2.1.15.* Notice that the 2-out-of-6 property is stronger than the 2-out-of-3 property. To see this, let either  $f, g$  or  $h$  be the identity, and then conclude with the 2-out-of-3 property.



**Remark 2.1.16.** The collection of weak equivalences contains every isomorphism. To see this pick an isomorphism  $f$  and  $f^{-1}$ , then the compositions are the identity on the domain and codomain, which are assumed to be in  $Ac$ .

Given such a homotopical category  $\mathcal{C}$ , we want to invert every weak equivalence and create the homotopy category of  $\mathcal{C}$ . This construction is developed in Gabriel and Zisman [17] called calculus of fractions. This method essentially tries to mimic localization for commutative rings in a category theoretic fashion.

**Definition 2.1.17.** Let  $\mathcal{C}$  be a homotopical category. It's homotopy category  $Ho\mathcal{C} = \mathcal{C}[Ac^{-1}]$ , together with a localization functor  $q : \mathcal{C} \rightarrow Ho\mathcal{C}$ . Recall that the localization are determined by the following universal property: If  $F : \mathcal{C} \rightarrow \mathcal{D}$  is a functor sending weak equivalences to isomorphisms, then it uniquely factors through the homotopy category up to a unique natural isomorphism  $\eta$ .

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{D} \\ & \searrow q \quad \Downarrow \eta \quad \nearrow F' & \\ & Ho\mathcal{C} & \end{array}$$

**Definition 2.1.18.** Suppose that  $\mathcal{C}$  is a homotopical category. Two objects of  $\mathcal{C}$  are said to be weakly equivalent if they are isomorphic in  $Ho\mathcal{C}$ . I.e. there is some zig-zag relation between the objects, consisting only of weak equivalences.

**Remark 2.1.19.** A renown problem with localizations is that even if  $\mathcal{C}$  is a locally small category, any localization  $\mathcal{C}[S^{-1}]$  does not need to be. Thus, without a good theory of classes or higher universes, we cannot in general ensure that the localization still exists as a locally small category.

We see from the definition of the homotopy category that a functor  $F$  admits a lift  $F'$  to the homotopy category whenever weak equivalences are sent to isomorphisms. Moreover, if we have a functor  $F$  between homotopical categories which preserves weak equivalences, it then induces a functor between the homotopy categories.

**Definition 2.1.20** (Homotopical functors). A functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  between homotopical categories is homotopical if it preserves weak equivalences. Moreover, there is a lift of functors as in the following diagram.

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{D} \\ \downarrow q_{\mathcal{C}} & \nearrow \eta & \downarrow q_{\mathcal{D}} \\ Ho\mathcal{C} & \xrightarrow{F'} & Ho\mathcal{D} \end{array}$$

Derived functors come into play whenever this is not the case. These lifts are however the closest approximation which we can make functorial. The general exposition of derived functors is

beyond the scope of this thesis, but an account of it may be found in [16]. As we will see, model categories are a nice environment to work with these concepts. Firstly we will amend the problem with localizations, where the homotopy category may not exist. Secondly, we will obtain a simple description of some important derived functors.

**Proposition 2.1.21.** *Any model category  $\mathcal{C}$  is a homotopical category.*

*Proof.* To show that a model category is homotopical it suffices to show that  $A_{\mathcal{C}}$  satisfies the 2-out-of-6 property. Assume there are 3 composable morphisms  $f, g, h$  such that  $gf, hg \in A_{\mathcal{C}}$ . By the 2-out-of-3 property for  $A_{\mathcal{C}}$  it is enough to show that at least one of  $f, g, h, fgh$  is a weak equivalence to deduce that every other morphism is a weak equivalence.

$$\begin{array}{ccccc}
 A & \xrightarrow{f} & B & & \\
 & \searrow gf & \downarrow g & \searrow hg & \\
 & & C & \xrightarrow{h} & D
 \end{array}$$

To be able to use the model structure, we will first show that we may assume  $f, g$  to be cofibrant and  $g, h$  to be fibrant. We know by MC4 that  $f, g, gf$  may be factored into a cofibration composed with an acyclic fibration, e.g.  $f = f_{\beta}f_{\alpha}$ . Since  $gf$  is a weak equivalence, so is  $(gf)_{\alpha}$  by the 2-out-of-3 property.

$$\begin{array}{ccc}
 \begin{array}{ccc} A & \xrightarrow{f} & B \\ & \searrow f_{\alpha} & \nearrow f_{\beta} \\ & B' & \end{array} & 
 \begin{array}{ccc} B & \xrightarrow{g} & C \\ & \searrow g_{\alpha} & \nearrow g_{\beta} \\ & C' & \end{array} & 
 \begin{array}{ccc} A & \xrightarrow{gf} & C \\ & \searrow (gf)_{\alpha} & \nearrow (gf)_{\beta} \\ & C'' & \end{array}
 \end{array}$$

Notice that the "cofibrant approximation" of the map from  $A$  to  $C$  either goes through  $C'$  or  $C''$ . We conjoin these by taking the pullback. Since acyclic fibrations are stable over pullbacks, we get a pullback square where every morphism is an acyclic fibration. Thus the map  $A \rightarrow \tilde{C}$  is a weak equivalence by 2-out-of-3.

$$\begin{array}{ccccc}
 A & & & & \\
 & \searrow (gf)_{\alpha} & & \searrow g_{\alpha} & \\
 & & \tilde{C} & \xrightarrow{r} & C' \\
 & & \downarrow & & \downarrow g_{\beta} \\
 & & C'' & \xrightarrow{(gf)_{\beta}} & C
 \end{array}$$

To replace  $f$  with  $f_\alpha$  we must lift the composition into our "new"  $C$ , which is  $\tilde{C}$ . This is done by using MC3, as  $f_\alpha$  is a cofibration and the pullback square above consists entirely of acyclic fibrations.

$$\begin{array}{ccc} A & \longrightarrow & \tilde{C} \\ \downarrow f_\alpha & \nearrow \text{dashed} & \downarrow \\ B' & \longrightarrow & C \end{array}$$

To summarize we have the following diagram, where every squiggly arrow is a weak equivalence.

$$\begin{array}{ccccccc} A & \xrightarrow{f_\alpha} & B' & \rightsquigarrow & B & & \\ & \searrow t & \downarrow s & & \downarrow & \searrow & \\ & & \tilde{C} & \rightsquigarrow & C & \longrightarrow & D \end{array}$$

We now wish to promote the arrow  $s : B' \rightarrow \tilde{C}$  into a cofibration. We do this by factoring both  $s$  and  $t$  with MC4. Notice that  $s_\beta, t_\beta$  and  $t_\alpha$  are weak equivalences.

$$\begin{array}{ccc} B' & \xrightarrow{s} & \tilde{C} \\ & \searrow s_\alpha & \nearrow s_\beta \\ & \tilde{C}' & \end{array} \quad \begin{array}{ccc} A & \xrightarrow{t} & \tilde{C} \\ & \searrow t_\alpha & \nearrow t_\beta \\ & \tilde{C}'' & \end{array}$$

To obtain our final factorization we use RLP of  $s_\beta$  on  $t_\alpha$ .

$$\begin{array}{ccc} & & B' \\ & \nearrow f_\alpha & \downarrow s_\alpha \\ A & \longrightarrow & \tilde{C}' \\ \downarrow t_\alpha & \nearrow \text{dashed} & \downarrow s_\beta \\ \tilde{C}'' & \xrightarrow{t_\beta} & \tilde{C} \end{array}$$

We have now obtained a factorization of  $gf$  which are two cofibrations followed by an acyclic fibration, in such a manner that it is compatible with the original composition. The dual to this claim is that we may also factor  $hg$  into two fibrations which is preceded by an acyclic cofibration. In other words, we may assume without loss of generality that  $f$  and  $g$  are cofibrations, and that  $g$  and  $h$  are fibrations.

In this case we will show that  $g$  is an isomorphism. Consider the diagram below with lifts  $i$  and  $j$ , these exist since we assume  $gf$  and  $hg$  to be weak equivalences.

$$\begin{array}{ccccc}
 A & \xrightarrow{f} & B & \xrightarrow{id_B} & B \\
 \downarrow gf & \nearrow i & \downarrow g & \nearrow j & \downarrow hg \\
 C & \xrightarrow{id_C} & C & \xrightarrow{h} & D
 \end{array}$$

Since the diagram is commutative, we get that  $i = j$ , and that  $g$  is both split mono and split epi, with  $i$  as its splitting.  $\square$

Since every model category is homotopical, it also has an associated homotopy category  $Ho\mathcal{C}$ . Let  $\mathcal{C}_c$ ,  $\mathcal{C}_f$  and  $\mathcal{C}_{cf}$  denote the full subcategories consisting of cofibrant, fibrant and bifibrant objects respectively.

**Proposition 2.1.22.** *Let  $\mathcal{C}$  be a model category. The following categories are equivalent:*

- $Ho\mathcal{C}$
- $Ho\mathcal{C}_c$
- $Ho\mathcal{C}_f$
- $Ho\mathcal{C}_{cf}$

*Proof.* We show that  $Ho\mathcal{C} \simeq Ho\mathcal{C}_c$ . The inclusion  $i : \mathcal{C}_c \rightarrow \mathcal{C}$  clearly preserves weak equivalences, thus  $i$  is homotopical and admits a lift. Moreover, since the cofibrant replacement is also homotopical, it also has a lift.

$$\begin{array}{ccc}
 \mathcal{C}_c & \xrightarrow{i} & \mathcal{C} \\
 \downarrow & \swarrow Ho\,Q & \downarrow \\
 Ho\mathcal{C}_c & \xleftarrow{Ho\,i} & Ho\mathcal{C}
 \end{array}$$

It is clear that  $Ho\,Q$  is the quasi-inverse of  $Ho\,i$ .  $\square$

As of now we still don't see how model categories will fix the size issues. To do this we will develop the notion of homotopy equivalence,  $\sim$ . On the subcategory of bifibrant objects  $\mathcal{C}_{cf}$ , this homotopy equivalence will be a congruence relation. This, together with the fact that there is an equivalence of categories  $Ho\mathcal{C}_{cf} \simeq \mathcal{C}_{cf}/\sim$ , is enough to solve the size issues.

**Definition 2.1.23** (Cylinder and path objects). Let  $\mathcal{C}$  be a model category. Given an object  $X$ , a cylinder object  $X \wedge I$  is a factorization of the fold map  $i : X \coprod X \rightarrow X$ , such that  $p_0$  is a cofibration and that  $p_1$  is a weak equivalence.

$$\begin{array}{ccc}
 X \amalg X & \xrightarrow{i} & X \\
 \searrow p_0 & & \nearrow p_1 \\
 & X \wedge I &
 \end{array}$$

Dually, a path object  $X^I$  is a factorization of the diagonal map  $i : X \rightarrow X \amalg X$ , such that  $p_0$  is a weak equivalence and that  $p_1$  is a fibration.

$$\begin{array}{ccc}
 X & \xrightarrow{i} & X \amalg X \\
 \searrow p_0 & & \nearrow p_1 \\
 & X^I &
 \end{array}$$

*Remark 2.1.24.* Even though we have written  $X \wedge I$  suggestively to be a functor, it is not. There may be many choices for a cylinder object. However, by using the functorial factorization from MC4, we get a canonical choice of a cylinder object, as it factors every map into a cofibration and an acyclic fibration. By letting the cylinder object be this object, we do indeed obtain a functor.

**Proposition 2.1.25.** *Let  $\mathcal{C}$  be a model category and  $X$  an object of  $\mathcal{C}$ . Given two cylinder objects  $X \wedge I$  and  $X \wedge I'$ , then they are weakly equivalent.*

*Proof.* It is enough to show that there is a weak equivalence from any cylinder object into one specified cylinder object. This is in fact true for the functorial cylinder object  $X \wedge I$ , as the final morphism  $p_1$  is an acyclic fibration, which enables a lift which is a weak equivalence by the 2-out-of-3 property.

$$\begin{array}{ccc}
 X \amalg X & \xrightarrow{p_0} & X \wedge I \\
 \downarrow p'_0 & \nearrow \text{dashed} & \downarrow p_1 \\
 X \wedge I' & \xrightarrow{p'_1} & X
 \end{array}$$

□

**Definition 2.1.26** (Homotopy equivalence). Let  $f, g : X \rightarrow Y$ . A left homotopy between  $f$  and  $g$  is a morphism  $H : X \wedge I \rightarrow Y$  such that  $H i_0 = f$  and  $H i_1 = g$ .  $f$  and  $g$  are left homotopic if a left homotopy exists, and it is denoted  $f \stackrel{l}{\sim} g$ .

$$\begin{array}{ccccc}
 X & & & & f \\
 \downarrow & \searrow^{i_0} & & & \nearrow \\
 X \amalg X & \xrightarrow{p_0} & X \wedge I & \xrightarrow{H} & Y \\
 \uparrow & \nwarrow_{i_1} & & & \nwarrow \\
 X & & & & g
 \end{array}$$

A right homotopy between  $f$  and  $g$  is a morphism  $H : X \rightarrow Y^I$  such that  $i_0 H = f$  and  $i_1 H = g$ .  $f$  and  $g$  are right homotopic if a right homotopy exists, and it is denoted  $f \stackrel{r}{\sim} g$ .

$$\begin{array}{ccccc}
 & & f & & Y \\
 & \searrow & & \nearrow & \uparrow \\
 X & \xrightarrow{H} & Y^I & \xrightarrow{p_1} & Y \amalg Y \\
 & \swarrow & \nwarrow_{i_1} & & \downarrow \\
 & & g & & Y
 \end{array}$$

$f$  and  $g$  are said to be homotopic if they are both left and right homotopic, it is denoted  $f \sim g$ .  $f$  is said to be a homotopy equivalence, if it has a homotopy inverse  $h : Y \rightarrow X$ , such that  $hf \sim id_X$  and  $fh \sim id_Y$ .

It is important to know that this is not a priori an equivalence relation. This is amended by taking both fibrant and cofibrant replacements. We see this in the following proposition.

**Proposition 2.1.27.** *Let  $\mathcal{C}$  be a model category, and  $f, g : X \rightarrow Y$  be morphisms. We have the following:*

1. If  $f \stackrel{l}{\sim} g$  and  $h : Y \rightarrow Z$ , then  $hf \stackrel{l}{\sim} hg$ .
2. If  $Y$  is fibrant,  $f \stackrel{l}{\sim} g$  and  $h : W \rightarrow X$ , then  $fh \stackrel{l}{\sim} gh$ .
3. If  $X$  is cofibrant, then left homotopy is an equivalence relation on  $\mathcal{C}(X, Y)$ .
4. If  $X$  is cofibrant and  $f \stackrel{l}{\sim} g$ , then  $f \stackrel{r}{\sim} g$ .

*Proof.* (1.) Assume that  $f \stackrel{l}{\sim} g$  and  $h : Y \rightarrow Z$ . Let  $H : X \wedge I \rightarrow Y$  denote the left homotopy between  $f$  and  $g$ . The left homotopy between  $hf$  and  $hg$  is given as  $hH$ .

(2.) Assume that  $Y$  is fibrant,  $f \stackrel{l}{\sim} g$  and that  $h : W \rightarrow X$ . Let  $H : X \wedge I \rightarrow Y$  be a left homotopy. We construct a new cylinder object for the homotopy. Factor  $p_1 : X \wedge I \rightarrow X$  as  $q_1 \circ q_0$  where  $q_0 : X \wedge I \rightarrow X \wedge I'$  is an acyclic cofibration and  $q_1 : X \wedge I' \rightarrow X$  is a fibration. By the 2-out-of-3

property  $q_1$  is an acyclic fibration, as  $p_1$  and  $q_0$  are weak equivalences.  $X \wedge I'$  is a cylinder object as  $q_0 \circ p_0$  is a cofibration and  $q_1$  is a weak equivalence. Since we assume  $Y$  to be fibrant we lift the left homotopy  $H : X \wedge I \rightarrow Y$  to the left homotopy  $H' : X \wedge I' \rightarrow Y$  with the following diagram:

$$\begin{array}{ccc} X \wedge I & \xrightarrow{H} & Y \\ \downarrow q_0 & \nearrow H' & \downarrow \\ X \wedge I' & \longrightarrow & * \end{array}$$

We can find the appropriate homotopy needed with lift given by the following diagram:

$$\begin{array}{ccc} W \coprod W & \xrightarrow{q_0 p_0 (h \coprod h)} & X \wedge I' \\ \downarrow p'_0 & \nearrow k & \downarrow q_1 \\ W \wedge I & \xrightarrow{h p'_1} & X \end{array}$$

Then the morphism  $H'k$  is the desired left homotopy witnessing  $fh \stackrel{l}{\sim} gh$ .

(3.) Assume that  $X$  is cofibrant. First observe that a left homotopy is reflexive and symmetric. We must show that in this case it is also transitive. Thus, assume that  $f, g, h : X \rightarrow Y$  and that  $H : X \wedge I \rightarrow Y$  is a left homotopy witnessing  $f \stackrel{l}{\sim} g$  and that  $H' : X \wedge I' \rightarrow Y$  is a left homotopy witnessing  $g \stackrel{l}{\sim} h$ . We first observe that  $i_0 : X \rightarrow X \wedge I$  is a weak equivalence, as  $id_X = p_1 i_0$  where  $id_X$  and  $p_1$  are weak equivalences. Since  $X$  is assumed to be cofibrant, we see that  $X \coprod X$  is cofibrant by the following pushout:

$$\begin{array}{ccc} * & \longrightarrow & X \\ \downarrow & & \downarrow inr \\ X & \xrightarrow{inl} & X \coprod X \end{array}$$

Moreover, both  $inl$  and  $inr$  are cofibrations. This shows that  $i_0$  is a cofibration as  $i_0 = p_0 \circ inr$  is a composition of two cofibrations.  $i_0$  is thus an acyclic cofibration. We define an almost cylinder object  $C$  by the pushout of  $i_1$  and  $i'_0$ . We define the maps  $t$  and  $H''$  by using the universal property in the following manner:

$$\begin{array}{ccc}
X & \xrightarrow{i_1} & X \wedge I \\
\downarrow i'_0 & & \downarrow \\
X \wedge I' & \longrightarrow & C \\
& \searrow p'_1 & \nearrow p_1 \\
& & X
\end{array}
\quad
\begin{array}{ccc}
X & \xrightarrow{i_1} & X \wedge I \\
\downarrow i'_0 & & \downarrow \\
X \wedge I' & \longrightarrow & C \\
& \searrow H' & \nearrow H \\
& & X
\end{array}$$

Observe that there is a factorization of the fold map  $X \coprod X \xrightarrow{s} C \xrightarrow{t} X$ . However,  $s$  may not be a cofibration, so we replace  $C$  with the cylinder object  $X \wedge I''$  such that we have the factorization  $X \coprod X \xrightarrow{s_\alpha} X \wedge I'' \xrightarrow{ts_\beta} X$ . The morphism  $Hs_\beta$  is then our required homotopy for  $f \stackrel{l}{\sim} g$ .

(4.) Suppose that  $X$  is cofibrant and that  $H : X \wedge I \rightarrow Y$  is a left homotopy for  $f \stackrel{l}{\sim} g$ . Pick a path object for  $Y$ , such that we have the factorization  $Y \xrightarrow{q_0} Y^I \xrightarrow{q_1} Y \coprod Y$  where  $q_0$  is a weak equivalence and  $q_1$  is a fibration. Again, as  $X$  is cofibrant we get that  $i_0$  is an acyclic cofibration, so we have the following lift of the homotopy:

$$\begin{array}{ccc}
X & \xrightarrow{q_0 f} & Y^I \\
\downarrow i_0 & \nearrow J & \downarrow q_1 \\
X \wedge I & \xrightarrow{(fp_1, H)} & Y \coprod Y
\end{array}$$

The right homotopy is given by injecting away from  $f$ , i.e.  $H' = Ji_1$ . □

**Corollary 2.1.27.1.** *We collect the dual results of the above proposition, and thus have the following.*

1. If  $f \stackrel{r}{\sim}$  and  $h : W \rightarrow X$ , then  $fh \stackrel{r}{\sim} gh$ .
2. If  $X$  is cofibrant,  $f \stackrel{r}{\sim} g$  and  $h : Y \rightarrow Z$ , then  $hf \stackrel{r}{\sim} hg$ .
3. If  $Y$  is fibrant, then left homotopy is an equivalence relation on  $\mathcal{C}(X, Y)$ .
4. If  $Y$  is fibrant and  $f \stackrel{r}{\sim} g$ , then  $f \stackrel{l}{\sim} g$ .

**Corollary 2.1.27.2.** *Homotopy is a congruence relation on  $\mathcal{C}_{cf}$ . In this manner, the category  $\mathcal{C}_{cf}/\sim$  is well-defined, exists and inverts every homotopy equivalence.*

**Lemma 2.1.28** (Weird Whitehead). *Let  $\mathcal{C}$  be a model category. Suppose that  $C$  is cofibrant and  $h : X \rightarrow Y$  is an acyclic fibration or a weak equivalence between fibrant objects, then  $h$  induces an isomorphism:*

$$\mathcal{C}(C, X)/\sim \xrightarrow{\cong} \mathcal{C}(C, Y)/\sim$$



Dually, if  $X$  is fibrant and  $h : C \rightarrow D$  is an acyclic cofibration or a weak equivalence between cofibrant objects, then  $h$  induces an isomorphism:

$$\mathcal{C}(D, X) / \sim \xrightarrow{h^*} \mathcal{C}(C, X) / \sim$$

*Proof.* We assume  $C$  to be cofibrant and  $h : X \rightarrow Y$  to be an acyclic fibration. We first prove that  $h$  is surjective. Let  $f : C \rightarrow Y$ . By RLP of  $h$  there is a morphism  $f' : C \rightarrow X$  such that  $f = hf'$ .

$$\begin{array}{ccc} \emptyset & \longrightarrow & X \\ \downarrow & \nearrow f' & \downarrow h \\ C & \xrightarrow{f} & Y \end{array}$$

To show injectivity we assume  $f, g : C \rightarrow X$  such that  $hf \stackrel{L}{\sim} hg$ , in particular there is a left homotopy  $H : C \wedge I \rightarrow Y$ . Remember that since  $C$  is cofibrant, the map  $p_0$  is a cofibration. We find a left homotopy  $H : C \wedge I \rightarrow X$  witnessing  $f \stackrel{L}{\sim} g$  by the following lift.

$$\begin{array}{ccc} C \amalg C & \xrightarrow{f+g} & X \\ \downarrow p_0 & \nearrow H' & \downarrow h \\ C \wedge I & \xrightarrow{H} & Y \end{array}$$

Moreover, if we assume both  $X$  and  $Y$  to be fibrant, the functor  $\mathcal{C}(C, -) / \sim$  sends acyclic fibrations to isomorphisms, i.e. to weak equivalences. By Ken Brown's lemma, lemma 2.1.13, the aforementioned functor sends weak equivalences between fibrant objects to isomorphisms.  $\square$

**Theorem 2.1.29** (Generalized Whiteheads theorem). *Let  $\mathcal{C}$  be a model category. Suppose that  $f : X \rightarrow Y$  is a morphism of bifibrant objects, then  $f$  is a weak equivalence if and only if  $f$  is a homotopy equivalence.*

*Proof.* Suppose first that  $f$  is a weak equivalence. Pick a bifibrant object  $A$ , then by lemma 2.1.28  $f_* : \mathcal{C}(A, X) / \sim \rightarrow \mathcal{C}(A, Y) / \sim$  is an isomorphism. Letting  $A = Y$  we know that there is a morphism  $g : Y \rightarrow X$ , such that  $f_*g = fg \sim id_Y$ . Furthermore, by proposition 2.1.27, since  $X$  is bifibrant composing on the right preserves homotopy equivalence, e.g.  $fgf \sim f$ . By letting  $A = X$ , we get that  $f_*gf = fgf \sim f = f_*id_X$ , thus  $gf \sim id_X$ .

For the opposite direction, assume that  $f$  is a homotopy equivalence. We factor  $f$  into an acyclic cofibration  $f_\gamma$  and a fibration  $f_\delta$ , i.e.  $X \xrightarrow{f_\gamma} Z \xrightarrow{f_\delta} Y$ . Observe that  $Z$  is bifibrant as  $X$  and  $Y$  is, in particular,  $f_\gamma$  is a weak equivalence of bifibrant objects, so it is a homotopy equivalence.

It is enough to show that  $f_\delta$  is a weak equivalence. Let  $g$  be the homotopy inverse of  $f$ , and  $H : Y \wedge I \rightarrow Y$  is a left homotopy witnessing  $fg \sim id_Y$ . Since  $Y$  is bifibrant, the following square has a lift.

$$\begin{array}{ccc} Y & \xrightarrow{f_\gamma g} & Z \\ \downarrow i_0 & \nearrow H' & \downarrow f_\delta \\ Y \wedge I & \xrightarrow{H} & Y \end{array}$$

Let  $h = H'i_1$ , then by definition we know that  $f_\delta H'i_1 = id_Y$ . Moreover,  $H$  is a left homotopy witnessing  $f_\gamma g \sim h$ . Let  $g' : Z \rightarrow X$  be the homotopy inverse of  $f_\gamma$ . We have the following relations  $f_\delta \sim f_\delta f_\gamma g' \sim fg'$ , and  $hf_\delta \sim (f_\gamma g)(fg') \sim f_\gamma g' \sim id_Z$ . Let  $H'' : Z \wedge I \rightarrow Z$  be a left homotopy witnessing this homotopy. Since  $Z$  is bifibrant,  $i_0$  and  $i_1$  are weak equivalences. By the 2-out-of-3 property  $H''$  and  $hf_\delta$  are weak equivalences. Since  $f_\delta h = id_Y$ , it follows that  $f_\delta$  is a retract of  $f_\delta h$ , and is thus a weak equivalence.  $\square$

**Corollary 2.1.29.1.** *The category  $\mathcal{C}_{cf}/\sim$  satisfy the universal property of the localization of  $\mathcal{C}_{cf}$  by the weak equivalences. I.e. there is a categorical equivalence  $Ho\mathcal{C}_{cf} \simeq \mathcal{C}_{cf}/\sim$ .*

*Proof.* By generalized Whiteheads theorem, theorem 2.1.29 weak equivalences and homotopy equivalences coincide. The corollary follows steadily from both the universal property of the localization category and the quotient category.  $\square$

We collect the results from above in the following theorem.

**Theorem 2.1.30** (Fundamental theorem of model categories). *Let  $\mathcal{C}$  be a model category and denote  $q : \mathcal{C} \rightarrow Ho\mathcal{C}$  the localization functor. Let  $X$  and  $Y$  be objects of  $\mathcal{C}$ .*

1. *There is an equivalence of categories  $Ho\mathcal{C} \simeq \mathcal{C}_{cf}/\sim$ .*
2. *There are natural isomorphisms  $\mathcal{C}_{cf}/\sim(QRX, QRY) \simeq Ho\mathcal{C}(X, Y) \simeq \mathcal{C}_{cf}/\sim(RQX, RQY)$ . Additionally,  $Ho\mathcal{C}(X, Y) \simeq \mathcal{C}_{cf}/\sim(QX, RY)$ .*
3. *The localization  $q$  identifies left or right homotopic morphisms.*
4. *A morphism  $f : X \rightarrow Y$  is a weak equivalence if and only if  $qf$  is an isomorphism.*

*Proof.* This is clear by the results above.  $\square$

### 2.1.3 Quillen adjoints

We now want to study morphisms, or certain functors, between model categories. Like in the case of homotopical functors we want these morphisms to induce a functor between the homotopy categories. However, we also want them to respect the cofibration and fibration structure, not

just weak equivalences. In this way we will instead look towards derived functors to be able to define this extension to the homotopy category. We recall the definition of a total (left/right) derived functor. In the case of model categories, we get a simple description for some of these derived functors which are of special interest.

**Definition 2.1.31** (Total derived functors). Let  $\mathcal{C}$  and  $\mathcal{D}$  be homotopical categories, and  $F : \mathcal{C} \rightarrow \mathcal{D}$  a functor. Whenever it exists, a total left derived functor of  $F$ , is a functor  $\mathbb{L}F : Ho\mathcal{C} \rightarrow Ho\mathcal{D}$  with a natural transformation  $\varepsilon : \mathbb{L}F \circ q \Rightarrow q \circ F$  satisfying the universal property: If  $G : Ho\mathcal{C} \rightarrow Ho\mathcal{D}$  is a functor and there is a natural transformation  $\alpha : G \circ q \Rightarrow q \circ F$ , then it factors uniquely up to unique isomorphism through  $\varepsilon$ .

$$\begin{array}{ccc}
 \mathcal{C} & \xrightarrow{F} & \mathcal{D} \\
 \downarrow q & \nearrow \varepsilon & \downarrow q \\
 Ho\mathcal{C} & \xrightarrow{\mathbb{L}F} & Ho\mathcal{D}
 \end{array}
 \qquad
 \begin{array}{ccc}
 \mathcal{C} & \xrightarrow{F} & \mathcal{D} \\
 \downarrow q & \nearrow \varepsilon & \downarrow q \\
 Ho\mathcal{C} & \xrightarrow{\mathbb{L}F} & Ho\mathcal{D} \\
 \exists! \uparrow \downarrow & & \\
 G & \xrightarrow{\quad} & 
 \end{array}$$

Dually, whenever it exists, a total right derived functor of  $F$ , is a functor  $\mathbb{R}F : Ho\mathcal{C} \rightarrow Ho\mathcal{D}$  with a natural transformation  $\eta : q \circ F \Rightarrow \mathbb{R}F \circ q$  having the opposite universal property.

$$\begin{array}{ccc}
 \mathcal{C} & \xrightarrow{F} & \mathcal{D} \\
 \downarrow q & \nwarrow \eta & \downarrow q \\
 Ho\mathcal{C} & \xrightarrow{\mathbb{R}F} & Ho\mathcal{D}
 \end{array}
 \qquad
 \begin{array}{ccc}
 \mathcal{C} & \xrightarrow{F} & \mathcal{D} \\
 \downarrow q & \nwarrow \eta & \downarrow q \\
 Ho\mathcal{C} & \xrightarrow{\mathbb{R}F} & Ho\mathcal{D} \\
 \exists! \uparrow \downarrow & & \\
 G & \xrightarrow{\quad} & 
 \end{array}$$

**Definition 2.1.32** (Deformation). A left (right) deformation on a homotopical category  $\mathcal{C}$  is an endofunctor  $Q$  together with a natural weak equivalence  $q : Q \Rightarrow Id_{\mathcal{C}}$  ( $q : Id_{\mathcal{C}} \Rightarrow Q$ ).

A left (right) deformation on a functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  between homotopical categories, is a left (right) deformation  $Q$  on  $\mathcal{C}$  such that weak equivalences in the image of  $Q$  is preserved by  $F$ .

*Remark 2.1.33* (Cofibrant and fibrant replacement). If  $\mathcal{C}$  is a model category, then we have a left and a right deformation. The cofibrant replacement  $Q$  defines a left deformation, and the fibrant replacement defines a right deformation. Notice that this is only due to the fact that the factorization system is functorial.

**Proposition 2.1.34.** Let  $F : \mathcal{C} \rightarrow \mathcal{D}$  be a functor between homotopical categories. If  $F$  has a left deformation  $Q$ , then the total left derived functor  $\mathbb{L}F$  exists. Moreover, the functor  $FQ$  is homotopical, and  $\mathbb{L}F$  is the unique extension of  $FQ$ .

*Proof.* Since we already have a candidate for the derived functor, the proof must just check that it has the universal property. A proof may be found in Reihl [16] under proposition 6.4.11.  $\square$

*Remark 2.1.35.* There is a somewhat weaker statement by Dwyer and Spalinski [15]. If we instead ask for functors  $F$  which have the cofibrant replacement  $Q$  (fibrant replacement  $R$ ) as a left (right) deformation we may make this proof more explicit. This is theorem 9.3.

Equipped with the above proposition and remark, it makes sense to define Quillen functors as left and right Quillen functors. A left Quillen functor should be left deformable by the cofibrant replacement. Moreover, for the composition of two left Quillen functors to make sense, we also need weak equivalences between cofibrant objects to be mapped to weak equivalences between cofibrant objects. We make the following definition.

**Definition 2.1.36** (Quillen adjunction). Let  $\mathcal{C}$  and  $\mathcal{D}$  be model categories.

1. A left Quillen functor is a functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  such that it preserves cofibrations and acyclic cofibrations.
2. A right Quillen functor is a functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  such that it preserves fibrations and acyclic fibrations.
3. Suppose that  $(F, U)$  is an adjunction where  $F : \mathcal{C} \rightarrow \mathcal{D}$  is left adjoint to  $U$ .  $(F, U)$  is called a Quillen adjunction if  $F$  is a left Quillen functor and  $U$  is a right Quillen functor.

*Remark 2.1.37.* By Ken Brown's lemma, lemma 2.1.13, we see that a left Quillen functor  $F$  is left deformable to the cofibrant replacement functor  $Q$ . Thus the total left derived functor exists and is given by  $\mathbb{L}F = HoFQ$ .

In order to eliminate the choice of left or right derivedness, we will think of a morphism of model categories as a Quillen adjunction. The direction of the arrow can be chosen to be along either the left or right adjoints, we make the convention of following the left adjoint functors. We summarize the following properties.

**Lemma 2.1.38.** Let  $\mathcal{C}$  and  $\mathcal{D}$  be model categories, and suppose there is an adjunction  $F : \mathcal{C} \rightleftharpoons \mathcal{D} : U$ . The following are equivalent:

1.  $(F, U)$  is a Quillen adjunction.
2.  $F$  is a left Quillen functor.
3.  $U$  is a right Quillen functor.

*Proof.* This follows from naturality of the adjunction. I.e. any square in  $\mathcal{C}$ , with the right side from  $\mathcal{D}$  is commutative if and only if any square in  $\mathcal{D}$  with the left side from  $\mathcal{C}$  is commutative. Now,  $f$  has LLP with respect to  $Ug$  if and only if  $Ff$  has LLP with respect to  $g$ .

$$\begin{array}{ccc}
 A & \xrightarrow{k} & UX \\
 f \downarrow & \nearrow h & \downarrow Ug \\
 B & \xrightarrow{l} & UY
 \end{array}
 \rightsquigarrow
 \begin{array}{ccc}
 FA & \xrightarrow{k^T} & X \\
 Ff \downarrow & \nearrow h^T & \downarrow g \\
 FB & \xrightarrow{l^T} & Y
 \end{array}$$

□

**Proposition 2.1.39.** Suppose that  $(F, U) : \mathcal{C} \rightarrow \mathcal{D}$  is a Quillen adjunction. The functors  $\mathbb{L}F : Ho\mathcal{C} \rightarrow Ho\mathcal{D}$  and  $\mathbb{R}U : Ho\mathcal{D} \rightarrow Ho\mathcal{C}$  forms an adjoint pair.

*Proof.* We must show that  $H\mathcal{O}\mathcal{D}(\mathbb{L}FX, Y) \simeq H\mathcal{O}\mathcal{D}(X, \mathbb{R}UY)$ . By using the fundamental theorem of model categories, theorem 2.1.30, we have the following isomorphisms:  $H\mathcal{O}\mathcal{D}(\mathbb{L}FX, Y) \simeq \mathcal{C}(FQX, RY)/\sim$  and  $H\mathcal{O}\mathcal{D}(X, \mathbb{R}UY) \simeq \mathcal{D}(QX, URY)/\sim$ . In other words, if we assume  $X$  to be cofibrant, and  $Y$  to be fibrant, we must show that the adjunction preserves homotopy equivalences.

We show it for one direction. Suppose that the morphisms  $f, g : FA \rightarrow B$  are homotopic, witnessed by a right homotopy  $H : FA \rightarrow B^I$ . Since we assume  $U$  to preserve products, fibrations and weak equivalences between fibrant objects,  $U(B^I)$  is a path object for  $UB$ . Thus the transpose  $H^T : A \rightarrow U(B^I)$  is the desired homotopy witnessing  $f^T \sim g^T$   $\square$

**Definition 2.1.40** (Quillen equivalence). Let  $\mathcal{C}$  and  $\mathcal{D}$  be model categories, and  $(F, U) : \mathcal{C} \rightarrow \mathcal{D}$  be a Quillen adjunction.  $(F, U)$  is called a Quillen equivalence if for any cofibrant  $X$  in  $\mathcal{C}$ , fibrant  $Y$  in  $\mathcal{D}$  and any morphism  $f : FX \rightarrow Y$  is a weak equivalence if and only if its transpose  $f^T : X \rightarrow UY$  is a weak equivalence.

**Proposition 2.1.41.** Suppose that  $(F, U) : \mathcal{C} \rightarrow \mathcal{D}$  is a Quillen adjunction. The following are equivalent:

1.  $(F, U)$  is a Quillen equivalence.
2. Let  $\eta : Id_{\mathcal{C}} \Rightarrow UF$  denote the unit, and  $\varepsilon : FU \Rightarrow Id_{\mathcal{D}}$  denote the counit. The composite  $Ur_F\eta : Id_{\mathcal{C}} \Rightarrow UR_F|_{\mathcal{C}}$ , and  $\varepsilon_{FQU}Fq_U : FQU|_{\mathcal{D}} \Rightarrow Id_{\mathcal{D}}$  are natural weak equivalences.
3. The derived adjunction  $(\mathbb{L}F, \mathbb{R}U)$  is an equivalence of categories.

*Proof.* Firstly observe that 2.  $\implies$  3. by definition. Secondly observe that equivalences both preserves and reflects isomorphisms, from this we get 3.  $\implies$  1.. We now show 1.  $\implies$  2.. Pick  $X$  in  $\mathcal{C}$  such that  $X$  is cofibrant. Since  $(F, U)$  is assumed to be a Quillen adjunction we know that  $FX$  is still cofibrant. The fibrant replacement  $r_{FX} : FX \rightarrow RFX$  gives us a weak equivalence. Furthermore, since  $(F, U)$  is assumed to be a Quillen equivalence, its transpose  $r_{FX}^T : X \rightarrow URFX$  is a weak equivalence. Unwinding the definition of the transpose we get that  $r_{FX}^T = Ur_{rFX}\eta_X$ .

$\square$

We have the following refinement.

**Corollary 2.1.41.1.** Suppose that  $(F, U) : \mathcal{C} \rightarrow \mathcal{D}$  is a Quillen adjunction. The following are equivalent:

1.  $(F, U)$  is a Quillen equivalence.
2.  $F$  reflects weak equivalences between cofibrant objects, and  $\varepsilon_{FQU}Fq_U : FQU|_{\mathcal{D}} \Rightarrow Id_{\mathcal{D}}$  is a natural weak equivalence.
3.  $U$  reflects weak equivalences between fibrant objects, and  $Ur_F\eta : Id_{\mathcal{C}} \Rightarrow UR_F|_{\mathcal{C}}$  is a natural weak equivalence.

*Proof.* We start by showing  $1. \implies 2.$  and  $3..$  We already know that the derived unit and counit are isomorphism in homotopy, so we only need to show that  $F(U)$  reflects weak equivalences between cofibrant (fibrant) objects. Suppose that  $Ff : FX \rightarrow FY$  is a weak equivalence between cofibrant objects. Since  $F$  preserves weak equivalences between cofibrant objects, we get that  $FQf$  is a weak equivalence, or that  $\mathbb{L}Ff$  is an isomorphism. By assumption,  $\mathbb{L}F$  is an equivalence of categories, so  $f$  is a weak equivalence as needed.

We will show  $2. \implies 1.$ , the case  $3. \implies 1.$  is dual. We assume that the counit map is an isomorphism in homotopy. By assumption, the derived unit  $\mathbb{L}\eta$  is split-mono on the image of  $\mathbb{L}F$ . Moreover, the derived counit  $\mathbb{R}\varepsilon$  is assumed to be an isomorphism, in particular the derived unit  $\mathbb{L}F\mathbb{L}\eta$  is an isomorphism. Unpacking this, we have a morphism, call it  $\eta'_X : FQX \rightarrow FQURFQX$ , which is a weak equivalence. Since  $F$  and  $Q$  reflects weak equivalences, we get that  $\eta_X : X \rightarrow URFQX$  is a weak equivalence.  $\square$

## 2.2 Model structures on Algebraic Categories

In order to understand  $\infty$ -quasi-isomorphism of strongly homotopy associative algebras we will study different homotopy theories of various categories. Munkholm [18] successfully showed that the derived category of augmented algebras is equivalent to the derived category of augmented algebras equipped with  $\infty$ -morphisms. Well, to be more precise, he showed that certain subcategories of augmented algebras had this property. Lefevre-Hasagawas phd thesis [12] builds upon this identification, but with help of further devolpment within the field. We will follow the approach of Lefevre-Hasagawa, by comparing the model structure for algebras and coalgebras,

### 2.2.1 DG-Algebras as a Model Category

Bousefield and Gugenheim [19] proved that the category of commutative dg-algebras had a model structure whenever the base field was a field of characteristic 0. In a joint project, Jardines paper from 1997 [20] shows that this construction may be extended to dg-algebras over any commutative ring. On the other hand, Munkholm expanded on the ideas from Bousfield and Gugenheim to get an identification of derived categories. Also, Hinichs paper from 1997 [21] details another method to obtain the model category which we want. We will follow the approach of Hinich, as it will be usefull later on. Notice that where Hinich use theory of algebraic operads to show that the category of algebras is a model category, we will however give a more explicit formulation.

Let  $\mathbb{K}$  be a field, and  $\mathcal{C}$  be a category such that there is an adjunction  $F : \mathcal{C} \rightleftharpoons Ch(\mathbb{K}) : \#$ , where  $F$  is left adjoint to  $\#$ . Furthermore, suppose that  $\mathcal{C}$  satisfies the 2 conditions:

- (H0)  $\mathcal{C}$  admits finite limits and every small colimit. The functor  $\#$  commutes with filtered colimits.
- (H1) Let  $M$  be the complex below, concentrated in 0 and 1.

$$\dots \longrightarrow 0 \longrightarrow \mathbb{K} \xrightarrow{id} \mathbb{K} \longrightarrow 0 \longrightarrow \dots$$

For any  $d \in \mathbb{Z}$  and for any  $A \in \mathcal{C}$  the injection  $A \rightarrow A \coprod F(M[d])$  induces a quasi-isomorphism  $A^\# \rightarrow (A \coprod F(M[d]))^\#$ .

With this adjunction in mind, we define weak equivalences, fibrations and cofibrations as follows: Let  $f \in \mathcal{C}$  be a morphism

- $f \in Ac$  if  $f^\#$  is a quasi-isomorphism.
- $f \in Fib$  if  $f^\#$  is surjective on each component.
- $f \in Cof$  if  $f$  has LLP to acyclic fibrations.

**Theorem 2.2.1.** *The category  $\mathcal{C}$  equipped with the weak equivalences, fibrations and cofibrations as defined above is a model category.*

Before we show this theorem we need to understand the cofibrations better. Let  $A \in \mathcal{C}$ ,  $M \in Ch(\mathbb{K})$  and  $\alpha : M \rightarrow A^\#$  a morphism in  $Ch(\mathbb{K})$ . We define a functor

$$h_{A,\alpha}(B) = \{(f, t) \mid f \in \mathcal{C}(A, B), t \in Hom_{\mathbb{K}}^{-1}(M, B^\#) \text{ s.t. } \partial t = f^\# \circ \alpha\}.$$

Note that  $t$  is not a morphism of chain maps. This is a homogenous morphism of degree  $-1$ . The differential then promotes this morphism to a chain map, and  $t$  is thus a homotopy for the comoposite  $f^\# \circ \alpha$ .

This functor is represented by an object of  $\mathcal{C}$ . We define this representing object  $A\langle M, \alpha \rangle$  as the pushout:

$$\begin{array}{ccc} F(A^\#) & \xrightarrow{\varepsilon_A} & A \\ \downarrow & & \downarrow a \\ F(\text{cone}(\alpha)) & \xrightarrow{e} & A\langle M, \alpha \rangle \end{array}$$

Let  $i : M[1] \rightarrow \text{cone}(\alpha)$  be a homogenous morphism which is the injection when considered as graded modules. Notice that we have a pair of morphisms  $(a, e^T i) \in h_{A,\alpha}(A\langle M, \alpha \rangle)$ .

**Proposition 2.2.2.** *The functor  $h_{A,\alpha}$  is represented by  $A\langle M, \alpha \rangle$ , i.e.  $h_{A,\alpha} \simeq \mathcal{C}(A\langle M, \alpha \rangle, -)$  is a natural isomorphism. Moreover, the pair  $(a, e^T i)$  is the universal element of the functor  $h_{A,\alpha}$ , i.e. the natural isomorphism is induced by this element under Yoneda's lemma.*

*Proof.* Let  $(f, t) \in h_{A,\alpha}(B)$  for some  $B \in \mathcal{C}$ . The condition that  $\partial t = f^\# \circ \alpha$  is equivalent to say that  $f^\#$  extends to a morphism  $f' : \text{cone}(\alpha) \rightarrow B^\#$  along  $t$ , i.e.  $f' = (f^\# \ t)$ . This concludes the isomorphism part, as being an element  $(f, t)$  is equivalent to the existence of the diagram below, where  $\tilde{f}$  is uniquely determined.

$$\begin{array}{ccc}
F(A^\#) & \xrightarrow{\varepsilon_A} & A \\
\downarrow & & \downarrow a \\
F(\text{cone}(\alpha)) & \xrightarrow{e} & A\langle M, \alpha \rangle
\end{array}
\quad
\begin{array}{c}
\searrow f \\
\downarrow \tilde{f} \\
\searrow f'
\end{array}
\quad
\begin{array}{c}
\\
\\
B
\end{array}$$

To obtain naturality, we use the adjunction to observe that the element  $(a, e^T i)$  is in fact universal.  $\square$

We are now in a position to explicitly find some important cofibrations. We collect these morphisms into the "standard" cofibrations.

**Definition 2.2.3.** Let  $f : A \rightarrow B$  be a morphism in  $\mathcal{C}$ . Suppose that  $f$  factors as a transfinite composition of morphisms on the form  $A_i \rightarrow A_i\langle M, \alpha \rangle$ , i.e.  $f$  factors into the diagram below, where  $A_{i+1} = A_i\langle M, \alpha \rangle$ .

$$A \longrightarrow A_1 \longrightarrow A_2 \longrightarrow \dots \longrightarrow B$$

- If every such  $M$  is a complex consisting of free  $\mathbb{K}$ -modules and has a 0-differential, we call  $f$  a standard cofibration.
- If every such  $M$  is a contractible complex (and  $\alpha = 0$ ), we call  $f$  a standard acyclic cofibration.

**Proposition 2.2.4.** Every standard cofibration is a cofibration, and every standard acyclic cofibration has LLP with respect to fibrations. Moreover, if  $\alpha = 0$ , then every standard acyclic cofibration is also a weak equivalence. We will see that these morphisms in some sense generate every (acyclic) cofibration.

*Proof.* First observe that every standard cofibration may be made iteratively from the chain complexes  $\mathbb{K}[n]$ , and likewise, every standard acyclic cofibration may be made iteratively from  $M$  as in H1.

We first prove that if  $M \simeq \mathbb{K}[n]$ , and  $\alpha : M \rightarrow A^\#$  is any map, then the map  $A \rightarrow A\langle M, \alpha \rangle$  is a cofibration. This amounts to show that it has LLP to every acyclic fibration. Suppose that  $h : B \rightarrow C$  is an acyclic fibration and that there is a commutative square as below.

$$\begin{array}{ccc}
A & \xrightarrow{f} & B \\
\downarrow a & & \downarrow h \\
A\langle M, \alpha \rangle & \xrightarrow{g} & C
\end{array}$$



By the universal property of  $h_{A,\alpha}$  2.2.2 it suffices to find a pair  $(f', t')$  such that  $f : A \rightarrow B$ ,  $t' : M \rightarrow B^\#$  is homogenous of degree  $-1$ ,  $\partial t = f^\# \alpha$  and that  $h$  induces a morphism  $h : (f', t') \rightarrow g$ . We see that we are forced to choose  $f' = f$  as  $hf = ga$ . We know there exists a  $t : M \rightarrow C^\#$  such that  $\partial t = g^\# a^\# \alpha = h^\# f^\# \alpha$ . Since  $h$  is an acyclic fibration  $h^\#$  is a surjective quasi-isomorphism. Since  $M \simeq \mathbb{K}[n]$ , the morphism  $t$  is really an element of  $C^{\#n-1}$ . By surjectivity of  $h^\#$  there is an element  $u$  of  $B^{\#n-1}$  such that  $h^\#(u) = t$ . Moreover, the difference  $h^\#(\partial u - f^\# \alpha) = 0$ , so  $\partial u - f^\# \alpha$  factors through the kernel  $\text{Ker } h^\#$ , which is acyclic. This element is furthermore a cycle, so by acyclicity there is another element  $u'$  such that  $\partial u' = \partial u - f^\# \alpha$ . We may now see that  $(f, u - u')$  is our desired factorization.

Secondly we prove that if  $M$  is in as  $H1$  and  $\alpha = 0$ , then the map  $A \rightarrow A\langle M, \alpha \rangle$  is an acyclic cofibration. This amounts to show that it has LLP to every fibration. Suppose that  $h : B \rightarrow C$  is a fibration and that there is a commutative square as below.

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow a & & \downarrow h \\ A\langle M, \alpha \rangle & \xrightarrow{g} & C \end{array}$$

We will again use 2.2.2, so it suffices to find a  $(t')$  such that  $\partial t' = f^\# \alpha$ . Let  $t : M \rightarrow C^\#$  such that  $\partial t = h^\# f^\# \alpha$ .

Finally we will assume that  $\alpha = 0$ . In this case the cone is a direct sum  $\text{cone}(\alpha) = A^\# \oplus M$ . Since  $F$  is left adjoint to  $\#$ , we know that  $F\text{cone}(\alpha) \simeq F(A^\#) \coprod FM$ . By  $H1$  we get that the map  $A \rightarrow A\langle M, \alpha \rangle \simeq A \coprod FM$  is a weak equivalence.

□

In light of the above proposition we would like to make some more convenient notation. If  $M \simeq \mathbb{K}[n]$  and  $\alpha : M \rightarrow Z^n(A^\#)$ , s.t.  $\alpha(1) = a$ , we write  $A\langle M, \alpha \rangle$  as  $A\langle T; dT = a \rangle$  instead. Hinich calls this "adding a variable to kill a cycle". If  $M$  is the acyclic complex as below and  $\alpha = 0$ , we write  $A\langle T, S; dT = S \rangle$ . This could be thought of "adding a variable and a cycle to kill itself".

$$\dots \longrightarrow 0 \longrightarrow \mathbb{K} \xrightarrow{id} \mathbb{K} \longrightarrow 0 \longrightarrow \dots$$

*proof of theorem.* **MC1** and **MC2** are satisfied. By definition we also have the first part of **MC3**. We start by checking **MC4**.

Let  $f : A \rightarrow B$  be a morphism in  $\mathcal{C}$ . Given any  $b \in B^\#$ , let  $C_b = A\langle T_b, S_b; dT_b = S_b \rangle$ . We define  $g_b : C_b \rightarrow B$  by the conditions that it acts on  $A$  as  $f$ ,  $g_b^\#(T_b) = b$  and  $g_b^\#(S_b) = db$ . Iterating this construction for every  $b \in B$ , we obtain an object  $C$ , such that the injection  $A \rightarrow C$  is an

acyclic standard cofibration, and the map  $g : C \rightarrow B$  is a fibration. This gives us a factorization  $f = f_\delta \circ f_\gamma$ , where  $f_\gamma$  is the injection and  $f_\delta = g$ .

To obtain the other factorization we want to make a standard cofibration. We already know that the map  $A \rightarrow C$  is a standard cofibration, so let  $C_0 = C$ . From here on, we will make each  $C_i$  inductively, such that  $\varinjlim C_i$  has the factorization property which we desire. Notice that from  $C_0$  there is a morphism  $g_0 : C_0 \rightarrow B$ , which is surjective and surjective on every kernel. This morphism may fail to be a quasi-isomorphism, so it is not an acyclic fibration.

To construct  $C_1$  we assign to every pair of elements  $(c, b)$ , such that  $c \in ZC_0^\#$  and  $g_0^\#(c) = db$ , a variable to kill a cycle. If  $(c, b)$  is such a pair, then we add a variable  $T$  such that  $dT = c$  and  $g_1^\#(T) = b$ .  $C_1$  is then the complex where each cycle  $c$  has been killed by adding a variable  $T$ . Now if we suppose that we have constructed  $C_i$ , then  $C_{i+1}$  is constructed in the same manner, by adding a variable to kill each cycle which is a boundary in the image.

When adding a variable, we have also updated the morphism  $g_i$  by letting  $g_{i+1}^\#(T) = b$ . Thus in each step we have also made a new morphism  $g_{i+1}$ . If  $g$  denote the morphism at the colimit, then it is clear that this is still a fibration and it has also become a quasi-isomorphism. This is because every cycle which failed to be in the homology of  $B$  have been killed.

It remains to check the last part of **CM3**. Suppose that  $f : A \rightarrow B$  is an acyclic cofibration. By **CM4**, we know that it factors as  $f = f_\delta \circ f_\gamma$ , where  $f_\delta$  is an acyclic fibration and  $f_\gamma$  is a standard acyclic fibration. We thus obtain that  $f$  is a retract of  $f_\gamma$  by the commutative diagram below.

$$\begin{array}{ccc} A & \xrightarrow{f_\gamma} & C \\ \downarrow f & \nearrow \text{---} & \downarrow f_\delta \\ B & \xlongequal{\quad} & B \end{array}$$

□

The following corollary will concretize what it means that the standard cofibrations generate every cofibration. This corollary is really a step used within in the proof.

**Corollary 2.2.4.1.** *Any (acyclic) cofibration is a retract of a standard (acyclic) cofibration.*

We may immedietly apply this theorem to some familiar examples.

**Corollary 2.2.4.2.** *Let  $A$  be a dg-algebra over the field  $\mathbb{K}$ . The category  $\text{mod}_A$  of left modules is a model category.*

*sketch of proof.* We establish the adjunction by letting  $F = A \otimes_{\mathbb{K}} \_$ . **H0** is satisfied as this category is bicomplete, and filtered colimits may be thought of as unions of sets. Moreover, since  $\text{mod}_A$  is an Abelian category, the forgetful functor  $\#$  commutes with coproducts, or direct sums, which makes **H1** trivially satisfied. □

**Corollary 2.2.4.3.** *The categories  $Alg_{\mathbb{K}}^{\bullet}$  ( $AugAlg_{\mathbb{K}}^{\bullet}$ ) are model categories.*

*Proof.* We establish the adjunction by letting  $F = \bar{T}(T)$ , the reduced tensor algebra of a cochain complex. For the same reasons as above,  $H\bar{0}$  is trivially satisfied. To show H1 we must find the coproduct and what  $\bar{T}(M)$  is. ...  $\square$

We summarize the last result:

The category of augmented dg-algebras  $AugAlg_{\mathbb{K}}^{\bullet}$  is a model category. Let  $f : X \rightarrow Y$  be a homomorphism of augmented algebras.

- $f \in Ac$  if  $f^{\#}$  is a quasi-isomorphism.
- $f \in Fib$  if  $f^{\#}$  is an epimorphism (surjective onto every component).
- $f \in Cof$  if  $f$  has LLP wrt. to every acyclic fibration.

The category of augmented dg-algebras has an initial and a terminal object. The initial object is the stalk  $\mathbb{K}$  and the terminal object is the 0-ring. We see that every object is fibrant, as 0 is preserved by the forgetful functor and every map into 0 is surjective. Every dg-algebra which is isomorphic to a tensor algebra when considered as a graded algebra is cofibrant.

This is wrong

## 2.2.2 A Model Structure on DG-Coalgebras

We now want to equip the category of dg-coalgebras with a suitable model structure. This model structure should be suitable in the sense that it give rise to the same homotopy theory of dg-algebras. The bar-cobar construction will be crucial in this construction, as it is in fact a Quillen adjunction. To this end we will follow the setup as presented by Lefevre-Hasegawa [12]. His method is a modification of Hinichs paper [22] which describes a model structure on dg-coalgebras, but in relation to dg-lie algebras.

Let  $f : C \rightarrow D$ , the category of dg-coalgebras will be equipped with the three following classes of morphisms:

- $f \in Ac$  if  $\Omega f$  is a quasi-isomorphism.
- $f \in Fib$  if  $f$  has RLP wrt. to every acyclic cofibration.
- $f \in Cof$  if  $f^{\#}$  is a monomorphism (injective in every component).

Recall that a filtration on a chain complex  $M$  is a sequence of inclusions  $M_0 \subseteq M_1 \subseteq \dots$ . The filtration is called exhaustive if  $\varinjlim M_i \simeq M$ , and admissible if  $Fr_0 M \simeq \mathbb{K}$  as well. Since each inclusion respect the differential, we may find the cokernel of each inclusion. The associated graded  $grM$  is then the graded complex given by  $gr_0 M = M_0$  and  $gr_i M = M_i/M_{i-1}$ . A dg-(co)algebra is filtered if the filtration respects the (co)multiplication. If  $f : M \rightarrow N$  is a morphism

of filtered cochain complexes, then it defines a morphism  $grf : grM \rightarrow grN$  on the associated graded. We call  $f$  a graded quasi-isomorphism if  $grf$  is a quasi-isomorphism.

Let  $C$  be a conilpotent dg-coalgebra. The coradical filtration  $Fr_0C \subseteq Fr_1C \subseteq \dots$  is an exhaustive filtration by assumption and  $Fr_0C \simeq \mathbb{K}$ , so it is admissible. Observe that  $\Delta Fr_{i+1}C \subseteq Fr_iC \otimes Fr_iC$ . To obtain quasi-isomorphisms of chain complexes we do not need the extra structure of the coalgebra. If not specified otherwise, whenever  $C$  is assumed to be a conilpotent dg-coalgebra then  $grC$  should be the associated graded of the coradical filtration.

**Lemma 2.2.5.** *Let  $f : C \rightarrow C'$  be a graded quasi-isomorphism between conilpotent dg-coalgebras, then  $\Omega f : \Omega C \rightarrow \Omega C'$  is a quasi-isomorphism.*

*Proof.* We do this by considering a spectral sequence. Endow  $C$  with a grading (as a vector space) induced by the coradical filtration, i.e.  $c \in C$  has degree  $|c| = n$  if  $n$  is the smallest number such that  $\bar{\Delta}^n c = 0$ . We define a filtration on  $\Omega C$  by

$$F_p \Omega C = \{sc_1 \otimes \dots \otimes sc_n \mid |c_1| + \dots + |c_n| \leq p\}$$

Since  $C$  is a dg-coalgebra, the coradical filtration respects the differential. In other words,  $F_p \Omega C$  is still a chain complex, which is a subcomplex of  $\Omega C$ . This filtration is clearly bounded below and exhaustive. Thus by the classical convergence theorem of spectral sequences, theorem 5.5.1 [23], the spectral sequence converges to the homology  $E\Omega C \Rightarrow H^* \Omega C$ .

By definition, the 0'th page is defined as  $E_{p,q}^0 \Omega C = (F_p \Omega C)_{p+q} / (F_{p-1} \Omega C)_{p+q}$ . Furthermore, notice that at this page we have the following isomorphism  $E_{p,q}^0 \Omega C \simeq (\Omega grC)_{p+q}^{(p)}$ , where  $(\Omega grC)^{(p)} = \{sc_1 \otimes \dots \otimes sc_n \mid |c_1| + \dots + |c_n| = p\}$ .

Evaluating  $f$  at the 0'th page would look like  $E^0 \Omega f \simeq \Omega grf$ . By the mapping lemma, exercise 5.2.3 [23], it is enough to check that  $\Omega grf$  is a quasi-isomorphism, to see that  $\Omega f$  is a quasi-isomorphism. We show that  $\Omega grf$  is a quasi-isomorphism by inspecting every  $E_{p,\bullet}^0 \Omega C$ .

We define a filtration  $G_k$  on  $E_{p,\bullet}^0 \Omega C$  as

$$G_k = \{sc_1 \otimes \dots \otimes sc_n \mid n \geq -k\}.$$

We see that  $G_0 = E_{p,\bullet}^0 \Omega C$  by definition and  $G_{-p-1} \simeq 0$  on the coaugmentation quotient  $\bar{C}$ . Again, by the classical convergence theorem of spectral sequences, this defines a spectral sequence such that  $EG \Rightarrow H^* E_{p,\bullet}^0 \Omega C$ .

To see that  $\Omega grf$  is a quasi-isomorphism, it is now enough to see that  $E^0 Gf$  is a quasi-isomorphism for any  $p$ . Notice that  $E_{l,\bullet}^0 G \subseteq (grC[-1])^{\otimes l}$  where the total grading is  $p$ . Since  $f$  is a graded quasi-isomorphism and by the Kunneth-formula, theorem 3.6.3 [23], it follows that  $E^0 Gf$  is a quasi-isomorphism.  $\square$

For completeness we include the following statement.

**Lemma 2.2.6.** *Let  $f : A \rightarrow A'$  be a quasi-isomorphism between dg-algebras, then  $Bf : BA \rightarrow BA'$  is both a filtered and unfiltered quasi-isomorphism.*

*Proof.* Notice that the homology of  $BA$  may be calculated from the double complex used to define  $BA$ . In fact, at the 0'th page we have  $E_{p,\bullet}^0 f \simeq f^{\otimes p}$ . It follows that  $f$  is a quasi-isomorphism on the 0'th page from the Kunneth formula, theorem 3.6.3 [23].  $\square$

Let  $A$  ( $C$ ) be a filtered dg-algebra (coalgebra). Given an element  $a \in A$  ( $c \in C$ ) we say that its filtered degree  $fdeg(a)$  ( $fdeg(c)$ ) is the number such that  $a \in F_{fdeg(a)}A$  ( $c \in F_{fdeg(c)}C$ ) but not  $a \in F_{fdeg(a)+1}A$  ( $c \in F_{fdeg(c)+1}C$ ). There is then an associated filtration on the bar (cobar) construction of this complex, defined as  $F_p BA = \{\omega a_1 \otimes \dots \otimes \omega a_n \mid \sum fdeg(a_i) \leq p\}$  ( $F_p \Omega C = \{sc_1 \otimes \dots \otimes sc_n \mid \sum fdeg(c_i) \leq p\}$ ).

**Proposition 2.2.7.** *Let  $A$  be an augmented dg-algebra and  $C$  a conilpotent dg-coalgebra. The counit  $\varepsilon_A : \Omega BA \rightarrow A$  is a quasi-isomorphism. The unit  $\eta_C : C \rightarrow B\Omega C$  is a filtered quasi-isomorphism, moreover  $B\eta_C$  is a quasi-isomorphism.*

The following proof is due to [12], but with corrections given by [24]. Some minor modifications are given to the proof as it resembles a proof given earlier.

*Proof.* We will show that the counit is quasi-invertible. The proof technique for quasi-invertability of the unit is analogous.

Define the following filtration for  $A$ .

$$\begin{aligned} F_0 A &= \mathbb{K} \\ F_1 A &= A \\ F_p A &= F_1 A \end{aligned}$$

We clearly see that this filtration endows  $A$  with the structure of a filtered dg-algebra. For  $\Omega BA$  we will use the induced filtration from the coradical filtration of  $BA$ .

The counit acts on  $\Omega BA$  as tensorwise projection, following with multiplication in  $A$ . This morphism respects the filtration, so it is a filtered morphism. Notice that both of these filtrations are bounded below, and exhaustive, so the classical convergence theorem of spectral sequences apply.

Let  $E_r \Omega BA$  and  $E_r A$  be the spectral sequences given by the filtration. We have that  $E_1^p \Omega BA \simeq gr_p \Omega BA$  and  $E_1^p A \simeq gr_p A$ . For  $p = 1$  both complexes are isomorphic to the same complex,  $\bar{A}$ . Moreover,  $E_1^1 \varepsilon_A = id_{\bar{A}}$ . Whenever  $p \neq 1$  we get that  $E_1^p A \simeq 0$ , so it remains to show that  $E_1^p \Omega BA \simeq gr_p \Omega BA$  is contractible for any  $p \geq 2$ .

The differential of  $\Omega BA$  is generated by three actions: the differential on  $A$ , the multiplication on  $A$  and the comultiplication on  $BA$ . With the induced filtration on  $\Omega BA$ , we see that the multiplication on  $A$  is the only action which maps  $F_p \Omega BA \rightarrow F_{p-1} \Omega BA$ . Thus this action is 0 in the associated graded and the spectral sequence.

There is a homotopy of the identity given as  $h : gr_i \Omega BA \rightarrow gr_i \Omega BA$ , which is 0 unless if there is an element on the form  $s^{-1}(sa) \otimes s^{-1}(\dots) \otimes \dots$ . In this case  $h$  is

$$r(s(\omega a) \otimes s(\dots) \otimes \dots) = (-1)^{|a|+1} s(\omega a \otimes \dots) \otimes \dots$$

We will show that this is a homotopy by induction on  $i$ .

Let  $i = 2$ . Then there are two cases we must handle, either an element is on the form  $s(\omega a_1) \otimes s(\omega a_2)$  or  $s(\omega a_1 \otimes \omega a_2)$ . We consider the latter case first. If we apply  $r$  to this element we are returned 0.

$$(r \circ d_{\Omega BA} + d_{\Omega BA} \circ r)(s(\omega a_1 \otimes \omega a_2)) = r((-1)^{|a_1|+1} s(\omega a_1) \otimes s(\omega a_2)) = s(\omega a_1 \otimes \omega a_2)$$

The we treat the former case

$$\begin{aligned} (r \circ d_{\Omega BA} + d_{\Omega BA} \circ r)(s(\omega a_1) \otimes s(\omega a_2)) \\ = r(s(\omega d_A a_1) \otimes s(\omega a) + (-1)^{|a_1|} s(\omega a_1) \otimes s(\omega d_A a_2)) + d_{\Omega BA}((-1)^{|a_1|+1} s(\omega a_1 \otimes \omega a_2)) \\ = (-1)^{|a_1|} s(\omega d_A a_1 \otimes \omega a_2) - s(\omega a_1 \otimes \omega d_A a_2) + s(\omega a_1) \otimes s(\omega a_2) \\ + (-1)^{|a|+1} s(\omega d_A a_1 \otimes \omega a_2) + s(\omega a_1 \otimes \omega d_A a_2) = s(\omega a_1) \otimes s(\omega a_2) \end{aligned}$$

We see that this is in fact a homotopy, which makes  $id_{gr_2 \Omega BA}$  a null-homotopy.

To extend this argument by induction, we will hypothesize that the terms where the differential is applied will have opposite signs, such that they cancel. Then the result follows for any  $i$ , by observing that the tensors far enough out to the right is not affected by  $r$ .

If  $C$  is a dg-coalgebra, we use the same technique as in 2.2.5. Consider the filtration on  $B\Omega C$  given as

$$F_p B\Omega C = \{\omega(sc_{1,1} \otimes \dots \otimes sc_{1,n_1}) \otimes \dots \otimes \omega(sc_{m,1} \otimes \dots \otimes sc_{m,n_m}) \mid |c_{1,1}| + \dots + |c_{m,n_m}| \leq p\}.$$

This filtration is both bounded below and exhaustive, so the classical convergence theorem says that the associated spectral sequence converges. If we denote this sequence as  $EF$ , then  $EF \implies H^* B\Omega C$ . Let  $EC$  denote the spectral sequence associated to  $C$ . Since  $C$  is conilpotent  $EC \implies H^* C$ . The unit  $\eta_C : C \rightarrow B\Omega C$  is now a map acting on  $EC^0$  as the identity, sending each element in  $EC_{p,q}^0$  to itself in  $EF_{p,q}^0$ .

On each row  $EF_{p,\bullet}^0$ , we make another filtration called  $G$ .

$$G_k EF_{p,\bullet}^0 = \{\omega(\dots)_1 \otimes \dots \otimes \omega(\dots)_n \mid n \geq -k\}$$

Similarly as in 2.2.5, this filtration is bounded below and exhaustive, so we may again apply the classical convergence theorem to obtain a spectral sequence  $E_p G$  such that  $E_p G \implies H^* EF_{p,\bullet}^0 \simeq EF_{p,\bullet}^1$ . Since the unit acts as the identity on  $EC^0$ , it descends to a morphism  $gr_p C \rightarrow E_p G_{k,\bullet}^0$  which is the identity when  $k = -1$  and  $0$  otherwise. Notice that every string of length  $\geq 2$  is not hit by this morphism. However, by employing  $r$  as above we may show that these summands are contractible.

To show that this map is a quasi-isomorphism it is sufficient to show that  $E_p G_{k,\bullet}^0$  is contractible for  $k \neq 1$ . This looks like the same situation as for algebras. We use the same homotopy  $r$  as in the first part to get a null-homotopy of the identity.  $\square$

**Lemma 2.2.8.** *Let  $f : C \rightarrow D$  be a morphism of dg-coalgebras, then:*

- *if  $f$  is a cofibration, then  $\Omega f$  is a standard cofibration.*
- *if  $f$  is a weak equivalence, then  $\Omega f$  is as well.*

*Almost dually, let  $f : A \rightarrow B$  be a morphism of dg-algebras, then:*

- *if  $f$  is a fibration, then  $Bf$  is a fibration.*
- *if  $f$  is a weak equivalence, then  $Bf$  is as well.*

*Proof.* First suppose that  $f : C \rightarrow D$  is a cofibration. We define a filtration on  $D$  as the sum of the image of  $f$  and the coradical filtration on  $D$ :  $D_i = \text{Im} f + Fr_i D$ .  $f$  being a cofibration ensures us that  $D_0 \simeq C$ . Since  $D$  is conilpotent we know that  $D \simeq \varinjlim D_i$ , and that  $\Omega$  commutes with colimits, there is a sequence of algebras  $\Omega C \rightarrow \Omega D_1 \rightarrow \dots \rightarrow \Omega D$ . It is enough to show that each morphism  $\Omega D_i \rightarrow \Omega D_{i+1}$  is a standard cofibration. The quotient coalgebra  $D_{i+1}/D_i$  only has a trivial comultiplication, thus every element is primitive. This means that as a cochain complex  $D_{i+1}$  is constructed from  $D_i$  by attaching possibly very many copies of  $\mathbb{K}$ . We treat the case when there is only one such  $\mathbb{K}$ , here  $D_{i+1} \simeq D_i \oplus \mathbb{K}\{x\}$  where  $dx = y$  for some  $y \in D_i$ . We observe that this is exactly the condition for that the morphism  $\Omega D_i \rightarrow \Omega D_{i+1}$  is a standard cofibration.

If  $f$  is a weak equivalence, then  $\Omega f$  is a quasi-isomorphism by definition.

By lemma 2.1.38, or adjointness more specifically,  $B$  preserving fibrations is a consequence of  $\Omega$  preserving cofibrations.

It remains to show that if  $f : A \rightarrow B$  is a quasi-isomorphism, then  $Bf$  is a weak equivalence. Now,  $Bf$  is a weak equivalence if and only if  $\Omega Bf$  is a quasi-isomorphism. By 2.2.7, the counit  $A \rightarrow \Omega B A$  is a quasi-isomorphism, so  $Bf$  is a weak equivalence by 2-out-of-3 property.

$$\begin{array}{ccc}
A & \xrightarrow{f} & B \\
\varepsilon_A \uparrow & & \varepsilon_B \uparrow \\
\Omega BA & \xrightarrow{\Omega Bf} & \Omega BB
\end{array}$$

□

We will need one more technical lemma.

**Lemma 2.2.9.** *Let  $A$  be a dg-algebra,  $D$  a dg-coalgebra and  $p : A \rightarrow \Omega D$  a fibration of algebras. The projection morphism  $\pi : BA \prod_{B\Omega D} D \rightarrow BA$  is an acyclic cofibration.*

$$\begin{array}{ccc}
BA \prod_{B\Omega D} D & \longrightarrow & D \\
\downarrow \pi & & \downarrow \eta_D \\
BA & \xrightarrow{Bp} & B\Omega D
\end{array}$$

*Proof.*  $\pi$  being a cofibration is immediate by corollary 2.1.9.1. To see that  $\pi$  is a quasi-isomorphism it is enough to understand that it is a quasi-isomorphism as chain complexes. This is checked by Lefevre-Hasegawa [12]. Keller has pointed out a slight mistake in this proof which he has proposed a fix to [25]. □

**Theorem 2.2.10.** *The category  $\text{ConilCoalg}_{\mathbb{K}}^{\bullet}$  is a model category with the classes  $Ac$ ,  $Fib$  and  $Cof$  as defined above.*

*Proof.* The axioms **MC1** and **MC2** are immediet. Also, fibrations having RLP wrt. acyclic cofibrations is by definition.

We show **MC4** first. Let  $f : C \rightarrow D$  be a morphism of coalgebras. There is a factorization  $\Omega f = pi$  of morphisms between algebras, where  $i$  is a cofibration,  $p$  is a fibration and at least one of  $i$  and  $p$  are quasi-isomorphisms. Applying bar we get a factorization  $B\Omega f = BiBp$ , where  $Bp$  is a fibration and at least one of  $Bi$  and  $Bp$  are weak equivalences.

$$\begin{array}{ccc}
\Omega C & \xrightarrow{\Omega f} & \Omega D \\
\searrow i & & \nearrow p \\
& A &
\end{array}
\rightsquigarrow
\begin{array}{ccc}
B\Omega C & \xrightarrow{B\Omega f} & B\Omega D \\
\searrow Bi & & \nearrow Bp \\
& BA &
\end{array}$$

We construct a pullback with  $Bp$  and  $\eta_D$ . By 2.2.9 the morphism  $\pi$  is an acyclic cofibration. We collect our morphisms in a big diagram. The dashed arrow exists since the rightmost square is a pullback.



$$\begin{array}{ccccc}
 & & BA \prod_{B\Omega D} D & & \\
 & \nearrow j & \downarrow \tau & \searrow q & \\
 C & \xrightarrow{f} & & & D \\
 \downarrow \eta_C & & \downarrow & & \downarrow \eta_D \\
 & \nearrow Bi & BA & \searrow Bp & \\
 B\Omega C & \xrightarrow{B\Omega f} & & & B\Omega D
 \end{array}$$

First notice that  $q$  is a fibration, since fibrations are stable under pullbacks.  $j$  is a cofibration, or a monomorphism, as the composition  $Bi\eta_C$  is a monomorphism. Thus it remains to see that if  $Bi(Bp)$  is a weak equivalence, then  $j(q)$  is as well. This is evident from the 2-out-of-3 property, as  $\eta$  is a natural weak equivalence,  $\pi$  is a weak equivalence and  $Bi(Bp)$  is a weak equivalence.

We now show **CM3**. Suppose that there is a square as below, where  $i$  is a cofibration and  $t$  is an acyclic cofibration.

$$\begin{array}{ccc}
 E & \longrightarrow & C \\
 \downarrow i & & \downarrow t \\
 F & \longrightarrow & D
 \end{array}$$

We can factor  $t$  as  $t = qj$  by **CM4**. Notice that  $t$  is a retract of  $q$ , i.e. there is a commutative diagram as below.

$$\begin{array}{ccc}
 C & \xlongequal{\quad} & C \\
 \downarrow j & \nearrow & \downarrow t \\
 BA \prod_{B\Omega A} D & \xrightarrow{q} & D
 \end{array}$$

So in order to find a lift to  $C$ , we may instead find a lift to  $BA \prod_{B\Omega A} D$ . Since  $p$  is an acyclic fibration by construction and  $\Omega i$  is a cofibration by 2.2.8, there is a lift  $h : \Omega E \rightarrow A$  of algebras. We obtain our desired lift from the bar-cobar adjunction and the universal property of the pullback.

$$\begin{array}{ccccc}
E & \longrightarrow & BA \coprod_{B\Omega D} D & \xrightarrow{\pi} & BA \\
\downarrow i & \nearrow & \downarrow q & \nearrow h^F & \downarrow Bp \\
F & \longrightarrow & D & \xrightarrow{\eta_D} & B\Omega D
\end{array}
\quad \longleftrightarrow \quad
\begin{array}{ccc}
\Omega E & \longrightarrow & A \\
\downarrow \Omega i & \nearrow h & \downarrow p \\
\Omega F & \longrightarrow & \Omega D
\end{array}$$

□

We restate the corollary of the adjunction.

**Corollary 2.2.10.1.** *The bar-cobar construction  $B : AugAlg_{\mathbb{K}}^{\bullet} \rightleftharpoons ConilCoalg_{\mathbb{K}}^{\bullet} : \Omega$  as a Quillen equivalence.*

*Proof.* We first observe that  $(B, \Omega)$  is a Quillen adjunction by lemma 2.2.8. Moreover, since the unit and counit are weak equivalences by proposition 2.2.7, it follows by either proposition 2.1.41 or its corollary 2.1.41.1 that  $(B, \Omega)$  is a Quillen equivalence. □

### 2.2.3 Homotopy theory of $A_{\infty}$ -algebras

This section aims to finalize the discussion of the homotopy theory of  $A_{\infty}$ -algebras. We will look at the homotopy invertability of every strongly homotopy associative quasi-isomorphism, and the relation to associative algebras. This discussion will end with mentioning different results which gives a clearer description of fibrations, cofibrations and homotopy equivalences. This section follows Lefevre-Hasegawa [12]. Before we get to the main theorem, we start by discussing a non-closed model structure on the category of  $Alg_{\infty}$ .

Let  $f : A \rightsquigarrow B$  be a morphism between  $A_{\infty}$ -algebras, the category of  $A_{\infty}$ -algebras will be equipped with the three following classes of morphisms:

- $f \in Ac$  if  $f$  is an  $\infty$ -quasi-isomorphism, i.e.  $f_1$  is a quasi-isomorphism.
- $f \in Fib$  if  $f_1$  is an epimorphism.
- $f \in Cof$  if  $f_1$  is a monomorphism.

This category does not make a model category in the sense of a closed model category, as we are lacking many finite limits. It does however come quite close to be such a category.

**Theorem 2.2.11.** *The category  $Alg_{\infty}$  equipped with the three classes as defined above satisfies:*

- a The axioms **MC1** through **MC4**.
- b Given a diagram as below, where  $p$  is a fibration, then its limit exists.

$$\begin{array}{ccc} & & A \\ & & \downarrow p \\ B & \longrightarrow & C \end{array}$$

We will first prove one lemma which technique we will reuse. A seconde lemma used to simplify the theorem will be stated without proof.

**Lemma 2.2.12.** *let  $A$  be an  $A_\infty$ -algebra, and  $K$  a contractible complex considered as an  $A_\infty$ -algebra. If  $g : (A, m_1^A) \rightarrow (K, m_1^K)$  is a cochain map, then it extends to an  $\infty$ -morphism  $f : A \rightsquigarrow K$ .*

*Proof.* We construct each  $f_i$  inductively. The case  $i = 1$  is degenerate as we have assumed  $f_1 = g$ .

Assume that we have already constructed  $f_1$  through  $f_n$ . We observe that the sum below is a cycle of  $Hom_{\mathbb{K}}^*(A, K)$ .

$$\sum_{\substack{p+1+r=k \\ p+q+r=n}} (-1)^{pq+r} f_k \circ_{p+1} m_q^A - \sum_{\substack{k \geq 2 \\ i_1 + \dots + i_k = n}} (-1)^e m_k^B \circ (f_{i_1} \otimes f_{i_2} \otimes \dots \otimes f_{i_k})$$

Thus since  $K$  is contractible,  $Hom_{\mathbb{K}}^*(A, K)$  is acyclic and there exists some morphism  $f_{n+1}$  such that  $\partial(f_{n+1})$  is the sum above. This says that this extension does in fact satisfy  $(rel_{n+1})$ .  $\square$

**Lemma 2.2.13.** *Let  $j : A \rightsquigarrow D$  be a cofibration of  $A_\infty$ -algebras, then there is an isomorphism  $k : D \rightsquigarrow D'$  such that the composition  $k \circ j : A \rightsquigarrow D'$  is a strict morphism of  $A_\infty$ -algebras.*

*Dually, if  $j : A \rightsquigarrow D$  is a fibration, then there is an isomorphism  $l : A' \rightsquigarrow A$  such that the composition  $j \circ l : A' \rightsquigarrow D$  is a strict morphism of  $A_\infty$ -algebras.*

*Proof.* A proof is given as lemma 1.3.3.3 in [12].  $\square$

*proof of 2.2.11.* We start by showing b. Suppose that we have a diagram of  $A_\infty$ -algebras, such that  $g_1$  is an epimorphism.

$$\begin{array}{ccc} & & A \\ & & \downarrow g \\ A' & \xrightarrow{f} & A'' \end{array}$$

First notice that as dg-coalgebras, this pullback exists and defines a new dg-coalgebra  $BA \prod_{BA''} BA'$ .

Since  $g_1$  is an epimorphism,  $A[1]$  as a graded vector space splits into  $A''[1] \oplus K$ , where  $K = \text{Ker} g_1$ . The pullback is then naturally identified with  $BA \prod_{BA''} BA' \simeq \bar{T}^c(K) \prod \bar{T}^c(A'[1])$  as graded vector spaces. Since the cofree coalgebra is right adjoint to forget, it commutes with products and we get,  $\bar{T}^c(A'[1]) \prod \bar{T}^c(K) \simeq \bar{T}^c(A'[1] \oplus K)$ . Thus the pullback is isomorphic to a cofree coalgebra as a graded coalgebra, i.e. it is an  $A_\infty$ -algebra.

We now prove a. MC1 and MC2 are immediate, so we will not prove them.

We start by proving MC3. Suppose that there is a square of  $A_\infty$ -algebras as below, where  $j$  is a cofibration and  $q$  is a fibration.

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow j & & \downarrow q \\ C & \xrightarrow{g} & D \end{array}$$

By lemma 2.2.13, we may assume that both  $j$  and  $q$  are strict morphisms. We now assume that  $q$  is an  $\infty$ -quasi-isomorphism, the proof will be analogous if  $j$  is an  $\infty$ -quasi-isomorphism instead.

Our goal is to construct a lifting in this diagram inductively. Having a lift means finding an  $\infty$ -morphism  $a : C \rightsquigarrow B$ , such that the following hold for any  $n \geq 1$ :

- $a$  satisfy  $(rel_n)$ .
- $a_n \circ j_1 = f_n$ .
- $q_1 \circ a_n = g_n$ .

We start by showing there is such an  $a_1$ . Consider the diagram below of chain complexes over  $\mathbb{K}$ .

$$\begin{array}{ccc} A & \xrightarrow{f_1} & B \\ \downarrow j_1 & \nearrow a_1 & \downarrow q_1 \\ C & \xrightarrow{g_1} & D \end{array}$$

The lift exists since the category  $Ch\mathbb{K}$  is a model category. Here  $j_1$  is a cofibration, while  $q_1$  is an acyclic fibration, so the lift  $a_1$  exists.

We now wish to extend this. Suppose that we have been able to create morphisms  $a_1$  up to  $a_n$ , all satisfying the above points. A naive solution to make  $a_{n+1}$  is  $b = f_{n+1}r^{\otimes n+1} + sg_{n+1} - sq_1f_{n+1}r^{\otimes n+1}$ . Notice that this satisfy the two last points by definition. We will augment  $b$  to get an  $a_{n+1}$  which also satisfies  $(rel_{n+1})$ .

For our own convenience, let  $-c(f_1, \dots, f_n)$  denote the right hand side of  $(rel_{n+1})$  formula. Since both  $j$  and  $q$  are strict  $\infty$ -morphisms we get the following identities:

$$\begin{aligned} (\partial b + c(a_1, \dots, a_n)) \circ j_1 &= \partial(b \circ j_1) + c(a_1 \circ j_1, \dots, a_n \circ j_1) = \partial f_{n+1} + c(f_1, \dots, f_n) = 0 \\ q_1 \circ (\partial b + c(a_1, \dots, a_n)) &= \partial(q_1 \circ b) + c(q_1 \circ a_1, \dots, q_1 \circ a_n) = \partial g_{n+1} + c(g_1, \dots, g_n) = 0 \end{aligned}$$

We thus obtain that the cycle  $\partial b + c(a_1, \dots, a_n)$  factors through the cokernel of  $j$  and the kernel of  $q$ . Let us say that it factors like the diagram below:

$$C \xrightarrow{p} \text{Cok} j_1 \xrightarrow{c'} \text{Ker} q_1 \xrightarrow{i} D$$

Now,  $c'$  is a morphism between two  $A_\infty$ -algebras. Since  $q$  is assumed to be an  $\infty$ -quasi-isomorphism, it follows that  $\text{Ker} q$  is acyclic. Since  $c'$  is a cycle, it necessarily has to be in the image of the differential. Let  $h$  be a morphism such that  $\partial h = c'$ , and define  $a_{n+1} = b - i \circ h \circ p$ . One may check that this morphism satisfies all three properties.

We will now show MC4. Since the two properties have a similar proof, we will only show one direction. Let  $f : A \rightsquigarrow B$  be an  $\infty$ -morphism. Let  $C = \text{cone}(id_B[-1])$ . The complex  $C$  may be considered as an  $A_\infty$ -algebra. Let  $j : A \rightsquigarrow A \amalg C$  be the morphism induced by  $id_A$  and  $0 : A \rightarrow C$ . The canonical projection  $q_1 : A \oplus C \rightarrow B$  gives a lift of the following diagram.

$$\begin{array}{ccc} A & \xrightarrow{f_1} & B \\ \downarrow j_1 & \nearrow q_1 & \downarrow \\ A \oplus C & \longrightarrow & 0 \end{array}$$

Since we have a morphism of chain complexes, lodged between an acyclic cofibration and a fibration we use the same technique as above to construct an  $\infty$ -morphism  $q : A \amalg C \rightarrow B$ .  $q$  is a fibration by construction. The morphism  $f$  may be factored as  $f = q \circ j$ , where  $j$  is an acyclic cofibration and  $q$  is a fibration.  $\square$

With this model structure we are finally able to characterize the fibrant and cofibrant conilpotent dg-coalgebras.

**Proposition 2.2.14.** *Let  $C$  be a conilpotent dg-coalgebra. Then  $C$  is cofibrant, and  $C$  is fibrant if and only if there is a cochain complex  $V$ , such that  $C \simeq T^c(V)$  as complexes.*

*Proof.* To see that  $C$  is cofibrant is the same as to verify that the map  $\mathbb{K} \rightarrow C$  is a monomorphism, but this is clear.

We start by assuming that  $C$  is fibrant. Then there is a lift in the square below, making the unit split-mono.

$$\begin{array}{ccc}
C & \xlongequal{\quad} & C \\
\downarrow \eta_C & \nearrow r & \downarrow \varepsilon_C \\
B\Omega C & \xrightarrow{\varepsilon_{B\Omega C}} & \mathbb{K}
\end{array}$$

Consider the morphism  $p_1^C : C \rightarrow Fr_1 C$  which is defined as  $p_1^C = Fr_1 r \circ p_1 \circ \eta_C$ , where  $p_1 : B\Omega C \rightarrow Fr_1 B\Omega C$  is the canonical projection on the filtration induced by the coradical filtration on  $C$ . Clearly,  $r$  makes  $p_1$  into a universal arrow in the category of conilpotent coalgebras, so  $C \simeq T^c(Fr_1 C)$ .

Now, assume that  $C$  is isomorphic to  $T^c(V)$  as coalgebras for some cochain complex  $V$ . Note that, by definition,  $C$  is an  $A_\infty$ -algebra. By definition, we have a commutative square of  $A_\infty$ -algebras. Since every  $A_\infty$ -algebra is bifibrant, we know that this diagram has a lift, exhibiting  $C$  as a retract of  $B\Omega C$ .

$$\begin{array}{ccc}
C & \xlongequal{\quad} & C \\
\downarrow & \nearrow & \downarrow \\
B\Omega C & \longrightarrow & \mathbb{K}
\end{array}$$

We know that  $\Omega C$  is fibrant, since the map  $\Omega C \rightarrow 0$  is epi. By lemma 2.2.8, we know that the bar construction preserves fibrations, so  $B\Omega C$  is fibrant. Thus  $C$  is fibrant as well.  $\square$

The model structure of  $A_\infty$ -algebras is compatible with the model structure of conilpotent dg-coalgebras in the following sense. If  $f : A \rightsquigarrow B$  is an  $\infty$ -morphism, we denote its dg-coalgebra counterpart as  $Bf : BA \rightarrow BB$ . Remember that the bar construction is extended such that it is an equivalence of categories on its image. We use this to realize  $Alg_\infty$  as a subcategory of  $ConilCoalg_{\mathbb{K}}$  to essentially obtain 2 different model structure on this category. The following proposition tells us that these structures do not differ.

**Proposition 2.2.15.** *Let  $f : A \rightsquigarrow B$  be an  $\infty$ -morphism. Then we have the following:*

- $f$  is an  $\infty$ -quasi-isomorphism if and only if  $Bf$  is a weak equivalence.
- $f_1$  is an epimorphism if and only if  $Bf$  is a fibration.
- $f_1$  is a monomorphism if and only if  $Bf$  is a monomorphism.

*Proof.* This is proposition 1.3.3.5 in [12].  $\square$

## 2.3 The Homotopy Category of $Alg_\infty$

With the results we have establish, we are now ready to talk about homotopies in  $Alg_\infty$ .

**Theorem 2.3.1.** *In the category  $Alg_\infty$  we have the following:*

- *Homotopy equivalence is an equivalence relation.*
- *A morphism is an  $\infty$ -quasi-isomorphism if and only if it is a homotopy equivalence.*
- *Let  $dash \subseteq Alg_\infty$  be the full subcategory consisting of dg-algebras considered as  $A_\infty$ -algebras.  $dash$  has an induced homotopy equivalence from  $Alg_\infty$ , and the inclusion  $Alg \rightarrow dash$  induces an equivalence in homotopy  $Alg[Qis^{-1}] \simeq dash/\sim$ .*

*Proof.* The first point is observed from corollary 2.1.27.2, and the second point is Whiteheads theorem, theorem 2.1.29.

To see the final point, observe that the inclusion functor is given by the bar construction  $B$ . By corollary 2.2.10.1, we know that the bar construction induces an equivalence on the homotopy categories, i.e.  $HoAlg \simeq HoCoalg$ . Moreover, we know that by theorem 2.1.30 that  $HoCoalg \simeq Alg_\infty/\sim$ . Notice that the image of  $B$  is  $dash$ , so in homotopy, we get that the image  $dash/\sim$  is equivalent to the essential image  $HoAlg_\infty$ .  $\square$

Homotopy equivalence in the algebraic sense and the model categorical sense differ in how they are defined. In homological algebra, two morphisms are called homotopic if their difference is a boundary of some homotopy, i.e.  $f - g = \partial h$ . In the model categorical sense, a homotopy is a morphism through either a cylinder or a path object. The following proposition tells us how they differ in the category of conilpotent dg-coalgebras.

**Proposition 2.3.2.** *Let  $C$  and  $D$  be two conilpotent dg-coalgebras, where  $f, g : C \rightarrow D$  are two morphisms. Then:*

- *If  $f - g$  is null homotopic, then they are left homotopic.*
- *If  $D$  is fibrant, then  $f - g$  is null homotopic if and only if  $f$  and  $g$  are left homotopic.*

This tells us that in general, these concepts are usually not the same. However, we know that the subcategory of bifibrant objects in this category is exactly the subcategory  $Alg_\infty$ . Thus for  $A_\infty$ -algebras homological homotopy is the same as model categorical homotopy. In this sense we obtain that the homotopy category in the homological sense is equivalent to the derived category.





## Chapter 3

# Derived Categories of Strongly Homotopy Associative Algebras

In this chapter we wish to study the derived categories of  $A_\infty$ -algebras. At the heart of homological algebra is the derived category of algebras, so it is only natural to ask how this category looks like in the  $A_\infty$  case. In the last chapter we studied the relationship between the category of algebras and coalgebras to understand how quasi-isomorphisms between  $A_\infty$ -algebras worked. In this chapter we will instead study the relationship between module and comodule categories in order to understand how quasi-isomorphisms between  $A_\infty$ -modules will work. At the heart of this discussion are twisting morphisms  $\alpha : C \rightarrow A$ , which allows us to study the relationship between  $Mod^A$  and  $CoMod^C$ .

From twisting morphisms we will obtain functors  $L_\alpha : CoMod^C \rightarrow Mod^A$  and  $R_\alpha : Mod^A \rightarrow CoMod^C$  which create an adjoint pair of functors. Whenever the twisting morphism  $\alpha$  is acyclic, this will in fact become a Quillen Equivalence.

We wish to reuse all of the methods we have gained and acquired throughout this thesis. This chapter will mostly be reformulation and recontextualization of previous definitions, concepts and techniques.

### 3.1 Twisting Morphisms

Twisting morphisms were already introduced in chapter 1. There, they were used mostly to be represented by the bar and cobar construction. Now we want twisting morphisms and twisting tensors to play a bigger role. In order to define the functors  $L_\alpha$  and  $R_\alpha$ , these constructions will be crucial.

### 3.1.1 Twisted Tensor Products

Let  $A$  be an augmented dg-algebra,  $C$  a conilpotent dg-coalgebra and  $\alpha : C \rightarrow A$  a twisting morphism. The right (left) twisted tensor product is the complex  $C \otimes_\alpha A$  ( $A \otimes_\alpha C$ ) together with the differential  $d_\alpha^\bullet = d_{C \otimes A}^\bullet + d_\alpha^r$ . The perturbation is defined as

$$d_\alpha^r = (\nabla_A \otimes id_C) \circ (id_A \otimes \alpha \otimes id_C) \circ (id_A \otimes \Delta_C).$$

If  $M$  is a right  $A$ -module and  $N$  is a left  $C$ -comodule then the tensor product  $M \otimes_{\mathbb{K}} N$  exists and is a  $\mathbb{K}$ -module with differential  $d_{M \otimes N}$ . We may define a perturbation to this differential as

$$d_\alpha^r = (\mu_M \otimes id_N) \circ (id_M \otimes \alpha \otimes id_N) \circ (id_M \otimes \nu_N).$$

By using the same line of thought as proposition 1.2.4, there is a twisted tensor product  $M \otimes_\alpha N$  with differential  $d_\alpha^\bullet = d_{M \otimes N} + d_\alpha^r$ .

*Remark 3.1.1.* Koszuls sign rule forces us to define the differential of the left twisted tensor product as  $d_\alpha^\bullet = d_{N \otimes M} - d_\alpha^l$ .

**Definition 3.1.2.** Suppose that  $M \in Mod^A$  ( $M \in Mod_A$ ) and  $N \in CoMod_C$  ( $N \in CoMod^C$ ), then the left (right) twisted tensor product is the  $\mathbb{K}$ -module  $M \otimes_\alpha N$  ( $N \otimes_\alpha M$ ).

In this setting right handedness and left handedness for the twisted tensor product is more clear in this setting, as we only have an action or coaction from one of the chosen sides. Trying to force the other handedness on the twisted tensors would just be ill-defined.

**Definition 3.1.3.** Let  $A$  be an augmented dg-algebra and  $C$  a conilpotent dg-coalgebra, such that there is a twisting morphism  $\alpha : C \rightarrow A$ . Given a linear map  $f : N \rightarrow M$  between a right  $C$ -comodule  $N$  and a right  $A$ -module  $M$  we say that it is an  $\alpha$  right twisted linear homomorphism, or just an  $\alpha$  twisted morphism, if it satisfies the following equation:

$$\partial f - f \star \alpha = 0$$

This definition gives us a functor  $Tw_\alpha : CoMod^C \times Mod^A \rightarrow Ab$  which is the collection of right twisting linear homomorphisms between a comodule and module.

Suppose that  $\alpha : C \rightarrow A$  is a twisting morphism. We define the functor  $L_\alpha = \otimes_\alpha A : CoMod^C \rightarrow Mod^A$  as an arbitrary right twisted tensor product with  $A$ . This functor does indeed hit  $Mod^A$  by using the free right  $A$ -module structure on  $A$ . Likewise, we define a functor  $R_\alpha = \otimes_\alpha C : Mod^A \rightarrow CoMod^C$  as an arbitrary left twisted tensor product with  $C$ . This does also hit right  $C$ -comodules by using the free right  $C$ -comodule structure on  $C$ .

**Proposition 3.1.4.** Suppose that  $\alpha : C \rightarrow A$  is a twisting morphism. The functor  $L_\alpha$  and  $R_\alpha$  form an adjoint pair of categories.

$$\begin{array}{ccc}
& L_\alpha & \\
\text{CoMod}^C & \xrightarrow{\quad} & \text{Mod}^A \\
& \perp & \\
& R_\alpha &
\end{array}$$

*Proof.* This proof breaks down to showing  $\text{CoMod}^C(N, L_\alpha(M)) \simeq \text{Tw}_\alpha(N, M) \simeq \text{Mod}^A(R_\alpha(N), M)$ . This is a routine calculation, much like the proof for 1.2.13.  $\square$

Let  $A$  be a dg-algebra, and  $M$  a right  $A$ -module. Recall that by the Cobar-Bar adjunction 1.2.13 there exists a universal twisting morphism  $\pi_A : BA \rightarrow A$ . We define the bar-construction of  $M$  as  $B_A M = R_{\pi_A} M = M \otimes_{\pi_A} BA$ . Likewise, given a conilpotent dg-coalgebra  $C$  and  $N$  a right  $C$ -comodule we define the cobar-construction as  $\Omega_C N = L_{\iota_C} N = N \otimes_{\iota_C} \Omega C$ . In these cases we obtain adjunctions  $\Omega_{BA} \dashv B_A$  and  $\Omega_C \dashv B_{\Omega C}$ .

Let  $A$  and  $B$  be two algebras and  $f : A \rightarrow B$  is an algebra morphism. Then  $f$  induces a functor between the module categories by restriction:  $f^* : \text{Mod}^B \rightarrow \text{Mod}^A$ . Since  $A$  and  $B$  considered as algebroids are small, and the category of abelian groups is cocomplete, so the left Kan extension (induction) along this functor exists.

$$\begin{array}{ccc}
& f_! & \\
\text{Mod}^B & \xrightarrow{\quad} & \text{Mod}^A \\
& f^* &
\end{array}$$

Dually, if  $C$  and  $D$  are two coalgebras and  $g : C \rightarrow D$  is an coalgebra morphism. Then  $g$  induces a functor between the module categories by composing:  $g_* : \text{CoMod}^C \rightarrow \text{CoMod}^D$ . Since  $C$  and  $D$  considered as coalgebroids are small, and the category of abelian groups is complete, so the right Kan extension (co-induction) along this functor exists.

$$\begin{array}{ccc}
\text{CoMod}^C & \xrightarrow{g_*} & \text{CoMod}^D \\
& g^! &
\end{array}$$

**Lemma 3.1.5.** *Let  $\tau : C \rightarrow A$  be a twisting morphism. The adjunction  $(L_\tau, R_\tau)$  factors as  $(f_{\tau!}, f_\tau^*) \circ (L_{\iota_C}, R_{\iota_C})$  or  $(L_{\pi_A}, R_{\pi_A}) \circ (g_{\tau*}, g_\tau^!)$ .*

*Proof.* This follows from corollary 1.2.14.1, that is  $\tau = f_\tau \circ \iota_C = \pi_A \circ g_\tau$ .  $\square$

**Definition 3.1.6.** A twisting morphism  $f : C \rightarrow A$  is called acyclic if the counit of the adjunction  $L_\alpha \dashv R_\alpha$  is a pointwise quasi-isomorphism.

**Lemma 3.1.7.** *Let  $A$  be an augmented dg-algebra and  $C$  a conilpotent dg-coalgebra. The universal twisting morphisms  $\pi_A$  and  $\iota_C$  are acyclic.*

*Proof.* We start with  $\pi_A$ . Recall that  $\pi_A$  is constructed as the twisting morphism corresponding to  $id_{BA}$ . This morphism is thus given as the projection onto the first dimension of  $BA$ , that is:

$$\begin{aligned}\pi_A sa &= a \\ \pi_A(sa \otimes \dots) &= 0\end{aligned}$$

We say that  $\pi_A$  is acyclic if the counit  $\varepsilon : L_{\pi_A} R_{\pi_A} \Rightarrow Id_{Mod^A}$  at each object  $M$  is a quasi-isomorphism.

For each  $M$  in  $Mod^A$ ,  $L_{\pi_A} R_{\pi_A} M = M \otimes_{\pi_A} BA \otimes_{\pi_A} A$ . We may split up the differential into two summands,  $d_v$  and  $d_h$ .  $d_v$  is the ordinary differential on the tensor product, while  $d_h = (-d_{\pi_A}^l \otimes A) + M \otimes d_2 \otimes A + d_{\pi_A}^r$ . Since  $(d_v + d_h)^2 = 0$  and  $d_v^2 = 0$  we can observe that  $d_v d_h = -d_h d_v$  and  $d_h^2 = 0$ . This is evident as  $d_v$  changes the homological degree while  $d_h$  does not, so in order for both of the first equations to hold, the last two must hold as well. We almost obtain a double complex.

$$\dots \xrightarrow{d_h} M \otimes BA_{(i)} \otimes A \xrightarrow{d_h} \dots \xrightarrow{d_h} M \otimes BA_{(1)} \otimes A \xrightarrow{d_h} M \otimes A \xrightarrow{0} 0$$

$\begin{array}{c} \curvearrowright d_v \\ \curvearrowright d_v \\ \curvearrowright d_v \end{array}$

It is clear that the total complex of this "double complex" is in fact  $L_{\pi_A} R_{\pi_A} M$ . Moreover, the counit induces an augmentation to this complex resolution of  $M$ , denoted as  $cone(\varepsilon_M)$ .

$$\dots \xrightarrow{d_h} M \otimes BA_{(i)} \otimes A \xrightarrow{d_h} \dots \xrightarrow{d_h} M \otimes BA_{(1)} \otimes A \xrightarrow{d_h} M \otimes A \xrightarrow{\varepsilon_M} M \xrightarrow{0} 0$$

$\begin{array}{c} d_{M \otimes BA_{(i)} \otimes A} \\ \curvearrowright \\ d_{M \otimes BA_{(1)} \otimes A} \\ \curvearrowright \\ d_{M \otimes A} \\ \curvearrowright \\ d_M \end{array}$

To see that this is in fact a resolution we define a morphism  $h : cone(\varepsilon_M) \rightarrow cone(\varepsilon_M)$  of degree  $-1$ . It works by the following formula:

$$h(m \otimes (sa_1 \otimes \dots \otimes sa_n) \otimes a) = m \otimes (sa_1 \otimes \dots \otimes sa_n \otimes sa) \otimes 1$$

It is clear that  $id_{cone(\varepsilon_M)} = d_h h - h d_h$  and  $d_v h = h d_v$ . Thus to see that the cone is acyclic we let  $c \in cone(\varepsilon_M)$  be a cycle, that is  $(d_v + d_h)(c) = 0$ . Our goal is to show that  $h(c)$  is a preimage of  $c$  along  $d_v + d_h$ .

$$(d_v + d_h) \circ h(c) = d_v \circ h(c) + d_h \circ h(c) = h \circ d_v(c) + c + h \circ d_h(c) = h \circ (d_v + d_h)(c) + c = c$$

proof

To treat the case of  $\iota_C$  we will use a double spectral sequence argument.

□

### 3.1.2 Model Structure on Module Categories

Let  $A$  be an augmented dg-algebra, then we know that  $Mod^A$  is a model category. By corollary 2.2.4.2 we have a model structure where the fibrations, cofibrations and weak equivalences are given as follows:

- $f \in Ac$  is a weak equivalence if  $f$  is a quasi-isomorphism.
- $f \in Fib$  is a fibration if  $f^\#$  is an epimorphism.
- $f \in Cof$  is a cofibration if it has LLP to acyclic fibrations.

Every object in this category is fibrant as the morphism  $0 : M \rightarrow 0$  is always an epimorphism.

### 3.1.3 Model Structure on Comodule Categories

Unless stated otherwise, for this section we fix  $A$  to be an augmented dg-algebra,  $C$  as a conilpotent dg-coalgebra and  $\tau : C \rightarrow A$  as an acyclic twisting morphism. We endow  $CoMod_{conil}^C$  with three classes of morphisms:

- $f \in Ac$  is a weak equivalence if  $L_\tau f$  is a quasi-isomorphism.
- $f \in Cof$  is a cofibration if  $f^\#$  is a monomorphism.
- $f \in Fib$  is a fibration if it has RLP to acyclic cofibrations.

**Theorem 3.1.8.** *The category  $CoMod_{conil}^C$  with the three classes as above forms a model category. Every object is cofibrant, and those objects which is a direct summand of  $R_\tau M$  for some  $M \in Mod^A$  are fibrant. The adjoint pair  $(L_\tau, R_\tau)$  is a Quillen equivalence.*

We will call this model structure for the canonical model structure on  $CoMod_{conil}^C$ . Under the hypothesis of this theorem, we may observe that every object of  $CoMod_{conil}^C$  are cofibrant. Since every  $M \in Mod^A$  is fibrant, and  $R_\tau$  preserves fibrant objects we know that  $R_\tau M$  is fibrant as well. Clearly every direct summand of  $R_\tau M$  is fibrant. If  $N \in CoMod_{conil}^C$  is fibrant, then it is a direct summand of  $R_\tau L_\tau N$ . This shows that the bifibrant objects of  $CoMod_{conil}^C$  are exactly the thick image of  $R_\tau$ .

To be able to prove this we will need some lemmata. This proof is essentially the same as the case for dg-coalgebras. The main difference is to show independence of the choice of a twisting morphism  $\tau$ . To this end we must establish the relationship between graded quasi-isomorphisms and weak equivalences, as well as a technical lemma.

Recall that given a coaugmented coalgebra  $C$  we have a filtration called the coradical filtration, defined as  $Fr_i C = Ker(\bar{\Delta}_C)^i$ . If  $N$  is a right  $C$ -comodule we may define the coradical filtration of  $N$  as  $Fr_i N = Ker(\bar{\omega}_N^i)$ . This filtration is admissible, meaning it is exhaustive and  $Fr_0 N = 0$ .

**Lemma 3.1.9.** *Let  $C$  be a conilpotent dg-coalgebra,  $M$  and  $N$  be right  $C$ -comodules. Then any graded quasi-isomorphism  $f : M \rightarrow N$  is a weak equivalence.*

*Proof.* This proof is identical to 2.2.5. □

**Lemma 3.1.10.** *Let  $M$  and  $N$  be two objects of  $Mod^A$ . The functor  $R_\tau$  sends a quasi-isomorphism  $f : M \rightarrow N$  to a weak equivalence  $R_\tau f : R_\tau M \rightarrow R_\tau N$ .*

*The unit of the adjunction  $\eta : Id \rightarrow R_\tau L_\tau$  is a pointwise weak equivalence.*

*Proof.*  $R_\tau f$  is a weak equivalence if  $L_\tau R_\tau f$  is a quasi-isomorphism. By naturality of the counit we have the following commutative diagram.

$$\begin{array}{ccc} M & \xleftarrow{\varepsilon_M} & L_\tau R_\tau M \\ \downarrow f & & \downarrow L_\tau R_\tau f \\ N & \xleftarrow{\varepsilon_N} & L_\tau R_\tau N \end{array}$$

By assumption we know that all three of  $f$ ,  $\varepsilon_M$  and  $\varepsilon_N$  are quasi-isomorphisms. It follows by the 2-out-of-3 property that  $L_\tau R_\tau f$  is a quasi-isomorphism as well.

To show that  $\eta : Id \rightarrow L_\tau R_\tau$  is a pointwise weak-equivalence, we must show that  $L_\tau \eta$  is a pointwise quasi-isomorphism. Since  $L_\tau$  is left adjoint to  $R_\tau$  we know that  $\eta$  is split on the image of  $L_\tau$ , i.e.

$$\varepsilon_{L_\tau} \circ L_\tau \eta = id_{L_\tau}$$

Since we know that the natural isomorphisms  $\varepsilon$  and  $id$  are pointwise quasi-isomorphisms, we get by the 2-out-of-3 property that  $L_\tau \eta$  is a pointwise quasi-isomorphism as well. □

**Lemma 3.1.11.** *The functor  $L_\tau$  preserves cofibrations and sends weak-equivalences to quasi-isomorphisms.*

*Proof.* This proof is essentially the same as 2.2.8. □

With the above lemmata we have now established that the adjunction  $(L_\tau, R_\tau)$  forms a Quillen equivalence if  $CoMod^C$  is a model category.

The next lemma is a technical lemma which we need. There will not be given a proof for it, but this is lemma 2.2.2.9 in [12].

**Lemma 3.1.12.** *Let  $M$  be a right  $A$ -module and  $N$  a right  $C$ -comodule. Let  $p : M \rightarrow L_\tau N$  be a fibration of modules. The projection  $j : R_\tau M \prod_{R_\tau L_\tau N} N \rightarrow R_\tau M$  is an acyclic cofibration of comodules.*

*Proof.* This proof is omitted. □

*Proof of 3.1.8.* With the above lemmata established, this proof is identical to the proof of 2.2.10. □

### 3.1.4 Triangulation of Homotopy Categories

In this section we will show that the homotopy categories are triangulated. If we look at the category  $Mod^A$  we will observe that the category  $HoMod^A$  is our beloved derived category  $\mathcal{D}(A)$ . It is not quite the same for the category  $CoMod^C$ . Here we want  $HoCoMod^C$  to be equivalent to the derived category of a ring, so we will see that the derived category is a further localization of  $HoCoMod^C$ .

Furthermore, by employing the theory of triangulated categories we will show that the model structure on  $CoMod^C$  is independent on the choice of acyclic twisting morphism. This breaks down to show that every acyclic twisting morphism induce an equivalence between derived categories, as done by Bernhard Keller in [26].

$Mod^A$  is an abelian category, where we employ the maximal exact structure  $\mathcal{E}'$  consisting of short exact sequences in  $Mod^A$ . This translates to short exact sequences which is short exact in each degree. However, this category also has an exact structure  $\mathcal{E}$  which makes  $Mod^A$  into a Frobenius category, which we will now describe.

Let  $f : M \rightarrow N$ , be a chain map from  $M$  to  $N$ . Then  $\mathcal{E}$  contains a conflation on the form:

$$N \twoheadrightarrow cone(f) \twoheadrightarrow M[1]$$

We define  $\mathcal{E}$  to be the smallest exact structure on  $Mod^A$  which contains every conflation arising from a chain map  $f$ . Observe that these conflations are exactly the short exact sequences of  $Mod^A$  such that they are split when regarded as graded modules, i.e. forgetting the differential. Thus the smallest such  $\mathcal{E}$  is exactly the collection of every conflation arising from a chain map  $f$ .

Recall that an object  $M$  is projective (injective) if the represented functor  $Mod^A(M, \_)$  ( $Mod^A(\_, M)$ ) is exact. For the category  $(Mod^A, \mathcal{E})$

**Proposition 3.1.13.** *Let  $M$  be an object of  $Mod^A$ . The following are equivalent:*

- $M$  is projective
- $M$  is injective
- $M$  is contractible

*Proof.* This is a well known statement from literature. See Krause [27], Happel [28], Buehler [29] or Thorbjørnsen [30] for an account of this result.  $\square$

To see that  $(Mod^A, \mathcal{E})$  has both enough projectives and injectives we consider the following conflation:

$$M \rightarrowtail cone(id_M) \twoheadrightarrow M[1]$$

It is known that the complex  $cone(id_M)$  is contractible for any complex  $M$  (and it is also universal with this property). In this way by letting  $M$  vary we can find an inflation or deflation from the identity cone from or to any complex. This concludes that  $(Mod^A, \mathcal{E})$  is a Frobenius category.

Let  $\underline{Mod}^A$  denote the injectively stable module category. Let  $I(M, N)$  denote the set of chain maps from  $M$  to  $N$  which factors through an injective object. We define the injectively stable category as the quotient of abelian groups  $\underline{Mod}^A(M, N) = Mod^A(M, N)/I(M, N)$ .

**Theorem 3.1.14.** *Suppose that  $(\mathcal{C}, \mathcal{E})$  is a Frobenius category, then the injectively stable category  $\underline{\mathcal{C}}$  is triangulated. The additive auto-equivalence is given by cozyzygy, and the standard triangles is the image of the conflations into the quotient.*

*Proof.* This is also well known in literature. An account for it may also be found in Krause [27], Happel [28], Buehler [29] or Thorbjørnsen [30].  $\square$

We thus obtain a triangulated category  $\underline{Mod}^A$  associated to the Frobenius pair  $(Mod^A, \mathcal{E})$ . This category is commonly denoted as  $K(A)$ , and we will do this as well. Notice that with the structure given by  $\mathcal{E}$ , the cozyzygy is defined by the shift functor  $[1]$ . Every standard triangle is also on the form:

$$M \xrightarrow{f} N \longrightarrow cone(f) \longrightarrow M[1]$$

To define the derived category  $D(A)$  of  $A$  we will consider the localization of  $K(A)$  at the quasi-isomorphisms,  $D(A) = K(A)[Qiso^{-1}]$ . To see that the derived category is still triangulated we may realize it as a Verdier quotient of  $K(A)$ .

**Proposition 3.1.15.** *The derived category of  $A$  is equivalent to the Verdier quotient  $K(A)/Ac$ , where  $Ac$  denotes the image of acyclic objects in  $K(A)$ .*

*Proof.* A proof may be found in Buehler [29] or Thorbjørnsen [30].  $\square$



There is another way of telling the story of the derived category  $D(A)$ . That is to directly localize it at the quasi-isomorphisms. We may directly see that  $D(A) \simeq Mod^A[Qiso^{-1}]$  which we know is  $HoMod^A$  by definition. This gives us our first important identification.

**Theorem 3.1.16.** *The homotopy category of  $Mod^A$  is triangulated, and moreover it is the derived category  $D(A)$ .*

*Proof.* Follows from discussion above. □

The triangulated construction for the category  $hoCoMod^C$  closely resembles that of  $HoMod^A$ . We start by studying the Frobenius pair  $(CoMod^C, \mathcal{E})$ , where  $\mathcal{E}$  is the same exact structure. Notice that this exact structure only takes the underlying category of chain complexes into account, so this follows from the above description.

We define the injectively stable category  $\underline{CoMod}^C = K(C)$  in the same manner. The standard triangles and the additive auto-equivalence stays the same.

At this point things start to differ. The definition for the homotopy category  $HoCoMod^C$  is  $CoMod^C[Ac^{-1}]$ , here  $Ac$  denotes the class of weak equivalences in  $CoMod^C$ . By abuse of notation we also let  $Ac \subset K(C)$  be the collection of objects which are cones of weak equivalences. This subcategory can be characterized by being the preimage of acyclic objects  $Ac \subset K(A)$  along  $L_\tau : CoMod^C \rightarrow Mod^A$ . This identification suffices to show that  $Ac \subset K(C)$  is a triangulated subcategory. In this manner  $HoCoMod^C$  is the category  $K(C)/Ac$ , which is a triangulated category.

**Remark 3.1.17.** We may show that  $Ac \subset K(C)$  is a subcategory of acyclic objects. In this manner we get that  $D(C) \simeq HoCoMod^C[Qiso^{-1}]$ . This is done in Lefevre-Hasegawa [12] as proposition 1.3.5.1 or lemma 2.2.2.11. This follows from the fact that we have an equivalence of categories  $CoMod^C[fQiso^{-1}] \simeq HoCoMod^C$ , here  $fQiso$  means the collection of filtered quasi-isomorphisms. Since every filtered quasi-isomorphism is in fact a quasi-isomorphism by a spectral sequence argument we get the inclusion of triangulated subcategories  $\langle cone(fQiso) \rangle \subseteq \langle cone(Qiso) \rangle \subseteq K(C)$ .

Let  $\tau : C \rightarrow A$  and  $v : C \rightarrow A'$  be two acyclic twisting morphisms. These defines independently two different model structures on  $CoMod^C$  by the adjunctions  $(L_\tau, R_\tau)$  and  $(L_v, R_v)$ . By lemma 3.1.5 we have the identification  $(L_\tau, R_\tau) = (f_{\tau!}, f_\tau^*)(L_{\iota_C}, R_{\iota_C}) = (f_{\tau!}L_{\iota_C}, R_{\iota_C}f_\tau^*)$ , and a likewise for  $v$ . In order to show that  $\tau$  and  $v$  defines equivalent module structures on  $CoMod^C$  it is enough that both define the same structure as  $\iota_C$ . By symmetry it is enough to assume that  $v = \iota_C$ . From lemma 3.1.7 we know that  $\iota_C$  is acyclic, so this technique is well-founded.

Since we already know that  $(L_\tau, R_\tau)$  and  $(L_{\iota_C}, R_{\iota_C})$  are Quillen equivalences it remains to show that  $(f_{\tau!}, f_\tau^*)$  is a Quillen equivalence. This is shown if  $f_\tau^*$  is a right Quillen functor, and that it induces a triangle equivalence between  $D(A)$  and  $D(\Omega C)$ .

We know that  $f_\tau^*$  preserves fibrations (epimorphisms). This is because on morphisms this functor acts as the identity, it only changes the ring action, so epimorphisms stay epimorphisms. It

remains to see that quasi-isomorphisms are preserved. We will show this by identifying the derived categories. This follows the methods given by Keller in [26]. We will however simplify this discussion by restricting our attention solely to dg-algebras.

Let  $A$  be a dg-algebra. We denote  $A$  as a free  $A$ -module as  $\hat{A} = \text{Hom}_A(\_, A)$ .  $\hat{A}$  is free in the enriched sense, i.e.  $\text{Hom}_A^*(\hat{A}, M) \simeq M$ . Recall that  $P$  is projective if it is a direct summand of  $\hat{A}^n$  for some  $n \in \mathbb{N}$ . Given a right bounded complex  $M$ , we know how to construct a projective resolution  $p : pM \rightarrow M$ . Associated to this resolution there is a triangle in  $K(A)$  consisting of the complexes  $M, pM$  and  $aM$ , where  $aM$  is an acyclic complex.

$$M \xrightarrow{p} pM \longrightarrow aM \longrightarrow M[1]$$

In this sense we obtain an identification  $M \simeq pM$  in  $D(A)^-$ . By following Kellers construction we are able to weaken this identification to all of  $D(A)$  by weakening the projective resolution. In Kellers paper he calls these complexes of property (P). We will however refer to them as homotopy projective complexes, since these complexes are built up from projective complexes in a manner respecting homotopy colimits.

**Definition 3.1.18.** Let  $P$  be a complex of  $\text{Mod}^A$ . We say that  $P$  is homotopy projective if there exists a complex  $P'$ , a homotopy equivalence  $P \simeq P'$  and a filtration of  $P'$ .

$$0 = F_0 \subseteq F_1 \subseteq \dots \subseteq F_n \subseteq \dots \subseteq P'$$

The filtration should satisfy these properties:

- F1  $P'$  is the colimit of the filtration.
- F2 Each inclusion  $i_n : F_n \subseteq F_{n+1}$  is split as graded modules.
- F3 The quotient  $F_{n+1}/F_n$  is projective.

**Remark 3.1.19.** The properties F1 and F2 may be reformulated to that  $P$  should be the homotopy colimit of the filtration. Thus there is a canonical triangle in  $K(A)$ :

$$\bigoplus F_n \xrightarrow{\Phi} \bigoplus F_n \longrightarrow P \longrightarrow \bigoplus F_p[1]$$

$\Phi$  is given as the unique morphism which acts as the identity and the inclusion on each summand of  $\bigoplus F_p$ :

$$\Phi_n = \begin{pmatrix} id_{F_n} \\ -i_n \end{pmatrix}$$

In the definition of a homotopy projective complex we have required that each quotient is strictly projective. If this was true, then these objects would be ill-behaved in the homotopy category. We can weaken this assumption to (F3') the quotient  $F_{n+1}/F_n$  is homotopy equivalent to a projective complex.

**Lemma 3.1.20.** *If  $P$  is the colimit of a filtration admitting (F2) and (F3'), then  $P$  is homotopy projective.*

*Proof.* Let  $\{F_n\}$  denote the filtration on  $P$ . To show that  $P$  is homotopy projective is to find a homotopy equivalence  $P'$  such that  $P'$  is the homotopy colimit of a filtration admitting (F3).

Suppose that  $F_{n+1}/F_n \simeq Q_{n+1}$ , where each  $Q_{n+1}$  is projective. We wish to inductively define a filtration  $\{F'_n\}$  which has (F2) and (F3) and a pointwise homotopy equivalence of filtrations  $f : \{F_n\} \rightarrow \{F'_n\}$ . The object  $P'$  is the (homotopy) colimit of the new filtration.

Define  $F'_0 = Q_0$ , and let  $f_0 : F_0 \rightarrow F'_0$  be the projection onto  $Q_0$ . By assumption  $f_0$  is a homotopy equivalence and we have a commutative square where the vertical arrows are homotopy equivalences. Moreover, each horizontal arrow splits as a graded arrow.

$$\begin{array}{ccc} 0 & \xrightarrow{0} & F_0 \\ \downarrow 0 & & \downarrow f_0 \\ 0 & \xrightarrow{0} & Q_0 \end{array}$$

Suppose that we have been able to construct this filtration up to  $F'_p$ . By using our known homotopy equivalences there is an isomorphism of Ext groups:

$$Ext_A(F_p/F_{p-1}, F_{p-1}) \simeq Ext_A(Q_p, F'_{p-1})$$

Given the triangle consisting of  $F_{p-1}$ ,  $F_p$  and  $F_p/F_{p-1}$  there is an associated triangle with the morphisms as follows:

$$\begin{array}{ccccccc} F_{p-1} & \longrightarrow & F_p & \longrightarrow & F_p/F_{p-1} & \longrightarrow & F_{p-1}[1] \\ \downarrow f_{p-1} & & \downarrow & & \downarrow \sim & & \downarrow f_{p-1}[1] \\ F'_{p-1} & \longrightarrow & F'_p & \longrightarrow & Q_p & \longrightarrow & F'_{p-1} \end{array}$$

By the morphism axiom there is a morphism  $f_p : F_p \rightarrow F'_p$  which is also a homotopy equivalence by the 2-out-of-3 property.

This defines a filtration  $\{F'_p\}$ , with (F3) and  $P'$  as its homotopy colimit. To see that  $P$  is homotopy equivalent to  $P'$  we use the maps  $f_p$  constructed to obtain a homotopy equivalence by the morphism axiom and the 2-out-of-3 property.

$$\begin{array}{ccccccc}
\bigoplus F_p & \xrightarrow{\Phi} & \bigoplus F_p & \longrightarrow & P & \longrightarrow & \bigoplus F_p[1] \\
\downarrow \oplus f_p & & \downarrow \oplus f_p & & \downarrow \sim & & \downarrow \oplus f_p[1] \\
\bigoplus F'_p & \xrightarrow{\Phi'} & \bigoplus F'_p & \longrightarrow & P' & \longrightarrow & \bigoplus F'_p[1]
\end{array}$$

□

The projective complexes are the complexes which are generated by the free module  $\hat{A}$  in the sense that they are all in the smallest thick triangulated subcategory of  $K(A)$  containing  $\hat{A}$ . By definition, we may see that the homotopy projective complexes are the complexes in the smallest thick triangulated subcategory of  $K(A)$  which is closed under well-ordered homotopy colimits and contains  $K(A)$ . By devissage we may extend fully-faithfulness of functors on the set  $\{\hat{A}\}$  to the class of homotopy projective objects.

**Lemma 3.1.21** (Devissage). *Let  $F : \mathcal{T} \rightarrow \mathcal{U}$  be a triangulated functor between triangulated categories. Suppose  $S \subseteq \mathcal{T}$  is a class of objects closed under shift, and denote  $\hat{S}$  for the smallest thick triangulated subcategory (closed under well-ordered homotopy colimits). If  $F|_S$  is fully faithful, then  $F|_{\hat{S}}$  is fully faithful as well.*

*Proof.* This is straightforward by using the Yoneda embeddings and 5-lemma. More details may be found in [27]. To get closed under homotopy colimits we also need that  $F$  commutes with infinite direct sums, and that the set  $\{S\}$  only contains small objects. □

**Lemma 3.1.22.** *Suppose we have  $F$  and  $S$  as above. If  $F|_S = 0$ , then it is 0 on all of  $\hat{S}$ .*

*Proof.* The same argument as above, except we have to squeeze out zeros from exact sequences. □

The final ingredient to construct a homotopy projective resolution for our complexes is the acyclic assembly lemma [23].

**Lemma 3.1.23** (acyclic assembly). *Suppose that  $C$  is a double complex of  $R$ -modules. Then  $\text{Tot}^\oplus C$  is acyclic if either:*

- $C$  is a lower half-plane complex with exact rows.
- $C$  is a left half-plane complex with exact columns.

*Proof.* This is proposition 2.7.3 in [23]. We omit the proof as the next proof is in some sense very similar. □

**Corollary 3.1.23.1.** *Suppose that  $C$  is a double complex of  $R$ -modules such that every column is exact and that the kernels along the rows give rise to exact columns, then  $\text{Tot}^\oplus C$  is acyclic.*

*Proof.* We want to realize the images along the rows as the coimage along the horizontal differential. Write  $Z^n(C)$  for the  $n$ -th horizontal kernel and  $B^n(C)$  for the  $n$ -th horizontal image. We have a short exact sequence of complexes:

$$Z^n(C)^* \longrightarrow C^{n,*} \longrightarrow B^n(C)^*$$

Given that  $C^{n,*}$  is acyclic we get that  $Z^n(C)^*$  is acyclic if and only if  $B^n(C)^*$  is acyclic.

Assuming that all of these three constructions are acyclic we make a filtration on  $C$ . Let  $F_n C^{p,*} = C$  if  $p \in [-n, n-1]$ ,  $F_n C^{n,*} = Z^n C$  and  $F_n C^{p,*} = 0$  otherwise.

This filtration is bounded below and exhaustive as colimits commute with colimits.

$$Tot^\oplus C = Tot^\oplus \varinjlim F_n C \simeq \varinjlim Tot^\oplus F_n C$$

We should be a bit careful here as the total complex is not really a coproduct, but since coproducts and cokernels are calculated pointwise we obtain the commutativity.

Now we apply the classical convergence theorem to the filtration to obtain a converging spectral sequence  $EF_2 C \implies H^*(Tot^\oplus C)$ . But since we assume each column to be exact in the filtration, the second page is 0, so  $H^*(Tot^\oplus C) \simeq 0$  as desired.  $\square$

**Theorem 3.1.24.** *Suppose that  $P$  is homotopy projective,  $N$  is acyclic. Then  $K(A)(P, N) \simeq 0$ .*

*Given any module  $M$ , there is a homotopy projective object  $pM$  and an acyclic object  $aM$  giving rise to a triangle in  $K(A)$ .*

$$pM \longrightarrow M \longrightarrow aM \longrightarrow pM[1]$$

*Proof.* We assume that  $P \simeq \hat{A}$ . By a devissage argument we may extend the isomorphism to all homotopy projective  $P$ .

$$K(A)(\hat{A}, N) \simeq H^0 Hom_A^*(\hat{A}, N) \simeq H^0 N \simeq 0$$

We want to construct two complexes  $pM$  and  $aM$  by taking total complexes. We show that  $aM$  is acyclic by using 3.1.23.1. To use it we will construct an exact sequence of complexes satisfying the assumptions. Described by MacLane [31] there is an exact structure  $\mathcal{E}$  on  $Mod^R$  such that the collections on conflation are the short exact sequences such that the kernel functor is exact.

$$L \xrightarrow{f} M \xrightarrow{g} N$$

$$Z^*L \xrightarrow{Z^*f} Z^*M \xrightarrow{Z^*g} Z^*N$$

Since limits commute with limits, the kernel functor preserves any limit. Thus the kernel is left exact, and its only obstruction for exactness is to preserve cokernels. We may thus characterize the conflations by inflations and deflations, which are monomorphisms and epimorphisms which are preserved by the kernel functor. Mac Lane calls these deflations for proper epimorphisms instead.

Mac Lane also shows that there are enough  $\mathcal{E}$ -projectives with this exact structure. We want to construct  $\mathcal{E}$ -projectives be on the form of homotopy projective complexes.  $\hat{A}[-n]$  is  $\mathcal{E}$ -projective by the following isomorphism.

$$Hom_A^\bullet(\hat{A}[-n], M) \simeq Z^0 Hom_A^*(\hat{A}, M[n]) \simeq M^n$$

Define the trivialization  $trivM$  of  $M$  be the underlying graded module  $M$  endowed with a trivial differential. This trivial differential is in some sense the inclusion of graded modules into chain complexes. Thus we have the following isomorphism on hom-sets:

$$Hom_A^\bullet(trivM, trivN) \simeq Hom_A^*(M, N)$$

$triv$  is then well-defined as a functor as every morphism between chain complexes uniquely defines a morphism between their trivializations. By using the isomorphisms from Keller [26] 2.2. we get that:

$$Hom_A^\bullet(cone(id_{triv\hat{A}}), M) \simeq Hom_A^\bullet(cone(id_{triv\hat{A}[-1]})[1], M)$$

$$\simeq Hom_A^*(triv\hat{A}, trivM[-1])^0 \simeq Hom_A^*(\hat{A}, M)^{-1} \simeq M^{-1}$$

This shows that if  $P$  is homotopy projective, then  $P$  and  $cone(id_{trivP})$  are  $\mathcal{E}$ -projective. To see that there are enough  $\mathcal{E}$ -projectives pick an arbitrary module  $M$ . Since we know that there are enough projectives, let  $P$  be a projective such that there is an epimorphism  $p : P \rightarrow M$ . We don't know if this morphism is a deflation, so pick another projective  $Q$  such that there is an epimorphism  $q : Q \rightarrow Z^*M$ . Since  $Z^*M$  has a trivial differential we know that  $d_{QQ} = 0$ . Thus this morphism extends to  $q' = [q \ 0] : cone(id_{trivQ}) \rightarrow M$  such that  $Z^*q'$  is an epimorphism. The morphism  $[p \ q'] : P \oplus cone(id_{trivQ}) \rightarrow M$  is thus a deflation.  $P' = P \oplus cone(id_{trivQ})$  shows that we have enough projectives. Moreover every  $cone(id_{trivQ})$  has homotopy type 0, so  $P' \simeq P$  in  $K(A)$ .

Since we have enough  $\mathcal{E}$ -projective, we may construct a  $\mathcal{E}$ -projective resolution  $P'^{*,*}$  of  $M$  in the standard way. See Keller [32] for details. Such resolutions are then double complexes, and the augmented resolution below is  $\mathcal{E}$ -acyclic.

$$\dots \longrightarrow P'_1 \longrightarrow P'_0 \twoheadrightarrow M \xrightarrow{0} 0$$

Having an  $\mathcal{E}$ -acyclic resolution means that each row is exact, and taking kernels along the columns preserve exactness of the rows.

Denote the augmentation of  $P'^{*,*}$  by  $m : P'^{*,*} \rightarrow M$ . We define the complexes  $pM = Tot^\oplus(P'^{*,*})$  and  $aM = Tot^\oplus(cone(m))$ .

$pM$  carries a natural filtration  $F_n pM$  from the double complex structure. Let  $F_n pM$  be the truncated complex:

$$\dots \longrightarrow 0 \longrightarrow P'^{n,*} \longrightarrow \dots \longrightarrow P'^{1,*} \longrightarrow P'^{0,*} \longrightarrow 0 \longrightarrow \dots$$

The filtration  $F_n pM$  satisfies F1 and F2 by construction. The quotients  $F_{n+1} pM / F_n pM \simeq P'_n$  which is homotopy equivalent to a projective. By lemma 3.1.20  $pM$  is homotopy projective.

The complex  $cone(m)$  satisfies the conditions for 3.1.23.1, thus  $aM$  is acyclic. Thus we have a triangle in  $K(A)$  as desired.  $\square$

**Corollary 3.1.24.1.** *Let  $M$  be an arbitrary module. If  $P$  is homotopy projective, then  $K(A)(P, M) \simeq K(A)(P, pM)$ . If  $N$  is acyclic, then  $K(A)(M, N) \simeq (aM, N)$ .*

$a$  and  $p$  are well-defined functors which commutes with infinite direct sums.

**Corollary 3.1.24.2.** *Let  $\{\hat{A}\}$  denote the smallest thick triangulated subcategory of  $D(A)$  which is closed under homotopy colimits. Then  $D(A) \simeq \{\hat{A}\}$ .*

**Corollary 3.1.24.3.** *Suppose that  $f : A \rightarrow B$  is a dg-ring homomorphism and a quasi-isomorphism between dg-algebras, then  $D(A) \simeq D(B)$ .*

*Proof.*  $f$  endows  $B$  with both a left and right  $A$ -module structure. We will think of  $\hat{B}$  as a left  $A$ -module and right  $B$  module. There is then a natural hom-tensor adjunction between the differential graded enriched categories.

$$\begin{array}{ccc} & \xrightarrow{H} & \\ Diff(B) & \top & Diff(A) \\ & \xleftarrow{T} & \end{array}$$

We define  $T = \_ \otimes_A \hat{B}$  and  $H = Diff(B)(B, \_)$ . We see that  $Diff(B)(T\hat{A}, M) \simeq Diff(A)(\hat{A}, HM) \simeq HM \simeq Diff(B)(\hat{B}, M)$ . Thus  $T\hat{A} \simeq \hat{B}$ , and the morphism  $T : Diff(A)(\hat{A}, \hat{A}) \rightarrow Diff(B)(\hat{B}, \hat{B})$  is given by  $f$ . Since we assume  $f$  to be a quasi-isomorphism, it follows that  $\mathbb{L}T : D(A) \rightarrow D(B)$  is fully faithful on the set  $\{\hat{A}\}$ .

By devissage the functor  $\mathbb{L}T$  is fully-faithful on all of  $D(A)$ , since  $D(A)$  is generated by  $\hat{A}$ . Since  $T$  hits all of  $D(B)$ 's generators,  $\mathbb{L}T$  is essentially surjective as well.  $\square$

*Remark 3.1.25.* We have ignored smallness conditions for objects. This technique does not always work, since it depends on some unstated isomorphisms which we have used. We have these since the objects  $\hat{A}$  and  $\hat{B}$  are small. This detail is given more care in Keller [26].

With this result we are able to show that  $HoMod^A$  and  $HoMod^{\Omega C}$  are equivalent. Since we assumed the morphism  $\tau : C \rightarrow A$  to be acyclic, we would expect the morphism  $f_\tau^* : \Omega C \rightarrow A$  to be a quasi-isomorphism. If this is the case, we know that  $D(\Omega C) \simeq D(A)$ .

### 3.1.5 The Fundamental Theorem of Twisting Morphisms

In this section we aim to finish what we started the previous section. We will prove a characterization for the acyclic twisting morphisms.

**Theorem 3.1.26** (Fundamental Theorem of Twisting Morphisms). *Let  $\tau : C \rightarrow A$  be a twisting morphism between augmented objects. The following are equivalent:*

1.  $\tau$  is acyclic, i.e. the natural transformation  $\varepsilon : L_\tau R_\tau \implies Id_{Mod^A}$  is a pointwise quasi-isomorphism.
2. The unit transformation  $\eta : Id_{CoMod^C} \implies R_\tau L_\tau$  is a pointwise weak equivalence.
3. The counit at  $A$  is a quasi-isomorphism, i.e.  $\varepsilon_A : L_\tau R_\tau A \rightarrow A$  is a quasi-isomorphism.
4. The unit at  $\mathbb{K}$  is a weak-equivalence, i.e. the algebra unit  $v_A$  and coaugmentation  $v_C$  assemble into a weak-equivalence:  $v_A \otimes v_C : \mathbb{K} \rightarrow A \otimes_\tau C$ .
5. The morphism of algebras  $f_\tau : \Omega C \rightarrow A$  is a quasi-isomorphism.
6. The morphism of coalgebras  $g_\tau : C \rightarrow BA$  is a weak-equivalence.

*Proof.* Notice that 1. is equivalent to 2. since  $\mathbb{L}L$  and  $\mathbb{R}R$  are quasi-inverse. 3. is a special case of 1. and 4. is a special case of 2. Observe that 5. and 6. are equivalent since the cobar-bar-adjunction is a Quillen equivalence, 2.2.10.1.

We show 3.  $\implies$  1. Let  $\mathcal{T} \subseteq D(A)$  be the full subcategory consisting of objects  $M$  where  $\varepsilon_M$  is a quasi-isomorphism. This subcategory is by assumption non-empty and contains  $A$ . By the 5-lemma, making triangles (and smallness of  $A$ ), this subcategory contains the smallest thick triangulated subcategory closed under homotopy colimits which contains  $A$ . We know this to be all of  $D(A)$ .

To show 4. implies 5. we consider the twisting morphism  $\iota_C$ . Since  $\iota_C$  is acyclic we know that the counit at  $A$  is a quasi-isomorphism.

$$L_{\iota_C} R_{\iota_C} f_\tau^* A \rightarrow f_\tau^* A$$



By assumption the unit morphism  $\eta_{\mathbb{K}} : \mathbb{K} \rightarrow A \otimes_{\tau} C$  is a weak equivalence, so the morphism  $L_{\iota_C} \eta_{\mathbb{K}} : \Omega C \rightarrow L_{\iota_C} R_{\tau} A = L_{\iota_C} R_{\iota_C} f_{\tau}^* A$  is a quasi-isomorphism. Let  $\varepsilon'$  denote the counit of  $L_{\iota_C} \dashv R_{\iota_C}$ , then we see that  $f_{\tau} = \varepsilon'_A \circ L_{\iota_C} \eta_{\mathbb{K}}$ , so  $f_{\tau}$  is a quasi-isomorphism by the 2-out-of-3 property.

It remains to show that 5. implies 1. Let the counit of  $f_{\tau*} \dashv f_{\tau}^*$  be denoted as  $\tilde{\varepsilon}$ . Since  $f_{\tau}$  is a quasi-isomorphism,  $f_{\tau}^*$  descends to an equivalence between the derived categories 3.1.24.3. Thus  $\tilde{\varepsilon} : f_{\tau!} f_{\tau}^* \Rightarrow Id$  is a pointwise quasi-isomorphism. Observe that the counit factors as

$$\varepsilon = \tilde{\varepsilon} \circ f_{\tau!} \varepsilon'_{f_{\tau}^*}$$

By the 2-out-of-3 property it follows that  $\varepsilon$  is a quasi-isomorphism.  $\square$

**Corollary 3.1.26.1.** *There is one canonical model structure on  $CoMod^C$  defined by the acyclic twisting morphisms  $\tau : C \rightarrow A$ , for any algebra  $A$ . I.e. each acyclic twisting morphism defines the same model structure for  $CoMod^C$ .*

*Proof.* Apply the fundamental theorem of twisting morphisms to the discussion of the last section.  $\square$

## 3.2 Polydules

### 3.2.1 The Bar Construction

In section 1.3 we saw that we could extend the domain of the bar construction to obtain an equivalence of categories. This converse led us to the definition of an  $A_{\infty}$ -algebra, as well as recognizing them as almost free dg-coalgebras. By employing the adjunction  $(L_{\tau}, R_{\tau}) : CoMod^C \rightleftarrows Mod^A$  we will do something similar for modules.

Let  $A$  be an augmented algebra. The bar construction of  $A$  gives us a universal adjunction  $(L_{\pi_A}, R_{\pi_A}) : CoMod^{BA} \rightleftarrows Mod^A$ . We will call  $R_{\pi_A}(-[1]) = -[1] \otimes_{\pi_A} BA$  for  $B_A$ , the bar construction on  $Mod^A$ . In this manner every  $A$ -module  $M$  gives rise to an almost free  $BA$ -comodule  $B_A M$ , but does the converse of this construction works?

Let us first look at what  $B_A$  does to a module  $M$ .  $B_A M$  is the dg-comodule which as a graded comodule is the free comodule  $M[1] \otimes BA$ . The differential of  $B_A M$  is given by the  $A$ -module structure of  $M$ . That is, every elementary element  $m'$  of  $B_A M$  is an element of  $M$  together with a finite string of elements of  $A$ .

$$m' = \omega m \otimes (\omega a_1 \otimes \dots \otimes \omega a_n)$$

The differential acts on  $m'$  by using the differential of  $d_{M[1] \otimes BA}$  and multiplication from the right.

$$d_{B_A M}(m') = d_{M[1] \otimes BA}(m') + (-1)^{|m|+|a|} \omega(m \cdot a_1) \otimes (\omega a_2 \otimes \dots \otimes \omega a_n)$$

By using delooping, we see that in turn that  $d_{B_AM}$  defines an  $A$ -module structure for  $M$ . We may decompose  $B_AM$  as:

$$B_AM = M[1] \oplus M[1] \otimes \bar{A} \oplus M[1] \otimes \bar{A}^{\otimes 2} \oplus \dots$$

Let  $\pi_M : R_{\pi_A} M \rightarrow M$  be the map which kills anything not on the form  $m$ . We denote  $(d_{B_AM})_i$  by  $d_{B_AM} \circ \iota_i$ , where  $\iota_i : M[-1] \otimes \bar{A}^{\otimes i-1} \hookrightarrow B_AM$ . Proposition 1.1.32 tells us that we may recover the structure of  $M$  from the differential  $d_{B_AM}$ . This is done by conjugating the components of  $d_{B_AM}$  with desuspension and applying projections appropriately. We recover the maps as:

1. The differential of  $M$  is  $d_M = s \circ \pi_{M[1]} \circ (d_{B_AM})_1 \omega$
2. The right multiplication from  $A$  is  $\mu_M = s \circ \pi_{M[-1]} \circ (d_{B_AM})_2 \circ \omega^{\otimes 2}$
3. For  $i \geq 3$  we have  $0 = s \circ \pi_{M[1]} \circ (d_{B_AM})_i \circ \omega^{\otimes i}$

Now, let  $\tilde{N}$  be an almost free  $BA$ -comodule. That is,  $\tilde{N} = N[1] \otimes BA$  as a graded comodule. We would now wish for that  $N$  carries an  $A$ -module structure. Unfortunately we are not that lucky, however, this defines a notion of  $A_\infty$ -module to the algebra  $A$ . If we try to recover the same structure we obtain the following structure morphisms for  $N$ :

$$\text{A differential of degree 1: } m_1 = d_N = s \circ \pi_N(d_{\tilde{N}})_1 \circ \omega$$

$$\text{A 2-ary operation of degree 0: } m_2 = s \circ \pi_N(d_{\tilde{N}})_2 \circ \omega^{\otimes 2}$$

$$\text{A 3-ary operation of degree } -1: m_3 = s \circ \pi_N(d_{\tilde{N}})_3 \circ \omega^{\otimes 3}$$

$$\text{A 4-ary operation of degree } -2: \dots$$

Let  $\tilde{m}_i$  be the looped versions of the  $m_i$ s. Then the sum  $\sum \tilde{m}_i : \tilde{N} \rightarrow N$  extends to  $d_{B_AM}$ . Since  $d_{B_AM}^2 = 0$  we get the relations  $(rel_n)$  defined in section 1.3 imposed on the morphisms  $m_i$ .

To summarize the datum of  $M[1] \otimes BA$  as a  $BA$ -comodule is equivalent to a chain complex  $M$  having maps

$$m_i : M \otimes \bar{A}^{\otimes i-1} \rightarrow M$$

of degree  $2 - i$  for any  $i \geq 2$ . The maps should satisfy the relations:

$$(rel_n) \quad \partial(m_n) = - \sum_{\substack{n=p+q+r \\ k=p+1+r \\ k>1, q>1}} (-1)^{pq+r} m_k \circ_{p+1} m_q$$

This gives  $M$  the structure of an  $A$ -module, where associativity is only well-defined up to strong homotopy. In other words,  $m_3$  is a homotopy for the associator for  $m_2$ ,  $m_4$  is a homotopy for  $m_3$ s associator and so on. Following Lefevre-Hasegawa [12], we call the chain-complex  $M$  an  $A$ -polydule, given it has maps  $m_i$  as above.

We have defined the objects of a category  $Mod_\infty^A$ . Our goal is to have that the converse bar construction defines an equivalence of categories, i.e.  $B_A$  extends to a functor  $B_A : Mod_\infty^A \rightarrow$

$CoMod^{BA}$  is fully-faithful. This makes sense as every  $A$ -module  $M$  is a non-full  $A$ -polydule by letting  $m_1 = d_M$ ,  $m_2 = \mu_M$  and  $m_i = 0$  for any  $i \geq 3$ .

Since we say that  $B_A$  is fully-faithful any  $\infty$ -morphisms between polydules is determined by a morphism of almost free  $BA$ -comodules. Let  $f : M \rightsquigarrow N$  be an  $\infty$ -morphism, then  $B_A f : B_A M \rightarrow B_A N$  is the associated morphism of dg-comodules. By cofreeness, proposition 1.1.27, we see that  $Bf$  is uniquely determined by morphisms  $f_i : M \otimes \bar{A}^{\otimes i-1} \rightarrow N$  of degree  $1-i$ . Since we know that  $\partial Bf = 0$ , we obtain the relations:

$$(rel_n) \quad \sum_{p+q+r=n} (-1)^{pq+r} f_{p+1+r} \circ_{p+1} m_q^M = \sum_{p+q=n} m_{p+1}^N \circ_1 f_q$$

If  $f : M \rightsquigarrow N$  and  $g : N \rightsquigarrow P$ , then their composition is:

$$(gf)_n = \sum_{p+q=n} g_{p+1} \circ_1 f_q$$

This defines the morphisms of  $Mod_{\infty}^A$ . By definition  $B_A : Mod_{\infty}^A \rightarrow CoMod^{BA}$  is fully-faithful. An  $\infty$ -morphism  $f$  is called strict if  $f_i = 0$  for any  $i \geq 2$ . We denote the restriction of  $Mod_{\infty}^A$  to strict  $\infty$ -morphisms as  $Mod_{\infty,strict}^A$ . Observe that we obtain an equivalence of categories  $Mod^A \simeq Mod_{\infty,strict}^A$ .

We will give some examples of  $A$ -polydules given an augmented algebra  $A$ .

### 3.2.2 Polydules of SHA-algebras

In the last section we developed the notion of a polydule to an augmented/unital algebra. By applying the converse of the bar construction, we are able to extend this notion to  $A_{\infty}$ -algebras.

Suppose that  $A$  is an  $A_{\infty}$ -algebra. By the bar construction,  $BA$  is an almost free coalgebra. In the same manner, we may consider the almost free dg-coalgebras of  $CoMod^{BA}$ . This is again the collections of comodules of the form  $M' = M[1] \otimes BA$ . Since there is no obstruction to the above arguments the differential  $d_{M'}$  is determined by a collection of morphisms  $m_n^M : M \otimes A^{\otimes n-1} \rightarrow M$  satisfying  $(rel_n)$ . Moreover, an  $\infty$ -morphism  $f : M \rightsquigarrow N$  is a collection of morphisms  $f_n : M \otimes A^{\otimes n-1} \rightarrow N$  satisfying  $(rel_n)$ .

**Definition 3.2.1.** Let  $A$  be an  $A_{\infty}$ -algebra. The category  $Mod_{\infty}^A$  has  $A$ -polydules as objects and  $\infty$ -morphisms as morphisms.

The quasi-isomorphisms in  $Mod_{\infty}^A$  are the  $\infty$ -morphisms  $f$  such that  $f_1$  is a quasi-isomorphism.

*Remark 3.2.2.* The isomorphisms of  $Mod_{\infty}^A$  are the  $\infty$ -morphisms  $f$  where  $f_1$  is an isomorphism.

We say that an  $\infty$ -morphism is strict if  $f_i = 0$  for any  $i \geq 2$ . The category  $Mod_{\infty,strict}^A$  is the non-full subcategory of  $Mod_{\infty}^A$  restricted to strict  $\infty$ -morphisms.

We may also lift homotopies between almost free  $BA$ -comodules and  $A$ -polydules. A homotopy  $B_A h : B_A M \rightarrow B_A M$  is a morphism of degree  $-1$ . Thus the collection  $h_n : M \otimes A^{\otimes n-1} \rightarrow N$  has morphisms of degree  $-i$ . Moreover,  $h : M \rightsquigarrow N$  defines a homotopy of  $f, g : M \rightsquigarrow N$  if we have

$$f_n - g_n = \sum_{p+q} (-1)^p m_{p+1}^N \circ_1 h_q - \sum_{p+q+r=n} (-1)^{pq+r} h_{p+1} \circ_{p+1} m_q^M$$

Suppose now that  $A$  is instead a strictly unital  $A_\infty$ -algebra (1.3.12). We may define strictly unital  $A$ -polydules as an  $A$ -polydule  $M$  such that

$$\begin{aligned} m_2^M \circ (id_M \otimes v_A) &= id_M \\ \forall i \geq 3 \quad m_i^M \circ (id_M \otimes \dots \otimes v_A \otimes \dots \otimes id_A) &= 0 \end{aligned}$$

An  $\infty$ -morphism  $f : M \rightsquigarrow N$  is strictly unital if

$$\forall i \geq 2 \quad f_i(id_M \otimes \dots \otimes v_A \otimes \dots \otimes id_A) = 0$$

This definition also extends to homotopies. We may then define the categories of strictly unital polydules  $iMod_\infty^A$  and  $iMod_{\infty,strict}^A$ .

Given an augmented  $A_\infty$ -algebra  $A$  (1.3.13) we obtain an equivalence of categories. Recall that the categories  $Alg_\infty$  and  $Alg_{\infty,+}$  were equivalent by taking the kernel of augmentation and applying the free augmentation as its quasi-inverse. In the same manner, given a strictly unital  $A$ -polydule  $M$ , then it defines a strictly unital  $\bar{A}$ -polydule  $\bar{M}$  by restricting the structure maps to  $\bar{A}^{\otimes n}$ . This defines an equivalence of categories.

$$\begin{array}{ccc} & \xrightarrow{-} & \\ iMod_\infty^A & & Mod_\infty^{\bar{A}} \\ & \xleftarrow{+} & \end{array}$$

We may call its quasi-inverse for the free strict unitization. This takes an  $\bar{A}$ -polydule  $M$  and turns it into a strictly unital  $A$ -polydule by defining the structure morphism as 0 on the unit.

We will for now restrict our attention to augmented  $A_\infty$ -algebras. The reason for this is that if  $A$  is an arbitrary  $A_\infty$ -algebra, then studying  $Mod_\infty^A$  would be the same as studying  $iMod_\infty^{A^+}$ . We extend the bar construction along this equivalence to a fully faithful functor  $B_A : iMod_\infty^A \rightarrow CoMod^{BA}$ .

### 3.2.3 Universal Enveloping Algebra

Given any augmented  $A_\infty$ -algebra  $A$ , there is a universal enveloping algebra  $UA$ . This algebra is universal in the sense that given any algebra  $A'$  and an  $\infty$ -morphism  $B \rightarrow A$ , then this factors through  $UA$  by an algebra map  $B \rightarrow UA$ . By the cobar-bar adjunction there is really only one way to define this algebra.

**Definition 3.2.3.** Let  $A$  be an  $A_\infty$ -algebra. The universal enveloping algebra is the algebra defined by  $\Omega BA$ .

*Remark 3.2.4.* In this definition we have used the extended bar construction to  $A_\infty$ -algebras and the cobar construction on dg-coalgebras.

**Lemma 3.2.5.** *There is an isomorphism of categories  $i : \text{Mod}^{UA} \rightarrow i\text{Mod}_{\infty, \text{strict}}^{BA}$  given by delooping.*

*Proof.* This is immediate by the definition of a  $UA$ -module. To have a  $UA$ -module  $SM$  we must have structure maps  $m_i^M : M \otimes A^{\otimes i-1} \rightarrow M$  of degree  $2 - i$  for any  $i \geq 2$ . Unwinding this definition and using the adjunction data establishes this isomorphism.  $\square$

### 3.2.4 The Derived Category $D_\infty A$ of Augmented $A_\infty$ Algebras

In this section we wish to define the derived category of strictly unitary polydules to an augmented  $A_\infty$ -algebra. If  $[Qis]$  denote the class of  $\infty$ -quasi-isomorphisms, we want the derived category to be the localization at  $\infty$ -quasi-isomorphisms, e.g.

$$\mathcal{D}_\infty A = i\text{Mod}_\infty^A[Qis^{-1}].$$

Like in the case of algebras, we may understand the quasi-isomorphisms better. The category  $i\text{Mod}_\infty^A$  is not complete, but we may in the same sense give it a model structure without limits. Within this structure we already know that every object is cofibrant, and the goal is to show that every object is fibrant as well. This will allow us to lift every  $\infty$ -quasi-isomorphism to a homotopy equivalence. With this we may see that the localization from  $K_\infty A \rightarrow D_\infty A$  is given by the identity.

Within the category  $i\text{Mod}_\infty^A$  we define three classes of morphisms:

- $f \in Ac$  is a weak equivalence if  $f_1$  is a quasi-isomorphism.
- $f \in Cof$  is a cofibration if  $f_1$  is a monomorphism.
- $f \in Fib$  is a fibration if  $f_1$  is an epimorphism.

**Theorem 3.2.6.** *The category  $i\text{Mod}_\infty^A$  is a model category without enough limits. Moreover, every object are bifibrant.*

*Proof.* This is more or less identical to the proof of 2.3.1.  $\square$

**Corollary 3.2.6.1.** *Homotopy equivalence defined in  $i\text{Mod}_\infty^A$  is an equivalence relation, and every  $\infty$ -quasi-isomorphism is a homotopy equivalence.*

If  $A$  is an ordinary associative algebra, then  $Mod^A$  fully-faithfully embeds into  $iMod_\infty^A$ . This inclusion defines an equivalence:

$$DA \simeq Mod^A / \sim_\infty$$

$iMod_\infty^A$  admits a "model" structure, but we want this model structure to respect the model structure of the category  $CoMod_{conil}^{BA}$ . In other words we want the functor  $B_A : iMod_\infty^A \rightarrow CoMod_{conil}^{BA}$  to preserve and reflect the model structure of both categories.

**Lemma 3.2.7.** *Let  $M$  be an object of  $iMod_\infty^A$ . The unit  $BM \rightarrow RLBM$  induces a quasi-isomorphism on the primitives.*

*Proof.* Fix this proof by using a double spectral sequence argument. □

**Proposition 3.2.8.** *Let  $M$  and  $M'$  be objects of  $iMod_\infty^A$ , together with an  $\infty$ -morphism  $f : M \rightarrow M'$ .*

- *$f$  is an  $\infty$ -quasi-isomorphism if and only if  $Bf$  is a weak equivalence.*
- *$f$  is a fibration if and only if  $Bf$  is a fibration.*
- *$f$  is a cofibration if and only if  $Bf$  is a cofibration.*

*Proof.* Recall from theorem 3.1.8 that the morphism  $\iota_{BA} : BA \rightarrow UA$  is an acyclic twisting morphism. Thus the adjoint pair  $(L_{\iota_{BA}}, R_{\iota_{BA}})$  defines a Quillen equivalence.

We show only the first bullet point. The last two are identical to the proof of 2.2.15.

If  $f_1$  is a quasi-isomorphism, then  $Bf$  is a filtered quasi-isomorphism by definition. So suppose that  $Bf$  is a weak equivalence instead. The unit transformation gives us a natural square.

$$\begin{array}{ccc} BM & \longrightarrow & RLBM \\ \downarrow Bf & & \downarrow RLBf \\ BM' & \longrightarrow & RLBM' \end{array}$$

In this case  $R = Bi$ , so this diagram is in the image of  $B$ . We pull it back by functoriality.

$$\begin{array}{ccc} M & \longrightarrow & iLBM \\ \downarrow f & & \downarrow iLBf \\ M' & \longrightarrow & iLBM' \end{array}$$

Since  $Bf$  is a weak equivalence, we know that  $iLBf$  is an  $\infty$ -quasi-isomorphism.

By the above proposition the horizontal maps are  $\infty$ -quasi-isomorphisms as well.  $\square$

Associated to each augmented  $A_\infty$ -algebra there is also a homotopy category. Since homotopy equivalence  $\sim_\infty$  in  $iMod_\infty^A$  defines a congruence relation we may construct the homotopy category  $K_\infty A$ .

**Corollary 3.2.8.1.** *The localization  $K_\infty A \rightarrow D_\infty A$  is given by the identity. Moreover,  $K_\infty A = D_\infty A$ .*

*Remark 3.2.9.* The name homotopy category comes from homological algebra and has a priori nothing to do with the homotopy category  $Ho(iMod_\infty^A)$ . However, in this particular case these naming conventions coincide.

**Lemma 3.2.10.** *The composition  $J : Mod^{UA} \rightarrow iMod_{\infty,strict}^A \rightarrow iMod_\infty^A$  given by  $J = \iota \circ i$ , the equivalence then non-full inclusion induces an equivalence of categories:*

$$DUA \simeq D_\infty A.$$

*Proof.* Consider the commutative square:

$$\begin{array}{ccc} Mod^{UA} & \xrightarrow{i} & iMod_{\infty,strict}^A \\ \downarrow R_{\iota_{BA}} & & \downarrow \iota \\ CoMod^{BA} & \xleftarrow{B} & iMod_\infty^A \end{array}$$

Since the three functors  $R_{\iota_{BA}}$ ,  $i$  and  $B$  all induces equivalences on the derived categories, then  $\iota$  has to as well.  $\square$

To summarize, we have established an equivalence between 4 different categories:

- $D_\infty A$ , derived category of  $A$
- $iMod_{\infty,strict}^A[Qis^{-1}]$ , derived category of  $A$  with only strict morphisms
- $DBA$ , derived category of  $BA$  as a dg-coalgebra
- $DUA$ , derived category of universal enveloping algebra

In this sense, we may also see that in the derived category, all of the higher homotopic data of each morphism have been collapsed by the homotopy.

The triangulated structure on  $D_\infty A$  may be lifted along these equivalences making them triangulated as well. Note that  $R_{\iota_{BA}}$  is already triangulated, and forcing there is only one way of forcing the triangulated structure on  $iMod_\infty^A$ . Since  $iMod_\infty^A$  isn't complete it isn't easy to obtain

a description of the triangles along any  $\infty$ -morphism  $f$ . However, this problem does not appear in  $iMod_{\infty,strict}^A$ , so one should think of only strict morphisms instead, but in this case we are already working in the category  $Mod^{UA}$ .

### 3.3 The Derived Category $D_{\infty}A$



# Bibliography

- [1] J. D. Stasheff, "Homotopy associativity of h-spaces. i," *Transactions of the American Mathematical Society*, vol. 108, pp. 275–292, 1963.
- [2] J. D. Stasheff, "Homotopy associativity of h-spaces. ii," *Transactions of the American Mathematical Society*, vol. 108, pp. 293–312, 1963.
- [3] J.-L. Loday and B. Vallette, *Algebraic Operads*. Springer Verlag, 2012.
- [4] G. M. Kelly, "Basic concepts of enriched category theory," *Elements of  $\infty$ -Category Theory*, 2005.
- [5] L. W. Tu, "An introduction to manifolds," 2007.
- [6] S. Eilenberg and S. Mac Lane, "On the groups  $h((\pi, n), i)$ ," *Annals of Mathematics*, vol. 58, pp. 55–106, 1953.
- [7] J. Adams, "On the cobar construction," *PNAS*, vol. 42, pp. 409–412, 1956.
- [8] S. Eilenberg and J. C. Moore, "Homology and fibrations. i. coalgebras, cotensor product and its derived functors," *Commentarii mathematici Helvetici*, vol. 40, pp. 199–236, 1965/66.
- [9] E. Riehl, *Categorical Homotopy Theory*. Cambridge University Press, 2014.
- [10] S. MacLane, *Categories for the working mathematician*, ser. Graduate Texts in Mathematics, Vol. 5. Springer-Verlag, New York-Berlin, 1971, pp. ix+262.
- [11] D. -M. Lu, J. H. Palmieri, Q. -S. Wu, and J. J. Zhang, *A-infinity structure on ext-algebras*, 2006. DOI: 10.48550/ARXIV.MATH/0606144. [Online]. Available: <https://arxiv.org/abs/math/0606144>.
- [12] K. Lefevre-Hasegawa, "Sur les a [infini]-catégories," *arXiv: Category Theory*, 2003.
- [13] M. Hovey, *Model Categories*. American Mathematical Society, 1999.
- [14] D. G. Quillen, *Homotopical Algebra*, A. Dold, Heidelberg, and B. Eckmann, Eds. Springer-Verlag, 1967.
- [15] W. G. Dwyer and J. Spalinsky, "Homotopy theories and model categories," in *Handbook of Algebraic Topology*, I. M. James, Ed. Elsevier Science, 1995, ch. 2, pp. 73–126.
- [16] E. Riehl, *Category Theory in Context*. Dover Publications, 2016.

- [17] P. Gabriel and M. Zisman, *Calculus of Fractions and Homotopy Theory*. Springer-Verlag, 1967, pp. 6–20.
- [18] H. J. Munkholm, “Dga algebras as a quillen model category and relations to shm maps,” *Journal of Pure and Applied Algebra*, vol. 13, pp. 221–232, 1978.
- [19] A. K. Bousfield and V. K. A. M. Gugenheim, “On pl de rham theory and rational homotopy type,” *Memoirs of the American Mathematical Society*, vol. 8, no. 179, 1976.
- [20] J. F. Jardine, “A closed model structure for differential graded algebras,” *Fields Institute Communications*, vol. 17, pp. 55–58, 1997.
- [21] V. Hinich, “Homology algebra of homotopy algebras,” *Communications in Algebra*, vol. 25(10), pp. 2391–3323, 1997.
- [22] V. Hinich, “Dg coalgebras as formal stacks,” *Journal of Pure and Applied Algebra*, vol. 162, no. 2, pp. 209–250, 2001, ISSN: 0022-4049. DOI: [https://doi.org/10.1016/S0022-4049\(00\)00121-3](https://doi.org/10.1016/S0022-4049(00)00121-3). [Online]. Available: <https://www.sciencedirect.com/science/article/pii/S0022404900001213>.
- [23] C. A. Weibel, *An Introduction to Homological Algebra*, ser. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1994. DOI: 10.1017/CB09781139644136.
- [24] B. Keller, *Corrections to 'sur les a-infini categories'*, <https://webusers.imj-prg.fr/~bernhard.keller/lefevre/TheseFinale/corrainf.pdf>, 2005.
- [25] B. Keller, “A-infinity algebras, modules and functor categories,” in *Trends in representation theory of algebras and related topics*, ser. Contemp. Math. Vol. 406, Amer. Math. Soc., Providence, RI, 2006, pp. 67–93. DOI: 10.1090/conm/406/07654. [Online]. Available: <https://doi.org/10.1090/conm/406/07654>.
- [26] B. Keller, “Deriving DG categories,” *Ann. Sci. École Norm. Sup. (4)*, vol. 27, no. 1, pp. 63–102, 1994, ISSN: 0012-9593. [Online]. Available: [http://www.numdam.org/item?id=ASENS\\_1994\\_4\\_27\\_1\\_63\\_0](http://www.numdam.org/item?id=ASENS_1994_4_27_1_63_0).
- [27] H. Krause, *Homological Theory of Representations -Draft Version of a Book Project-*. Cambridge University Press, Aug. 2021. [Online]. Available: <https://www.math.uni-bielefeld.de/~hkrause/HomTheRep.pdf>.
- [28] D. Happel, *Triangulated Categories in the Representation of Finite Dimensional Algebras*, ser. London Mathematical Society Lecture Note Series. Cambridge University Press, 1988. DOI: 10.1017/CB09780511629228.
- [29] T. Bühler, “Exact categories,” *Expositiones Mathematicae*, vol. 28, no. 1, pp. 1–69, 2010, ISSN: 0723-0869. DOI: <https://doi.org/10.1016/j.exmath.2009.04.004>. [Online]. Available: <https://www.sciencedirect.com/science/article/pii/S0723086909000395>.
- [30] T. W. Thorbjørnsen, “An introduction to triangulated categories,” Bachelor’s thesis, Norwegian University of Science and Technology, 2021. [Online]. Available: <https://ntnuopen.ntnu.no/ntnu-xmlui/handle/11250/2980275>.
- [31] S. Mac Lane, *Homology*, 1st ed., ser. Classics in Mathematics. Springer Berlin, 1994. DOI: <https://doi.org/10.1007/978-3-642-62029-4>.

- [32] B. Keller, “Chain complexes and stable categories,” *manuscripta mathematica*, vol. 67, pp. 379–417, 1990.