

Assignment 4 - MAT257

David Knott
Student #999817685

October 7, 2013

1. Munkres Question 1

Since $f'(a; u) = Df(a) \cdot u$, then $f'(a; cu) = Df(a) \cdot cu = c(Df(a) \cdot u) = cf'(a; u)$.

2. Munkres Question 2

a) Note that the function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ can be rewritten like so:

$$f((x, y)) = \begin{cases} \frac{xy}{\|(x, y)\|^2} & \text{if } (x, y) \neq 0 \\ 0 & \text{if } (x, y) = 0 \end{cases}$$

Let $u = (h, k)$ be an arbitrary, non-zero vector. The directional derivative of f at origin with respect to u will then be:

$$\begin{aligned} f'(0; u) &= \lim_{t \rightarrow 0} \frac{f(0 + tu) - f(0)}{t} \\ &= \lim_{t \rightarrow 0} \frac{thtk}{\|t(h, k)\|^2} \frac{1}{t} \\ &= \lim_{t \rightarrow 0} \frac{t^2hk}{t^3\|(h, k)\|^2} \\ &= \lim_{t \rightarrow 0} \frac{hk}{t\|(h, k)\|^2} \end{aligned}$$

This limit will only exist when h or k are zero. When this is the case, the limit will evaluate to zero.

- b) Since D_1f and D_2f are the directional derivatives of elementary unit vectors then k or h will be zero, and thus both of these will be zero.
- c) The function is not differentiable since some of the directional derivatives don't exist.
- d) The function is not continuous, since if $x = y$ then the limit of the function $\phi(x) = f((x, x)) = \frac{1}{2}$ as $x \rightarrow 0$ is $\frac{1}{2}$, however if $-x = y$ then the limit of the function $\phi(x) = f((x, -x)) = -\frac{1}{2}$ as $x \rightarrow 0$ is $-\frac{1}{2}$.

3. Munkres Question 3

a) Note the function can be written as follows

$$f((x, y)) = \begin{cases} \frac{x^2y^2}{x^2y^2 + (y-x)^2} & \text{if } (x, y) \neq 0 \\ 0 & \text{if } (x, y) = 0 \end{cases}$$

Let $u = (h, k)$ be an arbitrary, non-zero vector. The directional derivative of f at origin with respect to u will then be:

$$f'(0; u) = \lim_{t \rightarrow 0} \frac{f(0 + tu) - f(0)}{t}$$

$$= \lim_{t \rightarrow 0} \frac{th^2k^2}{t^2h^2k^2 + (k-h)^2}$$

Now assuming that $(k-h)^2 \neq 0$ then $\lim_{t \rightarrow 0} t^2h^2k^2 + (k-h)^2 = (k-h)^2 \neq 0$ so the limit becomes:

$$\begin{aligned} &= \lim_{t \rightarrow 0} \frac{th^2k^2}{t^2h^2k^2 + (k-h)^2} \\ &= \frac{\lim_{t \rightarrow 0} th^2k^2}{\lim_{t \rightarrow 0} t^2h^2k^2 + (k-h)^2} \\ &= 0 \end{aligned}$$

Assuming that $(k-h)^2 = 0$ then $k = h$ so the limit becomes:

$$\begin{aligned} &= \lim_{t \rightarrow 0} \frac{th^2k^2}{t^2h^2k^2 + (k-h)^2} \\ &= \lim_{t \rightarrow 0} \frac{tk^4}{t^2k^4} \\ &= \lim_{t \rightarrow 0} \frac{1}{t} \end{aligned}$$

Which does not exist. Therefore the directional derivatives only exist when $k \neq h$.

- b) Since D_1f and D_2f are the directional derivatives of elementary unit vectors then $k \neq h$, and thus both of these will be zero.
- c) The function is not differentiable since some of the directional derivatives don't exist.
- d) The function is continuous.

4. Munkres Question 4

- a) Note the function is to be written as follows

$$f((x, y)) = \begin{cases} \frac{x^3}{x^2+y^2} & \text{if } (x, y) \neq 0 \\ 0 & \text{if } (x, y) = 0 \end{cases}$$

Let $u = (h, k)$ be an arbitrary, non-zero vector. The directional derivative of f at origin with respect to u will then be:

$$\begin{aligned} f'(0; u) &= \lim_{t \rightarrow 0} \frac{f(0 + tu) - f(0)}{t} \\ &= \lim_{t \rightarrow 0} \frac{t^3h^2}{t^2(h^2 + k^2)} \\ &= \lim_{t \rightarrow 0} \frac{th^2}{h^2 + k^2} \end{aligned}$$

Which is always zero

- b) All directional derivatives exist, therefore D_1f and D_2f must exist.
- c) The function is differentiable since all of the partial derivatives are continuous.
- d) The function is continuous since it's differentiable.

5. Additional work, question 1

a) Note that the function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ can be rewritten like so:

$$f((x, y)) = \begin{cases} \frac{x|y|}{\|(x, y)\|} & \text{if } (x, y) \neq 0 \\ 0 & \text{if } (x, y) = 0 \end{cases}$$

Let $u = (h, k)$ be an arbitrary, non-zero vector. The directional derivative of f at origin with respect to u will then be:

$$\begin{aligned} f'(0; u) &= \lim_{t \rightarrow 0} \frac{f(0 + tu) - f(0)}{t} \\ &= \lim_{t \rightarrow 0} \frac{th|tk|}{\|tu\|} \frac{1}{t} \\ &= \lim_{t \rightarrow 0} \frac{t|t|h|k|}{t|t|\|u\|} \\ &= \frac{h|k|}{\|u\|} \end{aligned}$$

Which is defined for all $u \neq 0$. Therefore all the directional derivatives exist for f .

b) If f were differentiable at the origin then $f'(0; u) = ah + bk$ for some constant $a, b \in \mathbb{R}$, however there are no such constants that satisfy $ah + bk = \frac{h|k|}{\|u\|}$. Therefore f is not differentiable at the origin.

6. Additional work, question 2

a) $f(1, 1) = 1^3 + 1^3 + 1 - 3 = 0$

b)