

II. Working with data in R (solution)

Data Science Lab, University of Copenhagen

01 September, 2021

Loading the core **tidyverse** packages, as well as the **readxl** package for importing data from .xlsx

```
library(tidyverse)
library(readxl)
```

Importing data

1-3. Use the *Import data facility*.

```
library(readxl)
climate <- read_excel("climate.xlsx")
climate
```

```
## # A tibble: 60 x 7
##   station year month   af rain  sun device
##   <chr>   <dbl> <dbl> <dbl> <dbl> <dbl> <chr>
## 1 armagh  2016     1     5 132.  44.5 Campbell Stokes
## 2 armagh  2016     2    10  62.6  71.3 Campbell Stokes
## 3 armagh  2016     3     4  43.8 117. Campbell Stokes
## 4 armagh  2016     4     5   54  140. Campbell Stokes
## 5 armagh  2016     5     0  41.4 210. Campbell Stokes
## 6 armagh  2016     6     0  75.1 114. Campbell Stokes
## 7 armagh  2016     7     0  80.6 113. Campbell Stokes
## 8 armagh  2016     8     0  52.5 135. Campbell Stokes
## 9 armagh  2016     9     0  65.4  91.1 Campbell Stokes
## 10 armagh 2016    10     0  37.1  89.8 Campbell Stokes
## # ... with 50 more rows
```

Working with the data

```
library(tidyverse)
```

4. Trying out some basic commands.

```
# Only data fra Oxford
filter(climate, station == "oxford")
```

```
## # A tibble: 12 x 7
##   station year month   af rain  sun device
##   <chr>   <dbl> <dbl> <dbl> <dbl> <dbl> <chr>
## 1 oxford  2016     1     5  83.9  59.1 Campbell Stokes
## 2 oxford  2016     2     6  47.6 113. Campbell Stokes
## 3 oxford  2016     3     4  74.2 124. Campbell Stokes
## 4 oxford  2016     4     1  53.1 164. Campbell Stokes
```

```
## 5 oxford 2016 5 0 86.1 203. Campbell Stokes
## 6 oxford 2016 6 0 95.7 100. Campbell Stokes
## 7 oxford 2016 7 0 3.4 228. Campbell Stokes
## 8 oxford 2016 8 0 41.2 204. Campbell Stokes
## 9 oxford 2016 9 0 44.6 113. Campbell Stokes
## 10 oxford 2016 10 0 26.5 112. Campbell Stokes
## 11 oxford 2016 11 3 76.1 88.3 Campbell Stokes
## 12 oxford 2016 12 10 25.8 62.3 Campbell Stokes
```

Only certain variables

```
select(climate, station, year, month, af)
```

```
## # A tibble: 60 x 4
```

```
##   station year month   af
##   <chr>   <dbl> <dbl> <dbl>
## 1 armagh 2016    1    5
## 2 armagh 2016    2   10
## 3 armagh 2016    3    4
## 4 armagh 2016    4    5
## 5 armagh 2016    5    0
## 6 armagh 2016    6    0
## 7 armagh 2016    7    0
## 8 armagh 2016    8    0
## 9 armagh 2016    9    0
## 10 armagh 2016   10    0
```

```
## # ... with 50 more rows
```

New variable with rain measured in cm

```
mutate(climate, rain_cm = rain/10)
```

```
## # A tibble: 60 x 8
```

```
##   station year month   af rain  sun device      rain_cm
##   <chr>   <dbl> <dbl> <dbl> <dbl> <dbl> <chr>      <dbl>
## 1 armagh 2016    1    5 132.  44.5 Campbell Stokes 13.2
## 2 armagh 2016    2   10 62.6  71.3 Campbell Stokes 6.26
## 3 armagh 2016    3    4 43.8 117. Campbell Stokes 4.38
## 4 armagh 2016    4    5 54 140. Campbell Stokes 5.4
## 5 armagh 2016    5    0 41.4 210. Campbell Stokes 4.14
## 6 armagh 2016    6    0 75.1 114. Campbell Stokes 7.51
## 7 armagh 2016    7    0 80.6 113. Campbell Stokes 8.06
## 8 armagh 2016    8    0 52.5 135. Campbell Stokes 5.25
## 9 armagh 2016    9    0 65.4 91.1 Campbell Stokes 6.54
## 10 armagh 2016   10    0 37.1 89.8 Campbell Stokes 3.71
```

```
## # ... with 50 more rows
```

No of obs at each station

```
count(climate, station)
```

```
## # A tibble: 5 x 2
```

```
##   station      n
##   <chr>   <int>
## 1 armagh    12
## 2 camborne 12
## 3 lerwick  12
## 4 oxford   12
## 5 sheffield 12
```

```
# Total sum over all stations
summarize(climate, total_sun = sum(sun))
```

```
## # A tibble: 1 x 1
##   total_sun
##   <dbl>
## 1    6829.
```

```
# Sort after sun
arrange(climate, sun)
```

```
## # A tibble: 60 x 7
##   station year month   af rain  sun device
##   <chr>   <dbl> <dbl> <dbl> <dbl> <dbl> <chr>
## 1 lerwick 2016   12    0 159.  11.5 Kipp Zonen
## 2 lerwick 2016    1    7 187.  34.6 Kipp Zonen
## 3 sheffield 2016    1    3 84.8  40.3 Kipp Zonen
## 4 armagh 2016    1    5 132.  44.5 Campbell Stokes
## 5 camborne 2016   12    0 58.4  47.3 Kipp Zonen
## 6 camborne 2016    1    0 222.  48   Kipp Zonen
## 7 lerwick 2016   11    1 133.  48   Kipp Zonen
## 8 armagh 2016   12    1 51.4  50.5 Campbell Stokes
## 9 camborne 2016   11    0 137.  56.5 Kipp Zonen
## 10 sheffield 2016   12    3 31.8  57.9 Kipp Zonen
## # ... with 50 more rows
```

5. Assigning the data set of observations from Oxford with 0 days of air frost to `oxford_af`.

```
oxford_af <- filter(climate, station == "oxford", af == 0)
oxford_af
```

```
## # A tibble: 6 x 7
##   station year month   af rain  sun device
##   <chr>   <dbl> <dbl> <dbl> <dbl> <dbl> <chr>
## 1 oxford 2016    5    0 86.1  203. Campbell Stokes
## 2 oxford 2016    6    0 95.7  100. Campbell Stokes
## 3 oxford 2016    7    0  3.4  228. Campbell Stokes
## 4 oxford 2016    8    0 41.2  204. Campbell Stokes
## 5 oxford 2016    9    0 44.6  113. Campbell Stokes
## 6 oxford 2016   10    0 26.5  112. Campbell Stokes
```

Counting the number of observations assigned to `oxford_af`.

```
count(oxford_af)
```

```
## # A tibble: 1 x 1
##       n
##   <int>
## 1     6
```

Alternative solution without using `oxford_af`.

```
climate %>%
  filter(station == "oxford", af == 0) %>%
  count()
```

```
## # A tibble: 1 x 1
##       n
```

```
## <int>
## 1 6
```

6. Dataset with observations from Camborne and Oxford.

```
filter(climate, station == "camborne" | station == "oxford")
# Or: filter(climate, station %in% c("camborne", "oxford"))
```

```
## # A tibble: 24 x 7
##   station year month   af rain sun device
##   <chr>   <dbl> <dbl> <dbl> <dbl> <dbl> <chr>
## 1 camborne 2016     1     0 222.  48 Kipp Zonen
## 2 camborne 2016     2     0 162.  64.1 Kipp Zonen
## 3 camborne 2016     3     0  88.4 140. Kipp Zonen
## 4 camborne 2016     4     0  81.4 184. Kipp Zonen
## 5 camborne 2016     5     0  45.6 206. Kipp Zonen
## 6 camborne 2016     6     0  65.8 132. Kipp Zonen
## 7 camborne 2016     7     0  23.2 161. Kipp Zonen
## 8 camborne 2016     8     0  57.4 171. Kipp Zonen
## 9 camborne 2016     9     0 154. 103. Kipp Zonen
## 10 camborne 2016    10     0  53.2 125. Kipp Zonen
## # ... with 14 more rows
```

7. Make the new variable and give the nes dataset a name

```
climate2 <- mutate(climate, sqrtSum=sqrt(sun))
```

8. Assigning the rainfall observations to the vector `rain_vector`.

```
rain_vector <- climate$rain
```

Extracting elements from the rainfall vector.

```
rain_vector
```

```
## [1] 131.9 62.6 43.8 54.0 41.4 75.1 80.6 52.5 65.4 37.1 40.8 51.4
## [13] 221.6 161.6 88.4 81.4 45.6 65.8 23.2 57.4 153.7 53.2 137.0 58.4
## [25] 187.0 119.6 79.2 55.1 46.0 48.4 115.2 108.4 110.0 56.2 133.4 159.4
## [37] 83.9 47.6 74.2 53.1 86.1 95.7 3.4 41.2 44.6 26.5 76.1 25.8
## [49] 84.8 68.6 87.2 65.8 58.2 130.4 30.0 62.0 44.2 29.2 95.6 31.8
```

```
# First six elements
```

```
rain_vector[1:6]
```

```
## [1] 131.9 62.6 43.8 54.0 41.4 75.1
```

```
# Element number five
```

```
rain_vector[5]
```

```
## [1] 41.4
```

```
# Element numbers 2, 4 and 6
```

```
rain_vector[c(2,4,6)]
```

```
## [1] 62.6 54.0 75.1
```

Some summary statistics for the rainfall observations.

```
mean(rain_vector)
```

```
## [1] 75.79667
```

```
sd(rain_vector)
```

```
## [1] 43.18915
```

```
sum(rain_vector)
```

```
## [1] 4547.8
```

9. Summary statistics for the rainfall observations computed using `summarize`. It is most easily made with the pipe operator, but can also be done in several steps (not shown).

```
summarize(climate,  
  avg_rain = mean(rain),  
  sd_rain = sd(rain),  
  sum_rain = sum(rain))
```

```
## # A tibble: 1 x 3
```

```
##   avg_rain sd_rain sum_rain
```

```
##   <dbl>   <dbl>   <dbl>
```

```
## 1    75.8    43.2   4548.
```

10. Summary statistics for rainfall data by weather station, including number of observations for each station, and sorted in descending order according to annual rainfall.

```
climate %>%  
  group_by(station) %>%  
  summarize(avg_rain = mean(rain),  
    sd_rain = sd(rain),  
    sum_rain = sum(rain),  
    n=n()) %>%  
  arrange(desc(sum_rain))
```

```
## # A tibble: 5 x 5
```

```
##   station avg_rain sd_rain sum_rain    n
```

```
##   <chr>    <dbl>   <dbl>   <dbl> <int>
```

```
## 1 lerwick    101.    45.6   1218.   12
```

```
## 2 camborne    95.6    59.4   1147.   12
```

```
## 3 sheffield    65.6    30.5    788.   12
```

```
## 4 armagh     61.4    26.1    737.   12
```

```
## 5 oxford     54.8    28.5    658.   12
```

11. Summary statistics for rainfall data by weather station, sorted in ascending order according to average monthly sunshine duration.

```
climate %>%  
  group_by(station) %>%  
  summarize(avg_rain = mean(rain),  
    sd_rain = sd(rain),  
    sum_rain = sum(rain),  
    avg_sun = mean(sun)) %>%  
  arrange(avg_sun)
```

```
## # A tibble: 5 x 5
```

```
##   station avg_rain sd_rain sum_rain avg_sun
```

```
##   <chr>    <dbl>   <dbl>   <dbl>   <dbl>
```

```
## 1 lerwick    101.    45.6   1218.   101.
```

```
## 2 armagh     61.4    26.1    737.   104.
```

```
## 3 sheffield    65.6    30.5    788.   113.
```

```
## 4 camborne      95.6    59.4    1147.    120.
## 5 oxford        54.8    28.5     658.    131.
```

12. Weather station with largest median number of monthly sunshine hours over the months April to September was Oxford:

```
climate %>%
  filter(month %in% 4:9) %>%
  group_by(station) %>%
  summarize(med_sun = median(sun)) %>%
  arrange(desc(med_sun))
```

```
## # A tibble: 5 x 2
##   station med_sun
##   <chr>    <dbl>
## 1 oxford    183.
## 2 camborne   166
## 3 sheffield  160.
## 4 lerwick   132.
## 5 armagh    124.
```

13. For each weather station apart from Armagh, the total rainfall (in cm) and duration of sunshine (in days) in the months with no days of air frost.

```
climate %>%
  group_by(station) %>%
  filter(station != "armagh", af == 0) %>%
  summarize(total_rain = sum(rain)/10,
            total_sun = sum(sun)/24)
```

```
## # A tibble: 4 x 3
##   station total_rain total_sun
##   <chr>    <dbl>    <dbl>
## 1 camborne    115.     59.9
## 2 lerwick     64.4     33.3
## 3 oxford      29.8     40.0
## 4 sheffield   35.4     35.0
```

14. For each month: Number of stations with at least two days of air frost or more than 95 mm rain, and the average sunshine duration for these stations.

```
climate %>%
  filter(af >= 2 | rain > 95) %>%
  group_by(month) %>%
  summarize(count = n(), avg_sun = mean(sun))
```

```
## # A tibble: 10 x 3
##   month count avg_sun
##   <dbl> <int>   <dbl>
## 1     1     5    45.3
## 2     2     5    86.2
## 3     3     4   106.
## 4     4     2   148.
## 5     6     2   103.
## 6     7     1    80.5
## 7     8     1    92.2
## 8     9     2   105.
## 9    11     5    68.0
```

```
## 10    12    3    43.9
```

15. Many solutions were already made with the pipe operator.