

Mortality of Indiana for February, 1912. (Stillbirths excluded.)

| POPULATION BY GEOGRAPHICAL SECTIONS AND AS URBAN AND RURAL. | Population Estimated 1912. | Total Deaths Reported for | | | | Annual Death Rate Per 100,000 Population. | | | | | | Important Ages. | | | | | | | | | | | | |
|---|----------------------------------|--|-----------|---|-----------|--|---|-----------|--|-----------|-------------------------------------|-----------------|-----------|---------|-----------|---------|-----------|-----------|-----------|-----------|-----------|--------------|-----------|------|
| | | February 1912. | | January 1912. | | Total Deaths for 1912 to Date. | February, 1912. | | January, 1912. | | Rate for Year 1911 to Same Date. | Under 1. | | 1 to 4. | | 5 to 9. | | 10 to 14. | | 15 to 19. | | 65 and Over. | | |
| | | Total Deaths Reported for February 1912. | | Total Deaths Reported for January 1912. | | | Total Deaths Reported for February, 1911. | | Total Deaths Reported for January 1911 to Same Date. | | | Number. | | Number. | | Number. | | Number. | | Number. | | Number. | | |
| | | Number. | Per Cent. | Number. | Per Cent. | | Number. | Per Cent. | Number. | Per Cent. | | Number. | Per Cent. | Number. | Per Cent. | Number. | Per Cent. | Number. | Per Cent. | Number. | Per Cent. | Number. | Per Cent. | |
| State | 2,730,506 | 2,985 | 3.188 | 2,908 | 3.171 | 6,052 | 13.7 | 13.7 | 14.0 | 13.7 | 14.1 | 295 | 9.8 | 132 | 4.4 | 48 | 1.6 | 42 | 1.4 | 62 | 2.0 | 1,042 | 34.9 | |
| Northern Counties | 939,532 | 955 | 1.091 | 948 | 2.048 | 2,010 | 12.8 | 13.7 | 13.2 | 13.2 | 13.4 | 114 | 11.9 | 44 | 4.6 | 11 | 1.1 | 14 | 1.4 | 14 | 1.4 | 330 | 36.6 | |
| Central Counties | 1,127,217 | 1,289 | 1.370 | 1,221 | 2.668 | 2,594 | 14.4 | 14.4 | 14.2 | 14.4 | 14.3 | 120 | 9.3 | 49 | 3.8 | 27 | 2.0 | 18 | 1.3 | 35 | 2.5 | 448 | 34.7 | |
| Southern Counties | 663,757 | 741 | 716 | 740 | 1.457 | 1,448 | 14.0 | 12.7 | 14.5 | 13.3 | 14.6 | 61 | 8.2 | 39 | 5.2 | 10 | 1.3 | 10 | 1.3 | 15 | 2.0 | 244 | 32.9 | |
| All Cities | 1,164,294 | 1,418 | 1,585 | 1,359 | 2,983 | 2,876 | 15.3 | 15.8 | 15.4 | 15.6 | 15.5 | 170 | 11.9 | 63 | 4.4 | 26 | 1.8 | 19 | 1.3 | 36 | 2.6 | 429 | 30.2 | |
| Over 100,000 | 240,698 | 317 | 328 | 301 | 645 | 606 | 16.6 | 16.1 | 16.7 | 16.3 | 16.0 | 31 | 9.7 | 10 | 3.1 | 3 | 0 | 5 | 1.5 | 7 | 2.2 | 81 | 25.5 | |
| 45,000 to 100,000 | 253,337 | 292 | 334 | 297 | 646 | 633 | 14.5 | 16.4 | 15.7 | 15.5 | 15.9 | 45 | 15.4 | 20 | 6.8 | 8 | 2.7 | 3 | 1.0 | 7 | 2.3 | 71 | 24.3 | |
| 20,000 to 45,000 | 132,435 | 146 | 167 | 147 | 313 | 310 | 13.9 | 14.8 | 14.6 | 14.3 | 14.6 | 16 | 10.9 | 10 | 6.8 | 1 | 6 | 2 | 3 | 4 | 6 | 4.1 | 44 | 30.1 |
| 10,000 to 20,000 | 208,083 | 232 | 292 | 240 | 544 | 520 | 15.2 | 16.5 | 14.9 | 15.9 | 15.3 | 36 | 14.2 | 7 | 7 | 3 | 1.9 | 2 | 7 | 4 | 1.5 | 55 | 33.7 | |
| Under 10,000 | 320,341 | 411 | 424 | 374 | 835 | 809 | 15.7 | 15.1 | 15.7 | 15.3 | 15.2 | 32 | 7.7 | 16 | 3.8 | 9 | 2.1 | 4 | 9 | 14 | 3.4 | 145 | 36.0 | |
| Country | 1,566,212 | 1,567 | 1,621 | 1,550 | 3,188 | 3,174 | 12.6 | 12.2 | 12.9 | 12.4 | 13.0 | 125 | 7.9 | 69 | 4.3 | 22 | 1.4 | 23 | 1.4 | 24 | 1.5 | 613 | 39.1 | |

| POPULATION BY GEOGRAPHICAL SECTIONS AND AS URBAN AND RURAL. | Deaths and Annual Death Rates Per 100,000 Population From Important Causes. | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|---|---|-------------|--------------------------------------|-------------|-------------------|-------------|----------------------------------|-------------|-------------------|-------------|----------|-------------|-------------------------|-------------|---|-------------|---|-------------|------------------------------|-------------|---|-------------|-----------------|-------------|------------------------------------|-------------|---------|-------------|---------------------|-------------|----------------|-------------|-----|
| | Pulmonary Tuber- culosis. | | Other Forms Tuber- culosis. | | Typhoid Fever. | | Diph- theria and Croup. | | Scarlet Fever. | | Measles. | | Whoop- ing Cough. | | Lobar and Broncho- Pneu- monia. | | Diarrhea and Enteritis (Under 2 Years.) | | Cerebro- spinal Fever. | | Acute Anterior Poli- myelitis. | | Influen- za. | | Puer- peral Septi- cemia. | | Cancer. | | External Causes. | | Small- pox. | | |
| | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | |
| | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | Number. | Death Rate. | |
| State | 326 | 150.6 | 37 | 17.1 | 41 | 18.6 | 29 | 13.4 | 14 | 6.4 | 3 | 1.3 | 21 | 9.7 | 391 | 180.7 | 37 | 17.1 | 1 | 4 | | | | 112 | 51.7 | 23 | 10.6 | 150 | 69.3 | 187 | 86.4 | 2 | 9 |
| Northern Counties | 91 | 122.2 | 9 | 12.0 | 16 | 21.4 | 8 | 10.7 | 2 | 2.6 | | | 6 | 8.0 | 133 | 178.6 | 27 | 36.2 | | | | | | 31 | 41.6 | 5 | 6.7 | 39 | 52.3 | 63 | 84.6 | 1 | 1.3 |
| Central Counties | 137 | 153.4 | 23 | 25.7 | 13 | 14.5 | 17 | 19.0 | 6 | 6.7 | 3 | 3.3 | 6 | 6.7 | 139 | 155.8 | 6 | 7.7 | | | | | | 33 | 36.9 | 15 | 16.8 | 79 | 88.4 | 39 | 99.6 | 1 | 1.1 |
| Southern Counties | 98 | 186.3 | 5 | 9.5 | 12 | 22.8 | 4 | 7.6 | 6 | 11.4 | | | 9 | 17.1 | 116 | 226.3 | 4 | 7.6 | 1 | 1.9 | | | | 48 | 91.2 | 3 | 5.7 | 32 | 60.8 | 35 | 96.5 | | |
| All Cities | 161 | 174.5 | 21 | 22.7 | 25 | 27.1 | 17 | 18.4 | 5 | 5.4 | 1 | 1.0 | 5 | 5.4 | 157 | 170.1 | 27 | 29.2 | | | | | | 35 | 37.9 | 15 | 16.2 | 63 | 88.9 | 79 | 85.6 | 1 | 1.0 |
| Over 100,000 | 43 | 226.0 | 10 | 52.5 | 1 | 5.2 | 4 | 21.0 | | | | | | | 27 | 141.9 | 1 | 5.2 | | | | | | 4 | 21.0 | 3 | 15.7 | 22 | 115.6 | 18 | 94.6 | | |
| 45,000 to 100,000 | 38 | 189.3 | 3 | 14.9 | 8 | 39.8 | 4 | 19.9 | 1 | 4.9 | | | | | 35 | 174.4 | 5 | 24.9 | | | | | | 6 | 29.8 | 3 | 14.9 | 17 | 84.7 | 20 | 96.6 | | |
| 20,000 to 45,000 | 11 | 104.8 | 1 | 9.5 | 4 | 38.1 | 2 | 19.0 | 2 | 19.0 | 1 | 9.5 | | | 12 | 114.3 | 3 | 25.5 | | | | | | 4 | 38.1 | 2 | 19.0 | 11 | 104.8 | 13 | 123.9 | | |
| 10,000 to 20,000 | 21 | 127.3 | 3 | 18.1 | 8 | 48.5 | 9 | 12.1 | | | | | 2 | 12.1 | 32 | 104.0 | 13 | 78.8 | | | | | | 6 | 38.3 | 1 | 6.0 | 18 | 109.1 | 17 | 103.1 | | |
| Under 10,000 | 48 | 183.3 | 4 | 15.2 | 4 | 15.2 | 5 | 19.1 | 2 | 7.6 | | | 3 | 11.4 | 51 | 194.8 | 5 | 19.1 | | | | | | 15 | 57.3 | 6 | 22.8 | 15 | 57.3 | 11 | 42.0 | 1 | 3.8 |
| Country | 165 | 132.9 | 16 | 12.8 | 16 | 12.8 | 12 | 9.6 | 9 | 7.2 | 2 | 1.6 | 16 | 12.8 | 234 | 188.6 | 10 | 8.0 | 1 | 8 | | | | 77 | 62.0 | 8 | 6.4 | 67 | 54.0 | 108 | 87.0 | 1 | 8 |

U. S. Department of Agriculture, Weather Bureau. Condensed Summary for Month of February, 1912.

V. H. CHURCH, SECTION DIRECTOR, INDIANAPOLIS, IND.

TEMPERATURE—IN DEGREES FAHRENHEIT.

| Section Average. | Departure from the normal. | Extremes. | | | | | |
|------------------|----------------------------|-----------|----|----------|---------|-------|---|
| | | Station. | | Highest. | | Date. | |
| | | Station. | | Lowest. | | Date. | |
| 22.9 | -5.1 | Rome | 60 | 17 | Laporte | -20 | 3 |

PRECIPITATION—IN INCHES AND HUNDREDTHS.

| Section Average. | Departure from the normal. | Extremes. | | | |
|------------------|----------------------------|-----------|------|--------------------------|------|
| | | Station. | | Greatest monthly amount. | |
| | | Station. | | Least monthly amount. | |
| 2.60 | -0.25 | Vincennes | 4.08 | Richmond | 1.30 |