
Emulating Satellite Tangential Scale Distortion

Remote sensing systems, provide an instantaneous “snapshot” view of the Earth from directly overhead. Each detector essentially takes a “snapshot” of each ground resolution cell.

However, a distortion occurs due to the rotation of the scanning optics. As the sensor scans across each line, the distance from the sensor to the ground increases further away from the centre of the swath. Although the scanning mirror rotates at a constant speed, the relative motion of the sensor is faster (relative to the ground) and scans a larger area as it moves closer to the edges. This effect results in the compression of image features at points away from the nadir and is called **tangential scale distortion**.




Figure 1: The ground resolution cell size along a single across-track scan

To reflect this deformation when generating synthetic imagery samples, we propose to apply an unidimensional sigmoid application to coordinates along a virtual scanning line. Explicitly, let σ_λ be :

$$\sigma_\lambda : \begin{cases} \mathbb{R} & \longrightarrow [0, 1] \\ x & \longmapsto \frac{1}{1+e^{-\lambda x}} \end{cases}, \quad \lambda > 0$$

Let now $\ell > 0$ denote the image size along the deformation line and the center position $c = \frac{\ell}{2}$ corresponding to a virtual nadir. To reflect invariance at nadir and compression at edges, we apply along the swath line $[0, \ell]$ the application $x \in [0, \ell] \mapsto \ell \sigma_\lambda(x - c)$.

In practice nearest method can be used for shifted pixels assignation and interpolation to cope with missing positions.

Similarly,

$$(??) \Rightarrow \forall x \in]c, \ell[\quad x - c \leq \sigma_\lambda^{-1} \left(\frac{x}{\ell} \right) \quad (5)$$

$$\Rightarrow \forall x \in]c, \ell[\quad \lambda \leq \sigma_{x-c}^{-1} \left(\frac{x}{\ell} \right) \quad (6)$$

Hence $\forall x \in]0, \ell[\setminus \{c\}$ we have

$$\lambda \leq \frac{1}{x-c} \log \left(\frac{x/\ell}{1-x/\ell} \right) = \underbrace{\frac{1}{x-c} \log \left(\frac{x}{\ell-x} \right)}_{\psi_\ell(x)}$$

ψ_ℓ decreases on $]0, c[$ and increases on $]c, \ell[$. It thus admits a lower bound in c , i.e.

$$\liminf_{]0, \ell[\setminus \{c\}} \psi_\ell = \lim_{x \rightarrow c} \psi_\ell(x)$$

To compute this limit, we observe that :

$$\begin{aligned} \log \left(\frac{x}{\ell-x} \right) &= \log(x) - \log(\ell-x) \\ &= \log(c + (x-c)) - \log(c - (x-c)) \\ &\underset{x \rightarrow c}{=} \frac{x-c}{c} + \frac{x-c}{c} + o(x-c) \\ &\underset{x \rightarrow c}{=} \frac{4}{\ell} (x-c) + o(x-c) \end{aligned}$$

Which entails :

$$\psi_\ell(x) \underset{x \rightarrow c}{=} \frac{1}{x-c} \left[\frac{4}{\ell} (x-c) + o(x-c) \right] \underset{x \rightarrow c}{=} \frac{4}{\ell} + o(1) \Rightarrow \lim_{x \rightarrow c} \psi_\ell(x) = \frac{4}{\ell}$$

Reversely, we verify that choosing $\lambda \leq \frac{4}{\ell}$ allows to verify (??) and (??), hence making $x \mapsto \ell \sigma_\lambda(x-c)$ contractive. \square