

# Computational Graphics: Lecture 8

The Cvdlab Team

Tue, Mar 17, 2015

# Outline: Algebra reminders

- 1 Linear spaces
- 2 Linear combinations
- 3 Subspaces
- 4 Spans
- 5 Bases
- 6 Affine spaces
- 7 Affine combinations
- 8 Convex combinations

# Linear spaces

# Definition

A **linear** (or **vector**) **space**  $\mathcal{V}$  over a field  $\mathcal{F}$  is a set with two composition rules, such that, for each  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$  and for each  $\alpha, \beta \in \mathcal{F}$ , the rules  $+$ ,  $\cdot$  satisfy the following axioms:

$$\textcircled{1} \quad \mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v}; \quad (\text{commutativity of addition})$$

# Definition

A **linear** (or **vector**) **space**  $\mathcal{V}$  over a field  $\mathcal{F}$  is a set with two composition rules, such that, for each  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$  and for each  $\alpha, \beta \in \mathcal{F}$ , the rules  $+$ ,  $\cdot$  satisfy the following axioms:

- ①  $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v};$

(commutativity of addition)

- ②  $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w};$

(associativity of addition)

# Definition

A **linear** (or **vector**) **space**  $\mathcal{V}$  over a field  $\mathcal{F}$  is a set with two composition rules, such that, for each  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$  and for each  $\alpha, \beta \in \mathcal{F}$ , the rules  $+$ ,  $\cdot$  satisfy the following axioms:

- ①  $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v};$  (commutativity of addition)
- ②  $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w};$  (associativity of addition)
- ③ there is a  $\mathbf{0} \in \mathcal{V}$  such that  $\mathbf{v} + \mathbf{0} = \mathbf{v};$  (neutral el. of addition)

# Definition

A **linear** (or **vector**) **space**  $\mathcal{V}$  over a field  $\mathcal{F}$  is a set with two composition rules, such that, for each  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$  and for each  $\alpha, \beta \in \mathcal{F}$ , the rules  $+$ ,  $\cdot$  satisfy the following axioms:

- ①  $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v};$  (commutativity of addition)
- ②  $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w};$  (associativity of addition)
- ③ there is a  $\mathbf{0} \in \mathcal{V}$  such that  $\mathbf{v} + \mathbf{0} = \mathbf{v};$  (neutral el. of addition)
- ④ there is a  $-\mathbf{v} \in \mathcal{V}$  such that  $\mathbf{v} + (-\mathbf{v}) = \mathbf{0};$  (inverse of add.)

# Definition

A **linear** (or **vector**) **space**  $\mathcal{V}$  over a field  $\mathcal{F}$  is a set with two composition rules, such that, for each  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$  and for each  $\alpha, \beta \in \mathcal{F}$ , the rules  $+$ ,  $\cdot$  satisfy the following axioms:

- ①  $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v};$  (commutativity of addition)
- ②  $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w};$  (associativity of addition)
- ③ there is a  $\mathbf{0} \in \mathcal{V}$  such that  $\mathbf{v} + \mathbf{0} = \mathbf{v};$  (neutral el. of addition)
- ④ there is a  $-\mathbf{v} \in \mathcal{V}$  such that  $\mathbf{v} + (-\mathbf{v}) = \mathbf{0};$  (inverse of add.)
- ⑤  $\alpha \cdot (\mathbf{v} + \mathbf{w}) = \alpha \cdot \mathbf{v} + \alpha \cdot \mathbf{w};$  (distrib. of addition w.r.t. product)



# Definition

A **linear** (or **vector**) **space**  $\mathcal{V}$  over a field  $\mathcal{F}$  is a set with two composition rules, such that, for each  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$  and for each  $\alpha, \beta \in \mathcal{F}$ , the rules  $+$ ,  $\cdot$  satisfy the following axioms:

- ①  $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v};$  (commutativity of addition)
- ②  $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w};$  (associativity of addition)
- ③ there is a  $\mathbf{0} \in \mathcal{V}$  such that  $\mathbf{v} + \mathbf{0} = \mathbf{v};$  (neutral el. of addition)
- ④ there is a  $-\mathbf{v} \in \mathcal{V}$  such that  $\mathbf{v} + (-\mathbf{v}) = \mathbf{0};$  (inverse of add.)
- ⑤  $\alpha \cdot (\mathbf{v} + \mathbf{w}) = \alpha \cdot \mathbf{v} + \alpha \cdot \mathbf{w};$  (distrib. of addition w.r.t. product)
- ⑥  $(\alpha + \beta) \cdot \mathbf{v} = \alpha \cdot \mathbf{v} + \beta \cdot \mathbf{v};$  (distrib. of product w.r.t. addition)

# Definition

A **linear** (or **vector**) **space**  $\mathcal{V}$  over a field  $\mathcal{F}$  is a set with two composition rules, such that, for each  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$  and for each  $\alpha, \beta \in \mathcal{F}$ , the rules  $+$ ,  $\cdot$  satisfy the following axioms:

- ①  $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v};$  (commutativity of addition)
- ②  $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w};$  (associativity of addition)
- ③ there is a  $\mathbf{0} \in \mathcal{V}$  such that  $\mathbf{v} + \mathbf{0} = \mathbf{v};$  (neutral el. of addition)
- ④ there is a  $-\mathbf{v} \in \mathcal{V}$  such that  $\mathbf{v} + (-\mathbf{v}) = \mathbf{0};$  (inverse of add.)
- ⑤  $\alpha \cdot (\mathbf{v} + \mathbf{w}) = \alpha \cdot \mathbf{v} + \alpha \cdot \mathbf{w};$  (distrib. of addition w.r.t. product)
- ⑥  $(\alpha + \beta) \cdot \mathbf{v} = \alpha \cdot \mathbf{v} + \beta \cdot \mathbf{v};$  (distrib. of product w.r.t. addition)
- ⑦  $\alpha \cdot (\beta \cdot \mathbf{v}) = (\alpha\beta) \cdot \mathbf{v};$  (associativity of product)

# Definition

A **linear** (or **vector**) **space**  $\mathcal{V}$  over a field  $\mathcal{F}$  is a set with two composition rules, such that, for each  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$  and for each  $\alpha, \beta \in \mathcal{F}$ , the rules  $+$ ,  $\cdot$  satisfy the following axioms:

- ①  $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v};$  (commutativity of addition)
- ②  $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w};$  (associativity of addition)
- ③ there is a  $\mathbf{0} \in \mathcal{V}$  such that  $\mathbf{v} + \mathbf{0} = \mathbf{v};$  (neutral el. of addition)
- ④ there is a  $-\mathbf{v} \in \mathcal{V}$  such that  $\mathbf{v} + (-\mathbf{v}) = \mathbf{0};$  (inverse of add.)
- ⑤  $\alpha \cdot (\mathbf{v} + \mathbf{w}) = \alpha \cdot \mathbf{v} + \alpha \cdot \mathbf{w};$  (distrib. of addition w.r.t. product)
- ⑥  $(\alpha + \beta) \cdot \mathbf{v} = \alpha \cdot \mathbf{v} + \beta \cdot \mathbf{v};$  (distrib. of product w.r.t. addition)
- ⑦  $\alpha \cdot (\beta \cdot \mathbf{v}) = (\alpha\beta) \cdot \mathbf{v};$  (associativity of product)
- ⑧  $1 \cdot \mathbf{v} = \mathbf{v}.$  (neutral element of product)

## Example: vector space of real matrices

Let  $\mathcal{M}_n^m(\mathbb{R})$  be the set of  $m \times n$  matrices with elements in the field  $\mathbb{R}$ . An element  $A$  in such a set is denoted as

$$A = (\alpha_{ij})$$

## Example: vector space of real matrices

Let  $\mathcal{M}_n^m(\mathbb{R})$  be the set of  $m \times n$  matrices with elements in the field  $\mathbb{R}$ . An element  $A$  in such a set is denoted as

$$A = (\alpha_{ij})$$

**Addition** and **multiplication by a scalar** are defined component-wise:

$$A + B = (\alpha_{ij}) + (\beta_{ij}) = (\alpha_{ij} + \beta_{ij})$$

$$\gamma A = \gamma(\alpha_{ij}) = (\gamma\alpha_{ij})$$

# Linear combinations

# Linear combination

Let  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \in \mathcal{V}$  and  $\alpha_1, \alpha_2, \dots, \alpha_n \in \mathcal{F}$ ,

The vector

$$\alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n = \sum_{i=1}^n \alpha_i \mathbf{v}_i \in \mathcal{V}$$

is called a **linear combination** of vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$  with scalars  $\alpha_1, \alpha_2, \dots, \alpha_n \in \mathcal{F}$

# Subspaces



# Subspace

Let  $(\mathcal{V}, +, \cdot)$  be a vector space on the field  $\mathcal{F}$ .

$\mathcal{U} \subset \mathcal{V}$  is a **subspace** of  $\mathcal{V}$  if  $(\mathcal{U}, +, \cdot)$  is a vector space with respect to the same operations.

# Subspace

Let  $(\mathcal{V}, +, \cdot)$  be a vector space on the field  $\mathcal{F}$ .

$\mathcal{U} \subset \mathcal{V}$  is a **subspace** of  $\mathcal{V}$  if  $(\mathcal{U}, +, \cdot)$  is a vector space with respect to the same operations.

$\mathcal{U} \subset \mathcal{V}$  is a **subspace** of  $\mathcal{V}$  if and only if

$$\mathcal{U} \neq \emptyset;$$

$$\text{for each } \alpha \in \mathcal{F} \text{ and } \mathbf{u}_1, \mathbf{u}_2 \in \mathcal{U}, \alpha \mathbf{u}_1 + \mathbf{u}_2 \in \mathcal{U}$$

# Subspace

Let  $(\mathcal{V}, +, \cdot)$  be a vector space on the field  $\mathcal{F}$ .

$\mathcal{U} \subset \mathcal{V}$  is a **subspace** of  $\mathcal{V}$  if  $(\mathcal{U}, +, \cdot)$  is a vector space with respect to the same operations.

$\mathcal{U} \subset \mathcal{V}$  is a **subspace** of  $\mathcal{V}$  if and only if

$$\mathcal{U} \neq \emptyset;$$

$$\text{for each } \alpha \in \mathcal{F} \text{ and } \mathbf{u}_1, \mathbf{u}_2 \in \mathcal{U}, \alpha \mathbf{u}_1 + \mathbf{u}_2 \in \mathcal{U}$$

**codimension** of a subspace  $\mathcal{U} \subset \mathcal{V}$   
is defined as  $\dim \mathcal{V} - \dim \mathcal{U}$

# Subspace

Let  $(\mathcal{V}, +, \cdot)$  be a vector space on the field  $\mathcal{F}$ .

$\mathcal{U} \subset \mathcal{V}$  is a **subspace** of  $\mathcal{V}$  if  $(\mathcal{U}, +, \cdot)$  is a vector space with respect to the same operations.

$\mathcal{U} \subset \mathcal{V}$  is a **subspace** of  $\mathcal{V}$  if and only if

$$\mathcal{U} \neq \emptyset;$$

$$\text{for each } \alpha \in \mathcal{F} \text{ and } \mathbf{u}_1, \mathbf{u}_2 \in \mathcal{U}, \alpha \mathbf{u}_1 + \mathbf{u}_2 \in \mathcal{U}$$

**codimension** of a subspace  $\mathcal{U} \subset \mathcal{V}$   
is defined as  $\dim \mathcal{V} - \dim \mathcal{U}$

# Subspace

Let  $(\mathcal{V}, +, \cdot)$  be a vector space on the field  $\mathcal{F}$ .

$\mathcal{U} \subset \mathcal{V}$  is a **subspace** of  $\mathcal{V}$  if  $(\mathcal{U}, +, \cdot)$  is a vector space with respect to the same operations.

$\mathcal{U} \subset \mathcal{V}$  is a **subspace** of  $\mathcal{V}$  if and only if

$$\mathcal{U} \neq \emptyset;$$

$$\text{for each } \alpha \in \mathcal{F} \text{ and } \mathbf{u}_1, \mathbf{u}_2 \in \mathcal{U}, \alpha \mathbf{u}_1 + \mathbf{u}_2 \in \mathcal{U}$$

**codimension** of a subspace  $\mathcal{U} \subset \mathcal{V}$   
is defined as  $\dim \mathcal{V} - \dim \mathcal{U}$

## Question

*Examples of codimension? in 1D, 2D, 3D*

# Spans

# Span

- The set of all linear combinations of elements of a set  $S \subset \mathcal{V}$  is a subspace of  $\mathcal{V}$ .

# Span

- The set of all linear combinations of elements of a set  $S \subset \mathcal{V}$  is a subspace of  $\mathcal{V}$ .
- Such a subspace is called the **span of  $S$**  and is denoted as

$$\text{lin } S$$



# Span

- The set of all linear combinations of elements of a set  $S \subset \mathcal{V}$  is a subspace of  $\mathcal{V}$ .
- Such a subspace is called the **span of  $S$**  and is denoted as

$$\text{lin } S$$

- If a subspace  $\mathcal{U}$  of  $\mathcal{V}$  can be generated as the span of a set  $S$  of vectors in  $\mathcal{V}$ , then  $S$  is called a **generating set** or a **spanning set** for  $\mathcal{U}$ .

# Linear independence

- A set of vectors  $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$  is **linearly independent** if

$$\sum_{i=1}^n \alpha_i \mathbf{v}_i = \mathbf{0}$$

implies that  $\alpha_i = 0$  for each  $i$

# Linear independence

- A set of vectors  $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$  is **linearly independent** if

$$\sum_{i=1}^n \alpha_i \mathbf{v}_i = \mathbf{0}$$

implies that  $\alpha_i = 0$  for each  $i$

- As a consequence, **a set of vectors is linearly independent** when none of them belongs to the span of the others.

# Bases

# Bases and coordinates

When working with vector spaces, the concept of **basis**, a **discrete subset of linearly independent elements**, is probably the most useful to deal with.

- each element of the space can be **represented uniquely as linear combination of basis elements**

# Bases and coordinates

When working with vector spaces, the concept of **basis**, a **discrete subset of linearly independent elements**, is probably the most useful to deal with.

- each element of the space can be **represented uniquely as linear combination of basis elements**
- this leads to a **parametrization** of the space, i.e. to **represent each element by a sequence of scalars**, called its **coordinates** with respect to the chosen basis.

# Bases

A set of vectors  $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$  is a **basis** for the vector space  $\mathcal{V}$  iff

- 1 the set is linearly independent, and

# Bases

A set of vectors  $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$  is a **basis** for the vector space  $\mathcal{V}$  iff

- 1 the set is linearly independent, and
- 2  $\mathcal{V} = \text{lin} \{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$



# Bases

- Every two bases of  $\mathcal{V}$  have the same number of elements, that is called the **dimension** of  $\mathcal{V}$  and is denoted

$$\dim \mathcal{V}$$

# Bases

- Every two bases of  $\mathcal{V}$  have the same number of elements, that is called the **dimension** of  $\mathcal{V}$  and is denoted

$$\dim \mathcal{V}$$

- Some important **properties** of the bases of a vector space are:

# Bases

- Every two bases of  $\mathcal{V}$  have the same number of elements, that is called the **dimension** of  $\mathcal{V}$  and is denoted

$$\dim \mathcal{V}$$

- Some important **properties** of the bases of a vector space are:
  - ① each spanning set for  $\mathcal{V}$  contains a basis;

# Bases

- Every two bases of  $\mathcal{V}$  have the same number of elements, that is called the **dimension** of  $\mathcal{V}$  and is denoted

$$\dim \mathcal{V}$$

- Some important **properties** of the bases of a vector space are:
  - 1 each spanning set for  $\mathcal{V}$  contains a basis;
  - 2 each minimal spanning set is a basis;

# Bases

- Every two bases of  $\mathcal{V}$  have the same number of elements, that is called the **dimension** of  $\mathcal{V}$  and is denoted

$$\dim \mathcal{V}$$

- Some important **properties** of the bases of a vector space are:
  - 1 each spanning set for  $\mathcal{V}$  contains a basis;
  - 2 each minimal spanning set is a basis;
  - 3 each linearly independent set of vectors is contained in a basis;

# Bases

- Every two bases of  $\mathcal{V}$  have the same number of elements, that is called the **dimension** of  $\mathcal{V}$  and is denoted

$$\dim \mathcal{V}$$

- Some important **properties** of the bases of a vector space are:
  - 1 each spanning set for  $\mathcal{V}$  contains a basis;
  - 2 each minimal spanning set is a basis;
  - 3 each linearly independent set of vectors is contained in a basis;
  - 4 each maximal set of linearly independent vectors is a basis;

## Example: vector space of polynomials of degree $\leq n$

A linear space we make often use of in **Computer Graphics** and **Geometric modeling** is the space of dimension  $n + 1$ :

$$\mathbb{P}^n(\mathbb{R}) = \{p : \mathbb{R} \rightarrow \mathbb{R} : u \mapsto \sum_{i=1}^n a_i p^i, a_i \in \mathbb{R}\}$$

of univariate **polynomials of degree  $\leq n$**  on the real field (with real coefficients), with  $p^i \in P_n$ , where

$$P_n = (p^n, p^{n-1}, \dots, p^1, p^0) \quad \text{and} \quad p^i : u \mapsto u^i$$

is **the power basis**.

# Components

If  $(\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n)$  is an ordered basis for  $\mathcal{V}$ , then for each  $\mathbf{v} \in \mathcal{V}$  there exists a **unique**  $n$ -tuple of scalars  $\alpha_1, \alpha_2, \dots, \alpha_n \in \mathcal{F}$  such that

$$\mathbf{v} = \sum_{i=1}^n \alpha_i \mathbf{e}_i.$$



# Components

The  $n$ -tuple of scalars  $(\alpha_i)$  is called the **components** of  $\mathbf{v}$  with respect to the ordered basis  $(\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n)$ .

- If such a  $n$ -tuple were not unique, then  $\mathbf{v} = \sum \alpha_i \mathbf{e}_i = \sum \beta_i \mathbf{e}_i$

# Components

The  $n$ -tuple of scalars  $(\alpha_i)$  is called the **components** of  $\mathbf{v}$  with respect to the ordered basis  $(\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n)$ .

- If such a  $n$ -tuple were not unique, then  $\mathbf{v} = \sum \alpha_i \mathbf{e}_i = \sum \beta_i \mathbf{e}_i$
- But this one would imply  $\sum (\alpha_i - \beta_i) \mathbf{e}_i = \mathbf{0}$ , hence  $(\alpha_i - \beta_i) = 0$ ,

# Components

The  $n$ -tuple of scalars  $(\alpha_i)$  is called the **components** of  $\mathbf{v}$  with respect to the ordered basis  $(\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n)$ .

- If such a  $n$ -tuple were not unique, then  $\mathbf{v} = \sum \alpha_i \mathbf{e}_i = \sum \beta_i \mathbf{e}_i$
- But this one would imply  $\sum (\alpha_i - \beta_i) \mathbf{e}_i = \mathbf{0}$ , hence  $(\alpha_i - \beta_i) = 0$ ,
- i.e.  $\alpha_i = \beta_i$ , for every  $i$ .

# Change of basis

- Let  $B = (\mathbf{e}_1, \dots, \mathbf{e}_n) \subset \mathcal{V}$  be a basis for  $\mathcal{V}$ .

# Change of basis

- Let  $B = (\mathbf{e}_1, \dots, \mathbf{e}_n) \subset \mathcal{V}$  be a basis for  $\mathcal{V}$ .
- Of course, their coordinates are  $(1 \ 0 \ \dots \ 0), (0 \ 1 \ \dots \ 0), \dots, (0 \ 0 \ \dots \ 1)$ , and, in  $B$  coordinates, the basis is represented by the matrix

$$[B] = [I]$$

.

# Change of basis

- Let  $B = (\mathbf{e}_1, \dots, \mathbf{e}_n) \subset \mathcal{V}$  be a basis for  $\mathcal{V}$ .
- Of course, their coordinates are  $(1 \ 0 \ \dots \ 0), (0 \ 1 \ \dots \ 0), \dots, (0 \ 0 \ \dots \ 1)$ , and, in  $B$  coordinates, the basis is represented by the matrix

$$[B] = [I]$$

- If we take  $n$  (linearly independent) vectors  $V = (\mathbf{v}_1, \dots, \mathbf{v}_n) \subset \mathcal{V}$ , represented in  $B$  coordinates as  $[V]$ , and want to parametrize  $\mathcal{V}$  with respect to the new basis, we have, for transformation of coordinates:

$$[I] = [T][V]$$

# Change of basis

- Let  $B = (\mathbf{e}_1, \dots, \mathbf{e}_n) \subset \mathcal{V}$  be a basis for  $\mathcal{V}$ .
- Of course, their coordinates are  $(1 \ 0 \ \dots \ 0), (0 \ 1 \ \dots \ 0), \dots, (0 \ 0 \ \dots \ 1)$ , and, in  $B$  coordinates, the basis is represented by the matrix

$$[B] = [I]$$

- .
- If we take  $n$  (linearly independent) vectors  $V = (\mathbf{v}_1, \dots, \mathbf{v}_n) \subset \mathcal{V}$ , represented in  $B$  coordinates as  $[V]$ , and want to parametrize  $\mathcal{V}$  with respect to the new basis, we have, for transformation of coordinates:

$$[I] = [T][V]$$

- and hence:

$$[T] = [V]^{-1}$$

## Example: two polynomial bases

- Let  $P_3 = (u^3, u^2, u, 1)$



## Example: two polynomial bases

- Let  $P_3 = (u^3, u^2, u, 1)$
- and  $B_3 = ((1 - u)^3, 3u(1 - u)^2, 3u^2(1 - u), u^3)$  be two ordered bases

## Example: two polynomial bases

- Let  $P_3 = (u^3, u^2, u, 1)$
- and  $B_3 = ((1-u)^3, 3u(1-u)^2, 3u^2(1-u), u^3)$  be two ordered bases
- for the linear space  $\mathbb{P}^3(\mathbb{R})$  of polynomials with  $\deg \leq 3$ .

## Example: two polynomial bases

- Let  $P_3 = (u^3, u^2, u, 1)$
- and  $B_3 = ((1-u)^3, 3u(1-u)^2, 3u^2(1-u), u^3)$  be two ordered bases
- for the linear space  $\mathbb{P}^3(\mathbb{R})$  of polynomials with  $\deg \leq 3$ .
- the  $[B_3]$  matrix in the  $P_3$  basis is

$$[B_3]_{P_3} = \begin{pmatrix} -1 & 3 & -3 & 1 \\ 3 & -6 & 3 & 0 \\ -3 & 3 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$$

## Example: two polynomial bases

- Let  $P_3 = (u^3, u^2, u, 1)$
- and  $B_3 = ((1-u)^3, 3u(1-u)^2, 3u^2(1-u), u^3)$  be two ordered bases
- for the linear space  $\mathbb{P}^3(\mathbb{R})$  of polynomials with  $\deg \leq 3$ .
- the  $[B_3]$  matrix in the  $P_3$  basis is

$$[B_3]_{P_3} = \begin{pmatrix} -1 & 3 & -3 & 1 \\ 3 & -6 & 3 & 0 \\ -3 & 3 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$$

- the  $[P_3]$  matrix in the  $B_3$  basis is

$$[P_3]_{B_3} = [B_3]_{P_3}^{-1} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1/3 & 1 \\ 0 & 1/3 & 1/6 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix}$$

## Example: two polynomial bases

- Let  $P_3 = (u^3, u^2, u, 1)$
- and  $B_3 = ((1-u)^3, 3u(1-u)^2, 3u^2(1-u), u^3)$  be two ordered bases
- for the linear space  $\mathbb{P}^3(\mathbb{R})$  of polynomials with  $\deg \leq 3$ .
- the  $[B_3]$  matrix in the  $P_3$  basis is

$$[B_3]_{P_3} = \begin{pmatrix} -1 & 3 & -3 & 1 \\ 3 & -6 & 3 & 0 \\ -3 & 3 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$$

- the  $[P_3]$  matrix in the  $B_3$  basis is

$$[P_3]_{B_3} = [B_3]_{P_3}^{-1} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1/3 & 1 \\ 0 & 1/3 & 1/6 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix}$$

- WHY ?

# Example

*# Open a file*

```
fo = open("foo.txt", "r+")
```

```
str = fo.read(10);
```

```
print "Read String is : ", str
```

*# Check current position*

```
position = fo.tell();
```

```
print "Current file position : ", position
```

*# Reposition pointer at the beginning once again*

```
position = fo.seek(0, 0);
```

```
str = fo.read(10);
```

```
print "Again read String is : ", str
```

*# Close opened file*

```
fo.close()
```

# Affine spaces

# Affine space

The idea of affine space corresponds to that of a set of points where the **displacement** from a point  $\mathbf{x}$  to another point  $\mathbf{y}$  is obtained by summing a vector  $\mathbf{v}$  to the  $\mathbf{x}$  point.



# Definition

A set  $\mathcal{A}$  of points is called an **affine space** modeled on the vector space  $\mathcal{V}$  if there is a function

$$\mathcal{A} \times \mathcal{V} \rightarrow \mathcal{A} : (\mathbf{x}, \mathbf{v}) \mapsto \mathbf{x} + \mathbf{v}$$

called **affine action**, with the properties:

# Definition

A set  $\mathcal{A}$  of points is called an **affine space** modeled on the vector space  $\mathcal{V}$  if there is a function

$$\mathcal{A} \times \mathcal{V} \rightarrow \mathcal{A} : (\mathbf{x}, \mathbf{v}) \mapsto \mathbf{x} + \mathbf{v}$$

called **affine action**, with the properties:

- 1  $(\mathbf{x} + \mathbf{v}) + \mathbf{w} = \mathbf{x} + (\mathbf{v} + \mathbf{w})$  for each  $\mathbf{x} \in \mathcal{A}$  and each  $\mathbf{v}, \mathbf{w} \in \mathcal{V}$ ;

# Definition

A set  $\mathcal{A}$  of points is called an **affine space** modeled on the vector space  $\mathcal{V}$  if there is a function

$$\mathcal{A} \times \mathcal{V} \rightarrow \mathcal{A} : (\mathbf{x}, \mathbf{v}) \mapsto \mathbf{x} + \mathbf{v}$$

called **affine action**, with the properties:

- ①  $(\mathbf{x} + \mathbf{v}) + \mathbf{w} = \mathbf{x} + (\mathbf{v} + \mathbf{w})$  for each  $\mathbf{x} \in \mathcal{A}$  and each  $\mathbf{v}, \mathbf{w} \in \mathcal{V}$ ;
- ②  $\mathbf{x} + \mathbf{0} = \mathbf{x}$  for each  $\mathbf{x} \in \mathcal{A}$ , where  $\mathbf{0} \in \mathcal{V}$  is the null vector;

# Definition

A set  $\mathcal{A}$  of points is called an **affine space** modeled on the vector space  $\mathcal{V}$  if there is a function

$$\mathcal{A} \times \mathcal{V} \rightarrow \mathcal{A} : (\mathbf{x}, \mathbf{v}) \mapsto \mathbf{x} + \mathbf{v}$$

called **affine action**, with the properties:


- ①  $(\mathbf{x} + \mathbf{v}) + \mathbf{w} = \mathbf{x} + (\mathbf{v} + \mathbf{w})$  for each  $\mathbf{x} \in \mathcal{A}$  and each  $\mathbf{v}, \mathbf{w} \in \mathcal{V}$ ;
- ②  $\mathbf{x} + \mathbf{0} = \mathbf{x}$  for each  $\mathbf{x} \in \mathcal{A}$ , where  $\mathbf{0} \in \mathcal{V}$  is the null vector;
- ③ for each pair  $\mathbf{x}, \mathbf{y} \in \mathcal{A}$  there is a unique  $(\mathbf{y} - \mathbf{x}) \in \mathcal{V}$  such that

$$\mathbf{x} + (\mathbf{y} - \mathbf{x}) = \mathbf{y}.$$

# Dimension

The affine space  $\mathcal{A}$  is said of **dimension**  $n$  if modeled on a vector space  $\mathcal{V}$  of dimension  $n$ .

# Vector sum vs affine action



**Figure:** (a) Vector sum and difference are given by the parallelogram rule  
 (b) associativity of displacement (point and vector sum) in an affine space

# Operations on vectors and points

- The **addition** of vectors is a primitive operation in a vector space.

# Operations on vectors and points

- The **addition** of vectors is a primitive operation in a vector space.
- The **difference** of vectors is defined through the two primitive operations:

$$\mathbf{v}_1 - \mathbf{v}_2 = \mathbf{v}_1 + (-1)\mathbf{v}_2.$$



# Operations on vectors and points

- The **addition** of vectors is a primitive operation in a vector space.
- The **difference** of vectors is defined through the two primitive operations:

$$\mathbf{v}_1 - \mathbf{v}_2 = \mathbf{v}_1 + (-1)\mathbf{v}_2.$$

- Addition and difference of vectors are geometrically produced by the **parallelogram rule**

# Operations on vectors and points

- The **addition** of vectors is a primitive operation in a vector space.
- The **difference** of vectors is defined through the two primitive operations:

$$\mathbf{v}_1 - \mathbf{v}_2 = \mathbf{v}_1 + (-1)\mathbf{v}_2.$$

- Addition and difference of vectors are geometrically produced by the **parallelogram rule**
- notice also the **associative property** of the affine action on a point space.

# Operations on vectors and points

The sum of a set  $\{\mathbf{v}_i\}$  of vectors ( $i = 1, \dots, n$ ) can be geometrically obtained, in an affine space:

- by setting  $\mathbf{p}_0 = \mathbf{0}$

Remark: operations on points

# Operations on vectors and points

The sum of a set  $\{\mathbf{v}_i\}$  of vectors ( $i = 1, \dots, n$ ) can be geometrically obtained, in an affine space:

- by setting  $\mathbf{p}_0 = \mathbf{0}$
- $\mathbf{p}_i = \mathbf{p}_{i-1} + \mathbf{v}_i$ ,

Remark: operations on points

# Operations on vectors and points

The sum of a set  $\{\mathbf{v}_i\}$  of vectors ( $i = 1, \dots, n$ ) can be geometrically obtained, in an affine space:

- by setting  $\mathbf{p}_0 = \mathbf{0}$
- $\mathbf{p}_i = \mathbf{p}_{i-1} + \mathbf{v}_i$ ,
- so that

$$\sum_i \mathbf{v}_i = \mathbf{p}_n - \mathbf{p}_0$$

Remark: operations on points

# Operations on vectors and points

The sum of a set  $\{\mathbf{v}_i\}$  of vectors ( $i = 1, \dots, n$ ) can be geometrically obtained, in an affine space:

- by setting  $\mathbf{p}_0 = \mathbf{0}$
- $\mathbf{p}_i = \mathbf{p}_{i-1} + \mathbf{v}_i$ ,
- so that

$$\sum_i \mathbf{v}_i = \mathbf{p}_n - \mathbf{p}_0$$

Remark: operations on points

- 1 the addition of points is not defined;

# Operations on vectors and points

The sum of a set  $\{\mathbf{v}_i\}$  of vectors ( $i = 1, \dots, n$ ) can be geometrically obtained, in an affine space:

- by setting  $\mathbf{p}_0 = \mathbf{0}$
- $\mathbf{p}_i = \mathbf{p}_{i-1} + \mathbf{v}_i$ ,
- so that

$$\sum_i \mathbf{v}_i = \mathbf{p}_n - \mathbf{p}_0$$

Remark: operations on points

- 1 the addition of points is **not** defined;
- 2 the difference of two points is a **vector**;

# Operations on vectors and points

The sum of a set  $\{\mathbf{v}_i\}$  of vectors ( $i = 1, \dots, n$ ) can be geometrically obtained, in an affine space:

- by setting  $\mathbf{p}_0 = \mathbf{0}$
- $\mathbf{p}_i = \mathbf{p}_{i-1} + \mathbf{v}_i$ ,
- so that

$$\sum_i \mathbf{v}_i = \mathbf{p}_n - \mathbf{p}_0$$

Remark: operations on points

- 1 the **addition of points** is **not** defined;
- 2 the **difference of two points** is a **vector**;
- 3 the **sum of a point and a vector** is a **point**.



# Affine combinations

# Positive, affine and convex combinations

Three types of combinations of vectors or points can be defined. They lead to the concepts of **cones**, **hyperplanes** and **convex sets**, respectively.

# Positive combination

Let  $\mathbf{v}_0, \dots, \mathbf{v}_d \in \mathbb{R}^n$  and  $\alpha_0, \dots, \alpha_d \in \mathbb{R}^+ \cup \{0\}$ .

The vector

$$\alpha_0 \mathbf{v}_0 + \dots + \alpha_d \mathbf{v}_d = \sum_{i=0}^d \alpha_i \mathbf{v}_i$$

is called a **positive combination** of such vectors.

The set of all the positive combinations of  $\{\mathbf{v}_0, \dots, \mathbf{v}_d\}$  is called the **positive hull** of  $\{\mathbf{v}_0, \dots, \mathbf{v}_d\}$  and denoted **pos**  $\{\mathbf{v}_0, \dots, \mathbf{v}_d\}$ .

This set is also called the **cone** generated by the given vectors

# Affine combination

Let  $\mathbf{p}_0, \dots, \mathbf{p}_d \in \mathbb{E}^n$  and  $\alpha_0, \dots, \alpha_d \in \mathbb{R}$ , such that  $\alpha_0 + \dots + \alpha_d = 1$ .

The point

$$\sum_{i=0}^d \alpha_i \mathbf{p}_i := \mathbf{p}_0 + \sum_{i=1}^d \alpha_i (\mathbf{p}_i - \mathbf{p}_0)$$

is called an **affine combination** of the points  $\mathbf{p}_0, \dots, \mathbf{p}_d$ .

# Affine combination

The set of all affine combinations of  $\{\mathbf{p}_0, \dots, \mathbf{p}_d\}$  is an **affine subspace**, denoted by  $\text{aff}\{\mathbf{p}_0, \dots, \mathbf{p}_d\}$

It is easy to verify that:

$$\text{aff}\{\mathbf{p}_0, \dots, \mathbf{p}_d\} = \mathbf{p}_0 + \text{lin}\{\mathbf{p}_1 - \mathbf{p}_0, \dots, \mathbf{p}_d - \mathbf{p}_0\}.$$

# Affine combination

- 1 The **dimension** of an affine subspace is the dimension of the corresponding linear vector space.

# Affine combination

- 1 The **dimension** of an affine subspace is the dimension of the corresponding linear vector space.
- 2 Affine subspaces of  $\mathbb{E}^d$  with dimensions 0, 1, 2 and  $d - 1$  are called **points**, **lines**, **planes** and **hyperplanes**, respectively.

# Affine combination

- 1 The **dimension** of an affine subspace is the dimension of the corresponding linear vector space.
- 2 Affine subspaces of  $\mathbb{E}^d$  with dimensions 0, 1, 2 and  $d - 1$  are called **points**, **lines**, **planes** and **hyperplanes**, respectively.
- 3 Affine subspaces are also called **flats**.



# Affine combination

- 1 The **dimension** of an affine subspace is the dimension of the corresponding linear vector space.
- 2 Affine subspaces of  $\mathbb{E}^d$  with dimensions 0, 1, 2 and  $d - 1$  are called **points**, **lines**, **planes** and **hyperplanes**, respectively.
- 3 Affine subspaces are also called **flats**.

# Affine combination

- 1 The **dimension** of an affine subspace is the dimension of the corresponding linear vector space.
- 2 Affine subspaces of  $\mathbb{E}^d$  with dimensions 0, 1, 2 and  $d - 1$  are called **points**, **lines**, **planes** and **hyperplanes**, respectively.
- 3 Affine subspaces are also called **flats**.

## Double description

Every affine subspace can be described either as

- the **intersection** of affine **hyperplanes**, or as

# Affine combination

- 1 The **dimension** of an affine subspace is the dimension of the corresponding linear vector space.
- 2 Affine subspaces of  $\mathbb{E}^d$  with dimensions 0, 1, 2 and  $d - 1$  are called **points**, **lines**, **planes** and **hyperplanes**, respectively.
- 3 Affine subspaces are also called **flats**.

## Double description

Every affine subspace can be described either as

- the **intersection** of affine **hyperplanes**, or as
- the **affine hull** of a finite set of **points**.

# Convex combinations

# Convex combination

Let  $\mathbf{p}_0, \dots, \mathbf{p}_d \in \mathbb{E}^n$  and  $\alpha_0, \dots, \alpha_d \geq 0$ , with  $\alpha_0 + \dots + \alpha_d = 1$ .

The point

$$\alpha_0 \mathbf{p}_0 + \dots + \alpha_d \mathbf{p}_d = \sum_{i=0}^d \alpha_i \mathbf{p}_i$$

is called a **convex combination** of points  $\mathbf{p}_0, \dots, \mathbf{p}_d$ .

A **convex** combinations is both **positive** and **affine**.

# Convex hull

The set of **all** convex combinations of  $\{\mathbf{p}_0, \dots, \mathbf{p}_d\}$  is a convex set, called **convex hull** of  $\{\mathbf{p}_0, \dots, \mathbf{p}_d\}$ , and is denoted by **conv**  $\{\mathbf{p}_0, \dots, \mathbf{p}_d\}$ .

# Convex hull

The set of **all** convex combinations of  $\{\mathbf{p}_0, \dots, \mathbf{p}_d\}$  is a convex set, called **convex hull** of  $\{\mathbf{p}_0, \dots, \mathbf{p}_d\}$ , and is denoted by  $\text{conv}\{\mathbf{p}_0, \dots, \mathbf{p}_d\}$ .

## Properties

- the convex hull of a set of points is the **intersection of all convex sets** that contain them

# Convex hull

The set of **all** convex combinations of  $\{\mathbf{p}_0, \dots, \mathbf{p}_d\}$  is a convex set, called **convex hull** of  $\{\mathbf{p}_0, \dots, \mathbf{p}_d\}$ , and is denoted by  $\text{conv}\{\mathbf{p}_0, \dots, \mathbf{p}_d\}$ .

## Properties

- the convex hull of a set of points is the **intersection of all convex sets** that contain them
- the convex hull of a set of points is the **smallest set** that contains them



# ASSIGNMENT

- Produce (and draw) 100 random points within the unit square  $[0, 1]^2$ ;

# ASSIGNMENT

- Produce (and draw) 100 random points within the unit square  $[0, 1]^2$ ;
- Produce (and draw) 1000 random points within  $S_1$ , the 1D sphere (circle) of unit radius centered at the origin  $(0, 0)$ ;

# References

Linear Algebra Done Right book

NumPy tutorial