

Computational Graphics: Lecture 4

The Cvdlab Team

Mon, Mar 10, 2014

Outline: Algebra1

- 1 Linear spaces
- 2 Linear combinations
- 3 Subspaces
- 4 Spans
- 5 Bases

Linear spaces

Definition

A **linear** (or **vector**) **space** \mathcal{V} over a field \mathcal{F} is a set with two composition rules, such that, for each $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$ and for each $\alpha, \beta \in \mathcal{F}$, the rules $+$, \cdot satisfy the following axioms:

$$\textcircled{1} \quad \mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v}; \quad (\text{commutativity of addition})$$

Definition

A **linear** (or **vector**) **space** \mathcal{V} over a field \mathcal{F} is a set with two composition rules, such that, for each $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$ and for each $\alpha, \beta \in \mathcal{F}$, the rules $+$, \cdot satisfy the following axioms:

① $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v};$

(commutativity of addition)

② $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w};$

(associativity of addition)

Definition

A **linear** (or **vector**) **space** \mathcal{V} over a field \mathcal{F} is a set with two composition rules, such that, for each $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$ and for each $\alpha, \beta \in \mathcal{F}$, the rules $+$, \cdot satisfy the following axioms:

- ① $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v};$ (commutativity of addition)
- ② $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w};$ (associativity of addition)
- ③ there is a $\mathbf{0} \in \mathcal{V}$ such that $\mathbf{v} + \mathbf{0} = \mathbf{v};$ (neutral el. of addition)

Definition

A **linear** (or **vector**) **space** \mathcal{V} over a field \mathcal{F} is a set with two composition rules, such that, for each $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$ and for each $\alpha, \beta \in \mathcal{F}$, the rules $+$, \cdot satisfy the following axioms:

- ① $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v};$ (commutativity of addition)
- ② $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w};$ (associativity of addition)
- ③ there is a $\mathbf{0} \in \mathcal{V}$ such that $\mathbf{v} + \mathbf{0} = \mathbf{v};$ (neutral el. of addition)
- ④ there is a $-\mathbf{v} \in \mathcal{V}$ such that $\mathbf{v} + (-\mathbf{v}) = \mathbf{0};$ (inverse of add.)

Definition

A **linear** (or **vector**) **space** \mathcal{V} over a field \mathcal{F} is a set with two composition rules, such that, for each $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$ and for each $\alpha, \beta \in \mathcal{F}$, the rules $+$, \cdot satisfy the following axioms:

- ① $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v};$ (commutativity of addition)
- ② $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w};$ (associativity of addition)
- ③ there is a $\mathbf{0} \in \mathcal{V}$ such that $\mathbf{v} + \mathbf{0} = \mathbf{v};$ (neutral el. of addition)
- ④ there is a $-\mathbf{v} \in \mathcal{V}$ such that $\mathbf{v} + (-\mathbf{v}) = \mathbf{0};$ (inverse of add.)
- ⑤ $\alpha \cdot (\mathbf{v} + \mathbf{w}) = \alpha \cdot \mathbf{v} + \alpha \cdot \mathbf{w};$ (distrib. of addition w.r.t. product)

Definition

A **linear** (or **vector**) **space** \mathcal{V} over a field \mathcal{F} is a set with two composition rules, such that, for each $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$ and for each $\alpha, \beta \in \mathcal{F}$, the rules $+$, \cdot satisfy the following axioms:

- ① $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v};$ (commutativity of addition)
- ② $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w};$ (associativity of addition)
- ③ there is a $\mathbf{0} \in \mathcal{V}$ such that $\mathbf{v} + \mathbf{0} = \mathbf{v};$ (neutral el. of addition)
- ④ there is a $-\mathbf{v} \in \mathcal{V}$ such that $\mathbf{v} + (-\mathbf{v}) = \mathbf{0};$ (inverse of add.)
- ⑤ $\alpha \cdot (\mathbf{v} + \mathbf{w}) = \alpha \cdot \mathbf{v} + \alpha \cdot \mathbf{w};$ (distrib. of addition w.r.t. product)
- ⑥ $(\alpha + \beta) \cdot \mathbf{v} = \alpha \cdot \mathbf{v} + \beta \cdot \mathbf{v};$ (distrib. of product w.r.t. addition)

Definition

A **linear** (or **vector**) **space** \mathcal{V} over a field \mathcal{F} is a set with two composition rules, such that, for each $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$ and for each $\alpha, \beta \in \mathcal{F}$, the rules $+$, \cdot satisfy the following axioms:

- ① $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v};$ (commutativity of addition)
- ② $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w};$ (associativity of addition)
- ③ there is a $\mathbf{0} \in \mathcal{V}$ such that $\mathbf{v} + \mathbf{0} = \mathbf{v};$ (neutral el. of addition)
- ④ there is a $-\mathbf{v} \in \mathcal{V}$ such that $\mathbf{v} + (-\mathbf{v}) = \mathbf{0};$ (inverse of add.)
- ⑤ $\alpha \cdot (\mathbf{v} + \mathbf{w}) = \alpha \cdot \mathbf{v} + \alpha \cdot \mathbf{w};$ (distrib. of addition w.r.t. product)
- ⑥ $(\alpha + \beta) \cdot \mathbf{v} = \alpha \cdot \mathbf{v} + \beta \cdot \mathbf{v};$ (distrib. of product w.r.t. addition)
- ⑦ $\alpha \cdot (\beta \cdot \mathbf{v}) = (\alpha\beta) \cdot \mathbf{v};$ (associativity of product)

Definition

A **linear** (or **vector**) **space** \mathcal{V} over a field \mathcal{F} is a set with two composition rules, such that, for each $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathcal{V}$ and for each $\alpha, \beta \in \mathcal{F}$, the rules $+$, \cdot satisfy the following axioms:

- ① $\mathbf{v} + \mathbf{w} = \mathbf{w} + \mathbf{v};$ (commutativity of addition)
- ② $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w};$ (associativity of addition)
- ③ there is a $\mathbf{0} \in \mathcal{V}$ such that $\mathbf{v} + \mathbf{0} = \mathbf{v};$ (neutral el. of addition)
- ④ there is a $-\mathbf{v} \in \mathcal{V}$ such that $\mathbf{v} + (-\mathbf{v}) = \mathbf{0};$ (inverse of add.)
- ⑤ $\alpha \cdot (\mathbf{v} + \mathbf{w}) = \alpha \cdot \mathbf{v} + \alpha \cdot \mathbf{w};$ (distrib. of addition w.r.t. product)
- ⑥ $(\alpha + \beta) \cdot \mathbf{v} = \alpha \cdot \mathbf{v} + \beta \cdot \mathbf{v};$ (distrib. of product w.r.t. addition)
- ⑦ $\alpha \cdot (\beta \cdot \mathbf{v}) = (\alpha\beta) \cdot \mathbf{v};$ (associativity of product)
- ⑧ $1 \cdot \mathbf{v} = \mathbf{v}.$ (neutral element of product)

Example: vector space of real matrices

Let $\mathcal{M}_n^m(\mathbb{R})$ be the set of $m \times n$ matrices with elements in the field \mathbb{R} . An element A in such a set is denoted as

$$A = (\alpha_{ij})$$

Example: vector space of real matrices

Let $\mathcal{M}_n^m(\mathbb{R})$ be the set of $m \times n$ matrices with elements in the field \mathbb{R} . An element A in such a set is denoted as

$$A = (\alpha_{ij})$$

Addition and **multiplication by a scalar** are defined component-wise:

$$A + B = (\alpha_{ij}) + (\beta_{ij}) = (\alpha_{ij} + \beta_{ij})$$

$$\gamma A = \gamma(\alpha_{ij}) = (\gamma\alpha_{ij})$$

Linear combinations

Linear combination

Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \in \mathcal{V}$ and $\alpha_1, \alpha_2, \dots, \alpha_n \in \mathcal{F}$,

The vector

$$\alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n = \sum_{i=1}^n \alpha_i \mathbf{v}_i \in \mathcal{V}$$

is called a **linear combination** of vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ with scalars $\alpha_1, \alpha_2, \dots, \alpha_n \in \mathcal{F}$

Subspaces

Subspace

Let $(\mathcal{V}, +, \cdot)$ be a vector space on the field \mathcal{F} .

$\mathcal{U} \subset \mathcal{V}$ is a **subspace** of \mathcal{V} if $(\mathcal{U}, +, \cdot)$ is a vector space with respect to the same operations.

Subspace

Let $(\mathcal{V}, +, \cdot)$ be a vector space on the field \mathcal{F} .

$\mathcal{U} \subset \mathcal{V}$ is a **subspace** of \mathcal{V} if $(\mathcal{U}, +, \cdot)$ is a vector space with respect to the same operations.

$\mathcal{U} \subset \mathcal{V}$ is a **subspace** of \mathcal{V} if and only if

$$\mathcal{U} \neq \emptyset;$$

$$\text{for each } \alpha \in \mathcal{F} \text{ and } \mathbf{u}_1, \mathbf{u}_2 \in \mathcal{U}, \alpha \mathbf{u}_1 + \mathbf{u}_2 \in \mathcal{U}$$

Subspace

Let $(\mathcal{V}, +, \cdot)$ be a vector space on the field \mathcal{F} .

$\mathcal{U} \subset \mathcal{V}$ is a **subspace** of \mathcal{V} if $(\mathcal{U}, +, \cdot)$ is a vector space with respect to the same operations.

$\mathcal{U} \subset \mathcal{V}$ is a **subspace** of \mathcal{V} if and only if

$$\mathcal{U} \neq \emptyset;$$

$$\text{for each } \alpha \in \mathcal{F} \text{ and } \mathbf{u}_1, \mathbf{u}_2 \in \mathcal{U}, \alpha \mathbf{u}_1 + \mathbf{u}_2 \in \mathcal{U}$$

codimension of a subspace $\mathcal{U} \subset \mathcal{V}$
is defined as $\dim \mathcal{V} - \dim \mathcal{U}$

Subspace

Let $(\mathcal{V}, +, \cdot)$ be a vector space on the field \mathcal{F} .

$\mathcal{U} \subset \mathcal{V}$ is a **subspace** of \mathcal{V} if $(\mathcal{U}, +, \cdot)$ is a vector space with respect to the same operations.

$\mathcal{U} \subset \mathcal{V}$ is a **subspace** of \mathcal{V} if and only if

$$\mathcal{U} \neq \emptyset;$$

$$\text{for each } \alpha \in \mathcal{F} \text{ and } \mathbf{u}_1, \mathbf{u}_2 \in \mathcal{U}, \alpha \mathbf{u}_1 + \mathbf{u}_2 \in \mathcal{U}$$

codimension of a subspace $\mathcal{U} \subset \mathcal{V}$
is defined as $\dim \mathcal{V} - \dim \mathcal{U}$

Subspace

Let $(\mathcal{V}, +, \cdot)$ be a vector space on the field \mathcal{F} .

$\mathcal{U} \subset \mathcal{V}$ is a **subspace** of \mathcal{V} if $(\mathcal{U}, +, \cdot)$ is a vector space with respect to the same operations.

$\mathcal{U} \subset \mathcal{V}$ is a **subspace** of \mathcal{V} if and only if

$$\mathcal{U} \neq \emptyset;$$

$$\text{for each } \alpha \in \mathcal{F} \text{ and } \mathbf{u}_1, \mathbf{u}_2 \in \mathcal{U}, \alpha \mathbf{u}_1 + \mathbf{u}_2 \in \mathcal{U}$$

codimension of a subspace $\mathcal{U} \subset \mathcal{V}$
is defined as $\dim \mathcal{V} - \dim \mathcal{U}$

Question

Examples of codimension? in 1D, 2D, 3D

Spans

Span

- The set of all linear combinations of elements of a set $S \subset \mathcal{V}$ is a subspace of \mathcal{V} .

Span

- The set of all linear combinations of elements of a set $S \subset \mathcal{V}$ is a subspace of \mathcal{V} .
- Such a subspace is called the **span of S** and is denoted as

$$\text{lin } S$$

Span

- The set of all linear combinations of elements of a set $S \subset \mathcal{V}$ is a subspace of \mathcal{V} .
- Such a subspace is called the **span of S** and is denoted as

$$\text{lin } S$$

- If a subspace \mathcal{U} of \mathcal{V} can be generated as the span of a set S of vectors in \mathcal{V} , then S is called a **generating set** or a **spanning set** for \mathcal{U} .

Linear independence

- A set of vectors $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ is **linearly independent** if

$$\sum_{i=1}^n \alpha_i \mathbf{v}_i = \mathbf{0}$$

implies that $\alpha_i = 0$ for each i

Linear independence

- A set of vectors $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ is **linearly independent** if

$$\sum_{i=1}^n \alpha_i \mathbf{v}_i = \mathbf{0}$$

implies that $\alpha_i = 0$ for each i

- As a consequence, **a set of vectors is linearly independent** when none of them belongs to the span of the others.

Bases

Bases and coordinates

When working with vector spaces, the concept of **basis**, a **discrete subset of linearly independent elements**, is probably the most useful to deal with.

- each element of the space can be **represented uniquely as linear combination of basis elements**

Bases and coordinates

When working with vector spaces, the concept of **basis**, a **discrete subset of linearly independent elements**, is probably the most useful to deal with.

- each element of the space can be **represented uniquely as linear combination of basis elements**
- this leads to a **parametrization** of the space, i.e. to **represent each element by a sequence of scalars**, called its **coordinates** with respect to the chosen basis.

Bases

A set of vectors $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ is a **basis** for the vector space \mathcal{V} iff

- 1 the set is linearly independent, and

Bases

A set of vectors $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ is a **basis** for the vector space \mathcal{V} iff

- 1 the set is linearly independent, and
- 2 $\mathcal{V} = \text{lin} \{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$

Bases

- Every two bases of \mathcal{V} have the same number of elements, that is called the **dimension** of \mathcal{V} and is denoted

$$\dim \mathcal{V}$$

Bases

- Every two bases of \mathcal{V} have the same number of elements, that is called the **dimension** of \mathcal{V} and is denoted

$$\dim \mathcal{V}$$

- Some important **properties** of the bases of a vector space are:

Bases

- Every two bases of \mathcal{V} have the same number of elements, that is called the **dimension** of \mathcal{V} and is denoted

$$\dim \mathcal{V}$$

- Some important **properties** of the bases of a vector space are:
 - ① each spanning set for \mathcal{V} contains a basis;

Bases

- Every two bases of \mathcal{V} have the same number of elements, that is called the **dimension** of \mathcal{V} and is denoted

$$\dim \mathcal{V}$$

- Some important **properties** of the bases of a vector space are:
 - 1 each spanning set for \mathcal{V} contains a basis;
 - 2 each minimal spanning set is a basis;

Bases

- Every two bases of \mathcal{V} have the same number of elements, that is called the **dimension** of \mathcal{V} and is denoted

$$\dim \mathcal{V}$$

- Some important **properties** of the bases of a vector space are:
 - 1 each spanning set for \mathcal{V} contains a basis;
 - 2 each minimal spanning set is a basis;
 - 3 each linearly independent set of vectors is contained in a basis;

Bases

- Every two bases of \mathcal{V} have the same number of elements, that is called the **dimension** of \mathcal{V} and is denoted

$$\dim \mathcal{V}$$

- Some important **properties** of the bases of a vector space are:
 - 1 each spanning set for \mathcal{V} contains a basis;
 - 2 each minimal spanning set is a basis;
 - 3 each linearly independent set of vectors is contained in a basis;
 - 4 each maximal set of linearly independent vectors is a basis;

Example: vector space of polynomials of degree $\leq n$

A linear space we make often use of in **Computer Graphics** and **Geometric modeling** is the space of dimension $n + 1$:

$$\mathbb{P}^n(\mathbb{R}) = \{p : \mathbb{R} \rightarrow \mathbb{R} : u \mapsto \sum_{i=1}^n a_i p^i, a_i \in \mathbb{R}\}$$

of univariate **polynomials of degree $\leq n$** on the real field (with real coefficients), with $p^i \in P_n$, where

$$P_n = (p^n, p^{n-1}, \dots, p^1, p^0) \quad \text{and} \quad p^i : u \mapsto u^i$$

is **the power basis**.

Components

If $(\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n)$ is an ordered basis for \mathcal{V} , then for each $\mathbf{v} \in \mathcal{V}$ there exists a **unique** n -tuple of scalars $\alpha_1, \alpha_2, \dots, \alpha_n \in \mathcal{F}$ such that

$$\mathbf{v} = \sum_{i=1}^n \alpha_i \mathbf{e}_i.$$

Components

The n -tuple of scalars (α_i) is called the **components** of \mathbf{v} with respect to the ordered basis $(\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n)$.

- If such a n -tuple were not unique, then $\mathbf{v} = \sum \alpha_i \mathbf{e}_i = \sum \beta_i \mathbf{e}_i$

Components

The n -tuple of scalars (α_i) is called the **components** of \mathbf{v} with respect to the ordered basis $(\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n)$.

- If such a n -tuple were not unique, then $\mathbf{v} = \sum \alpha_i \mathbf{e}_i = \sum \beta_i \mathbf{e}_i$
- But this one would imply $\sum (\alpha_i - \beta_i) \mathbf{e}_i = \mathbf{0}$, hence $(\alpha_i - \beta_i) = 0$,

Components

The n -tuple of scalars (α_i) is called the **components** of \mathbf{v} with respect to the ordered basis $(\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n)$.

- If such a n -tuple were not unique, then $\mathbf{v} = \sum \alpha_i \mathbf{e}_i = \sum \beta_i \mathbf{e}_i$
- But this one would imply $\sum (\alpha_i - \beta_i) \mathbf{e}_i = \mathbf{0}$, hence $(\alpha_i - \beta_i) = 0$,
- i.e. $\alpha_i = \beta_i$, for every i .

Change of basis

- Let $B = (\mathbf{e}_1, \dots, \mathbf{e}_n) \subset \mathcal{V}$ be a basis for \mathcal{V} .

Change of basis

- Let $B = (\mathbf{e}_1, \dots, \mathbf{e}_n) \subset \mathcal{V}$ be a basis for \mathcal{V} .
- Of course, their coordinates are $(1 \ 0 \ \dots \ 0), (0 \ 1 \ \dots \ 0), \dots, (0 \ 0 \ \dots \ 1)$, and, in B coordinates, the basis is represented by the matrix

$$[B] = [I]$$

.

Change of basis

- Let $B = (\mathbf{e}_1, \dots, \mathbf{e}_n) \subset \mathcal{V}$ be a basis for \mathcal{V} .
- Of course, their coordinates are $(1 \ 0 \ \dots \ 0), (0 \ 1 \ \dots \ 0), \dots, (0 \ 0 \ \dots \ 1)$, and, in B coordinates, the basis is represented by the matrix

$$[B] = [I]$$

- If we take n (linearly independent) vectors $V = (\mathbf{v}_1, \dots, \mathbf{v}_n) \subset \mathcal{V}$, represented in B coordinates as $[V]$, and want to parametrize \mathcal{V} with respect to the new basis, we have, for transformation of coordinates:

$$[I] = [T][V]$$

Change of basis

- Let $B = (\mathbf{e}_1, \dots, \mathbf{e}_n) \subset \mathcal{V}$ be a basis for \mathcal{V} .
- Of course, their coordinates are $(1 \ 0 \ \dots \ 0), (0 \ 1 \ \dots \ 0), \dots, (0 \ 0 \ \dots \ 1)$, and, in B coordinates, the basis is represented by the matrix

$$[B] = [I]$$

- If we take n (linearly independent) vectors $V = (\mathbf{v}_1, \dots, \mathbf{v}_n) \subset \mathcal{V}$, represented in B coordinates as $[V]$, and want to parametrize \mathcal{V} with respect to the new basis, we have, for transformation of coordinates:

$$[I] = [T][V]$$

- and hence:

$$[T] = [V]^{-1}$$

Example: two polynomial bases

- Let $P_3 = (u^3, u^2, u, 1)$

Example: two polynomial bases

- Let $P_3 = (u^3, u^2, u, 1)$
- and $B_3 = ((1 - u)^3, 3u(1 - u)^2, 3u^2(1 - u), u^3)$ be two ordered bases

Example: two polynomial bases

- Let $P_3 = (u^3, u^2, u, 1)$
- and $B_3 = ((1-u)^3, 3u(1-u)^2, 3u^2(1-u), u^3)$ be two ordered bases
- for the linear space $\mathbb{P}^3(\mathbb{R})$ of polynomials with $\deg \leq 3$.

Example: two polynomial bases

- Let $P_3 = (u^3, u^2, u, 1)$
- and $B_3 = ((1-u)^3, 3u(1-u)^2, 3u^2(1-u), u^3)$ be two ordered bases
- for the linear space $\mathbb{P}^3(\mathbb{R})$ of polynomials with $\deg \leq 3$.
- the $[B_3]$ matrix in the P_3 basis is

$$[B_3]_{P_3} = \begin{pmatrix} -1 & 3 & -3 & 1 \\ 3 & -6 & 3 & 0 \\ -3 & 3 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$$

Example: two polynomial bases

- Let $P_3 = (u^3, u^2, u, 1)$
- and $B_3 = ((1-u)^3, 3u(1-u)^2, 3u^2(1-u), u^3)$ be two ordered bases
- for the linear space $\mathbb{P}^3(\mathbb{R})$ of polynomials with $\deg \leq 3$.
- the $[B_3]$ matrix in the P_3 basis is

$$[B_3]_{P_3} = \begin{pmatrix} -1 & 3 & -3 & 1 \\ 3 & -6 & 3 & 0 \\ -3 & 3 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$$

- the $[P_3]$ matrix in the B_3 basis is

$$[P_3]_{B_3} = [B_3]_{P_3}^{-1} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1/3 & 1 \\ 0 & 1/3 & 1/6 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix}$$

Example: two polynomial bases

- Let $P_3 = (u^3, u^2, u, 1)$
- and $B_3 = ((1-u)^3, 3u(1-u)^2, 3u^2(1-u), u^3)$ be two ordered bases
- for the linear space $\mathbb{P}^3(\mathbb{R})$ of polynomials with $\deg \leq 3$.
- the $[B_3]$ matrix in the P_3 basis is

$$[B_3]_{P_3} = \begin{pmatrix} -1 & 3 & -3 & 1 \\ 3 & -6 & 3 & 0 \\ -3 & 3 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$$

- the $[P_3]$ matrix in the B_3 basis is

$$[P_3]_{B_3} = [B_3]_{P_3}^{-1} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1/3 & 1 \\ 0 & 1/3 & 1/6 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix}$$

- WHY ?

Example

Open a file

```
fo = open("foo.txt", "r+")
```

```
str = fo.read(10);
```

```
print "Read String is : ", str
```

Check current position

```
position = fo.tell();
```

```
print "Current file position : ", position
```

Reposition pointer at the beginning once again

```
position = fo.seek(0, 0);
```

```
str = fo.read(10);
```

```
print "Again read String is : ", str
```

Close opened file

```
fo.close()
```

References

Linear Algebra Done Right book

NumPy tutorial