

# Computational Graphics: Lecture 7

Alberto Paoluzzi

Mon, Mar 17, 2014


# Outline: LAR1

- 1 Solid Modeling
- 2 Space decompositions
- 3 Cellular complex
- 4 Simplicial mapping
- 5 References

# Solid Modeling

# Representation scheme: definition

mapping  $s : M \rightarrow R$  from a space  $M$  of mathematical models to a space  $R$  of computer representations




- 1 The  $M$  set contains the mathematical models of the class of solid objects the scheme aims to represent

A. Requicha, [Representations for Rigid Solids: Theory, Methods, and Systems](#), *ACM Comput. Surv.*, 1980.

V. Shapiro, [Solid Modeling](#), In [Handbook of Computer Aided Geometric Design](#), 2001

# Representation scheme: definition

mapping  $s : M \rightarrow R$  from a space  $M$  of mathematical models to a space  $R$  of computer representations



- 1 The  $M$  set contains the mathematical models of the class of solid objects the scheme aims to represent
- 2 The  $R$  set contains the symbolic representations, i.e. the proper data structures, built according to a suitable grammar

A. Requicha, [Representations for Rigid Solids: Theory, Methods, and Systems](#), *ACM Comput. Surv.*, 1980.

V. Shapiro, [Solid Modeling](#), In [Handbook of Computer Aided Geometric Design](#), 2001

# Representation schemes

Most of such papers introduce or discuss one or more representation schemes ...

- |  |   |
|--|---|
| 1 Requicha, ACM Comput. Surv., 1980 [?]                | 16 Shapiro, Cornell Ph.D Th., 1991, [?]               |
| 2 Requicha & Voelcker, PEP TM-25, 1977, [?]            | 17 Paoluzzi et al., ACM Trans. Graph., 1993, [?]      |
| 3 Rossignac & Requicha, Comput. Aided Des., 1991, [?]  | 18 Pratt & Anderson, ICAP, 1994, [?]                  |
| 4 Bowyer, SVLIS, 1994, [?]                             | 19 Bowyer, Djinn, 1995, [?]                           |
| 5 Baumgart, Stan-CS-320, 1972, [?]                     | 20 Gomes et al., ACM SMA, 1999, [?]                   |
| 6 Braid, Commun. ACM, 1975, [?]                        | 21 Raghothama & Shapiro, ACM Trans. Graph., 1998, [?] |
| 7 Dobkin & Laszlo, ACM SCG, 1987, [?]                  | 22 Shapiro & Vossler, ACM SMA, 1995, [?]              |
| 8 Guibas & Stolfi, ACM Trans. Graph., 1985, [?]        | 23 Hoffmann & Kim, Comput. Aided Des., 2001, [?]      |
| 9 Woo, IEEE Comp. Graph. & Appl., 1985, [?]            | 24 Raghothama & Shapiro, ACM SMA, 1999, [?]           |
| 10 Yamaguchi & Kimura, Comp. Graph. & Appl., 1995, [?] | 25 DiCarlo et al., IEEE TASE, 2008, [?]               |
| 11 Gursoz & Choi & Prinz, Geom.Mod., 1990, [?]         | 26 Bajaj et al., CAD&A, 2006, [?]                     |
| 12 S.S.Lee & K.Lee, ACM SMA, 2001, [?]                 | 27 Pascucci et al., ACM SMA, 1995, [?]                |
| 13 Rossignac & O'Connor, IFIP WG 5.2, 1988, [?]        | 28 Paoluzzi et al., ACM Trans. Graph., 1995, [?]      |
| 14 Weiler, IEEE Comp. Graph. & Appl., 1985, [?]        | 29 Paoluzzi et al., Comput. Aided Des., 1989, [?]     |
| 15 Silva, Rochester, PEP TM-36, 1981, [?]              | 30 Ala, IEEE Comput. Graph. Appl., 1992, [?]          |

and much more ...

# Space decompositions

# Join of pointsets

The **join** of two sets  $P, Q \subset \mathbb{E}^n$  is the convex hull of their points:

$$PQ = \text{join}(P, Q) := \{\gamma p + \lambda q, p \in P, q \in Q\}$$
$$\gamma, \lambda \in \mathbb{R}, \gamma, \lambda \geq 0, \gamma + \lambda = 1$$

The join operation is **associative** and **commutative**.



# Join of pointsets: examples

```
pts = [[0,0],[.5,0],[0,.5],[.5,.5],
        [1,.5],[1.5,.5],[1.5,1],[.25,1]]
```

```
P = AA(MK)(pts)
```

```
S = AA(JOIN)([P[0:4],P[4:7],P[7]])
```

```
H = JOIN(S)
```

*# coords*

*# 0-polyhedra*

*# array of d-polyhedra*

*# 2-polyhedron*

```
VIEW(STRUCT(AA(SKELETON(1))(S)))
```

```
VIEW(H)
```




Figure : (a) 1-skeleton of pointsets in  $S$ ; (b) convex hull  $H$  of pointset  $P$

# Simplex

A **simplex**  $\sigma \subset \mathbb{E}^n$  of order  $d$ , or  $d$ -simplex, is the **join** of  $d + 1$  **affinely independent points**, called **vertices**.

The  $n + 1$  points  $p_0, \dots, p_n$  are **affinely independent** when the  $n$  vectors  $p_1 - p_0, \dots, p_n - p_0$  are **linearly independent**.

A  **$d$ -simplex** can be seen as a  **$d$ -dimensional triangle**: 0-simplex is a **point**, 1-simplex is a **segment**, 2-simplex is a **triangle**, 3-simplex is a **tetrahedron**, and so on.

# Simplex: examples

```
s0,s1,s2,s3 = [SIMPLEX(d) for d in range(4)] # array of standard d-simplices
VIEW(s1); VIEW(s2); VIEW(s3);

points = [[1,1,1],[0,1,1],[1,0,0],[1,1,0]] # coords of 4 points
tetra = JOIN(AA(MK)(points)) # 3-simplex
VIEW(tetra)
```




Figure : (a,b,c) 1-, 2-, and 3-standard simplex; (d) 3-simplex defined by 4 points

# Simplicial complex

Any subset of  $s + 1$  vertices ( $0 \leq s \leq d$ ) of a  $d$ -simplex  $\sigma$  defines an  $s$ -simplex, which is called  $s$ -face of  $\sigma$ .

# Simplicial complex

Any subset of  $s + 1$  vertices ( $0 \leq s \leq d$ ) of a  $d$ -simplex  $\sigma$  defines an  $s$ -simplex, which is called  $s$ -face of  $\sigma$ .

A simplicial complex is a set of simplices  $\Sigma$ , verifying the following conditions:

- 1 if  $\sigma \in \Sigma$ , then any  $s$ -face of  $\sigma$  belongs to  $\Sigma$ ;

# Simplicial complex

Any subset of  $s + 1$  vertices ( $0 \leq s \leq d$ ) of a  $d$ -simplex  $\sigma$  defines an  $s$ -simplex, which is called  $s$ -face of  $\sigma$ .

A simplicial complex is a set of simplices  $\Sigma$ , verifying the following conditions:

- 1 if  $\sigma \in \Sigma$ , then any  $s$ -face of  $\sigma$  belongs to  $\Sigma$ ;
- 2 if  $\sigma, \tau \in \Sigma$ , then  $\sigma \cap \tau \in \Sigma$ .

# Simplicial complex

Any subset of  $s + 1$  vertices ( $0 \leq s \leq d$ ) of a  $d$ -simplex  $\sigma$  defines an  $s$ -simplex, which is called  $s$ -face of  $\sigma$ .

A simplicial complex is a set of simplices  $\Sigma$ , verifying the following conditions:

- 1 if  $\sigma \in \Sigma$ , then any  $s$ -face of  $\sigma$  belongs to  $\Sigma$ ;
- 2 if  $\sigma, \tau \in \Sigma$ , then  $\sigma \cap \tau \in \Sigma$ .

# Simplicial complex

Any subset of  $s + 1$  vertices ( $0 \leq s \leq d$ ) of a  $d$ -simplex  $\sigma$  defines an  $s$ -simplex, which is called  $s$ -face of  $\sigma$ .

A simplicial complex is a set of simplices  $\Sigma$ , verifying the following conditions:

- 1 if  $\sigma \in \Sigma$ , then any  $s$ -face of  $\sigma$  belongs to  $\Sigma$ ;
- 2 if  $\sigma, \tau \in \Sigma$ , then  $\sigma \cap \tau \in \Sigma$ .

Geometric carrier  $|\Sigma|$  is the pointset union of simplices in  $\Sigma$ .



# Simplicial complex: examples

```
from larcc import *
V,CV = larSimplexGrid([5,5,5])
FV = larSimplexFacets(CV)
EV = larSimplexFacets(FV)
VIEW(EXPLODE(1.5,1.5,1.5)(MKPOLs((V,CV))))
VIEW(EXPLODE(1.5,1.5,1.5)(MKPOLs((V,FV))))
VIEW(EXPLODE(1.5,1.5,1.5)(MKPOLs((V,EV))))
```

```
BV = [FV[t] for t in boundaryCells(CV,FV)]
VIEW(EXPLODE(1.5,1.5,1.5)(MKPOLs((V,BV))))
```

```
# import LAR library
# structured simplicial grid
# 2-simplicial grid
# 1-simplicial grid
```

```
# boundary 2-simplices
```




Figure : (a) 3-complex; (b) 2-subcomplex; (c) 1-subcomplex; (d) 2-boundary.

# Simplicial complex: examples

(see Disk Point Picking)

```
from larcc import *; from random import random as rand
points = [[2*PI*rand(),rand()] for k in range(1000)]
V = [[SQRT(r)*COS(alpha),SQRT(r)*SIN(alpha)] for alpha,r in points]
cells = [[k+1] for k,v in enumerate(V)]
VIEW(MKPOL([V,cells,None]))

from scipy.spatial import Delaunay
FV = Delaunay(array(V)).vertices
VIEW(EXPLODE(1.2,1.2,1)(MKPOLS((V,FV))))
VIEW(SKELETON(1)(STRUCT(MKPOLS((V,FV)))))
```




Figure : (a) Points; (b) Delaunay triangulation; (c) 1-skeleton.

# Simplicial complex: examples

(see Disk Point Picking)

```
from larcc import *; from random import random as rand
points = [[2*rand()-1,2*rand()-1,2*rand()-1] for k in range(30000)]
V = [p for p in points if VECTNORM(p) <= 1]
VIEW(STRUCT(MKPOLS((V,AA(LIST)(range(len(V)))))))
```

```
from scipy.spatial import Delaunay
CV = Delaunay(array(V)).vertices
def test(tetra): return AND([v[-1] < 0 for v in tetra])
CV = [cell for cell in CV if test([V[v] for v in cell])]
VIEW(STRUCT(MKPOLS((V,CV))))
```




Figure: (a) Points; (b) Delaunay triangulation; (c) 1-skeleton

# Cellular complex

# CW-complex

- Roughly speaking, a CW-complex is made of basic building blocks called cells.

# CW-complex

- Roughly speaking, a **CW-complex** is made of **basic building blocks** called **cells**.
- The C stands for **closure-finite**, and the W for **weak topology**.

# CW-complex

- Roughly speaking, a **CW-complex** is made of **basic building blocks** called **cells**.
- The C stands for **closure-finite**, and the W for **weak topology**.
- **Closure-finite** means that **each closed cell** is covered by a **finite union** of **open cells**.

# CW-complex

- Roughly speaking, a **CW-complex** is made of **basic building blocks** called **cells**.
- The C stands for **closure-finite**, and the W for **weak topology**.
- **Closure-finite** means that **each closed cell** is covered by a **finite union** of **open cells**.



# CW-complex

- Roughly speaking, a **CW-complex** is made of **basic building blocks** called **cells**.
  - The C stands for **closure-finite**, and the W for **weak topology**.
  - **Closure-finite** means that **each closed cell** is covered by a **finite union** of **open cells**.
- ① An  $n$ -dimensional closed cell is the image of an  $n$ -disk under an attaching map.

# CW-complex

- Roughly speaking, a **CW-complex** is made of **basic building blocks** called **cells**.
  - The C stands for **closure-finite**, and the W for **weak topology**.
  - **Closure-finite** means that **each closed cell** is covered by a **finite union** of **open cells**.
- 1 An  $n$ -dimensional closed cell is the image of an  $n$ -disk under an attaching map.
  - 2 A **simplex** is a **closed cell**, and more generally, a **convex polytope** is a **closed cell**.

# CW-complex

- Roughly speaking, a **CW-complex** is made of **basic building blocks** called **cells**.
  - The C stands for **closure-finite**, and the W for **weak topology**.
  - **Closure-finite** means that **each closed cell** is covered by a **finite union** of **open cells**.
- 
- 1 An  $n$ -dimensional closed cell is the image of an  $n$ -disk under an attaching map.
  - 2 A **simplex** is a **closed cell**, and more generally, a **convex polytope** is a **closed cell**.
  - 3 An  $n$ -dimensional open cell is a topological space that is homeomorphic to the open disk.

# CW-complex

- Roughly speaking, a **CW-complex** is made of **basic building blocks** called **cells**.
  - The C stands for **closure-finite**, and the W for **weak topology**.
  - **Closure-finite** means that **each closed cell** is covered by a **finite union** of **open cells**.
- 
- 1 An  $n$ -dimensional closed cell is the image of an  $n$ -disk under an attaching map.
  - 2 A **simplex** is a **closed cell**, and more generally, a **convex polytope** is a **closed cell**.
  - 3 An  $n$ -dimensional open cell is a topological space that is homeomorphic to the open disk.
  - 4 A 0-dimensional open (and closed) cell is a **singleton** space.

# CW-complex

- Roughly speaking, a **CW-complex** is made of **basic building blocks** called **cells**.
  - The C stands for **closure-finite**, and the W for **weak topology**.
  - **Closure-finite** means that **each closed cell** is covered by a **finite union** of **open cells**.
- 
- 1 An  $n$ -dimensional closed cell is the image of an  $n$ -disk under an attaching map.
  - 2 A **simplex** is a **closed cell**, and more generally, a **convex polytope** is a **closed cell**.
  - 3 An  $n$ -dimensional open cell is a topological space that is homeomorphic to the open disk.
  - 4 A 0-dimensional open (and closed) cell is a **singleton** space.

# CW-complex

- Roughly speaking, a **CW-complex** is made of **basic building blocks** called **cells**.
  - The C stands for **closure-finite**, and the W for **weak topology**.
  - **Closure-finite** means that **each closed cell** is covered by a **finite union** of **open cells**.
- 1 An  $n$ -dimensional closed cell is the image of an  $n$ -disk under an attaching map.
  - 2 A **simplex** is a **closed cell**, and more generally, a **convex polytope** is a **closed cell**.
  - 3 An  $n$ -dimensional open cell is a topological space that is homeomorphic to the open disk.
  - 4 A 0-dimensional open (and closed) cell is a **singleton** space.

$n$ -Dimensional disk:

$$D^n = \{x \in \mathbb{R}^n : |x| \leq 1\}$$

Interior of  $D^n \subset \mathbb{R}^n$ :

$$\text{int}(D^n) = \{x \in \mathbb{R}^n : |x| < 1\}$$

Boundary of  $D^n \subset \mathbb{R}^n$ :

$$S^{n-1} = \{x \in \mathbb{R}^n : |x| = 1\}$$

# CW-complex: inductive definition

Let  $X$  be a topological space.

Let  $\Lambda(X) = \cup_p X_p \in \mathbb{N}$  be a partition of  $X$ , with  $X_p$  the set of  $p$ -cells.

# CW-complex: inductive definition

Let  $X$  be a **topological space**.

Let  $\Lambda(X) = \cup_p X_{p \in \mathbb{N}}$  be a **partition** of  $X$ , with  $X_p$  the set of  $p$ -cells.

**Definition (CW-structure)**

A **CW-structure** on the space  $X$  is a **filtration**

$$\emptyset = X^{-1} \subset X_0 \subset X_1 \subset \cdots \subset X_n = \Lambda(X),$$

such that, for each  $n$ , the space  $X_n$  is **homeomorphic** to a space obtained from  $X_{n-1}$  by attachment of  $n$ -cells.



# CW-complex: inductive definition

Let  $X$  be a **topological space**.

Let  $\Lambda(X) = \cup_p X_{p \in \mathbb{N}}$  be a **partition** of  $X$ , with  $X_p$  the set of  $p$ -cells.

**Definition (CW-structure)**

A **CW-structure** on the space  $X$  is a **filtration**

$$\emptyset = X^{-1} \subset X_0 \subset X_1 \subset \cdots \subset X_n = \Lambda(X),$$

such that, for each  $n$ , the space  $X_n$  is **homeomorphic** to a space obtained from  $X_{n-1}$  by attachment of  $n$ -cells.

**Definition (Cellular complex)**

A **cellular complex** is a space endowed with a CW-structure.

# CW-complex: inductive definition

Let  $X$  be a **topological space**.

Let  $\Lambda(X) = \cup_p X_{p \in \mathbb{N}}$  be a **partition** of  $X$ , with  $X_p$  the set of  $p$ -cells.

**Definition (CW-structure)**

A **CW-structure** on the space  $X$  is a **filtration**

$$\emptyset = X^{-1} \subset X_0 \subset X_1 \subset \cdots \subset X_n = \Lambda(X),$$

such that, for each  $n$ , the space  $X_n$  is **homeomorphic** to a space obtained from  $X_{n-1}$  by attachment of  $n$ -cells.

**Definition (Cellular complex)**

A **cellular complex** is a space endowed with a CW-structure.

A cellular complex is **finite** when it contains a finite number of cells.

# Polytopal, simplicial, cuboidal complexes

Cellular complexes characterised by different types of cells:

- 1 In polytopal complexes cells are polytopes, i.e. ~bounded convex sets;

# Polytopal, simplicial, cuboidal complexes

Cellular complexes characterised by different types of cells:

- 1 In polytopal complexes cells are polytopes, i.e.  $\sim$  bounded convex sets;
- 2 In simplicial complexes cells are simplices, i.e.  $\sim$   $d$ -polyedra with  $d + 1$  facets ( $(d - 1)$ -faces) and  $d + 1$  vertices.

# Politopal, simplicial, cuboidal complexes

Cellular complexes characterised by different types of cells:

- 1 In polytopal complexes cells are polytopes, i.e.  $\sim$  bounded convex sets;
- 2 In simplicial complexes cells are simplices, i.e.  $\sim d$ -polyedra with  $d + 1$  facets ( $(d - 1)$ -faces) and  $d + 1$  vertices.
- 3 In cuboidal complexes cells are cuboids, (in general, sets homeomorphic to) Cartesian products of intervals, i.e.  $\sim d$ -polyedra with  $2d$  facets and  $2d$  vertices.

# Numbers of vertices and facets

A  $d$ -simplex, or  $d$ -dimensional simplex, has  $d + 1$  extremal points called vertices and  $d + 1$  facets.

- a point (0-simplex) has  $0 + 1 = 1$  vertices and 1 facet ( $\emptyset$ );

# Numbers of vertices and facets

A  $d$ -simplex, or  $d$ -dimensional simplex, has  $d + 1$  extremal points called vertices and  $d + 1$  facets.

- a point (0-simplex) has  $0 + 1 = 1$  vertices and 1 facet ( $\emptyset$ );
- an edge (1-simplex) has  $1 + 1 = 2$  vertices and 2 facets;

# Numbers of vertices and facets

A  $d$ -simplex, or  $d$ -dimensional simplex, has  $d + 1$  extremal points called vertices and  $d + 1$  facets.

- a point (0-simplex) has  $0 + 1 = 1$  vertices and 1 facet ( $\emptyset$ );
- an edge (1-simplex) has  $1 + 1 = 2$  vertices and 2 facets;
- a triangle (2-simplex) has  $2 + 1 = 3$  vertices and 3 facets;



# Numbers of vertices and facets

A  $d$ -simplex, or  $d$ -dimensional simplex, has  $d + 1$  extremal points called vertices and  $d + 1$  facets.

- a point (0-simplex) has  $0 + 1 = 1$  vertices and 1 facet ( $\emptyset$ );
- an edge (1-simplex) has  $1 + 1 = 2$  vertices and 2 facets;
- a triangle (2-simplex) has  $2 + 1 = 3$  vertices and 3 facets;
- a tetrahedron (3-simplex) has  $3 + 1 = 4$  vertices and 4 facets, etc.

# Numbers of vertices and facets

A  $d$ -simplex, or  $d$ -dimensional simplex, has  $d + 1$  extremal points called vertices and  $d + 1$  facets.

- a point (0-simplex) has  $0 + 1 = 1$  vertices and 1 facet ( $\emptyset$ );
- an edge (1-simplex) has  $1 + 1 = 2$  vertices and 2 facets;
- a triangle (2-simplex) has  $2 + 1 = 3$  vertices and 3 facets;
- a tetrahedron (3-simplex) has  $3 + 1 = 4$  vertices and 4 facets, etc.

# Numbers of vertices and facets

A  $d$ -simplex, or  $d$ -dimensional simplex, has  $d + 1$  extremal points called vertices and  $d + 1$  facets.

- a point (0-simplex) has  $0 + 1 = 1$  vertices and 1 facet ( $\emptyset$ );
- an edge (1-simplex) has  $1 + 1 = 2$  vertices and 2 facets;
- a triangle (2-simplex) has  $2 + 1 = 3$  vertices and 3 facets;
- a tetrahedron (3-simplex) has  $3 + 1 = 4$  vertices and 4 facets, etc.

A  $d$ -cuboid has conversely  $2^d$  vertices and  $2d$  facets:

- a point (0-cuboid) has  $2^0 = 1$  vertices and 0 facets;

# Numbers of vertices and facets

A  $d$ -simplex, or  $d$ -dimensional simplex, has  $d + 1$  extremal points called vertices and  $d + 1$  facets.

- a point (0-simplex) has  $0 + 1 = 1$  vertices and 1 facet ( $\emptyset$ );
- an edge (1-simplex) has  $1 + 1 = 2$  vertices and 2 facets;
- a triangle (2-simplex) has  $2 + 1 = 3$  vertices and 3 facets;
- a tetrahedron (3-simplex) has  $3 + 1 = 4$  vertices and 4 facets, etc.

A  $d$ -cuboid has conversely  $2^d$  vertices and  $2d$  facets:

- a point (0-cuboid) has  $2^0 = 1$  vertices and 0 facets;
- an edge (1-cuboid) has  $2^1 = 2$  vertices and 2 facets;

# Numbers of vertices and facets

A  $d$ -simplex, or  $d$ -dimensional simplex, has  $d + 1$  extremal points called vertices and  $d + 1$  facets.

- a point (0-simplex) has  $0 + 1 = 1$  vertices and 1 facet ( $\emptyset$ );
- an edge (1-simplex) has  $1 + 1 = 2$  vertices and 2 facets;
- a triangle (2-simplex) has  $2 + 1 = 3$  vertices and 3 facets;
- a tetrahedron (3-simplex) has  $3 + 1 = 4$  vertices and 4 facets, etc.

A  $d$ -cuboid has conversely  $2^d$  vertices and  $2d$  facets:

- a point (0-cuboid) has  $2^0 = 1$  vertices and 0 facets;
- an edge (1-cuboid) has  $2^1 = 2$  vertices and 2 facets;
- a quadrilateral (2-cuboid) has  $2^2 = 4$  vertices and 4 facets;

# Numbers of vertices and facets

A  $d$ -simplex, or  $d$ -dimensional simplex, has  $d + 1$  extremal points called vertices and  $d + 1$  facets.

- a point (0-simplex) has  $0 + 1 = 1$  vertices and 1 facet ( $\emptyset$ );
- an edge (1-simplex) has  $1 + 1 = 2$  vertices and 2 facets;
- a triangle (2-simplex) has  $2 + 1 = 3$  vertices and 3 facets;
- a tetrahedron (3-simplex) has  $3 + 1 = 4$  vertices and 4 facets, etc.

A  $d$ -cuboid has conversely  $2^d$  vertices and  $2d$  facets:

- a point (0-cuboid) has  $2^0 = 1$  vertices and 0 facets;
- an edge (1-cuboid) has  $2^1 = 2$  vertices and 2 facets;
- a quadrilateral (2-cuboid) has  $2^2 = 4$  vertices and 4 facets;
- a hexahedron (3-cuboids) has  $2^3 = 8$  vertices and 8 facets, etc.

## Cellular complex: other definitions

Support  $|K|$  of a cellular complex  $K$  is the union of points of its cells

A triangulation of a polytope  $P$  is a simplicial complex  $K$  whose support is  $|K| = P$

For example, a triangulation of a polygon is a subdivision in triangles

Simplices and cuboids are polytopes.

A polytope is always triangulable;

For example, a quadrilateral by be divided in two triangles, and a cube in either 5 or 6 tetrahedra without adding new vertices

# Simplicial mapping



# Simplicial mapping: definition

## Definition

A **simplicial map** is a map **between simplicial complexes** with the property that the images of the vertices of a simplex always span a simplex.

## Remarks

Simplicial maps are **determined by their effects on vertices**  
for a precise definition of **Simplicial Map** look at [Wolfram MathWorld](#)

# MAP operator in plasm

## Map operator

`MAP(fun)(domain)`

## Semantics

- 1 **domain** (HPC value) is decomposed into a **simplicial complex**

# MAP operator in plasm

## Map operator

MAP(fun)(domain)

## Semantics

- 1 **domain** (HPC value) is decomposed into a **simplicial complex**
- 2 **fun** (a simplicial function) is applied to the **domain vertices**

# MAP operator in plasm

## Map operator

MAP(fun)(domain)

## Semantics

- 1 **domain** (HPC value) is decomposed into a **simplicial complex**
- 2 **fun** (a simplicial function) is applied to the **domain vertices**
- 3 the **mapped domain** is returned

# MAP examples: 1-sphere ( $S^1$ ) and 2-disk ( $D^2$ )

```
def sphere1(p): return [COS(p[0]), SIN(p[0])] # point function
def domain(n): return INTERVALS(2*PI)(n)     # generator of domain decomp
VIEW( MAP(sphere1)(domain(32)) )              # geometric value (HPC type)

def disk2D(p):                                # point function
    u,v = p
    return [v*COS(u), v*SIN(u)]               # coordinate functions
domain2D = PROD([INTERVALS(2*PI)(32), INTERVALS(1)(3)]) # 2D domain decompos
VIEW( MAP(disk2D)(domain2D) )
VIEW( SKELETON(1)(MAP(disk2D)(domain2D)) )
```




Figure : (a) sphere  $S^1$  (b) disk  $D^2$ ; (c) 1-skeleton.

# References

# References

A. DiCarlo, V. Shapiro, and A. Paoluzzi, Linear Algebraic Representation for Topological Structures, Computer-Aided Design, Volume 46, Issue 1 , January 2014, Pages 269-274 (doi:10.1016/j.cad.2013.08.044)