

1 Estimating individual subjective values of emotion regulation strategies

2 Christoph Scheffel<sup>†,1</sup>, Josephine Zerna<sup>†,1</sup>, Anne Gärtner<sup>1</sup>, Denise Dörfel<sup>1</sup>, & Alexander  
3 Strobel<sup>1</sup>

4 <sup>1</sup> Faculty of Psychology, Technische Universität Dresden, 01069 Dresden, Germany

5 Author Note

6 The authors made the following contributions. Christoph Scheffel: Conceptualization,  
7 Methodology, Funding acquisition, Formal analysis, Investigation, Project administration,  
8 Software, Visualization, Writing - original draft preparation, Writing - review & editing;  
9 Josephine Zerna: Conceptualization, Methodology, Funding acquisition, Investigation,  
10 Project administration, Software, Writing - review & editing; Anne Gärtner: Formal  
11 analysis, Writing - review & editing; Denise Dörfel: Conceptualization, Writing - review &  
12 editing; Alexander Strobel: Conceptualization, Writing - review & editing. <sup>†</sup> Christoph  
13 Scheffel and Josephine Zerna contributed equally to this work.

14 Correspondence concerning this article should be addressed to Christoph Scheffel,  
15 Zellescher Weg 17, 01069 Dresden, Germany. E-mail: christoph\_scheffel@tu-dresden.de

## Abstract

Individuals have a repertoire of emotion regulation (ER) strategies at their disposal, which they can use more or less flexibly. In ER flexibility research, strategies that facilitate goal achievement are considered adaptive and therefore are subjectively valuable. Individuals are motivated to reduce their emotional arousal effectively and to avoid cognitive effort. Perceived costs of ER strategies in the form of effort, however, are highly subjective. Subjective values (SVs) should therefore represent a trade-off between effectiveness and subjectively required cognitive effort. However, SVs of ER strategies have not been determined so far. We present a paradigm that is suitable for determining individual SVs of ER strategies. Using a multilevel modelling approach, it will be investigated whether individual SVs can be explained by effectiveness (subjective arousal, facial muscle activity) and subjective effort. Relations of SVs to personality traits will be explored.

*Keywords:* emotion regulation, regulatory effort, effort discounting, registered report, specification curve analysis

Word count: 139

Estimating individual subjective values of emotion regulation strategies

## 1. Introduction

Every day we are confronted with stressful or emotionally demanding situations. The ability to modify emotional experiences, expressions, and physiological reactions<sup>1</sup> to regulate emotions is an important cognitive skill. It is therefore not surprising that emotion regulation (ER) has substantial implications for well-being and adaptive functioning.<sup>2</sup> Different strategies can be used to regulate emotions, namely situation selection, situation modification, attentional deployment, cognitive change, and response modification,<sup>1</sup> and, following the taxonomy of Powers and LaBar,<sup>3</sup> individuals can implement ER strategies by means of different tactics. So called antecedent-focused strategies, e.g., attentional deployment and cognitive change, take effect early in the emotion generation process.<sup>1</sup> In contrast, response modification takes place late in the process and is therefore conceptualized as a response-focused strategy.<sup>1</sup> This postulated temporal sequence of ER strategies influences their effectiveness. It is meta-analytically proven that all mentioned strategies reduce subjective emotional arousal.<sup>4</sup> Distraction as a tactic of attentional deployment and (expressive) suppression as a tactic of response modulation showed small to medium effect sizes on measures of emotional experience (distraction:  $d_+ = 0.27$ ; suppression:  $d_+ = 0.27$ ). Distancing as tactic of cognitive change showed the highest effectiveness with an effect size of  $d_+ = 0.45$ .<sup>4</sup> Moreover, it is known that ER strategies reduce physiological responses of emotions, especially emotional valence as measured via facial electromyography (EMG)<sup>5</sup>. So these tactics from three different strategies proved to be effective in the short term. However, in order to be able to make a statement on their general benefit, longer-term consequences must also be considered. In particular, strategies that do not put the emotional content of the situation into a neutral perspective (i.e., distraction and suppression) are presumed to be disadvantageous in the

longer term. Longer lasting consequences of self-reported habitual use of ER strategies for affect and well-being have been discussed.<sup>6</sup> Especially suppression is generally associated with poorer outcomes (i.e., more negative affect, lower general well-being), which led to the postulation of *adaptive* and *maladaptive* ER strategies. For example, it was shown that maladaptive ER strategies mediate the effect between neuroticism and depressive symptoms.<sup>7</sup> In addition, a number of ER strategies have been associated with mental disorders (for meta-analytic review, see Aldao et al.).<sup>8</sup>

The postulation of *adaptive* and *maladaptive* ER strategies was put in a new perspective with the concept of ER flexibility. Similar to other psychological domains, e.g., attention and goal pursuit, *maladaptive* now refers to inflexible ER strategy use or use of strategies that are hindering goal achievement.<sup>9</sup> Adaptive flexible ER requires a large repertoire of ER strategies.<sup>9</sup> For example, greater ER flexibility is related to reduced negative affect and therefore beneficial in daily life.<sup>10</sup> Strategies have to be chosen from the repertoire that are useful for goal achieving. Evidence from other contexts (e.g., intertemporal choice,<sup>11</sup> shows that subjective values (SVs) are attributed to the choice options on the basis of which the decision is made. Research on ER choice has identified numerous factors that influence the choice of ER strategies, which can be seen as indirect evidence for factors influencing SVs. For example, it was shown that the intensity of a stimulus or situation plays a role in the choice.<sup>12</sup> Higher intensity of the stimulus leads to a choice of rather disengaging strategies, like distraction.<sup>12,13</sup> Further influencing factors are for example extrinsic motivation (e.g., monetary incentives), motivational determinants (i.e., hedonic regulatory goals), and effort.<sup>13,14</sup> Especially for effort, in our previous work we could show that the choice for an ER strategy is mainly influenced by the effort required to implement a given strategy.<sup>14</sup> In this study, participants used the strategies distancing and suppression while inspecting emotional pictures. Afterwards, they choose which strategy they wanted to use again. Participants tended to re-apply the strategy that was subjectively less effortful, even though it was subjectively not the most effective one - in

this case: suppression. Interestingly, the choice was independent of self-reported habitual use of suppression and reappraisal. What has been missing in research on ER choice so far is information regarding the strategy *not* chosen. People choose a strategy that they prefer for different, relatively well-known reasons. However, nothing is revealed about the strategy that is *not chosen*.

We assume that people choose the strategy that has the highest value for them at the moment. The value is determined against the background of goal achievement in the specific situation: A strategy is highly valued if it facilitates goal achievement.<sup>9</sup> One certainly central goal is the regulation of negative affect. A second, intrinsic and less obvious goal is the avoidance of effort.<sup>15</sup> When given the choice, most individuals prefer tasks that are less effortful.<sup>16</sup> We assume that both aspects are traded off against each other by individuals to determine individual subjective values (SVs) of ER strategies: A strategy is more valuable if it can reduce emotional arousal *and* is less effortful. SVs of ER Strategies could be helpful to describe the ER repertoire<sup>9</sup> more comprehensively. Depending on the flexibility of a person, different patterns of SVs could be conceivable: A person with high flexibility would show relatively high SVs for a number of strategies. This would mean that all strategies are a good option for goal achievement. A second person with less flexibility, however, would show high SVs only for one strategy, or for no strategy at all. This in turn would mean that no strategy is a good choice to achieve ER goals.


However, so far we have not seen any attempt in ER choice research to determine individual SVs of ER strategies. To investigate this question, the individual SVs of each strategy available for selection would have to be determined. Promising approaches can be found in studies on difficulty levels of effortful cognitive tasks.

Individual SVs of effortful cognitive tasks have been quantified using the Cognitive Effort Discounting Paradigm (COG-ED).<sup>17</sup>

In the original study by Westbrook et al.,<sup>17</sup> cognitive load was varied using the

*n*-back task, a working memory task that requires fast and accurate responses to sequentially presented stimuli. Participants had to decide in an iterative procedure whether they wanted to repeat a higher *n*-back level for a larger, fixed monetary reward, or a lower level for a smaller, varying reward, with the implicit assumption that the objectively easiest *n*-back level has the highest SV. In the present study, we want to use this paradigm to determine SVs of ER strategies. In doing so, we need to make an important change: We have to adapt the assumption that the easiest *n*-back level has the highest SV. As we have shown in previous studies, there are large inter-individual differences in the preference and perceived subjective effort of ER strategies.<sup>14</sup> Moreover, there is nothing like an objectively easiest ER strategy. Therefore, we have to add an additional step, which precedes the other steps and where the option with the higher subjective value is determined. In this step, the same monetary value (i.e., 1 €) is assigned to both options. The assumption is that participants now choose the option that has the higher SV for them. In the next step we return to the original paradigm. The higher monetary value (i.e., 2 €) is assigned to the option that was not chosen in the first step and therefore is assumed to have the lower SV. In the following steps, the lower value is changed in every iteration according to Westbrook et al.<sup>17</sup> until the indifference point is reached. This procedure will be repeated until all strategies have been compared. The SV of each strategy is calculated as the mean of this strategy's SV from all comparisons. In case a participant has a clear preference for one strategy, the SV of this strategy will be 1. But our paradigm can also account for the case that a person does not have a clear preference. Then no SV will be 1, but still, the SVs of all strategies can be interpreted as absolute values and in relation to the other strategy's SVs (see Figure 1). Additionally, we will test our adapted paradigm in a *n*-back task and explore whether this paradigm can describe individuals that do not prefer the easiest *n*-back option (see Zerna et al., ...).

The aim of the present study is to evaluate whether this paradigm is suitable for determining SVs of ER strategies. As a manipulation check, we want to investigate



*Figure 1.* Exemplary visualization of two response patterns. In the top half, the person has a clear preference for one of the three strategies. In the lower half, they have no clear preference and therefore show an inconsistent response pattern. This pattern can be represented by our paradigm.

whether the ER strategies distraction, distancing, and suppression effectively reduce emotional arousal and physiological responses as well as whether they require cognitive effort. The following hypotheses are proposed:

- H1a) Subjective arousal (arousal ratings) is lower after using an ER strategy (distraction, distancing, suppression) compared to active viewing.
- H2a) Physiological responses in facial EMG that is sensitive to negative emotion (*corrugator* activity) are lower after using an ER strategy (distraction, distancing, suppression) compared to active viewing.
- H2b) Physiological responses in facial EMG that is sensitive to negative emotion (*levator* activity) are lower after using an ER strategy (distraction, distancing, suppression) compared to active viewing.

- H3a) Subjective effort (effort ratings) is greater after using an ER strategy (distraction, distancing, suppression) compared to active viewing.
- H3b) The majority of participants re-uses the strategy that was least effortful for them.

Furthermore, we want to investigate whether the following variables predict individual subjective values of ER strategies and whether effort is the best predictor for SVs of ER strategies with the following hypotheses:

- H4a) Subjective effort ratings negatively predict SVs of ER strategies.
- H4b) Subjective arousal ratings negatively predict SVs of ER strategies.
- H4c) EMG *corrugator* activity negatively predicts SVs of ER strategies.
- H4d) EMG *levator* activity negatively predicts SVs of ER strategies.
- H5a) SVs decline with increasing effort, even after controlling for task performance measured by subjective arousal ratings, *corrugator* and *levator* activity.

We also want to investigate whether SVs are related to flexible emotion regulation:

- H6a) The higher the SV, the more likely the respective strategy is chosen.
- H6b) SVs are lower and decline stronger when ER flexibility is lower.

Exploratorily, we want to investigate whether individual SVs are related to personality traits and how individual SVs of ER strategies relate to SVs of other tasks with different demand levels, namely *n*-back.

## 2. Method

We report how we determined our sample size, all data exclusions (if any), all manipulations, and all measures in the study.<sup>18</sup> The paradigm was written and presented



using *PsychoPy*.<sup>19</sup> We used *R* with *R Studio*<sup>20,21</sup> with the main packages *afex*<sup>22</sup> and *BayesFactor*<sup>23</sup> for all analyses. The R Markdown file used to analyze the data and write this document, as well as the raw data and the materials are freely available at <https://github.com/ChScheffel/CERED>. A complete list of all measures assessed in the study can be found at OSF (<https://osf.io/vnj8x/>) and GitHub (<https://github.com/ChScheffel/CERED>).

## 2.1 Ethics information

The study protocol complies with all relevant ethical regulations and was approved by the ethics committee of the Technische Universität Dresden (reference number EK...). Prior to testing, written informed consent will be obtained. Participants will receive 30 € in total or course credit for participation.

## 2.2 Pilot data

The newly developed ER paradigm was tested in a pilot study with  $N = 16$  participants (9 female; age:  $M = 24.1 \pm SD = 3.6$ ). Regarding self-reported arousal, results showed significant higher subjective arousal for active viewing of negative compared to neutral pictures. However, ER strategies did not lead to a reduction of subjective arousal compared to active viewing of negative pictures. Regarding physiological responses, ER strategies were associated with reduced facial muscle activity of the *corrugator* and *levator* compared to active viewing of negative pictures. In accordance with our previous study,<sup>14</sup> we found that the use of ER strategies compared to active viewing was associated with increased subjective effort. All results are detailed in the Supplementary Material

## 2.3 Design

Young healthy participants (aged 18 to 30 years) will be recruited using the software *ORSEE*<sup>24</sup> at the Technische Universität Dresden. Participants will be excluded from participation if they do not fluently speak German, have current or a history of psychological disorders or neurological trauma, or report to take medication. Participants will be invited to complete an online survey containing different questionnaires to assess broad and narrow personality traits and measures of well-being. The study consists of two lab sessions, which will take place in a shielded cabin with constant lighting. Before each session, participants will receive information about the respective experimental procedure and provide informed consent. At the beginning of the first session participants will fill out a demographic questionnaire and complete an n-back task with the levels one to four. Then, they will complete an effort discounting (ED) procedure on screen, followed by a random repetition of one n-back level. The second session will comprise the ER paradigm and take place exactly one week after session one. Participants will provide informed consent and receive written instructions on the ER paradigm and ER strategies that they should apply. A brief training will ensure that all participants are able to implement the ER strategies. Next, electrodes to measure facial EMG will be attached and the ER paradigm will be conducted. Study data will be collected and managed using REDCap electronic data capture tools hosted at Technische Universität Dresden.<sup>25,26</sup>

**2.3.1 Psychometric measures.** The online survey will contain a number of questionnaires: General psychological well-being will be assessed using the German version of the WHO-5 scale.<sup>27,28</sup> To measure resilience, the German version 10-item-form of the Connor-Davidson resilience Scale (CD-RISC)<sup>31</sup> will be used. Habitual use of ER will be assessed using the German version of the Emotion Regulation Questionnaire (ERQ).<sup>6,32</sup> For the assessment of ER flexibility we will use the Flexible Emotion Regulation Scale (FlexER).<sup>33</sup> Implicit theories of willpower in emotion control will be assessed using the

implicit theories questionnaire from.<sup>34</sup> To assess Need for Cognition, the German version short form of the Need for Cognition Scale<sup>35,36</sup> will be used. To assess self-control,<sup>37</sup> sum scores of the German versions of the following questionnaires will be used: the Self-Regulation Scale (SRS),<sup>38</sup> the Brief Self-Control Scale (BSCS),<sup>39,40</sup> and the Barratt Impulsiveness Scale (BIS-11).<sup>41,42</sup> Attentional control will be assessed using the Attentional Control Scale (ACS).<sup>43</sup> For more detailed information on psychometric properties of the questionnaires, please see supplementary material.

**2.3.2 Emotion regulation paradigm.** The ER paradigm will consist of three parts that will be described in the following.


*Part one: ER task.* Part one will be a standard ER task in a block design (see Figure 2), similar to paradigms previously used by our group.<sup>14</sup> Participants will be told to actively view neutral and negative pictures (see 2.3.3) or to regulate all upcoming emotions by means of distraction, distancing, and expressive suppression, respectively. Every participant first will have the condition “active viewing-neutral” that serves as a baseline condition. During this block, 20 neutral pictures will be presented. Participants will be asked to “actively view all pictures and permit all emotions that may arise.” In the second block, participants will actively view negative pictures. During the third, fourth, and fifth block, participants will see negative pictures and will be asked to regulate their emotions using distraction, distancing, and suppression. In order to achieve distraction, participants will be asked to think of a geometric object or an everyday activity, like brushing their teeth. During distancing, participants will be asked to “take the position of a non-involved observer, thinking about the picture in a neutral way.” Participants will be told not to re-interpret the situation or attaching a different meaning to the situation. During suppression, participants will be told to “suppress their emotional facial expression.” They should imagine being observed by a third person that should not be able to tell by looking at the facial expression whether the person is looking at an emotional picture. Participants will be instructed not to suppress their thoughts or change their facial expression to the

opposite.<sup>14</sup> All participants will receive written instruction and complete a training session. After the training session, participants will be asked about their applied ER strategies to avoid misapplication. The order of the three regulation blocks (distraction, distancing, and suppression) will be randomized between participants.

*Part two: ER effort discounting.* In the second part, ER effort discounting will take place. The procedure of the discounting will follow the COG-ED paradigm by Westbrook et al.<sup>17</sup> with a major change. We will use the following adaption that allows the computation of SVs for different strategies without presuming that all individuals would inherently evaluate the same strategy as the easiest one: For each possible pairing (distraction vs. distancing, distraction vs. suppression, and distancing vs. suppression), two strategies with monetary values will be presented. The order of the comparisons will be randomized. Because there is no strategy that is objectively more difficult, we will add an initial comparison that begins with the option “1 € for strategy A or 1 € for strategy B”. The strategy that is not chosen will be assigned the value of 2 €. From this point on, comparisons between strategies will follow the original COG-ED paradigm.<sup>17</sup> Participants will be instructed to decide as realistically as possible by imagining that the money displayed is actually available for choice.

*Part three: ER choice.* After the discounting part, participants will choose which of the three ER strategies (distraction, distancing or suppression) they want to re-apply. Importantly, there will be no further instruction on what basis they should make their decision. Participants should make their decision freely, according to criteria they consider important for themselves. However, participants will be asked to state the reasons for the decision afterwards. As soon as they have decided, they will see the respective instruction and the block with another 20 negative pictures starts.

**2.3.3 Stimuli.** Pictures that will be used in the paradigm are selected from the Emotional Picture Set (EmoPicS)<sup>44</sup> and the International Affective Picture System (IAPS).<sup>45</sup> The 20 neutral pictures (Valence (V):  $M \pm SD = 4.81 \pm 0.51$ ; Arousal (A):  $M \pm$



*Figure 2.* Block design of the paradigm. Every participant starts with two "active viewing" blocks containing neutral (Block 1) and negative (Block 2) pictures. Order of the regulation blocks (Blocks 3, 4, and 5) is randomized between participants. After, the discounting procedure takes place. All three regulation strategies are pairwise compared. Before the last block, participants can decide which regulation strategy they want to reapply. Subjective arousal and effort ratings are assessed after each block using a slider on screen with a continuous scale.

$SD = 3 \pm 0.65$ ) depicted content related to the categories persons, objects, and scenes. Further, 100 negative pictures, featuring categories animals, body, disaster, disgust, injury, suffering, violence, and weapons, will be used. An evolutionary algorithm<sup>46</sup> is used to cluster these pictures into five sets with comparable valence and arousal values (set one: V:  $M \pm SD = 2.84 \pm 0.57$ , A:  $M \pm SD = 5.62 \pm 0.34$ ; set two: V:  $M \pm SD = 2.64 \pm 0.46$ , A:  $M \pm SD = 5.58 \pm 0.35$ ; set three: V:  $M \pm SD = 2.82 \pm 0.62$ , A:  $M \pm SD = 5.60 \pm 0.39$ ; set four: V:  $M \pm SD = 2.65 \pm 0.75$ , A:  $M \pm SD = 5.61 \pm 0.41$ ; set five: V:  $M \pm SD = 2.74 \pm 0.70$ , A:  $M \pm SD = 5.63 \pm 0.37$ ). A complete list of all pictures and their classification into sets can be found in supplementary material 1.

**2.3.4 Facial electromyography.** Bipolar facial electromyography (EMG) will be measured for *corrugator supercilii* and *levator labii* as indices of affective valence,<sup>47</sup> similar to previous work by our group.<sup>48</sup> Two passive surface Ag/AgCl electrodes (8 mm inner diameter, 10 mm distance between electrodes) will be placed over each left muscle according to the guidelines of.<sup>49</sup> The ground electrode will be placed over the left *Mastoid*. Before electrode placement, the skin will be abraded with Every abrasive paste, cleaned with alcohol, and filled with Lectron III electrolyte gel. Raw signals will be amplified by a BrainAmp amplifier (Brain Products Inc., Gilching, Germany). Impedance level will be kept below 10  $k\Omega$ . Data will be sampled at 1000 Hz, filtered, rectified and integrated. A 20 Hz high pass (order 8), a 300 Hz low pass (order 8), and a 50 Hz notch filter will be applied to both signals. Corrugator and Levator EMG will be analyzed during the 6 s of picture presentation. EMG data will be baseline-corrected using a time window of 2 s prior to stimulus onset.<sup>47</sup> Last, the sampling rate will be changed to 100 Hz, and EMG data will be averaged for each condition and each participant.

## 2.4 Sampling plan

Sample size calculation is done using *G\*Power*.<sup>50,51</sup> In a meta-analysis of Zaehring and colleagues,<sup>5</sup> effect sizes of ER on peripheral-physiological measures were reported: To

find an effect of  $d = -0.32$  of ER on *corrugator* muscle activity with  $\alpha = .05$  and  $\beta = .95$ , data of at least  $N = 85$  have to be analyzed. Power analyses of all other hypotheses yielded smaller sample sizes. However, if participants withdraw from study participation, technical failures occur, or experimenter considers the participant for not suitable for study participation (e.g., because the participant does not follow instructions or shows great fatigue), respective data will also be excluded from further analyses. Therefore, we aim to collect data of XX participants. Detailed information on power calculation for each hypothesis can be found in the design table.

## 2.5 Analysis plan

Data collection and analysis will not be performed blind to the conditions of the experiments. The level of significance will be set to  $\alpha = .05$ .

*Effects of emotion regulation on arousal, facial EMG and effort.* To examine the impact of valence of emotional pictures on subjective arousal ratings, a repeated measures analysis of variance (rmANOVA) with the factor valence (neutral and negative) for the strategy active viewing will be conducted. To investigate the effects of the three ER strategies on subjective arousal, another rmANOVA with the factor strategy (active viewing - negative, distraction, distancing, and suppression) for subjective arousal ratings will be conducted.

To examine the impact of valence on facial EMG, a rmANOVA with the factor valence (neutral and negative) for the strategy active viewing will be conducted for *corrugator* and *levator* activity. To examine the effects of the three ER strategies on facial EMG, another rmANOVA with the factor strategy (active viewing - negative, distraction, distancing, and suppression) for *corrugator* and *levator* activity will be conducted.

To examine the effect of ER strategies on cognitive effort, a rmANOVA with the factor strategy (active viewing, distraction, distancing, and suppression) for subjective

effort ratings will be conducted. Greenhouse-Geisser-corrected  $p$ -values and degrees of freedom will be reported when the assumption of sphericity is violated. Proportion of explained variance  $\eta_p^2$  will be reported as a measure of effect size. If indicated by the data, estimated marginal means will be computed as post-hoc contrasts. rmANOVAS will be run using the *afex* package<sup>22</sup>

*Subjective values of emotion regulation strategies.* For each ER strategy, SVs will be calculated as follows: first, the value 0.015625 will be added to or subtracted from the last monetary value of the flexible strategy, depending on the participant's last choice. Second, the resulting (monetary) value will be divided by 2.00 €. The final SV for each participant will be computed by averaging all final SVs of each strategy. The resulting values will be entered in a rmANOVA to compare the SVs of the three strategies (distraction, distancing, and suppression) and explore group effects. Again, estimated marginal means will be computed as post-hoc contrasts.

To investigate whether individual SVs predict ER choice, a Chi-squared test with predicted choice (highest SV of each participant) and actual choice will be computed. Furthermore, an ordinal logistic regression with the dependent variable choice and independent variables SVs of each strategy will be computed.

To explore the association between subjective arousal, physiological responses, and subjective effort on SVs, a multilevel model (MLM) will be specified using the *lmerTest* package.<sup>52</sup> First, ER strategies will be recoded and centered for each subject according to their individual SVs: The strategy with the highest SV will be coded as -1, the strategy with the second highest SV 0, and the strategy with the lowest SV will be coded as 1. Restricted maximum likelihood (REML) will be applied to fit the model. A random slopes model of SVs including subjective effort (effort ratings), subjective arousal (arousal ratings), and physiological responses (*corrugator* and *levator* activity) as level-1-predictors



will be specified.

$$SV \sim strategy + effort\ rating + arousal\ rating + corrugator\ activity + levator\ activity + (strategy|subject)$$

Level-1-predictors will be centered within cluster.<sup>53</sup> Residuals of the final model will be inspected visually. Intraclass correlation coefficient (ICC),  $\rho$ , will be reported for each model (null model, as well as full model).

The influence of personality traits on SVs will be investigated exploratorily. Therefore, the MLM specified above will be extended by the level-2-predictors NFC and self-control.

The association between flexible ER and SVs of ER strategies will be investigated with a regression using the *intercept* and *slope* of each participants' SVs to predict their FlexER score. Firstly, SVs will be ordered by magnitude. Secondly, for each participant a linear model will be built to estimate the individual *intercept* and *slope*.

For each result of the analyses, both  $p$ -values and Bayes factors  $BF_{10}$ , calculated using the *BayesFactor* package,<sup>23</sup> will be reported. Bayes factors are calculated using the default prior widths of the functions *anovaBF*, *lmBF* and *regressionBF*.

## Data availability

The data of this study can be downloaded from [osf.io/vnj8x/](https://osf.io/vnj8x/).

## Code availability

The paradigm code, as well as the R Markdown file used to analyze the data and write this document is available at our Github repository.

## References

1.

Gross, J. J. Antecedent- and response-focused emotion regulation: Divergent consequences for experience, expression, and physiology. *Journal of Personality and Social Psychology* **74**, 224–37 (1998).

2.

Gross, J. J. The emerging field of emotion regulation: An integrative review. *Review of General Psychology* **2**, 271–299 (1998).

3.

Powers, J. P. & LaBar, K. S. Regulating emotion through distancing: A taxonomy, neurocognitive model, and supporting meta-analysis. *Neuroscience and Biobehavioral Reviews* **96**, 155–173 (2019).

4.

Webb, T. L., Miles, E. & Sheeran, P. Dealing with feeling: A meta-analysis of the effectiveness of strategies derived from the process model of emotion regulation. *Psychological Bulletin* **138**, 775–808 (2012).

5.

Zaehringer, J., Jennen-Steinmetz, C., Schmahl, C., Ende, G. & Paret, C. Psychophysiological effects of downregulating negative emotions: Insights from a meta-analysis of healthy adults. *Front Psychol* **11**, 470 (2020).

6.

Gross, J. J. & John, O. P. Individual differences in two emotion regulation processes: Implications for affect, relationships, and well-being. *Journal of Personality and Social Psychology* **85**, 348–62 (2003).

385 7.

386 Yoon, K. L., Maltby, J. & Joormann, J. A pathway from neuroticism to depression:  
Examining the role of emotion regulation. *Anxiety, Stress, & Coping* **26**, 558–72  
387 (2013).

388 8.

389 Aldao, A., Nolen-Hoeksema, S. & Schweizer, S. Emotion-regulation strategies across  
psychopathology: A meta-analytic review. *Clinical Psychology Review* **30**, 217–237  
390 (2010).

391 9.

392 Aldao, A., Sheppes, G. & Gross, J. J. Emotion regulation flexibility. *Cognitive Ther-*  
393 *apy and Research* **39**, 263–278 (2015).

394 10.

395 Blanke, E. S. *et al.* Mix it to fix it: Emotion regulation variability in daily life.  
396 *Emotion* **20**, 473–485 (2020).

397 11.

398 Kable, J. W. & Glimcher, P. W. The neural correlates of subjective value during  
399 intertemporal choice. *Nat Neurosci* **10**, 1625–33 (2007).

400 12.

401 Sheppes, G., Scheibe, S., Suri, G. & Gross, J. J. Emotion-regulation choice. *Psycho-*  
402 *logical Science* **22**, 1391–6 (2011).

403 13.

404 Sheppes, G. *et al.* Emotion regulation choice: A conceptual framework and supporting  
405 evidence. *Journal of Experimental Psychology: General* **143**, 163–81 (2014).

406 14.

407 Scheffel, C. *et al.* Effort beats effectiveness in emotion regulation choice: Differences  
between suppression and distancing in subjective and physiological measures. *Psy-*  
408 *chophysiology* **00**, e13908 (2021).

409 15.

410 Inzlicht, M., Shenhav, A. & Olivola, C. Y. The effort paradox: Effort is both costly  
and valued. *Trends Cogn Sci* **22**, 337–349 (2018).

412 16.

413 Hull, C. L. *Principles of behavior: An introduction to behavior theory.* (Appleton-  
414 Century-Crofts, 1943).

415 17.

416 Westbrook, A., Kester, D. & Braver, T. S. What is the subjective cost of cognitive  
effort? Load, trait, and aging effects revealed by economic preference. *PLOS ONE*  
417 **8**, e68210 (2013).

418 18.

419 Simmons, J. P., Nelson, L. D. & Simonsohn, U. A. A 21 word solution. *SSRN*  
420 *Electronic Journal* (2012) doi:10.2139/ssrn.2160588.

421 19.

422 Peirce, J. *et al.* PsychoPy2: Experiments in behavior made easy. *Behavior Research*  
423 *Methods* **51**, 195–203 (2019).

424 20.

425 R Core Team. *R: A language and environment for statistical computing.* (R Founda-  
426 tion for Statistical Computing, 2021).

427 21.

428 RStudio Team. RStudio: Integrated development for R. (2020).

429  
430 22.

431 Singmann, H., Bolker, B., Westfall, J., Aust, F. & Ben-Shachar, M. S. *Afex: Analysis*  
432 *of factorial experiments*. (2021).

433 23.

434 Morey, R. D. & Rouder, J. N. *BayesFactor: Computation of Bayes factors for common*  
435 *designs*. (2021).

436 24.

437 Greiner, B. Subject pool recruitment procedures: Organizing experiments with  
438 ORSEE. *Journal of the Economic Science Association* **1**, 114–125 (2015).

439 25.

440 Harris, P. A. *et al.* Research electronic data capture (REDCap)—A metadata-driven  
methodology and workflow process for providing translational research informatics  
441 support. *Journal of Biomedical Informatics* **42**, 377–381 (2009).

442 26.

443 Harris, P. A. *et al.* The REDCap consortium: Building an international community  
444 of software platform partners. *Journal of Biomedical Informatics* **95**, 103208 (2019).

445 27.

446 Bech, P. Measuring the dimensions of psychological general well-being by the WHO-5.  
447 *Quality of life newsletter* **32**, 15–16 (2004).

448 28.

449 Brähler, E., Mühlen, H., Albani, C. & Schmidt, S. Teststatistische prüfung und  
normierung der deutschen versionen des EUROHIS-QOL lebensqualität-index und  
450 des WHO-5 wohlbefindens-index. *Diagnostica* **53**, 83–96 (2007).

451 29.

452 Connor, K. M. & Davidson, J. R. Development of a new resilience scale: The connor-  
453 davidson resilience scale (CD-RISC). *Depression and Anxiety* **18**, 76–82 (2003).

454 30.

455 Sarubin, N. *et al.* First analysis of the 10-and 25-item german version of the connor-  
davidson resilience scale (CD-RISC) regarding psychometric properties and compo-  
456 nents. *Zeitschrift Fur Gesundheitspsychologie* **23**, 112–122 (2015).

457 31.

458 Campbell-Sills, L. & Stein, M. B. Psychometric analysis and refinement of the connor-  
davidson resilience scale (CD-RISC): Validation of a 10-item measure of resilience.  
459 *Journal of Traumatic Stress* **20**, 1019–28 (2007).

460 32.

461 Abler, B. & Kessler, H. Emotion regulation questionnaire - a german version of the  
462 ERQ by gross and john. *Diagnostica* **55**, 144–152 (2009).

463 33.

464 Dörfel, D., Gärtner, A. & Strobel, A. A new self-report instrument for measuring  
emotion regulation flexibility. *Society for Affective Science (SAS) Annual Conference*  
465 (2019).

466 34.

467 Bernecker, K. & Job, V. Implicit theories about willpower in resisting temptations  
and emotion control. *Zeitschrift Fur Psychologie-Journal of Psychology* **225**, 157–166  
468 (2017).

469 35.

470 Cacioppo, J. T. & Petty, R. E. The need for cognition. *Journal of Personality and*  
471 *Social Psychology* **42**, 116–131 (1982).

472 36.

473 Bless, H., Wanke, M., Bohner, G., Fellhauer, R. F. & Schwarz, N. Need for cognition  
- a scale measuring engagement and happiness in cognitive tasks. *Zeitschrift Für*  
474 *Sozialpsychologie* **25**, 147–154 (1994).

475 37.

476 Paschke, L. M. *et al.* Individual differences in self-reported self-control predict suc-  
cessful emotion regulation. *Social Cognitive and Affective Neuroscience* **11**, 1193–204  
477 (2016).

478 38.

479 Schwarzer, R., Diehl, M. & Schmitz, G. S. Self-regulation scale. (1999).

481 39.

482 Tangney, J. P., Baumeister, R. F. & Boone, A. L. High self-control predicts good  
adjustment, less pathology, better grades, and interpersonal success. *Journal of Per-*  
483 *sonality* **72**, 271–324 (2004).

484 40.

485 Sproesser, G., Strohbach, S., Schupp, H. & Renner, B. Candy or apple? How self-  
control resources and motives impact dietary healthiness in women. *Appetite* **56**,  
486 784–787 (2011).

487 41.

488 Patton, J. H., Stanford, M. S. & Barratt, E. S. Factor structure of the barratt impul-  
siveness scale. *Journal of Clinical Psychology* **51**, 768–774 (1995).

490 42.

Hartmann, A. S., Rief, W. & Hilbert, A. Psychometric properties of the german version of the barratt impulsiveness scale, version 11 (BIS-11) for adolescents. *Perceptual and Motor Skills* **112**, 353–368 (2011).

43.

Derryberry, D. & Reed, M. A. Anxiety-related attentional biases and their regulation by attentional control. *Journal of abnormal psychology* **111**, 225–236 (2002).

44.

Wessa, M. *et al.* EmoPicS: Subjective und psychophysiologische evaluation neuen bildmaterials für die klinisch-biopsychologische forschung. *Zeitschrift für Klinische Psychologie und Psychotherapie* **39**, 77 (2010).

45.

Lang, P. J., Bradley, M. M. & Cuthbert, B. N. *International affective picture system (IAPS): Affective ratings of pictures and instruction manual*. (University of Florida, 2008).

46.

Yu, X. & Gen, M. *Introduction to evolutionary algorithms*. (Springer Science & Business Media, 2010).

47.

Bradley, M. M. & Lang, P. J. Measuring emotion: Behavior, feeling, and physiology. in *Cognitive neuroscience of emotion* (eds. Lane, R. D. & Nadel, L.) 242–276 (Oxford University Press, 2000).

48.

Gärtner, A., Jawinski, P. & Strobel, A. Individual differences in inhibitory control are not related to emotion regulation. *PsyArXiv* (2021) doi:10.31234/osf.io/cd8rx.



511 49.

512 Fridlund, A. J. & Cacioppo, J. T. Guidelines for human electromyographic research.  
513 *Psychophysiology* **23**, 567–89 (1986).

514 50.

515 Faul, F., Erdfelder, E., Lang, A.-G. & Buchner, A. G\*Power 3: A flexible statistical  
power analysis program for the social, behavioral, and biomedical sciences. *Behavior*  
516 *Research Methods* **39**, 175–191 (2007).

517 51.

518 Faul, F., Erdfelder, E., Buchner, A. & Lang, A.-G. Statistical power analyses using  
G\*Power 3.1: Tests for correlation and regression analyses. *Behavior Research Meth-*  
519 *ods* **41**, 1149–1160 (2009).

520 52.

521 Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. lmerTest package: Tests in  
522 linear mixed effects models. *Journal of Statistical Software* **82**, 1–26 (2017).

523 53.

524 Enders, C. K. & Tofighi, D. Centering predictor variables in cross-sectional multilevel  
525 models: A new look at an old issue. *Psychological Methods* **12**, 121–138 (2007).

526 54.

527 Preece, D. A., Becerra, R., Robinson, K. & Gross, J. J. The emotion regulation  
questionnaire: Psychometric properties in general community samples. *J Pers Assess*  
528 **102**, 348–356 (2020).

529 55.

530 Fleischhauer, M. *et al.* Same or different? Clarifying the relationship of need for  
cognition to personality and intelligence. *Personality & Social Psychology Bulletin*  
531 **36**, 82–96 (2010).

532 56.

533 Fleischhauer, M., Strobel, A. & Strobel, A. Directly and indirectly assessed Need  
for Cognition differentially predict spontaneous and reflective information processing  
534 behavior. *Journal of Individual Differences* **36**, 101–109 (2015).

### Acknowledgements

This research is partly funded by the German Research Foundation (DFG) as part of the Collaborative Research Center (CRC) 940. Additionally, we have applied for funding of the participants' compensation from centralized funds of the Faculty of Psychology at Technische Universität Dresden. Applications for the centralized funds will be reviewed in May. Regardless of whether or not this additional funding will be granted, the study can commence immediately. The funders have/had no role in study design, data collection and analysis, decision to publish or preparation of the manuscript.

### Competing Interests

The authors declare no competing interests.

**Figures and figure captions**



Figure 1

Figure 1. Exemplary visualization of two response patterns. In the top half, the person has a clear preference for one of the three strategies. In the lower half, they have no clear preference and therefore show an inconsistent response pattern. This pattern can also be represented by our paradigm.

Figure 2. Block design of the paradigm. Every participant starts with two “active viewing” blocks containing neutral (Block 1) and negative (Block 2) pictures. Order of the regulation blocks (Blocks 3, 4, and 5) is randomized between participants. After, the discounting procedure takes place. All three regulation strategies are pairwise compared. Before the last block, participants can decide which regulation strategy they want to reapply. Subjective arousal and effort ratings are assessed after each block using a slider on screen with a continuous scale.

Design Table

Question	Hypothesis	Sampling plan (e.g. power analysis)	Analysis Plan	Interpretation given to different outcomes
1.) Do ER strategies reduce emotional arousal? (Manipulation check)	1a) Subjective arousal (arousal rating) is lower after using an emotion regulation strategy (distraction, distancing, suppression) compared to active viewing.	<p>F tests - ANOVA: Repeated measures, within factors            Analysis: A priori: Compute required sample size  <u>Input:</u>            Effect size <math>f = 0.50</math> (<math>\eta_p^2 = 0.20</math>) (Scheffé et al., 2021)  <math>\alpha</math> err prob = 0.05            Power (<math>1 - \beta</math> err prob) = 0.95            Number of groups = 1            Number of measurements = 4            Corr among rep measures = 0.5            Nonsphericity correction <math>\epsilon = 1</math></p> <p><u>Output:</u>            Noncentrality parameter <math>\lambda = 20.0</math>            Critical F = 2.9603513            Numerator df = 3.0            Denominator df = 27.0            Total sample size = 10            Actual power = 0.95210128</p>	<p>Repeated measures ANOVA with four linear contrasts, comparing the subjective arousal ratings of four blocks (active viewing, distraction, distancing, suppression).</p> <p>ANOVA is calculated using <code>aov_ez()</code> function of the <code>afex</code>-package, estimated marginal means are calculated using <code>emmeans()</code> function from the <code>emmeans</code>-package, pairwise contrasts are calculated using <code>pairs()</code>.</p> <p>Bayes factors are computed for the ANOVA and each contrast using the <code>BayesFactor</code>-package.</p>	<p>ANOVA yields <math>p &lt; .05</math> is interpreted as arousal ratings changing significantly with blocks. Values of arousal ratings are interpreted as equal between blocks if <math>p &gt; .05</math>.</p> <p>Each contrast yielding <math>p &lt; .05</math> is interpreted as arousal ratings being different between those two blocks, magnitude and direction are inferred from the respective estimate. Values of arousal ratings are interpreted as equal between blocks if <math>p &gt; .05</math>.</p> <p>The Bayes factor <math>BF_{10}</math> is reported alongside every <math>p</math>-value to assess the strength of evidence.</p>
2.) Do ER strategies reduce physiological responding? (Manipulation check)	2a) Physiological responding ( <i>corrugator</i> muscle activity) is lower after using an emotion regulation strategy (distraction, distancing, suppression) compared to active viewing.	<p>F tests - ANOVA: Repeated measures, within factors            Analysis: A priori: Compute required sample size  <u>Input:</u>            Effect size <math>f = 0.1605</math> (Zaehring et al., 2020)  <math>\alpha</math> err prob = 0.05            Power (<math>1 - \beta</math> err prob) = 0.95            Number of groups = 1            Number of measurements = 4            Corr among rep measures = 0.5</p>	<p>Repeated measures ANOVA with four linear contrasts, comparing the <i>corrugator</i> muscle activity of four blocks (active viewing, distraction, distancing, suppression).</p> <p>ANOVA is calculated using <code>aov_ez()</code> function of the <code>afex</code>-package, estimated marginal means are calculated using <code>emmeans()</code> function from the</p>	<p>ANOVA yields <math>p &lt; .05</math> is interpreted as <i>corrugator</i> muscle activity changing significantly with blocks. Values of <i>corrugator</i> muscle activity are interpreted as equal between blocks if <math>p &gt; .05</math>.</p> <p>Each contrast yielding <math>p &lt; .05</math> is interpreted as <i>corrugator</i> muscle activity being different between those two blocks, magnitude and direction are inferred from the respective estimate.</p>

		<p>Nonsphericity correction <math>\epsilon = 1</math></p> <p><u>Output:</u>  Noncentrality parameter <math>\lambda = 17.5169700</math>  Critical F = 2.6404222  Numerator df = 3.0  Denominator df = 252  Total sample size = 85  Actual power = 0.9509128</p>	<p>emmeans-package, pairwise contrasts are calculated using pairs().</p> <p>Bayes factors are computed for the ANOVA and each contrast using the BayesFactor-package.</p>	<p>Values of <i>corrugator</i> muscle activity are interpreted as equal between blocks if <math>p &gt; .05</math>.</p> <p>The Bayes factor <i>BF10</i> is reported alongside every <math>p</math>-value to assess the strength of evidence.</p>
	<p>2b) Physiological responding (<i>levator</i> muscle activity) is lower after using an emotion regulation strategy (distraction, distancing, suppression) compared to active viewing.</p>	<p>F tests - ANOVA: Repeated measures, within factors  Analysis: A priori: Compute required sample size  <u>Input:</u>  Effect size <math>f = 0.1605</math> (Zaehringer et al., 2020)  <math>\alpha</math> err prob = 0.05  Power (<math>1 - \beta</math> err prob) = 0.95  Number of groups = 1  Number of measurements = 4  Corr among rep measures = 0.5  Nonsphericity correction <math>\epsilon = 1</math></p> <p><u>Output:</u>  Noncentrality parameter <math>\lambda = 17.5169700</math>  Critical F = 2.6404222  Numerator df = 3.0  Denominator df = 252  Total sample size = 85  Actual power = 0.9509128</p>	<p>Repeated measures ANOVA with four linear contrasts, comparing the <i>levator</i> muscle activity of four blocks (active viewing, distraction, distancing, suppression).</p> <p>ANOVA is calculated using aov_ez() function of the afex-package, estimated marginal means are calculated using emmeans() function from the emmeans-package, pairwise contrasts are calculated using pairs().</p> <p>Bayes factors are computed for the ANOVA and each contrast using the BayesFactor-package.</p>	<p>ANOVA yields <math>p &lt; .05</math> is interpreted as <i>levator</i> muscle activity changing significantly with blocks. Values of <i>levator</i> muscle activity are interpreted as equal between blocks if <math>p &gt; .05</math>.</p> <p>Each contrast yielding <math>p &lt; .05</math> is interpreted as <i>levator</i> muscle activity being different between those two blocks, magnitude and direction are inferred from the respective estimate. Values of <i>levator</i> muscle activity are interpreted as equal between blocks if <math>p &gt; .05</math>.</p> <p>The Bayes factor <i>BF10</i> is reported alongside every <math>p</math>-value to assess the strength of evidence.</p>

3.) Do ER strategies require cognitive effort? (Manipulation check)	3a) Subjective effort (effort ratings) is greater after using an emotion regulation strategy (distraction, distancing, suppression) compared to active viewing.	<p>F tests - ANOVA: Repeated measures, within factors Analysis: A priori: Compute required sample size <u>Input:</u> Effect size <math>f = 0.2041241</math> (<math>\eta_p^2 = 0.04</math>) (Scheffé et al., 2021) <math>\alpha</math> err prob = 0.05 Power (<math>1 - \beta</math> err prob) = 0.95 Number of groups = 1 Number of measurements = 4 Corr among rep measures = 0.5 Nonsphericity correction <math>\epsilon = 1</math></p> <p><u>Output:</u> Noncentrality parameter <math>\lambda = 17.6666588</math> Critical F = 2.6625685 Numerator df = 3.0 Denominator df = 156.0 Total sample size = 53 Actual power = 0.95206921</p>	<p>Repeated measures ANOVA with four linear contrasts, comparing the subjective effort ratings of four blocks (active viewing, distraction, distancing, suppression).</p> <p>ANOVA is calculated using <code>aov_ez()</code> function of the <code>afex</code>-package, estimated marginal means are calculated using <code>emmeans()</code> function from the <code>emmeans</code>-package, pairwise contrasts are calculated using <code>pairs()</code>.</p> <p>Bayes factors are computed for the ANOVA and each contrast using the <code>BayesFactor</code>-package.</p>	<p>ANOVA yields <math>p &lt; .05</math> is interpreted as effort ratings changing significantly with blocks. Values of effort ratings are interpreted as equal between blocks if <math>p &gt; .05</math>.</p> <p>Each contrast yielding <math>p &lt; .05</math> is interpreted as effort ratings being different between those two blocks, magnitude and direction are inferred from the respective estimate. Values of effort ratings are interpreted as equal between blocks if <math>p &gt; .05</math>.</p> <p>The Bayes factor <i>BF</i><sub>10</sub> is reported alongside every <math>p</math>-value to assess the strength of evidence.</p>
	3b) Majority of participants reuse the strategy that was least effortful for them.	-	Subjects are asked about the reasons for their choice in the follow-up survey. These answers are classified into categories and counted.	The percentage choice of strategies is described descriptively.
4.) Which variables can predict individual subjective values of ER strategies?	4a) Subjective effort ratings negatively predict subjective values of ER strategies.	<p>t tests - Linear multiple regression: Fixed model, single regression coefficient Analysis: A priori: Compute required sample size</p>	Multilevel model of SVs with level-1-predictors subjective effort, subjective arousal, <i>corrugator</i> , and <i>levator</i> muscle activity using subject specific	Fixed effects yield $p < .05$ are interpreted as subjective values are related to subjective effort. Subjective values are interpreted as not being related to subjective effort if $p > .05$ .



		<u>Input:</u> Tail(s) = One Effect size $f^2 = 0.34$ (Since there are no findings in this respect yet, we have inferred from the effect size in the closest-similar model: Westbrook et al., 2013) $\alpha$ err prob = 0.05 Power ( $1-\beta$ err prob) = 0.95 Number of predictors = 4 <u>Output:</u> Noncentrality parameter $\delta = 3.4$ Critical t = 1.6991270 Df = 29 Total sample size = 34 Actual power = 0.9529571	intercepts and allowing random slopes for ER strategies.  The null model and the random slopes model are calculated using lmer() of the lmerTest-package.   Bayes factors are computed for the MLM using the BayesFactor-package.	The Bayes factor <i>BF10</i> is reported alongside every <i>p</i> -value to assess the strength of evidence.  Fixed effects yield $p < .05$ are interpreted as subjective values are related to subjective arousal. Subjective values are interpreted as not being related to subjective arousal if $p > .05$ .  The Bayes factor <i>BF10</i> is reported alongside every <i>p</i> -value to assess the strength of evidence.
	4b) Subjective arousal ratings negatively predict subjective values of ER strategies.			
	4c) <i>Corrugator</i> muscle activity negatively predict subjective values of ER strategies.			Fixed effects yield $p < .05$ are interpreted as subjective values are related to <i>corrugator</i> activity. Subjective values are interpreted as not being related to <i>corrugator</i> activity if $p > .05$ .  The Bayes factor <i>BF10</i> is reported alongside every <i>p</i> -value to assess the strength of evidence.
	4d) <i>Levator</i> muscle activity negatively predict subjective values of ER strategies.			Fixed effects yield $p < .05$ are interpreted as subjective values are related to <i>levator</i> activity. Subjective values are interpreted as not being related to <i>levator</i> activity if $p > .05$ .  The Bayes factor <i>BF10</i> is reported alongside every <i>p</i> -value to assess the strength of evidence.

5.) Is the effort required for an ER strategy the best predictor for subjective values of ER strategies?	5a) Subjective values decline with increasing effort, even after controlling for task performance measured by subjective arousal ratings, <i>corrugator</i> and <i>levator</i> muscle activity.	t tests - Linear multiple regression: Fixed model, single regression coefficient Analysis: A priori: Compute required sample size <u>Input:</u> Tail(s) = One Effect size $f^2 = 0.34$ (Since there are no findings in this respect yet, we have inferred from the effect size in the closest-similar model: Westbrook et al., 2013) $\alpha$ err prob = 0.05 Power (1- $\beta$ err prob) = 0.95 Number of predictors = 4 <u>Output:</u> Noncentrality parameter $\delta = 3.4$ Critical t = 1.6991270 Df = 29 Total sample size = 34 Actual power = 0.9529571		Fixed effects yield $p < .05$ are interpreted as subjective values changing significantly with ER strategy. Subjective values are interpreted as equal between ER strategies if $p > .05$ .  The Bayes factor <i>BF10</i> is reported alongside every $p$ -value to assess the strength of evidence.
6.) Are subjective values related to flexible emotion regulation?	6a) The higher the subjective value, the more likely the respective strategy is chosen.	1) $\chi^2$ tests – Goodness-of-fit tests_ Contingency tables Analysis: A priori: Compute required sample size <u>Input:</u> Effect size $\omega = 0.5$ (Based on our theoretical considerations, we assume a large effect) $\alpha$ err prob = 0.05 Power (1- $\beta$ err prob) = 0.95 Df = 1 <u>Output:</u> Noncentrality parameter $\lambda = 19.8$ Critical $\chi^2 = 11.0704977$	1) Chi-squared test with the variables “predicted choice” (= highest SV of each participant) and “choice” (Strategy 1, 2, or 3)  2) Ordinal regression with dependent variable “Choice” (Strategy 1, 2, or 3) and independent variables “SV strategy 1”, “SV strategy 2” and “SV strategy 3”.	1) $\chi^2$ yields $p < .05$ is interpreted as predicted choice (highest SV of each participant) and actual choice show significant consistency. Predicted choice and actual choice are interpreted as independent if $p > .05$ .  The Bayes factor <i>BF10</i> is reported alongside every $p$ -value to assess the strength of evidence.  2) Ordinal logistic regression yields $p < .05$ is interpreted as the respective subjective value has a significant

		<p>Total sample size = 52 Actual power = 0.9500756</p> <p>2) z tests –Logistic regression Analysis: A priori: Compute required sample size <u>Input:</u> Tails: One <math>\Pr(Y=1 X=1)</math> H1 = 0.80 (Based on our theoretical considerations, that a higher SVs should lead almost certainly to the choice of the respective strategy) <math>\Pr(Y=1 X=1)</math> H0 = 0.333 (Based on theoretical considerations: if all SVs are equal, choice is on chance level) <math>\alpha</math> err prob = 0.05 Power (1-<math>\beta</math> err prob) = 0.95 <math>R^2</math> other X = 0 X distribution: normal X param <math>\mu</math> = 0 X param <math>\sigma</math> = 1 <u>Output:</u> Critical z = 1.6448536 Total sample size = 25 Actual power = 0.9528726</p>		<p>influence on the OR of the choice of a strategy. Respective SV is interpreted as not related to choice if <math>p &gt; .05</math>.</p> <p>The Bayes factor <i>BF10</i> is reported alongside every <math>p</math>-value to assess the strength of evidence.</p>
	6b) Subjective values are lower and decline stronger when ER flexibility is lower.	<p>t tests – Linear multiple regression: Fixed model, single regression coefficient Analysis: A priori: compute required sample size <u>Input:</u> Tail(s) = One</p>	<p>SVs will be ordered by magnitude. Values will be fitted in a GLM to estimate the individual intercept and slope.</p> <p>A linear regression will be computed with intercept and</p>	<p><math>\beta</math> yield <math>p &lt; .05</math> are interpreted as significant association between predictor (intercept, slope) and ER flexibility. The direction of effect is interpreted according to sign (negative or positive). <math>p</math> – values <math>&gt; .05</math> are interpreted as no association between predictor and ER flexibility.</p>

		<p>Effect size <math>f^2 = 0.15</math> (as there is no evidence in the literature, we assume a medium sized effect)</p> <p><math>\alpha</math> err prob = 0.05</p> <p>Power (1-<math>\beta</math> err prob) = 0.95</p> <p>Number of predictors = 2</p> <p><u>Output:</u></p> <p>Noncentrality parameter <math>\delta = 3.316662</math></p> <p>Critical t = 1.69665997</p> <p>Df = 71</p> <p>Total sample size = 74</p> <p>Actual power = 0.95101851</p>	slope as predictors and FlexER score as criterion.	The Bayes factor <i>BF10</i> is reported alongside every <i>p</i> -value to assess the strength of evidence.
Exploratory: Are individual subjective values of ER strategies related to personality traits?			<p>Multilevel model of SVs with level-1-predictors subjective effort, subjective arousal, <i>corrugator</i>, and <i>levator</i> muscle activity and level-2-predictors NFC and self-control using subject specific intercepts and allowing random slopes for ER strategies.</p> <p>The null model and the random slopes model are calculated using lmer() of the lmerTest-package.</p> <p>Bayes factors are computed for the MLM using the BayesFactor-package.</p>	<p>Fixed effects yield <math>p &lt; .05</math> are interpreted as subjective values are related to NFC and self-control. Subjective values are interpreted as not being related to subjective effort if <math>p &gt; .05</math>.</p> <p>The Bayes factor <i>BF10</i> is reported alongside every <i>p</i>-value to assess the strength of evidence.</p>




Figure 2

565

## **Supplementary Material**

## Supplementary Material 1

Table S1

List of IAPS (Lang, Bradley, and Cuthbert, 2008) and EmoPicS (Wessa et al., 2010) used in the ER paradigm.

	Neutral	Negative 1	Negative 2	Negative 3	Negative 4	Negative 5
	083 <sup>†</sup>	225 <sup>†</sup>	210 <sup>†</sup>	208 <sup>†</sup>	227 <sup>†</sup>	223 <sup>†</sup>
	107 <sup>†</sup>	230 <sup>†</sup>	218 <sup>†</sup>	219 <sup>†</sup>	252 <sup>†</sup>	238 <sup>†</sup>
	124 <sup>†</sup>	255 <sup>†</sup>	222 <sup>†</sup>	226 <sup>†</sup>	1051*	245 <sup>†</sup>
	140 <sup>†</sup>	327 <sup>†</sup>	228 <sup>†</sup>	253 <sup>†</sup>	2800*	2981*
	143 <sup>†</sup>	1111*	246 <sup>†</sup>	254 <sup>†</sup>	3061*	3016*
	7000*	3017*	251 <sup>†</sup>	326 <sup>†</sup>	3230*	3101*
	7002*	3022*	2703*	1301*	6561*	3181*
	7004*	3180*	3051*	3350*	6838*	3215*
	7006*	3280*	3160*	6242*	9120*	3220*
	7009*	6190*	3185*	6410*	9181*	3225*
	7021*	6244*	3301*	6555*	9185*	6020*
	7025*	6836*	6562*	6825*	9230*	6571*
	7041*	9180*	9031*	6940*	9254*	6831*
	7100*	9182*	9040*	8230*	9295*	8231*
	7150*	9253*	9042*	9041*	9332*	9373*
	7185*	9300*	9043*	9140*	9411*	9400*
	7211*	9326*	9145*	9340*	9420*	9402*
	7224*	9424*	9160*	9409*	9421*	9403*
	7233*	9425*	9184*	9570*	9599*	9405*
	7235*	9920*	9904*	9800*	9905*	9423*
<b>Valence</b>	4.86 ± 0.49	2.84 ± 0.57	2.64 ± 0.46	2.82 ± 0.62	2.65 ± 0.75	2.74 ± 0.70
<b>Arousal</b>	3.01 ± 0.61	5.62 ± 0.34	5.58 ± 0.38	5.60 ± 0.39	5.61 ± 0.41	5.63 ± 0.37

Note. \* Pictures taken from the IAPS (Lang, Bradley, and Cuthbert, 2008); <sup>†</sup> Pictures taken from the EmoPicS (Wessa et al., 2010).

## Detailed information on psychometric measures

*WHO-5.* General psychological well-being is assessed using the WHO-5 scale<sup>27,28</sup>. 5  
Items such as “Over the past 2 weeks I have felt calm and relaxed.” are rated on a 6-point  
Likert scale ranging from 0 (at no time) to 5 (all of the time). The German version of the  
scale showed a high internal consistency (Cronbach’s  $\alpha = .92$ )<sup>28</sup>.

*Connor-Davidson Resilience Scale.* Resilience is assessed using the Connor-Davidson  
Resilience Scale (CD-RISC)<sup>29–31</sup>. 10 items such as “I am able to adapt to change.” are  
rated on a scale from 0 (not true at all) to 4 (true nearly all the time). The 10-item version  
showed a high internal consistency (Cronbach’s  $\alpha = .84$ ) and a satisfactory retest-reliability  
of  $r_{tt} = .81$  across 6 months<sup>30</sup>.

*Emotion Regulation Questionnaire.* Habitual use of reappraisal and suppression is  
measured using the 10-item Emotion Regulation Questionnaire (ERQ)<sup>6,32</sup>. The scale has  
items such as “I keep my emotions to myself” (ERQ-suppression - 4 items) and “When I’m  
faced with a stressful situation, I make myself think about it in a way that helps me stay  
calm” (ERQ-reappraisal - 6 items), which are answered on a 7-point Likert scale ranging  
from 1 (strongly disagree) to 7 (strongly agree), and has acceptable to high internal  
consistency (Cronbach’s  $\alpha > .75$ ).<sup>54</sup>

*FlexER Scale.* Flexible use of ER strategies is assessed using the FlexER Scale<sup>33</sup> with  
items such as “If I want to feel less negative emotions, I have several strategies to achieve  
this.”, which are answered on a 4-point scale ranging from “strongly agree” to “strongly  
disagree”. Psychometric properties are currently under investigation.

*Implicit Theories Questionnaire.* Implicit theories of willpower in emotional control  
are assessed using the Implicit Theories Questionnaire of.<sup>34</sup> 4 items such as “Having to  
control a strong emotion makes you exhausted and you are less able to manage your feelings  
right afterwards.” are rated on a 6-point scale ranging from 1 (fully agree) to 6 (do not  
agree at all). The questionnaire showed an internal consistency of Cronbach’s  $\alpha = .87$ <sup>34</sup>.



*Need for Cognition Scale.* Need for Cognition (NFC) is assessed with the 16-item short version of the German NFC scale.<sup>36</sup> Responses to each item (e.g., “Thinking is not my idea of fun”, recoded) are recorded on a 7-point Likert scale ranging from -3 (completely disagree) to +3 (completely agree) and are summed to the total NFC score. The scale shows comparably high internal consistency (Cronbach’s  $\alpha > .80$ )<sup>36,55</sup> and a retest reliability of  $r_{tt} = .83$  across 8 to 18 weeks.<sup>56</sup>

*Self-Regulation Scale.* As one measure of self-control, the Self-Regulation Scale (SRS)<sup>38</sup> is used. The scale has 10 items (e.g., “It is difficult for me to suppress thoughts that interfere with what I need to do.”, recoded) on a 4-point scale ranging from 1 (not at all true) to 4 (exactly true). It has high internal consistency [Cronbach’s  $\alpha > .80$ ].<sup>38</sup>

*Brief Self-Control Scale.* As a second measure of self-control, the Brief Self-Control Scale (BSCS)<sup>39,40</sup> is used. It comprises 13 items (e.g., “I am good at resisting temptations”) with a 5-point rating scale ranging from 1 (not at all like me) to 5 (very much like me). The scale shows acceptable internal consistency (Cronbach’s  $\alpha = .81$ )<sup>40</sup>.

*Barratt Impulsiveness Scale.* As a third measure of self-control, the Barratt Impulsiveness Scale (BIS-11)<sup>41,42</sup> is used. Responses to each item (e.g., “I am self-controlled.”, recoded) are assessed on a 4-point scale ranging from 1 (never/rarely) to 4 (almost always/always). An internal consistency of Cronbach’s  $\alpha = .74$  and a retest reliability of  $r_{tt} = .56$  for General Impulsiveness and  $r_{tt} = .66$  for Total Score across 6 month were reported.<sup>42</sup>

*Attentional Control Scale.* Attentional control is measured using the Attentional Control Scale (ACS)<sup>43</sup> with items such as “My concentration is good even if there is music in the room around me”. The 20 items are rated on a 4-point scale ranging from 1 (almost never) to 4 (always). An internal consistency of Cronbach’s  $\alpha = .88$  was reported.<sup>43</sup>

**Pilot study: Subjective arousal in the conditions “Active viewing - neutral”  
and “Active viewing - negative”**

ANOVA:

Effect	df	MSE	F	ges	p.value
block	1, 15	3895.91	34.32 ***	.475	<.001

$BF10 = 1,244.99$

Paired contrasts:

Table 1

*Paired contrasts for the rmANOVA comparing subjective arousal of negative and neutral pictures in the condition "active viewing".*

Contrast	Estimate	<i>SE</i>	<i>df</i>	<i>t</i>	<i>p</i>	<i>BF10</i>	$\eta_p^2$	95% <i>CI</i>
$View_{neutral} - View_{negative}$	-129.28	22.07	15.00	-5.86	0.00	794.78	0.70	[0.43, 1.00]

*Note.* *SE* = standard error, *df* = degrees of freedom, *t* = *t*-statistic, *p* = *p*-value, CI = confidence interval.

Figure:




Figure 3. Subjective arousal ratings for the conditions "Active viewing - neutral" and "Active viewing - negative" visualized as boxplots. Each dot represents the effort rating of a single subject. Bold dots represent outliers.

Pilot study: Subjective arousal in the conditions “Active viewing - negative”, “Distraction”, “Distancing”, and “Suppression”

ANOVA:

Effect	df	MSE	F	ges	p.value
block	2.79, 41.89	2238.27	1.17	.011	.332

$BF10 = 0.11$

Paired contrasts:

Table 2  
*Paired contrasts for the rmANOVA comparing subjective arousal of conditions "Active viewing - negative", "Distraction", "Distancing", and "Suppression".*

Contrast	Estimate	SE	df	t	p	BF10	$\eta_p^2$	95%CI
$View_{negative} - Distraction$	-0.74	16.14	45.00	-0.05	1.00	0.26	4.68e-05	[0.00, 1.00]
$View_{negative} - Distancing$	-5.35	16.14	45.00	-0.33	1.00	0.27	2.43e-03	[0.00, 1.00]
$View_{negative} - Suppression$	-26.23	16.14	45.00	-1.63	0.67	1.25	0.06	[0.00, 1.00]
$Distraction - Distancing$	-4.61	16.14	45.00	-0.29	1.00	0.26	1.81e-03	[0.00, 1.00]
$Distraction - Suppression$	-25.49	16.14	45.00	-1.58	0.73	0.77	0.05	[0.00, 1.00]
$Distancing - Suppression$	-20.88	16.14	45.00	-1.29	1.00	0.52	0.04	[0.00, 1.00]

*Note.* SE = standard error, df = degrees of freedom, t = t-statistic, p = p-value, CI = confidence interval.

630

Figure:




Figure 4. Subjective arousal ratings for the conditions "Active viewing - negative", "Distraction", "Distancing", and "Suppression" visualized as boxplots. Each dot represents the effort rating of a single subject. Bold dots represent outliers.

**Pilot study: Physiological responding (*Corrugator* and *Levator* activity) in the conditions “Active viewing - neutral” and “Active viewing - negative”**

*Corrugator*: ANOVA:

Effect	df	MSE	F	ges	p.value
block	1, 15	1.01	9.70 **	.237	.007

$BF10 = 6,690,401.91$

Paired contrasts:

Table 3

*Paired contrasts for the rmANOVA comparing physiological responding (*Corrugator* activity) of negative and neutral pictures in the condition "active viewing".*

Contrast	Estimate	<i>SE</i>	<i>df</i>	<i>t</i>	<i>p</i>	<i>BF10</i>	$\eta_p^2$	95% <i>CI</i>
$View_{neutral} - View_{negative}$	-1.11	0.36	15.00	-3.11	0.01	5,019,313.20	0.39	[0.09, 1.00]

*Note.* *SE* = standard error, *df* = degrees of freedom, *t* = *t*-statistic, *p* = *p*-value, CI = confidence interval.

*Levator*: ANOVA:

Effect	df	MSE	F	ges	p.value
block	1, 15	0.17	7.72 *	.162	.014

$BF10 = 48.44$

Paired contrasts:

Table 4

*Paired contrasts for the rmANOVA comparing physiological responding (*Levator* activity) of negative and neutral pictures in the condition "active viewing".*

Contrast	Estimate	<i>SE</i>	<i>df</i>	<i>t</i>	<i>p</i>	<i>BF10</i>	$\eta_p^2$	95% <i>CI</i>
$View_{neutral} - View_{negative}$	-0.40	0.14	15.00	-2.78	0.01	41.02	0.34	[0.05, 1.00]

*Note.* *SE* = standard error, *df* = degrees of freedom, *t* = *t*-statistic, *p* = *p*-value, CI = confidence interval.

641

Figures:




Figure 5. Corrugator activity for the conditions "Active viewing - neutral" and "Active viewing - negative" visualized as boxplots. Each dot represents the corrugator activity of a single trial. Bold dots represent outliers.




Figure 6. Levator activity for the conditions "Active viewing - neutral" and "Active viewing - negative" visualized as boxplots. Each dot represents the levator activity of a single trial. Bold dots represent outliers.

Pilot study: Physiological responding (*Corrugator* and *Levator* activity) in the conditions “Active viewing - negative”, “Distraction”, “Distancing”, and “Suppression”

*Corrugator*: ANOVA:

Effect	df	MSE	F	ges	p.value
block	1.53, 22.98	1.16	5.71 *	.189	.015

$$BF10 = 5,257,689.54$$

Paired contrasts:

Table 5

Paired contrasts for the *rmANOVA* comparing physiological responding (*Corrugator* activity) of conditions "Active viewing - negative", "Distraction", "Distancing", and "Suppression".

Contrast	Estimate	SE	df	t	p	BF10	$\eta_p^2$	95%CI
<i>View<sub>negative</sub></i> – <i>Distraction</i>	0.88	0.27	45.00	3.22	0.01	4,962.89	0.19	[0.05, 1.00]
<i>View<sub>negative</sub></i> – <i>Distancing</i>	0.95	0.27	45.00	3.50	0.01	616.63	0.21	[0.06, 1.00]
<i>View<sub>negative</sub></i> – <i>Suppression</i>	0.92	0.27	45.00	3.40	0.01	11,678.82	0.20	[0.06, 1.00]
<i>Distraction</i> – <i>Distancing</i>	0.08	0.27	45.00	0.28	1.00	0.07	1.78e-03	[0.00, 1.00]
<i>Distraction</i> – <i>Suppression</i>	0.05	0.27	45.00	0.18	1.00	0.08	7.22e-04	[0.00, 1.00]
<i>Distancing</i> – <i>Suppression</i>	-0.03	0.27	45.00	-0.10	1.00	0.06	2.36e-04	[0.00, 1.00]

Note. SE = standard error, df = degrees of freedom, t = t-statistic, p = p-value, CI = confidence interval.

*Levator*: ANOVA:

Effect	df	MSE	F	ges	p.value
block	2.07, 31.00	0.20	8.27 **	.225	.001

$$BF10 = 672,341.29$$

Paired contrasts:

Table 6


*Paired contrasts for the rmANOVA comparing physiological responding (Levator activity) of conditions "Active viewing - negative", "Distraction", "Distancing", and "Suppression".*

Contrast	Estimate	SE	df	t	p	BF10	$\eta_p^2$	95%CI
$View_{negative} - Distraction$	0.42	0.13	45.00	3.24	0.01	58.02	0.19	[0.05, 1.00]
$View_{negative} - Distancing$	0.45	0.13	45.00	3.46	0.01	93.49	0.21	[0.06, 1.00]
$View_{negative} - Suppression$	0.62	0.13	45.00	4.79	0.00	6,253.91	0.34	[0.16, 1.00]
$Distraction - Distancing$	0.03	0.13	45.00	0.22	1.00	0.07	1.06e-03	[0.00, 1.00]
$Distraction - Suppression$	0.20	0.13	45.00	1.54	0.78	1.52	0.05	[0.00, 1.00]
$Distancing - Suppression$	0.17	0.13	45.00	1.32	1.00	0.52	0.04	[0.00, 1.00]

*Note.* SE = standard error, df = degrees of freedom, t = t-statistic, p = p-value, CI = confidence interval.

653

Figures:



*Figure 7.* Corrugator activity for the conditions "Active viewing - negative", "Distraction", "Distancing", and "Suppression" visualized as boxplots. Each dot represents the corrugator activity of a single trial. Bold dots represent outliers.






Figure 8. Levator activity for the conditions "Active viewing - negative", "Distraction", "Distancing", and "Suppression" visualized as boxplots. Each dot represents the levator activity of a single trial. Bold dots represent outliers.

Pilot study: Subjective effort in the conditions “Active viewing - negative”, “Distraction”, “Distancing”, and “Suppression”

ANOVA:

Effect	df	MSE	F	ges	p.value
block	2.38, 35.66	4388.19	11.13 ***	.185	<.001

$BF_{10} = 7.40$

Paired contrasts:

Table 7

Paired contrasts for the *rmANOVA* comparing subjective effort of conditions "Active viewing - negative", "Distraction", "Distancing", and "Suppression".

Contrast	Estimate	<i>SE</i>	<i>df</i>	<i>t</i>	<i>p</i>	<i>BF</i> 10	$\eta_p^2$	95% <i>CI</i>
<i>View</i> <sub>negative</sub> – <i>Distancing</i>	-110.72	20.85	45.00	-5.31	0.00	59.77	0.39	[0.20, 1.00]
<i>View</i> <sub>negative</sub> – <i>Distraction</i>	-89.72	20.85	45.00	-4.30	0.00	20.49	0.29	[0.12, 1.00]
<i>View</i> <sub>negative</sub> – <i>Suppression</i>	-88.15	20.85	45.00	-4.23	0.00	33.13	0.28	[0.11, 1.00]
<i>Distraction</i> – <i>Distancing</i>	21.00	20.85	45.00	1.01	1.00	0.50	0.02	[0.00, 1.00]
<i>Distraction</i> – <i>Suppression</i>	22.57	20.85	45.00	1.08	1.00	0.57	0.03	[0.00, 1.00]
<i>Distancing</i> – <i>Suppression</i>	1.57	20.85	45.00	0.08	1.00	0.26	1.27e-04	[0.00, 1.00]

Note. *SE* = standard error, *df* = degrees of freedom, *t* = *t*-statistic, *p* = *p*-value, *CI* = confidence interval.

660

Figure:




Figure 9. Subjective effort ratings for the conditions "Active viewing - negative", "Distraction", "Distancing", and "Suppression" visualized as boxplots. Each dot represents the effort rating of a single subject. Bold dots represent outliers.