

Mathematical Foundations of Computer Science

CS 499, Shanghai Jiaotong University, Dominik Scheder

- Homework assignment published on Monday, 2018-03-05.
- Work on it and submit a first solution or questions by Sunday, 2018-03-11, 12:00 by email to me and the TAs.
- You will receive feedback by Wednesday, 2018-03-14.
- Submit your final solution by Sunday, 2018-03-18 to me and the TAs.

2 Partial Orderings

2.1 Equivalence Relations as a Partial Ordering

An equivalence relation $R \subseteq V \times V$ is basically the same as a partition of V . A *partition* of V is a set $\{V_1, \dots, V_k\}$ where (1) $V_1 \cup \dots \cup V_k = V$ and (2) the V_i are pairwise disjoint, i.e., $V_i \cap V_j = \emptyset$ for $1 \leq i < j \leq k$. For example, $\{\{1\}, \{2, 3\}, \{4\}\}$ is a partition of $\{1, 2, 3, 4\}$ but $\{\{1\}, \{2, 3\}, \{1, 4\}\}$ is not.

Exercise 2.1. Let E_4 be the set of all equivalence relations on $\{1, 2, 3, 4\}$. Note that E_4 is ordered by set inclusion, i.e.,

$$(E_4, \{(R_1, R_2) \in E_4 \times E_4 \mid R_1 \subseteq R_2\})$$

is a partial ordering.

1. Draw the Hasse diagram of this partial ordering in a nice way.
2. What is the size of the largest chain?
3. What is the size of the largest antichain?

2.2 Chains and Antichains

Define the partially ordered set (\mathbb{N}_0^n, \leq) as follows: $x \leq y$ if $x_i \leq y_i$ for all $1 \leq i \leq n$. For example, $(2, 5, 4) \leq (2, 6, 6)$ but $(2, 5, 4) \not\leq (3, 1, 1)$.

Exercise 2.2. Consider the infinite partially ordered set (\mathbb{N}_0^n, \leq) .

1. Which elements are minimal? Which are maximal?
2. Is there a minimum? A maximum?
3. Does it have an infinite chain?
4. Does it have arbitrarily large antichains? That is, can you find an antichain A of size $|A| = k$ for every $k \in \mathbb{N}$?

***Exercise 2.3.** Does every infinite subset $S \subseteq \mathbb{N}_0^n$ contain an infinite chain?

Exercise 2.4. Show that (\mathbb{N}_0^n, \leq) has no infinite antichain. **Hint.** Use the previous exercise.

Consider the induced ordering on $\{0, 1\}^n$. That is, for $x, y \in \{0, 1\}^n$ we have $x \leq y$ if $x_i \leq y_i$ for every coordinate $i \in [n]$.

Exercise 2.5. Draw the Hasse diagrams of $(\{0, 1\}^n, \leq)$ for $n = 2, 3$.

Exercise 2.6. Determine the maximum, minimum, maximal, and minimal elements of $\{0, 1\}^n$.

Exercise 2.7. What is the longest chain of $\{0, 1\}^n$?

****Exercise 2.8.** What is the largest antichain of $\{0, 1\}^n$?

2.3 Infinite Sets

In the lecture (and the lecture notes) we have showed that $\mathbb{N} \times \mathbb{N} \cong \mathbb{N}$, i.e., there is a bijection $f : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$. From this, and by induction, it follows quite easily that $\mathbb{N}^k \cong \mathbb{N}$ for every k .

Exercise 2.9. Consider \mathbb{N}^* , the set of all finite sequences of natural numbers, that is, $\mathbb{N}^* = \{\epsilon\} \cup \mathbb{N} \cup \mathbb{N}^2 \cup \mathbb{N}^3 \cup \dots$. Here, ϵ is the empty sequence. Show that $\mathbb{N} \cong \mathbb{N}^*$ by defining a bijection $\mathbb{N} \rightarrow \mathbb{N}^*$.

Exercise 2.10. Show that $R \cong R \times R$. **Hint:** Use the fact that $R \cong \{0, 1\}^{\mathbb{N}}$ and thus show that $\{0, 1\}^{\mathbb{N}} \cong \{0, 1\}^{\mathbb{N}} \times \{0, 1\}^{\mathbb{N}}$.

Proof. We can define f :

$$f : \{0, 1\}^{\mathbb{N}} \times \{0, 1\}^{\mathbb{N}} \rightarrow \{0, 1\}^{\mathbb{N}}$$

$$((a_{11}, a_{12}, a_{13}, \dots), (a_{21}, a_{22}, a_{23}, \dots)) \mapsto (a_{11}, a_{21}, a_{12}, a_{22}, \dots)$$

For any element $\mathbf{a} \in \{0, 1\}^{\mathbb{N}}$, \mathbf{a} can be represented by a combination of unique ordered pair $(\mathbf{a}_1, \mathbf{a}_2)$ where $\mathbf{a}_1, \mathbf{a}_2 \in \{0, 1\}^{\mathbb{N}}$. In this way, f is bijective. Then we can get $R \cong \{0, 1\}^{\mathbb{N}} \cong \{0, 1\}^{\mathbb{N}} \times \{0, 1\}^{\mathbb{N}} \cong R \times R$. \square

Exercise 2.11. Consider $\mathbb{R}^{\mathbb{N}}$, the set of all infinite sequences (r_1, r_2, r_3, \dots) of real numbers. Show that $\mathbb{R} \cong \mathbb{R}^{\mathbb{N}}$. **Hint:** Again, use the fact that $\mathbb{R} \cong \{0, 1\}^{\mathbb{N}}$.

Proof. \square

Next, let us view $\{0, 1\}^{\mathbb{N}}$ as a partial ordering: given two elements $\mathbf{a}, \mathbf{b} \in \{0, 1\}^{\mathbb{N}}$, that is, sequences $\mathbf{a} = (a_1, a_2, \dots)$ and $\mathbf{b} = (b_1, b_2, \dots)$, we define $\mathbf{a} \leq \mathbf{b}$ if $a_i \leq b_i$ for all $i \in \mathbb{N}$. Clearly, $(0, 0, \dots)$ is the minimum element in this ordering and $(1, 1, \dots)$ the maximum.

Exercise 2.12. Give a countably infinite chain in $\{0, 1\}^{\mathbb{N}}$. Remember that a set A is countably infinite if $A \cong \mathbb{N}$.

Solution. For set $A \subseteq \{0, 1\}^{\mathbb{N}}$, let $A = \{\mathbf{a}_i \in \{0, 1\}^{\mathbb{N}} \mid \text{if } k < i, a_k = 1, \text{ else, } a_k = 0\}$, any two elements in A are comparable

We can define a function f :

$$f : N \rightarrow A, x \mapsto \mathbf{a}_x$$

which is obviously bijective. In this way, A is a countably infinite chain. ■

Exercise 2.13. Find a countably infinite antichain in $\{0, 1\}^{\mathbb{N}}$.

Solution. For set $A \subseteq \{0, 1\}^{\mathbb{N}}$, let $A = \{\mathbf{a}_i \in \{0, 1\}^{\mathbb{N}} | a_i = 1 \text{ and other bits is } 0\}$. Any two elements in A is uncomparable since each has one bit larger than the other.

Also, we can define a function f :

$$f : N \rightarrow A, x \mapsto \mathbf{a}_x$$

which is bijective.

In this scense, A is a countably infinite antichain in $\{0, 1\}^{\mathbb{N}}$. ■

Exercise 2.14. Find an uncountable antichain in $\{0, 1\}^{\mathbb{N}}$. That is, an antichain A with $A \cong \mathbb{R}$.

Solution. Define a function f :

$$f : \{0, 1\}^{\mathbb{N}} \rightarrow A, (a_1, a_2, a_3 \dots) \mapsto (a_1, 1 - a_1, a_2, 1 - a_2, \dots)$$

f is a bijective function, and $\{0, 1\}^{\mathbb{N}} \cong A$

For $\mathbf{a}, \mathbf{b} \in A$, if $\mathbf{a} \neq \mathbf{b}$ and \mathbf{a}, \mathbf{b} are comparable, let $\mathbf{a} < \mathbf{b}$.

There must exist $a_i < b_i$, but $1 - a_i > 1 - b_i$, so \mathbf{a} and \mathbf{b} are uncomparable. Then we get a contradiction.

In this way, A is a uncountable antichain. ■

****Exercise 2.15.** Find an uncountable chain in $\{0, 1\}^{\mathbb{N}}$. That is, an antichain A with $A \cong \mathbb{R}$.