

Mathematical Foundations of Computer Science

CS 499, Shanghai Jiaotong University, Dominik Scheder

Group Name: **NOIDEA**

1 Broken Chessboard and Jumping With Coins

1.1 Tiling a Damaged Checkerboard

Exercise 1.1. Re-write the proof in your own way, using simple English sentences.

Proof. Let's color the board black and beige, just like a typical chessboard. In this case, there are 32 black and 30 beige squares.




Figure 1: The Chessboard

Since a domino stone placed in the board will cover both a black square and a beige square, it's obviously that we can't fully tile the board with 1×2 domino stones.

□

Exercise 1.2. Look at the seriously damaged 8×8 checkerboard. For convenience I already colored it black and white (or rather black and beige):

Try to tile it with domino stones and you will fail. However, since there are 24 black and 24 beige squares, the simple argument from the lecture will fail.

Prove that the above board cannot be tiled. Try to find a short and simple argument!




Figure 2: The “Corridor”

Proof. Solution: In the left half there's 13 black squares and 11 white squares. cut it into 2 halves. A complete domino tiling “marries” one black block to one white tiling. In the left half, at least 2 white squares have to “marry” a black foreigner from the right half. But only 1 black can be married to a white block from the right half. Thus, you cannot cover the complete board.

□

1.2 Jumping with Coins

Exercise 1.3. You jump around with two coins. Show that you cannot increase the distance between the two coins.

Proof. Clearly, the distance between two coins is fixed in one jump.



Figure 3: Two Coins

□

Exercise 1.4. You jump around with three coins. Show that you cannot start with an equilateral triangle and end up with a bigger equilateral triangle. Give a simple proof!

Proof. During a jump, either the side or the height of the triangle will not change, thus the area stays the same.




Figure 4: Three Coins

□

You jump around with four coins which in the beginning form a square of side length 1.

Exercise 1.5. Show that you cannot form a square of side length 2.

Proof. We use the coordinate to express the position of coins. The initial coordinate will be

$$(odd, odd), (odd, even), (even, even), (even, odd).$$

If a coin at (a, b) jump over the coin at (c, d) , the new position (e, f) will be

$$e = 2c - a, f = 2d - b$$

Both $2c$ and $2d$ are even, so the parity of e, f will be the same as a, b . It means that the parity of coordinate of a coin will not change during a jump.

If a square of side length 2 is formed, the parity of the coordinate will be different with the initial one, which is impossible. \square

Exercise 1.6. Show that you cannot achieve a position in which two coins are at the same position.

Proof. We have proved that the parity of coordinate of a coin will not change during a jump. Hence two coins with different initial parity of the coordinate can not be at the same position.

And we know that all the four coins have different initial parity, so there won't be any coins at the same position. \square

Exercise 1.7. Show that you cannot form a larger square.

Proof. If we can make the four coins form a larger square with side length $n > 1$ in a few jumps. Then if the initial side length is $1/n$ we can use the same steps of jump to form a square of side length 1.

It means that with the converse steps, We can form a smaller square with side length $1/n < 1$ with initial side length 1, which requires that the coordinates of coins are not integer.

Obviously, it's impossible.

- 1. Everything is reversible.
- 2. If we can form a larger square, we can also form a smaller square.
- 3. This is impossible since you always stay on the grid
- P.S. Hexagon can be also proofed as above but not pentagon.

\square

2 Exclusion-Inclusion

2.1 Sets

Exercise 2.1.

1. *Proof.* As is shown in the Venn diagram below, $|A| + |B|$ add the common part $|A \cap B|$ twice. So it should be subtracted once if we want to count $|A \cup B|$. \square




Figure 5: Venn Diagram

2. *Solution.* $|A \cup B \cup C| = |A| + |B| + |C| - |A \cap B| - |A \cap C| - |B \cap C| + |A \cap B \cap C|$ \blacksquare
3. *Solution.* $|A \cup B \cup C \cup D| = |A| + |B| + |C| + |D| - |A \cap B| - |A \cap C| - |A \cap D| - |B \cap C| - |B \cap D| - |C \cap D| + |A \cap B \cap C| + |A \cap B \cap D| + |A \cap C \cap D| + |B \cap C \cap D| - |A \cap B \cap C \cap D|$ \blacksquare

Exercise 2.2.

Solution. $|A_1 \cup \dots \cup A_n| = \sum_{i=1}^n |A_i| - \sum_{i,j:1 \leq i < j \leq n} |A_i \cap A_j| + \sum_{i,j,k:1 \leq i < j < k \leq n} |A_i \cap A_j \cap A_k| - \dots + (-1)^{n-1} |A_1 \cap \dots \cap A_n|$ \blacksquare

Exercise 2.3.

Proof. \square

1. proof using induction on n

First, let $A_{n,k} = \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} |A_{i_1} \cap A_{i_2} \dots \cap A_{i_k}|$, which denotes the sum of all the possible k-wise intersections in $\{A_1, A_2, \dots, A_n\}$.

Then the Inclusion-exclusion principle which we want to prove is as follows:

$$|A_1 \cup \dots \cup A_n| = \sum_{k=1}^n (-1)^{k+1} A_{n,k}$$

The theorem holds for $n = 1$, obviously.

The theorem holds for $n = 2$, as is showed in the 2.1.1

For the induction step, we want to show if it holds for $n - 1$, then it holds for n .

$$\begin{aligned} |A_1 \cup \dots \cup A_n| &= |A_1 \cup \dots \cup A_{n-1}| + |A_n| - |(A_1 \cup \dots \cup A_{n-1}) \cap A_n| \\ &= \sum_{k=1}^{n-1} (-1)^{k+1} A_{n-1,k} + |A_n| - |(A_1 \cup \dots \cup A_{n-1}) \cap A_n| \\ &= \sum_{k=1}^{n-1} (-1)^{k+1} A_{n-1,k} + |A_n| - |(\cup_{i=1}^{n-1} (A_i \cap A_n))| \end{aligned} \tag{1}$$

Let $B_i = (A_i \cap A_n)$.

Similarly, let $B_{n-1,k} = \sum_{1 \leq i_1 < i_2 < \dots \leq n-1} |B_{i_1} \cap B_{i_2} \dots \cap B_{i_k}|$, which denotes the sum of all the possible k -wise intersections in $\{B_1, B_2, \dots, B_{n-1}\}$.

(1) now becomes

$$\sum_{k=1}^{n-1} (-1)^{k+1} A_{n-1,k} + |A_n| - |(\cup_{i=1}^{n-1} B_i)| \tag{2}$$

Similarly, it holds:

$$|B_1 \cup \dots \cup B_{n-1}| = \sum_{k=1}^{n-1} (-1)^{k+1} B_{n-1,k} \tag{3}$$

(2) now becomes

$$\sum_{k=1}^{n-1} (-1)^{k+1} A_{n-1,k} + |A_n| + \sum_{k=1}^{n-1} (-1)^k B_{n-1,k} \quad (4)$$

In addition,

$$|A_n| = (-1)^{1+1} |A_n| \quad (5)$$

Thus,

$$|A_n| + \sum_{k=1}^{n-1} (-1)^k B_{n-1,k} = \sum_{k=1}^n (-1)^{k+1} A_{n,k} - \sum_{k=1}^{n-1} (-1)^{k+1} A_{n-1,k} \quad (6)$$

Then equation(4) finally becomes: $\sum_{k=1}^n (-1)^{k+1} A_{n,k}$

2. proof not using induction on n

First, let $A = |A_1 \cup \dots \cup A_{n-1}|$.

Function $P_S(x)$ defined as if set S includes element x , then $P_S(x) = 1$, else $P_S(x) = 0$.

(1) If $P_A(x) = 1$, there must exist an i that $P_{A_i}(x) = 1$. In this way:
 $(P_A(x) - P_{A_1}(x))(P_A(x) - P_{A_2}(x)) \dots (P_A(x) - P_{A_n}(x)) = 0$

(2) According to the properties of set, $P_{A_i}(x)P_{A_j}(x) = P_{A_i \cap A_j}(x)$.

(3) Let $P_{n,k}$ denotes $P_{A_{i_1} \cap A_{i_2} \dots \cap A_{i_k}} (1 \leq i_1 < i_2 < \dots \leq n)$.

Then decompose the first equation, we can have:

$$P_A(x) = \sum_{k=1}^n P_{n,k}$$

which can be demonstrated as:

$$|A_1 \cup \dots \cup A_n| = \sum_{k=1}^n (-1)^{k+1} A_{n,k}$$

solution: Let $X = A_1 \cup \dots \cup A_n$. Then $LHS = \sum_{a \in X} 1, RHS = \sum_{\phi \neq I \in [n]} (-1)^{|I|+1} \sum_{a \in X} [a \subseteq A_I]$ **Claim:** LHS=RHS for each $a \in X$ indi-

vidually, namely $1 = \sum_{\phi \neq I \subseteq [n]} (-1)^{|I|+1} [a \in A_I]$

Page 100 in the book.

3 Feasible Intersection Patterns

3.1

Exercise 3.1.

Find sets A_1, A_2, A_3, A_4 such that all pairwise intersections have size 3 and all three-wise intersections have size 1.

Formally, 1. $|A_i \cap A_j| = 3$ for all $i, j \in \binom{[4]}{2}$, 2. $|A_i \cap A_j \cap A_k| = 1$ for all $\{i, j, k\} \in \binom{[4]}{3}$.

$$A_1 = \{1, 2, 3, 5, 6, 9\};$$

$$A_2 = \{1, 2, 3, 4, 7, 8\};$$

$$A_3 = \{1, 5, 7, 8, 9, 10\};$$

$$A_4 = \{2, 4, 5, 6, 8, 10\};$$

Exercise 3.2.

In the spirit of the previous questions, let us call a sequence $(a_1, a_2, \dots, a_n) \in \mathbb{N}_0$ feasible if there are sets A_1, \dots, A_n such that all k -wise intersections have size a_k . That is, $|A_i| = a_1$ for all i , $|A_i \cap A_j| = a_2$ for all $i \neq j$ and so on. The previous exercise would thus state that $(5, 3, 1, 0)$ is not feasible, but $(6, 3, 1, 0)$ is, as one solution of Exercise 3.1 shows.

Proof.

□

Assume that there exist such sets A, B, C, D to which $(5, 3, 1, 0)$ is feasible.

$$|A| = |B| = |C| = |D| = 5;$$

$$\text{Since } |A \cap B| = 3, |A \cap B \cap C| = 1,$$

A and B have 3 same elements. A , B and C have 1.

There are 2 elements in $A \cap B$ that are not in C ;

Since $|A \cap C| = 3$, there are 2 elements in A and C but not in B .

There are 2 elements in B and C but not in A .

So, now we have 1 elements in A , B , C , 2 in A, C but not B , 2 in B, C but not A .

Then, set D has the same requirement with C .

Similarly, We have 1 elements in A, B, D, 2 in A,D but not B, 2 in B,D but not A.

A has only 5 elements. So dose B.

Then there is a contradiction. D will have at least 5 same elements with C.

$$|C \cap D| = 5 \neq 3.$$

Exercise 3.3.

Given a sequence (a_1, \dots, a_n) , Find a way to determine whether such a sequence is feasible or not.

Proof.

□

Definition:

Given $A_1, A_2, A_3, \dots, A_n$,
define

$$A_{\{I\}} = \bigcap_{i \in I} A_i.$$

In other words,

$$\begin{aligned} A_{\{i,j,k,\dots\}} &= A_i \cap A_j \cap A_k \cap \dots \\ a_{|I|} &= |A_{\{I\}}|. \\ B_{\{I\}} &= \bigcap_{i \in I} A_i \setminus \bigcup_{j \notin I} A_j. \end{aligned}$$

In other words,

$$\begin{aligned} B_{\{I\}} &\text{ is the number of elements in } \bigcap_{i \in I} A_i \text{ but not in } A_j, j \notin I. \\ b_{|I|} &= |B_{\{I\}}|. \end{aligned}$$

Because every set B_i is not divided by existing boundaries, we can put any number of elements in set B_i .

a. Whether a given B-table is feasible:

Thus, (b_1, b_2, \dots, b_n) is feasible for all situations as long as b_i are nonnegative integers.

Obviously, (a_1, a_2, \dots, a_n) has something to do with (b_1, b_2, \dots, b_n) . We have

b. Given a feasible B-table, how to compute A-table:

$$A_{\{I\}} = \bigcup_{J \geq I} B_{\{J\}};$$

$$\text{Thus, } a_{|I|} = \sum_{J \geq I} b_{|J|} = \sum_{j=i}^{j=n} \binom{n-i}{j-i} b_{|J|}$$

c. Given a A-table, how to compute B-table:

$$\text{Then, } b_i = \sum_{k=0}^{n-i} (-1)^k \binom{n-i}{k} a_{i+k}$$

d. Then apply a:

For (a_1, a_2, \dots, a_n) , replace the a_i into the formula.

If all the results are nonnegative integers. The sequence is feasible.