

Analysis 1

Jiaqi Wang

November 29, 2023

Contents

1	Sets, Spaces and Function	6
1.1	Metric Space	6
1.2	Normed Vector Spaces	6
1.3	The reverse triangle inequality	7
2	Real Numbers	8
2.1	What are the real numbers?	8
2.2	The completeness axiom	8
2.3	Alternative characterizations of suprema and infima	8
2.4	Maxima and minima	9
2.5	The Archimedean property	10
2.6	Computation rules for suprema	10
2.7	Bernoulli's inequality	10
3	Sequences	11
3.1	Sequence	11
3.2	Terminology around sequences	11
3.3	Convergence of sequences	11
3.4	Examples and limits of simple sequences	12
3.5	Uniqueness of limits	12
3.6	More properties of convergent sequences	12
3.7	Limit theorems for sequences taking values in a normed vector space	13
3.8	Index shift	13
4	Real-valued sequences	14
4.1	Terminology	14
4.2	Monotone, bounded sequences and convergent	14
4.3	Limit theorems	15
4.4	The squeeze theorem	15
4.5	Divergence to ∞ and $-\infty$	15
4.6	Limit theorems for improper limits	16
4.7	Standard sequences	16

5	Series	18
5.1	Geometric series	18
5.2	The harmonic series	18
5.3	The hyperharmonic series	18
5.4	Only the tail matters for convergence	18
5.5	Divergence test	18
5.6	Limit laws for series	18
6	Series with positive terms	19
6.1	Comparison test	19
6.2	Limit comparison test	19
6.3	Ratio test	19
6.4	Root test	19
7	Series with general terms	20
7.1	Series with real terms: the Leibniz test	20
7.2	Series characterization of completeness in normed vector space	20
7.3	The Cauchy product	20
8	Subsequences, \limsup and \liminf	21
8.1	Index sequences and subsequences	21
8.2	(Sequential) accumulation points	21
8.3	Subsequences of a converging sequence	21
8.4	\limsup	21
8.5	\liminf	21
8.6	Relations between \lim , \limsup and \liminf	21
9	Point-set topology of metric spaces	22
9.1	Open sets	22
9.2	Closed sets	22
9.3	Cauchy sequences	22
9.4	Completeness	22
9.5	Series characterization of completeness in normed vector spaces	22
10	Compactness	23
10.1	Boundedness and total boundedness	23
10.2	Alternative characterization of compactness	23
11	Limits and continuity	24
11.1	Accumulation points	24
11.2	Limit in an accumulation point	24
11.3	Uniqueness of limits	24
11.4	Sequential characterization of limits	24
11.5	Limit laws	24
11.6	Continuity	24
11.7	Sequential characterization of continuity	24
11.8	Rules for continuous functions	24
11.9	Images of compact sets under continuous functions are compact	24
11.10	Uniform continuity	24

12 Real-valued functions	25
12.1 More limit laws	25
12.2 Building of standard functions	25
12.3 Continuity of standard functions	25
12.4 Limits from the left and from the right	25
12.5 The extended real line	25
12.6 Limits to ∞ or $-\infty$	25
12.7 Limits at ∞ and $-\infty$	25
12.8 The Intermediate Value Theorem	25
12.9 The Extreme Value Theorem	25
12.10 Equivalence of norms	25
12.11 Bounded linear maps and operator norms	25
13 Differentiability	26
13.1 The derivative as a function	26
13.2 Constant and linear maps are differentiable	26
13.3 Bases and coordinates	26
13.4 The matrix representation	26
13.5 The chain rule	26
13.6 Sum, product and quotient rules	26
13.7 Differentiability of components	26
13.8 Differentiability implies continuity	26
13.9 Derivative vanishes in local maxima and minima	26
13.10 The Mean Value Theorem	26
14 Differentiability of standard functions	27
14.1 Global context	27
14.2 Polynomials and rational functions are differentiable	27
14.3 Differentiability of the standard functions	27
15 Directional and partial derivatives	28
15.1 A recurring and very important construction	28
15.2 Directional derivatives	28
15.3 Partial derivatives	28
15.4 The Jacobian of a map	28
15.5 Linearization and tangent planes	28
15.6 The gradient of a function	28
16 The Mean-Value Inequality	29
16.1 The mean-value inequality for functions defined on an interval	29
16.2 The mean-value inequality for functions on general domains	29
16.3 Continuous partial derivatives imply differentiability	29
17 Higher order derivatives	30
17.1 Multilinear maps	30
17.2 Relation to n -fold directional derivatives	30
17.3 A criterion for higher differentiability	30
17.4 Symmetry of second order derivatives	30
17.5 Symmetry of higher-order derivatives	30

18 Polynomials and approximation by polynomials	31
18.1 Homogeneous polynomials	31
18.2 Taylor's theorem	31
18.3 Taylor approximations of standard functions	31
19 Banach fixed point theorem	32
20 Implicit function theorem	33
20.1 The objective	33
20.2 Notation	33
20.3 The implicit function theorem	33
20.4 The inverse function theorem	33
21 Function sequences	34
21.1 Point-wise convergence	34
21.2 Uniform convergence	34
21.3 Preservation of continuity under uniform convergence	34
21.4 Differentiability theorem	34
21.5 The normed vector space of bounded functions	34
22 Function series	35
22.1 The Weierstrass M-test	35
22.2 Conditions for differentiation of function series	35
23 Power series	36
23.1 Convergence of power series	36
23.2 Standard functions defined as power series	36
23.3 Operations with power series	36
23.4 Differentiation of power series	36
23.5 Taylor series	36
24 Riemann integration in one dimension	37
24.1 Riemann integrable functions and the Riemann integral	37
24.2 Sums, products of Riemann integrable functions	37
24.3 Continuous functions are Riemann integrable	37
24.4 The fundamental theorem of calculus	37
25 Riemann integration in multiple dimensions	38
25.1 Partitions in multiple dimensions	38
25.2 Riemann integral on rectangles in \mathbb{R}^n	38
25.3 Properties of the multidimensional Riemann integral	38
25.4 Continuous functions are Riemann integrable	38
25.5 Fubini's theorem	38
25.6 The (topological) boundary of a set	38
25.7 Jordan content	38
25.8 Integration over general domains	38
25.9 The volume of bounded sets	38

26 Change-of-variables Theorem	39
26.1 Polar coordinates	39
26.2 Cylindrical coordinates	39
26.3 Spherical coordinates	39

1 Sets, Spaces and Function

1.1 Metric Space

Definition 1.1.1 – distance Let X be a set. A function $d : X \times X \rightarrow \mathbb{R}$ is called a *distance* on X if it satisfies the following properties:

- (i) Positivity: For all $a, b \in X$, it holds that $d(a, b) \geq 0$.
- (ii) Non-degeneracy: For all $a, b \in X$, if $d(a, b) = 0$, then $a = b$.
- (iii) Symmetry: For all $a, b \in X$, it holds that $d(a, b) = d(b, a)$.
- (iv) Triangle inequality: For all $a, b, c \in X$, it holds that $d(a, c) \leq d(a, b) + d(b, c)$.
- (v) Reflexivity: For all $a \in X$, it holds that $d(a, a) = 0$.

Usually conditions (ii) and (v) are combined into one condition: For all $a, b \in X$, $d(a, b) = 0$ if and only if $a = b$.

Definition 1.1.2 – metric space A metric space is a pair $(X, dist)$, where X is a set and $dist$ is a distance function $dist : X \times X \rightarrow \mathbb{R}$ on X .

Example 1.1.3 Let $X = \{\text{Die Hard, Barbie, Oppenheimer}\}$

d	Die Hard	Barbie	Oppenheimer
Die Hard	0	5	2
Barbie	5	0	3
Oppenheimer	2	3	0

Then d is a distance function on X

Definition 1.1.4 – ball in a metric space Let (X, d) be a metric space. Let $c \in X$ and $r \in \mathbb{R}$. The ball of radius r centered at c is the set

$$B(c, r) = \{x \in X \mid d(c, x) < r\}$$

Example 1.1.5 If $(X, d) = (\mathbb{R}, d_{\mathbb{R}})$, then $B(1, 3) = (-2, 4) = \{x \in \mathbb{R} \mid |x - 1| < 3\}$

Example 1.1.6 Let $X := \{\text{Die Hard, Barbie, Oppenheimer}\}$, with distance defined before. Then $B(\text{Barbie}, 4) = \{\text{Barbie, Oppenheimer}\} = \{x \in X \mid d(x, \text{Barbie}) < 4\}$.

1.2 Normed Vector Spaces

Definition 1.2.1 – norm Let V be a vector space over \mathbb{R} . A norm on V is a function $\|\cdot\| : V \rightarrow \mathbb{R}$ such that

- Positivity: for all $u, v \in V$ we have $\|u\| \geq 0$ and $\|u\| = 0$ if and only if $u = 0$.
- Non-degeneracy: for all $u \in V$ if $\|u\| = 0$ then $u = 0$.
- Absolute Homogeneity: for all $u \in V$ and for all $\lambda \in \mathbb{R}$ we have $\|\lambda u\| = |\lambda| \|u\|$.
- Triangle inequality: for all $u, v \in V$ we have $\|u + v\| \leq \|u\| + \|v\|$.

Example 1.2.2 Let $V = \mathbb{R}^n$. Then $\|\cdot\|_2 : \mathbb{R}^n \rightarrow \mathbb{R}$ defined by $\|x\|_2 = \sqrt{x_1^2 + \cdots + x_n^2}$ is a norm on \mathbb{R}^n .

Proposition 1.2.3 – Let $(V, \|\cdot\|)$ be a normed vector space. Then the function $d : V \times V \rightarrow \mathbb{R}$ defined by $d(u, v) = \|u - v\|$ is a distance on V . And (V, d) is a metric space.

Remark 1.2.4 (Notation for Euclidean distance on \mathbb{R}^d and \mathbb{R}). We will usually write $\text{dist}_{\mathbb{R}^d}$ instead of $\text{dist}_{\|\cdot\|_2}$ for the standard (Euclidean) distance on \mathbb{R}^d . In particular, if $d \geq 2$, we have

$$\text{dist}_{\mathbb{R}^d}(v, w) = \|v - w\|_2 = \sqrt{\sum_{i=1}^d (v_i - w_i)^2}$$

and if $d = 1$ we just have

$$\text{dist}_{\mathbb{R}} = |v - w|$$

And if there is no room for confusion, we will just leave out the subscript altogether and write dist instead of $\text{dist}_{\mathbb{R}^d}$.

1.3 The reverse triangle inequality

Lemma 1.3.1 – Reverse triangle inequality Let $(V, \|\cdot\|)$ be a normed vector space. Then for all $u, v \in V$ we have,

$$|||v|| - ||w||| \leq \|v - w\|$$

2 Real Numbers

2.1 What are the real numbers?

Definition 2.1.1 – Real numbers The real numbers are a complete totally ordered field.

2.2 The completeness axiom

Definition 2.2.1 – Upper and Lower bound We say a number $M \in \mathbb{R}$ is an *upper bound* for a set $A \subseteq \mathbb{R}$ if

$$\forall a \in A [a \leq M].$$

We say a number $m \in \mathbb{R}$ is a *lower bound* for a set $A \subseteq \mathbb{R}$ if

$$\forall a \in A [a \geq m].$$

Given the definition of upper and lower bounds, we define what it means for a set to be bounded from above, bounded from below and just bounded.

Definition 2.2.2 – bounded from above, bounded from below, bounded A set $A \subseteq \mathbb{R}$ is *bounded from above* if there exists an upper bound for A .

A set $A \subseteq \mathbb{R}$ is *bounded from below* if there exists a lower bound for A .

A set $A \subseteq \mathbb{R}$ is *bounded* if it is bounded from above and bounded from below.

Definition 2.2.3 – Least upper bound (supremum) Precisely, M is a *least upper bound* of a subset A if both

1. M is an upper bound of A .
2. For every upper bound $L \in \mathbb{R}$ of A , it holds that $M \leq L$.

Proposition 2.2.4 – Suppose both M and W are a least upper bound of a subset $A \subseteq \mathbb{R}$. Then $M = W$.

Axiom 2.2.5 – Completeness axiom We say that a totally ordered field \mathbf{R} satisfies the *completeness axiom* if every nonempty subset of \mathbf{R} that is bounded from above has a least upper bound.

Lemma 2.2.6 – Every non-empty subset of the real line that is bounded from below has a *largest lower bound*.

Definition 2.2.7 – infimum We usually call the largest lower bound of a non-empty set $A \subseteq \mathbb{R}$ that is bounded from below the *infimum* of A , and we denote it by $\inf A$.

2.3 Alternative characterizations of suprema and infima

Proposition 2.3.1 – alternative characterizations of supremum Let $A \subseteq \mathbb{R}$ be non-empty and bounded from above. Let $M \in \mathbb{R}$. Then M is the supremum of A if and only if

1. M is an upper bound for A ,
2. and

$$\begin{aligned} &\text{for all } \varepsilon > 0, \\ &\text{there exists } a \in A, \\ &a > M - \varepsilon. \end{aligned}$$

Proposition 2.3.2 – alternative characterizations of infimum Let $A \subseteq \mathbb{R}$ be non-empty and bounded from below. Let $m \in \mathbb{R}$. Then m is the infimum of A if and only if

1. m is a lower bound for A ,
2. and

$$\begin{aligned} &\text{for all } \varepsilon > 0, \\ &\text{there exists } a \in A, \\ &a < m + \varepsilon. \end{aligned}$$

These alternative characterizations of the supremum and infimum really provide a standard way to determining the supremum and infimum of subsets of the real line.

2.4 Maxima and minima

Definition 2.4.1 – maximum and minimum Let $A \subseteq \mathbb{R}$ be a subset of the real numbers. We say that $y \in A$ is the *maximum* of A , and write $y = \max A$, if

$$\begin{aligned} &\text{for all } a \in A, \\ &a \leq y. \end{aligned}$$

We say that $x \in A$ is the *minimum* of A , and write $x = \min A$, if

$$\begin{aligned} &\text{for all } a \in A, \\ &a \geq x. \end{aligned}$$

Remark 2.4.2. Even if a set $A \subseteq \mathbb{R}$ is non-empty and bounded, it may not have a maximum or minimum. For example, the set $(0, 1)$ has no maximum or minimum.

Proposition 2.4.3 – Let A be a subset of \mathbb{R} . If A has a maximum, then A is non-empty and bounded from above, and $\sup A = \max A$. If A has a minimum, then A is non-empty and bounded from below, and $\inf A = \min A$.

Proposition 2.4.4 – Let A be a subset of \mathbb{R} . Assume that A is non-empty and bounded from above. If $\sup A \in A$ then A has a maximum and $\max A = \sup A$.

Proposition 2.4.5 – Let A be a subset of \mathbb{R} . Assume that A is non-empty and bounded from below. If $\inf A \in A$ then A has a minimum and $\min A = \inf A$.

2.5 The Archimedean property

Proposition 2.5.1 – Archimedean property For every real number $x \in \mathbb{R}$ there exists a natural number $n \in \mathbb{N}$ such that $x < n$.

Given this proposition, we can define the ceiling function.

Definition 2.5.2 – ceiling function The *ceiling function* $\lceil \cdot \rceil : \mathbb{R} \rightarrow \mathbb{Z}$ is defined as follows. For $x \in \mathbb{R}$, $\lceil x \rceil$ denotes the smallest integer $z \in \mathbb{Z}$ such that $x \leq z$.

Proposition 2.5.3 – For every two real numbers $a, b \in \mathbb{R}$ with $a < b$ there exists a $q \in \mathcal{Q}$ with $a < q < b$.

2.6 Computation rules for suprema

In the proposition below, we use the definitions

$$A + B = \{a + b \mid a \in A, b \in B\}$$

and

$$\lambda A = \{\lambda a \mid a \in A\}$$

for subsets $A, B \subseteq \mathbb{R}$ and a scalar $\lambda \in \mathbb{R}$.

Proposition 2.6.1 – Let A, B, C, D be non-empty subsets of \mathbb{R} . Assume that A and B are bounded from above and C and D are bounded from below. Then

1. $\sup(A + B) = \sup A + \sup B$.
2. $\inf(C + D) = \inf C + \inf D$.
3. For all $\lambda \geq 0$, $\sup(\lambda A) = \lambda \sup A$.
4. For all $\lambda \leq 0$, $\sup(\lambda A) = \lambda \inf A$.
5. $\sup(-C) = -\inf C$.
6. $\inf(-C) = -\sup C$.

2.7 Bernoulli's inequality

Proposition 2.7.1 – Bernoulli's inequality Let $x \in \mathbb{R}$ and $n \in \mathbb{N}$. Then

1. If $x \geq -1$, then $(1 + x)^n \geq 1 + nx$.
2. If $x \geq 0$ and $n \geq 2$, then $(1 + x)^n \geq 1 + nx$.

3 Sequences

3.1 Sequence

Definition 3.1.1 – Sequence A sequence is a function for which the domain is \mathbb{N} .

$$a : \mathbb{N} \rightarrow Y$$

Y can be any set.

Example 3.1.2 Here are some functions that are sequences:

1. $a : \mathbb{N} \rightarrow \mathbb{Q}$
2. $b : \mathbb{N} \rightarrow (\mathbb{N} \rightarrow Y)$
3. $c : \mathbb{N} \rightarrow \mathbb{N}$

And some functions that are not sequences:

1. $d : (\mathbb{N} \rightarrow \mathbb{N}) \rightarrow \mathbb{N}$
2. $e : \mathbb{Q} \rightarrow \mathbb{N}$

3.2 Terminology around sequences

3.2.1 Bounded sequences

Definition 3.2.2 – bounded sequence Let (X, dist) be a metric space. We say a sequence $a : \mathbb{N} \rightarrow X$ is bounded if

$$\begin{aligned} &\text{there exists } q \in X, \\ &\text{there exists } M > 0, \\ &\text{for all } n \in \mathbb{N}, \\ &\text{dist}(a_n, q) \leq M. \end{aligned}$$

In a normed linear space, we can use a simpler criterion to check whether a sequence is bounded. That is the content of the following proposition.

Proposition 3.2.3 – Let $(V, \|\cdot\|)$ be a normed vector space. Let $a : \mathbb{N} \rightarrow V$ be a sequence. The sequence a is bounded if and only if

$$\begin{aligned} &\text{there exists } M > 0, \\ &\text{for all } n \in \mathbb{N}, \\ &\|a_n\| \leq M. \end{aligned}$$

3.3 Convergence of sequences

Definition 3.3.1 – Convergence of sequences Let (X, dist) be a metric space. We say that a sequence $a : \mathbb{N} \rightarrow X$ converges to a point $p \in X$ if

$$\begin{aligned} &\text{for all } \varepsilon > 0, \\ &\text{there exists } N \in \mathbb{N}, \\ &\text{for all } n \geq N, \\ &\text{dist}(a_n, p) < \varepsilon. \end{aligned}$$

We sometimes write

$$\lim_{n \rightarrow \infty} a_n = p$$

to express that the sequence (a_n) converges to p .

Definition 3.3.2 – Divergence of sequences Let (X, dist) be a metric space. A sequence $a : \mathbb{N} \rightarrow X$ is called *divergent* if it is not convergent.

3.4 Examples and limits of simple sequences

Proposition 3.4.1 – The constant sequence Let (X, dist) be a metric space. Let $p \in X$ and assume that the sequence (a_n) is given by $a_n = p$ for every $n \in \mathbb{N}$. We also say that (a_n) is a constant sequence. Then $\lim_{n \rightarrow \infty} a_n = p$.

Example 3.4.2 A standard limit Let $a : \mathbb{N} \rightarrow \mathbb{R}$ be a real-valued sequence such that $a_n = 1/n$ for $n \geq 1$. Then $a : \mathbb{N} \rightarrow \mathbb{R}$ converges to 0.

Proof. Let $\varepsilon > 0$. Choose $N = \lceil 1/\varepsilon \rceil + 1$. Take $n \geq N$. Then

$$\text{dist}_{\mathbb{R}}(a_n, 0) = |a_n - 0| = |1/n| = 1/n \leq 1/N < \varepsilon.$$

□

3.5 Uniqueness of limits

Proposition 3.5.1 – Uniqueness of limits Let (X, dist) be a metric space and let $a : \mathbb{N} \rightarrow \mathbb{R}$ be a sequence in X . Assume that $p, q \in X$ and assume that

$$\lim_{n \rightarrow \infty} a_n = p \text{ and } \lim_{n \rightarrow \infty} a_n = q$$

Then $p = q$.

3.6 More properties of convergent sequences

Proposition 3.6.1 – Let (X, dist) be a metric space and suppose that $a : \mathbb{N} \rightarrow X$ is a sequence. Let $p \in X$. Then the sequence $a : \mathbb{N} \rightarrow X$ converges to p if and only if the real-valued sequence

$$n \mapsto \text{dist}(a_n, p)$$

converges to 0 in \mathbb{R} .

Proposition 3.6.2 – Convergent sequences are bounded Let (X, dist) be a metric space. Let $a : \mathbb{N} \rightarrow X$ be a sequence in X converging to $p \in X$. Then the sequence $a : \mathbb{N} \rightarrow X$ is bounded.

Proposition 3.6.3 – Let (X, dist) be a metric space and let $a : \mathbb{N} \rightarrow X$ and $b : \mathbb{N} \rightarrow X$ be two sequences. Let $p \in X$ and suppose that $\lim_{n \rightarrow \infty} a_n = p$. Then $\lim_{n \rightarrow \infty} b_n = p$ if and only if

$$\lim_{n \rightarrow \infty} \text{dist}(a_n, b_n) = 0$$

Corollary 3.6.4 – Eventually equal sequences have the same limit Let (X, dist) be a metric space and

let $a : \mathbb{N} \rightarrow X$ and $b : \mathbb{N} \rightarrow X$ be two sequences such that there exists an $N \in \mathbb{N}$ such that for all $n \geq N$,

$$a_n = b_n$$

Then the sequence $a : \mathbb{N} \rightarrow X$ converges if and only if the sequence $b : \mathbb{N} \rightarrow X$ converges. If the sequences converge, they have the same limit.

3.7 Limit theorems for sequences taking values in a normed vector space

Theorem 3.7.1 – Let $(V, \|\cdot\|)$ be a normed vector space and let $a : \mathbb{N} \rightarrow V$ and $b : \mathbb{N} \rightarrow V$ be two sequences. Assume that the $\lim_{n \rightarrow \infty} a_n$ exists and is equal to $p \in V$ and that the $\lim_{n \rightarrow \infty} b_n$ exists and is equal to $q \in V$. Let $\lambda : \mathbb{N} \rightarrow \mathbb{R}$ be a real-valued sequence. Let $\mu \in \mathbb{R}$. Assume that $\lim_{n \rightarrow \infty} \lambda_n = \mu$. Then

1. The $\lim_{n \rightarrow \infty} (a_n + b_n)$ exists and is equal to $p + q$.
2. The $\lim_{n \rightarrow \infty} (\lambda_n a_n)$ exists and is equal to μp .

3.8 Index shift

Proposition 3.8.1 – Index shift Let (X, dist) be a metric space and let $a : \mathbb{N} \rightarrow X$ be a sequence. Let $k \in \mathbb{N}$ and $p \in X$. Then the sequence $a : \mathbb{N} \rightarrow X$ converges to p if and only if the sequence $(a_{n+k})_n$ (i.e. the sequence $n \mapsto a_{n+k}$) converges to p .

4 Real-valued sequences

4.1 Terminology

Definition 4.1.1 – increasing, decreasing and monotone sequences We say a sequence (a_n) is

1. *increasing* if for every $n \in \mathbb{N}$, $a_{n+1} \geq a_n$
2. *strictly increasing* if for every $n \in \mathbb{N}$, $a_{n+1} > a_n$
3. *decreasing* if for every $n \in \mathbb{N}$, $a_{n+1} \leq a_n$
4. *strictly decreasing* if for every $n \in \mathbb{N}$, $a_{n+1} < a_n$
5. *monotone* if it is either increasing or decreasing
6. *strictly monotone* if it is either strictly increasing or strictly decreasing

Definition 4.1.2 – upper bound and lower bound for a sequence We say that a number $M \in \mathbb{R}$ is an *upper bound* for a sequence $a : \mathbb{N} \rightarrow \mathbb{R}$ if

for all $n \in \mathbb{N}$

$$a_n \leq M$$

We say that a number $m \in \mathbb{R}$ is a *lower bound* for a sequence $a : \mathbb{N} \rightarrow \mathbb{R}$ if

for all $n \in \mathbb{N}$

$$a_n \geq m$$

Definition 4.1.3 – bounded sequence We say that a sequence $a : \mathbb{N} \rightarrow \mathbb{R}$ is *bounded above* if there exists an $M \in \mathbb{R}$ such that M is an upper bound for a .

We say that a sequence $a : \mathbb{N} \rightarrow \mathbb{R}$ is *bounded below* if there exists an $m \in \mathbb{R}$ such that m is a lower bound for a .

Proposition 4.1.4 – Let $a : \mathbb{N} \rightarrow \mathbb{R}$ be a sequence. Then $a : \mathbb{N} \rightarrow \mathbb{R}$ is bounded if and only if it is both bounded above and bounded below.

4.2 Monotone, bounded sequences and convergent

Theorem 4.2.1 – Let (a_n) be an increasing sequence that is bounded from above. Then (a_n) convergent and

$$\lim_{n \rightarrow \infty} a_n = \sup_{n \in \mathbb{N}} a_n \quad (= \sup\{a_n \mid n \in \mathbb{N}\})$$

Theorem 4.2.2 – Let (a_n) be a decreasing sequence that is bounded from below. Then (a_n) is convergent and

$$\lim_{n \rightarrow \infty} a_n = \inf_{n \in \mathbb{N}} a_n \quad (= \inf\{a_n \mid n \in \mathbb{N}\})$$

4.3 Limit theorems

Theorem 4.3.1 – Limit theorems for real-valued sequences Let $a : \mathbb{N} \rightarrow \mathbb{R}$ and $b : \mathbb{N} \rightarrow \mathbb{R}$ be two converging sequences, and let $c, d \in \mathbb{R}$ be real numbers such that

$$\lim_{n \rightarrow \infty} a_n = c \text{ and } \lim_{n \rightarrow \infty} b_n = d.$$

Then

1. The $\lim_{n \rightarrow \infty} (a_n + b_n)$ exists and is equal to $c + d$.
2. The $\lim_{n \rightarrow \infty} (a_n b_n)$ exists and is equal to $c \cdot d$.
3. If $d \neq 0$, then $\lim_{n \rightarrow \infty} (\frac{a_n}{b_n})$ exists and is equal to $\frac{c}{d}$.
4. For every non-negative integer $m \in \mathbb{N}$, the limit $\lim_{n \rightarrow \infty} (a_n)^m$ exists and is equal to c^m .
5. If for every $n \in \mathbb{N}$, the number a_n is non-negative, then for every positive integer $k \in \mathbb{N} \setminus \{0\}$, the limit $\lim_{n \rightarrow \infty} (a_n)^{\frac{1}{k}}$ exists and is equal to $c^{\frac{1}{k}}$.

4.4 The squeeze theorem

Theorem 4.4.1 – The squeeze theorem Let $a, b, c : \mathbb{N} \rightarrow \mathbb{R}$ be three sequences. Suppose that there exists an $N \in \mathbb{N}$ such that for every $n \geq N$, we have

$$a_n \leq b_n \leq c_n$$

and assume $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} c_n = L$ for some $L \in \mathbb{R}$. Then $\lim_{n \rightarrow \infty} b_n$ exists and is equal to L .

4.5 Divergence to ∞ and $-\infty$

Definition 4.5.1 – We say a sequence $a : \mathbb{N} \rightarrow \mathbb{R}$ *diverges to ∞* and write

$$\lim_{n \rightarrow \infty} a_n = \infty$$

if

for all $M \in \mathbb{R}$,
there exists $N \in \mathbb{N}$,
for all $n \geq N$,
 $a_n > M$.

Similarly, we say a sequence (a_n) *diverges to $-\infty$* and write

$$\lim_{n \rightarrow \infty} a_n = -\infty$$

if

for all $M \in \mathbb{R}$,
there exists $N \in \mathbb{N}$,
for all $n \geq N$,
 $a_n < M$.

Proposition 4.5.2 – Let $a : \mathbb{N} \rightarrow \mathbb{R}$ be a sequence such that

$$\lim_{n \rightarrow \infty} a_n = \infty.$$

Then the sequence (a_n) is bounded from below.

Similarly, let $b : \mathbb{N} \rightarrow \mathbb{R}$ be a sequence such that

$$\lim_{n \rightarrow \infty} b_n = -\infty.$$

Then the sequence (b_n) is bounded from above.

4.6 Limit theorems for improper limits

Theorem 4.6.1 – Limit theorems for improper limits Let $a, b, c, d : \mathbb{N} \rightarrow \mathbb{R}$ be four sequences such that

$$\lim_{n \rightarrow \infty} a_n = \infty \text{ and } \lim_{n \rightarrow \infty} c_n = -\infty$$

the sequence (b_n) is bounded from below and the sequence (d_n) is bounded from above. Let $\lambda : \mathbb{N} \rightarrow \mathbb{R}$ be a sequence bounded below by some $\mu > 0$. Then

- i. $\lim_{n \rightarrow \infty} (a_n + b_n) = \infty$
- ii. $\lim_{n \rightarrow \infty} (c_n + d_n) = -\infty$
- iii. $\lim_{n \rightarrow \infty} (\lambda_n a_n) = \infty$
- iv. $\lim_{n \rightarrow \infty} (\lambda_n c_n) = -\infty$

Proposition 4.6.2 – Let $a : \mathbb{N} \rightarrow \mathbb{R}$ and $b : \mathbb{N} \rightarrow (0, \infty)$ be two sequences. Then

- 1. $\lim_{n \rightarrow \infty} a_n = \infty$ if and only if $\lim_{n \rightarrow \infty} (-a_n) = -\infty$.
- 2. $\lim_{n \rightarrow \infty} b_n = \infty$ if and only if $\lim_{n \rightarrow \infty} \frac{1}{b_n} = 0$.

4.7 Standard sequences

4.7.1 Geometric sequence

Proposition 4.7.2 – Standard limit of of geometric sequence Let $q \in \mathbb{R}$. The sequence (a_n) defined by $a_n := q^n$ for $n \in \mathbb{N}$

- converges to 0 if $q \in (-1, 1)$
- converges to 1 if $q = 1$
- diverges to ∞ if $q > 1$
- diverges, but not to ∞ or $-\infty$ if $q \leq -1$

4.7.3 The n^{th} root of n

Proposition 4.7.4 – Standard limit of the n^{th} root of n The sequence (a_n) defined by $a_n := \sqrt[n]{n}$ for $n \in \mathbb{N}$ converges to 1.

Corollary 4.7.5 – Let $a > 0$. Then the sequence (b_n) defined by $b_n := \sqrt[n]{a}$ converges to 1.

4.7.6 The number e

First let's define the sequence (a_n) by

$$a_n := \left(1 + \frac{1}{n}\right)^n.$$

We show that (a_n) is increasing and bounded from above by 3. Hence (a_n) converges to some $e \in \mathbb{R}$ by the monotone convergence theorem.

Lemma 4.7.7 – The sequence (a_n) defined by $a_n := \left(1 + \frac{1}{n}\right)^n$ for $n \in \mathbb{N} \setminus \{0\}$ and $a_0 = 1$ is increasing.

Lemma 4.7.8 – The sequence (a_n) defined by $a_n := \left(1 + \frac{1}{n}\right)^n$ for $n \in \mathbb{N} \setminus \{0\}$ and $a_0 = 1$ is bounded from above by 3.

By these two lemmas, the sequence

$$n \mapsto \left(1 + \frac{1}{n}\right)^n$$

converges.

Definition 4.7.9 – (Standard limit of e) We define the number e by

$$e := \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n.$$

4.7.10 Exponentials beat powers

5 Series

5.1 Geometric series

5.2 The harmonic series

5.3 The hyperharmonic series

5.4 Only the tail matters for convergence

5.5 Divergence test

5.6 Limit laws for series

6 Series with positive terms

6.1 Comparison test

6.2 Limit comparison test

6.3 Ratio test

6.4 Root test

7 Series with general terms

7.1 Series with real terms: the Leibniz test

7.2 Series characterization of completeness in normed vector space

7.3 The Cauchy product

8 Subsequences, \limsup and \liminf

8.1 Index sequences and subsequences

8.2 (Sequential) accumulation points

8.3 Subsequences of a converging sequence

8.4 \limsup

8.5 \liminf

8.6 Relations between \lim , \limsup and \liminf

9 Point-set topology of metric spaces

9.1 Open sets

9.2 Closed sets

9.3 Cauchy sequences

9.4 Completeness

9.5 Series characterization of completeness in normed vector spaces

10 Compactness

10.1 Boundedness and total boundedness

10.2 Alternative characterization of compactness

11 Limits and continuity

11.1 Accumulation points

11.2 Limit in an accumulation point

11.3 Uniqueness of limits

11.4 Sequential characterization of limits

11.5 Limit laws

11.6 Continuity

11.7 Sequential characterization of continuity

11.8 Rules for continuous functions

11.9 Images of compact sets under continuous functions are compact

11.10 Uniform continuity

12 Real-valued functions

12.1 More limit laws

12.2 Building of standard functions

12.3 Continuity of standard functions

12.4 Limits from the left and from the right

12.5 The extended real line

12.6 Limits to ∞ or $-\infty$

12.7 Limits at ∞ and $-\infty$

12.8 The Intermediate Value Theorem

12.9 The Extreme Value Theorem

12.10 Equivalence of norms

12.11 Bounded linear maps and operator norms

13 Differentiability

13.1 The derivative as a function

13.2 Constant and linear maps are differentiable

13.3 Bases and coordinates

13.4 The matrix representation

13.5 The chain rule

13.6 Sum, product and quotient rules

13.7 Differentiability of components

13.8 Differentiability implies continuity

13.9 Derivative vanishes in local maxima and minima

13.10 The Mean Value Theorem

14 Differentiability of standard functions

14.1 Global context

14.2 Polynomials and rational functions are differentiable

14.3 Differentiability of the standard functions

15 Directional and partial derivatives

15.1 A recurring and very important construction

15.2 Directional derivatives

15.3 Partial derivatives

15.4 The Jacobian of a map

15.5 Linearization and tangent planes

15.6 The gradient of a function

16 The Mean-Value Inequality

16.1 The mean-value inequality for functions defined on an interval

16.2 The mean-value inequality for functions on general domains

16.3 Continuous partial derivatives imply differentiability

17 Higher order derivatives

17.1 Multilinear maps

17.2 Relation to n -fold directional derivatives

17.3 A criterion for higher differentiability

17.4 Symmetry of second order derivatives

17.5 Symmetry of higher-order derivatives

18 Polynomials and approximation by polynomials

18.1 Homogeneous polynomials

18.2 Taylor's theorem

18.3 Taylor approximations of standard functions

19 Banach fixed point theorem

20 Implicit function theorem

20.1 The objective

20.2 Notation

20.3 The implicit function theorem

20.4 The inverse function theorem

21 Function sequences

21.1 Point-wise convergence

21.2 Uniform convergence

21.3 Preservation of continuity under uniform convergence

21.4 Differentiability theorem

21.5 The normed vector space of bounded functions

22 Function series

22.1 The Weierstrass M-test

22.2 Conditions for differentiation of function series

23 Power series

23.1 Convergence of power series

23.2 Standard functions defined as power series

23.3 Operations with power series

23.4 Differentiation of power series

23.5 Taylor series

24 Riemann integration in one dimension

24.1 Riemann integrable functions and the Riemann integral

24.2 Sums, products of Riemann integrable functions

24.3 Continuous functions are Riemann integrable

24.4 The fundamental theorem of calculus

25 Riemann integration in multiple dimensions

25.1 Partitions in multiple dimensions

25.2 Riemann integral on rectangles in \mathbb{R}^n

25.3 Properties of the multidimensional Riemann integral

25.4 Continuous functions are Riemann integrable

25.5 Fubini's theorem

25.6 The (topological) boundary of a set

25.7 Jordan content

25.8 Integration over general domains

25.9 The volume of bounded sets

26 Change-of-variables Theorem

26.1 Polar coordinates

26.2 Cylindrical coordinates

26.3 Spherical coordinates