

# Homework: Week 2

Jiaqi Wang

November 29, 2023

## 1 Exercise 6.8.1

**Problem 1.1** Let  $a : \mathbb{N} \rightarrow \mathbb{R}$  be a sequence. Then  $a : \mathbb{N} \rightarrow \mathbb{R}$  is bounded if and only if it is both bounded above and bounded below. Prove it.

*Proof.* We prove both directions of the if and only if statement.  $a : \mathbb{N} \rightarrow \mathbb{R}$  is bounded if and only if it is both bounded above and bounded below.

1. We prove the forward direction.

Suppose  $a : \mathbb{N} \rightarrow \mathbb{R}$  is bounded.

Then

there exists  $M_0 > 0$ ,  
for all  $n \in \mathbb{N}$ ,  
 $|a_n| \leq M_0$ .

Obtain such  $M_0$ .

It holds that

for all  $n \in \mathbb{N}$ ,  
 $-M_0 \leq a_n \leq M_0$ .

Choose  $m = -M_0$ , then  $m \in \mathbb{R}$ .

It holds that

for all  $n \in \mathbb{N}$ ,  
 $m \leq a_n$ .

We conclude that  $a : \mathbb{N} \rightarrow \mathbb{R}$  is bounded from below.

Choose  $M = M_0$ , then  $M \in \mathbb{R}$ .

It holds that

for all  $n \in \mathbb{N}$ ,  
 $a_n \leq M$ .

We conclude that  $a : \mathbb{N} \rightarrow \mathbb{R}$  is bounded from above.

2. Now we prove the reverse direction.

Assume  $a : \mathbb{N} \rightarrow \mathbb{R}$  is bounded from above and bounded from below.

By definition of bounded from above, we have

there exists  $M_1 \in \mathbb{R}$ ,  
for all  $n \in \mathbb{N}$ ,  
 $a_n \leq M_1$ .

By definition of bounded from below, we have

there exists  $M_2 \in \mathbb{R}$ ,  
for all  $n \in \mathbb{N}$ ,  
 $M_2 \leq a_n$ .

We need to show that

there exists  $M_0 > 0$ ,  
for all  $n \in \mathbb{N}$ ,  
 $|a_n| \leq M_0$ .

Choose  $M_0 = \max\{|M_1|, |M_2|\}$ , then  $M_0 \in \mathbb{R}$ .  
It holds that

for all  $n \in \mathbb{N}$ ,  
 $-M_0 \leq a_n$  and  $a_n \leq M_0$ .

Then it holds that  $|a_n| \leq M_0$ .

We conclude that  $a : \mathbb{N} \rightarrow \mathbb{R}$  is bounded.

Since both directions hold, we conclude that  $a : \mathbb{N} \rightarrow \mathbb{R}$  is bounded if and only if it is both bounded above and bounded below.  $\square$

## 2 Exercise 6.8.2

**Problem 2.1** Let  $a : \mathbb{N} \rightarrow \mathbb{R}$  and  $b : \mathbb{N} \rightarrow (0, \infty)$  be real-valued sequences. Prove that

$$\lim_{n \rightarrow \infty} a_n = \infty \iff \lim_{n \rightarrow \infty} (-a_n) = -\infty$$

*Proof.* We show both directions of the if and only if statement.

$$\lim_{n \rightarrow \infty} a_n = \infty \iff \lim_{n \rightarrow \infty} (-a_n) = -\infty$$

1. First we prove the forward direction.

Suppose  $\lim_{n \rightarrow \infty} a_n = \infty$ .

Then

for all  $M \in \mathbb{R}$ ,  
there exists  $N_0 \in \mathbb{N}$ ,  
for all  $n \geq N_0$ ,  
 $a_n > M$ .

We need to show that

for all  $M \in \mathbb{R}$ ,  
there exists  $N \in \mathbb{N}$ ,  
for all  $n \geq N$ ,  
 $-a_n \leq M$ .

Take  $M \in \mathbb{R}$  Choose  $N = N_0$ , then  $N \in \mathbb{N}$ .

It holds that

for all  $n \geq N$ ,  
 $a_n > M$ .

$\square$

### 3 Exercise 6.8.3

**Problem 3.1** Let  $a : \mathbb{N} \rightarrow \mathbb{R}$  and  $b : \mathbb{N} \rightarrow (0, \infty)$  be real-valued sequences. Prove that

$$\lim_{n \rightarrow \infty} b_n = \infty \iff \lim_{n \rightarrow \infty} \frac{1}{b_n} = 0$$

*Proof.*

□

### 4 Exercise 6.8.5

**Problem 4.1** Define the sequence  $x : \mathbb{N} \rightarrow \mathbb{R}$  recursively by

$$x_{n+1} := \frac{2 + x_n^2}{2x_n}$$

for  $n \in \mathbb{N}$  while  $x_0 = 2$ . Prove that the sequence  $x : \mathbb{N} \rightarrow \mathbb{R}$  converges and determine its limit.

*Proof.*

□

### 5 Exercise 6.8.6

**Problem 5.1** Determine whether the following sequences converge, diverge to  $+\infty$ , diverge to  $-\infty$ , or diverge in a different way. In case the sequence converges, determine the limit.

- $a_n := \frac{1}{n^3} - 3$
- $b_n := \frac{5n^5 + 2n^2}{3n^5 + 7n^3 + 4}$
- $c_n := n - \sqrt{n}$
- $d_n := \frac{2^n}{n^{100}}$
- $e_n := \sqrt{n^2 + n} - n$
- $f_n := \sqrt[n]{3n^2}$
- $g_n := \frac{2^n + 5n^{200}}{3^n + n^{10}}$
- $h_n := (-1)^n 3^n$
- $i_n := \sqrt[n]{5^n + n^2}$

*Proof.* By limit laws and standard limits, we have

$$\begin{aligned}\lim_{n \rightarrow \infty} a_n &= \lim_{n \rightarrow \infty} \left( \frac{1}{n^3} - 3 \right) \\ &= \lim_{n \rightarrow \infty} \frac{1}{n^3} - \lim_{n \rightarrow \infty} 3 \\ &= \left( \lim_{n \rightarrow \infty} \frac{1}{n} \right)^3 - 3 \\ &= 0^3 - 3 \\ &= -3\end{aligned}$$

$$\begin{aligned}\lim_{n \rightarrow \infty} b_n &= \lim_{n \rightarrow \infty} \left( \frac{5n^5 + 2n^2}{3n^5 + 7n^3 + 4} \right) \\ &= \lim_{n \rightarrow \infty} \frac{5 + \frac{2}{n^3}}{3 + \frac{7}{n^2} + \frac{4}{n^5}} \\ &= \frac{\lim_{n \rightarrow \infty} (5 + \frac{2}{n^3})}{\lim_{n \rightarrow \infty} (3 + \frac{7}{n^2} + \frac{4}{n^5})} \\ &= \frac{\lim_{n \rightarrow \infty} 5 + \lim_{n \rightarrow \infty} \frac{2}{n^3}}{\lim_{n \rightarrow \infty} 3 + \lim_{n \rightarrow \infty} \frac{7}{n^2} + \lim_{n \rightarrow \infty} \frac{4}{n^5}} \\ &= \frac{5}{3}\end{aligned}$$

$$\begin{aligned}
\lim_{n \rightarrow \infty} c_n &= \lim_{n \rightarrow \infty} (n - \sqrt{n}) \\
&= \lim_{n \rightarrow \infty} \left( \frac{(n - \sqrt{n})(n + \sqrt{n})}{n + \sqrt{n}} \right) \\
&= \lim_{n \rightarrow \infty} \left( \frac{n^2 - n}{n + \sqrt{n}} \right) \\
&= \lim_{n \rightarrow \infty} \left( \frac{n(n - 1)}{n(1 + \frac{1}{\sqrt{n}})} \right) \\
&= \lim_{n \rightarrow \infty} \left( \frac{n - 1}{1 + \frac{1}{\sqrt{n}}} \right) \\
&= \frac{\lim_{n \rightarrow \infty} (n - 1)}{\lim_{n \rightarrow \infty} (1 + \frac{1}{\sqrt{n}})} \\
&= \frac{\lim_{n \rightarrow \infty} n - \lim_{n \rightarrow \infty} 1}{\lim_{n \rightarrow \infty} 1 + \lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}}} \\
&= \frac{\lim_{n \rightarrow \infty} n - 1}{1 + \lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}}} \\
&= \frac{\lim_{n \rightarrow \infty} n - 1}{1 + 0} \\
&= \lim_{n \rightarrow \infty} n - 1 \\
&= \infty
\end{aligned}$$

$$\begin{aligned}
\lim_{n \rightarrow \infty} d_n &= \lim_{n \rightarrow \infty} \left( \frac{2^n}{n^{100}} \right) \\
&= \lim_{n \rightarrow \infty} \left( \frac{2^n}{n^{100}} \right)
\end{aligned}$$

□