


Convergence of Gradient Descent: Smooth and Convex Settings

Avishek Ghosh

Assistant Professor, Dept. of CSE., IIT Bombay

IIT Bombay

August 6, 2025





1. Gradient Descent and Decent lemma for Smooth functions
 - GD for Non-Convex Smooth Functions
 - GD for Convex and Smooth function
 - GD for Smooth and Strongly Convex functions

2. Projection



Gradient Descent(GD) Formulation

Problem:

$$\min_{x \in \mathbb{R}^d} f(x)$$

.



Gradient Descent(GD) Formulation

Problem:

$$\min_{x \in \mathbb{R}^d} f(x)$$

Gradient Descent Update:



Gradient Descent(GD) Formulation

Problem:

$$\min_{x \in \mathbb{R}^d} f(x)$$

Gradient Descent Update:

$$x_{k+1} = x_k - \alpha_k \nabla f(x_k), \quad \alpha_k > 0 \text{ (step size).}$$



Gradient Descent(GD) Formulation

Problem:

$$\min_{x \in \mathbb{R}^d} f(x)$$

Gradient Descent Update:

$$x_{k+1} = x_k - \alpha_k \nabla f(x_k), \quad \alpha_k > 0 \text{ (step size).}$$

- Choose α_k appropriately (for example, through line search methods)



Gradient Descent(GD) Formulation

Problem:

$$\min_{x \in \mathbb{R}^d} f(x)$$

Gradient Descent Update:

$$x_{k+1} = x_k - \alpha_k \nabla f(x_k), \quad \alpha_k > 0 \text{ (step size).}$$

- Choose α_k appropriately (for example, through line search methods)

Now from next slide, we'll see about GD convergence results in the following cases:

- Smooth functions



Gradient Descent(GD) Formulation

Problem:

$$\min_{x \in \mathbb{R}^d} f(x)$$

Gradient Descent Update:

$$x_{k+1} = x_k - \alpha_k \nabla f(x_k), \quad \alpha_k > 0 \text{ (step size).}$$

- Choose α_k appropriately (for example, through line search methods)

Now from next slide ,we'll see about GD convergence results in the following cases:

- Smooth functions
- Smooth + Convex functions



Gradient Descent(GD) Formulation

Problem:

$$\min_{x \in \mathbb{R}^d} f(x)$$

Gradient Descent Update:

$$x_{k+1} = x_k - \alpha_k \nabla f(x_k), \quad \alpha_k > 0 \text{ (step size).}$$

- Choose α_k appropriately (for example, through line search methods)

Now from next slide ,we'll see about GD convergence results in the following cases:

- Smooth functions
- Smooth + Convex functions
- Smooth + Strongly Convex functions

Descent lemma for smooth function



We consider the unconstrained optimization problem

$$\min_{x \in \mathbb{R}^d} f(x) \quad (1)$$

where:



Descent lemma for smooth function

We consider the unconstrained optimization problem

$$\min_{x \in \mathbb{R}^d} f(x) \quad (1)$$

where:

- $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is M -smooth, i.e. $\|\nabla f(x) - \nabla f(y)\| \leq M\|x - y\|$.



Descent lemma for smooth function

We consider the unconstrained optimization problem

$$\min_{x \in \mathbb{R}^d} f(x) \quad (1)$$

where:

- $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is M -smooth, i.e. $\|\nabla f(x) - \nabla f(y)\| \leq M\|x - y\|$.
- $f^* = \inf_x f(x) > -\infty$ is a global lower bound.



Descent lemma for smooth function

We consider the unconstrained optimization problem

$$\min_{x \in \mathbb{R}^d} f(x) \quad (1)$$

where:

- $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is M -smooth, i.e. $\|\nabla f(x) - \nabla f(y)\| \leq M\|x - y\|$.
- $f^* = \inf_x f(x) > -\infty$ is a global lower bound.

Lemma (Descent Lemma)

Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be M -smooth. Using gradient descent

$$x^{k+1} = x^k - \frac{1}{M} \nabla f(x^k),$$

we have



Descent lemma for smooth function

We consider the unconstrained optimization problem

$$\min_{x \in \mathbb{R}^d} f(x) \quad (1)$$

where:

- $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is M -smooth, i.e. $\|\nabla f(x) - \nabla f(y)\| \leq M\|x - y\|$.
- $f^* = \inf_x f(x) > -\infty$ is a global lower bound.

Lemma (Descent Lemma)

Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be M -smooth. Using gradient descent

$$x^{k+1} = x^k - \frac{1}{M} \nabla f(x^k),$$

we have

$$f(x^{k+1}) \leq f(x^k) - \frac{1}{2M} \|\nabla f(x^k)\|^2 \leq f(x^k).$$

Proof of Descent lemma





Since f is M -smooth, for all $x, y \in \mathbb{R}^d$,



Since f is M -smooth, for all $x, y \in \mathbb{R}^d$,

$$f(y) \leq f(x) + \langle \nabla f(x), y - x \rangle + \frac{M}{2} \|y - x\|^2.$$



Proof of Descent lemma

Since f is M -smooth, for all $x, y \in \mathbb{R}^d$,

$$f(y) \leq f(x) + \langle \nabla f(x), y - x \rangle + \frac{M}{2} \|y - x\|^2.$$

Setting $x = x^k$, $y = x^{k+1}$ and $\alpha^k = 1/M$ gives



Proof of Descent lemma

Since f is M -smooth, for all $x, y \in \mathbb{R}^d$,

$$f(y) \leq f(x) + \langle \nabla f(x), y - x \rangle + \frac{M}{2} \|y - x\|^2.$$

Setting $x = x^k$, $y = x^{k+1}$ and $\alpha^k = 1/M$ gives

$$f(x^{k+1}) \leq f(x^k) - \langle \nabla f(x^k), \alpha^k \nabla f(x^k) \rangle + \frac{M}{2} \|\alpha^k \nabla f(x^k)\|^2$$



Proof of Descent lemma

Since f is M -smooth, for all $x, y \in \mathbb{R}^d$,

$$f(y) \leq f(x) + \langle \nabla f(x), y - x \rangle + \frac{M}{2} \|y - x\|^2.$$

Setting $x = x^k$, $y = x^{k+1}$ and $\alpha^k = 1/M$ gives

$$\begin{aligned} f(x^{k+1}) &\leq f(x^k) - \langle \nabla f(x^k), \alpha^k \nabla f(x^k) \rangle + \frac{M}{2} \|\alpha^k \nabla f(x^k)\|^2 \\ &= f(x^k) - \frac{1}{M} \|\nabla f(x^k)\|^2 + \frac{M}{2} \frac{1}{M^2} \|\nabla f(x^k)\|^2 \end{aligned}$$



Proof of Descent lemma

Since f is M -smooth, for all $x, y \in \mathbb{R}^d$,

$$f(y) \leq f(x) + \langle \nabla f(x), y - x \rangle + \frac{M}{2} \|y - x\|^2.$$

Setting $x = x^k$, $y = x^{k+1}$ and $\alpha^k = 1/M$ gives

$$\begin{aligned} f(x^{k+1}) &\leq f(x^k) - \langle \nabla f(x^k), \alpha^k \nabla f(x^k) \rangle + \frac{M}{2} \|\alpha^k \nabla f(x^k)\|^2 \\ &= f(x^k) - \frac{1}{M} \|\nabla f(x^k)\|^2 + \frac{M}{2} \frac{1}{M^2} \|\nabla f(x^k)\|^2 \\ &= f(x^k) - \frac{1}{2M} \|\nabla f(x^k)\|^2 \end{aligned}$$



Proof of Descent lemma

Since f is M -smooth, for all $x, y \in \mathbb{R}^d$,

$$f(y) \leq f(x) + \langle \nabla f(x), y - x \rangle + \frac{M}{2} \|y - x\|^2.$$

Setting $x = x^k$, $y = x^{k+1}$ and $\alpha^k = 1/M$ gives

$$\begin{aligned} f(x^{k+1}) &\leq f(x^k) - \langle \nabla f(x^k), \alpha^k \nabla f(x^k) \rangle + \frac{M}{2} \|\alpha^k \nabla f(x^k)\|^2 \\ &= f(x^k) - \frac{1}{M} \|\nabla f(x^k)\|^2 + \frac{M}{2} \frac{1}{M^2} \|\nabla f(x^k)\|^2 \\ &= f(x^k) - \frac{1}{2M} \|\nabla f(x^k)\|^2 \\ &\leq f(x^k). \end{aligned} \tag{2}$$

Proof of Descent lemma

Since f is M -smooth, for all $x, y \in \mathbb{R}^d$,

$$f(y) \leq f(x) + \langle \nabla f(x), y - x \rangle + \frac{M}{2} \|y - x\|^2.$$

Setting $x = x^k$, $y = x^{k+1}$ and $\alpha^k = 1/M$ gives

$$\begin{aligned} f(x^{k+1}) &\leq f(x^k) - \langle \nabla f(x^k), \alpha^k \nabla f(x^k) \rangle + \frac{M}{2} \|\alpha^k \nabla f(x^k)\|^2 \\ &= f(x^k) - \frac{1}{M} \|\nabla f(x^k)\|^2 + \frac{M}{2} \frac{1}{M^2} \|\nabla f(x^k)\|^2 \\ &= f(x^k) - \frac{1}{2M} \|\nabla f(x^k)\|^2 \\ &\leq f(x^k). \end{aligned} \tag{2}$$

Which Complete the proof of Descent Lemma.



Theorem

Let $f: \mathbb{R}^n \rightarrow \mathbb{R}$ be M -smooth (but not necessarily convex).



Theorem

Let $f: \mathbb{R}^n \rightarrow \mathbb{R}$ be M -smooth (but not necessarily convex). Running gradient descent with step size $\alpha_k = \frac{1}{M}$

Case1: Non-Convex Smooth Functions

Theorem

Let $f: \mathbb{R}^n \rightarrow \mathbb{R}$ be M -smooth (but not necessarily convex). Running gradient descent with step size $\alpha_k = \frac{1}{M}$

$$x^{k+1} = x^k - \frac{1}{M} \nabla f(x^k) \quad (k = 0, 1, \dots, T-1)$$

Case1: Non-Convex Smooth Functions

Theorem

Let $f: \mathbb{R}^n \rightarrow \mathbb{R}$ be M -smooth (but not necessarily convex). Running gradient descent with step size $\alpha_k = \frac{1}{M}$

$$x^{k+1} = x^k - \frac{1}{M} \nabla f(x^k) \quad (k = 0, 1, \dots, T-1)$$

yields

$$\min_{0 \leq k \leq T-1} \|\nabla f(x^k)\| \leq \sqrt{\frac{2M(f(x^0) - \bar{f})}{T}},$$

where $\bar{f} = \inf_x f(x)$.



Case1: Non-Convex Smooth Functions

Theorem

Let $f: \mathbb{R}^n \rightarrow \mathbb{R}$ be M -smooth (but not necessarily convex). Running gradient descent with step size $\alpha_k = \frac{1}{M}$

$$x^{k+1} = x^k - \frac{1}{M} \nabla f(x^k) \quad (k = 0, 1, \dots, T-1)$$

yields

$$\min_{0 \leq k \leq T-1} \|\nabla f(x^k)\| \leq \sqrt{\frac{2M(f(x^0) - \bar{f})}{T}},$$

where $\bar{f} = \inf_x f(x)$. Moreover,

$$\lim_{T \rightarrow \infty} \|\nabla f(x^T)\| = 0.$$



- 1 The above result is a first-order necessary condition (FONC) guarantee.



- 1 The above result is a first-order necessary condition (FONC) guarantee.
- 2 In general for non-convex functions, this is the best one can establish for GD.



- 1 The above result is a first-order necessary condition (FONC) guarantee.
- 2 In general for non-convex functions, this is the best one can establish for GD.
- 3 In the above theorem, we don't have any condition on the final iterate but

$$\lim_{T \rightarrow \infty} \|\nabla f(x^T)\| = 0$$



Use Decent lemma with $k = T - 1$, and so on..



Use Decent lemma with $k = T - 1$, and so on..

$$f(x^T) \leq f(x^{T-1}) - \frac{1}{2M} \|\nabla f(x^{T-1})\|^2$$



Use Decent lemma with $k = T - 1$, and so on..

$$\begin{aligned} f(x^T) &\leq f(x^{T-1}) - \frac{1}{2M} \|\nabla f(x^{T-1})\|^2 \\ &\leq f(x^{T-2}) - \frac{1}{2M} \|\nabla f(x^{T-2})\|^2 - \frac{1}{2M} \|\nabla f(x^{T-1})\|^2 \end{aligned}$$

Use Decent lemma with $k = T - 1$, and so on..

$$\begin{aligned}
 f(x^T) &\leq f(x^{T-1}) - \frac{1}{2M} \|\nabla f(x^{T-1})\|^2 \\
 &\leq f(x^{T-2}) - \frac{1}{2M} \|\nabla f(x^{T-2})\|^2 - \frac{1}{2M} \|\nabla f(x^{T-1})\|^2 \\
 &\leq \dots \\
 &\leq f(x^0) - \frac{1}{2M} \sum_{k=0}^{T-1} \|\nabla f(x^k)\|^2.
 \end{aligned}$$

Use Decent lemma with $k = T - 1$, and so on..

$$\begin{aligned} f(x^T) &\leq f(x^{T-1}) - \frac{1}{2M} \|\nabla f(x^{T-1})\|^2 \\ &\leq f(x^{T-2}) - \frac{1}{2M} \|\nabla f(x^{T-2})\|^2 - \frac{1}{2M} \|\nabla f(x^{T-1})\|^2 \\ &\leq \dots \\ &\leq f(x^0) - \frac{1}{2M} \sum_{k=0}^{T-1} \|\nabla f(x^k)\|^2. \end{aligned}$$

Since $\bar{f} \leq f(x^T)$, it follows

$$\sum_{k=0}^{T-1} \|\nabla f(x^k)\|^2 \leq 2M(f(x^0) - \bar{f}).$$



(a) Asymptotic: Taking $T \rightarrow \infty$ gives



(a) Asymptotic: Taking $T \rightarrow \infty$ gives $\lim_{T \rightarrow \infty} \|\nabla f(x^T)\|^2 = 0$.



(a) Asymptotic: Taking $T \rightarrow \infty$ gives $\lim_{T \rightarrow \infty} \|\nabla f(x^T)\|^2 = 0$.

(b) Non-asymptotic: For finite T ,



(a) Asymptotic: Taking $T \rightarrow \infty$ gives $\lim_{T \rightarrow \infty} \|\nabla f(x^T)\|^2 = 0$.

(b) Non-asymptotic: For finite T ,

$$\frac{1}{T} \sum_{k=0}^{T-1} \|\nabla f(x^k)\|^2 \leq \frac{2M(f(x^0) - \bar{f})}{T},$$



(a) Asymptotic: Taking $T \rightarrow \infty$ gives $\lim_{T \rightarrow \infty} \|\nabla f(x^T)\|^2 = 0$.

(b) Non-asymptotic: For finite T ,

$$\frac{1}{T} \sum_{k=0}^{T-1} \|\nabla f(x^k)\|^2 \leq \frac{2M(f(x^0) - \bar{f})}{T},$$

Since, minimum of non-negative sum term is upper-bounded by average sum, hence



(a) Asymptotic: Taking $T \rightarrow \infty$ gives $\lim_{T \rightarrow \infty} \|\nabla f(x^T)\|^2 = 0$.

(b) Non-asymptotic: For finite T ,

$$\frac{1}{T} \sum_{k=0}^{T-1} \|\nabla f(x^k)\|^2 \leq \frac{2M(f(x^0) - \bar{f})}{T},$$

Since, minimum of non-negative sum term is upper-bounded by average sum, hence

$$\min_{0 \leq k \leq T-1} \|\nabla f(x^k)\|^2 \leq \frac{2M(f(x^0) - \bar{f})}{T}.$$

Remarks:

- Convergence rate: $\mathcal{O}\left(\frac{1}{\sqrt{T}}\right)$.



(a) Asymptotic: Taking $T \rightarrow \infty$ gives $\lim_{T \rightarrow \infty} \|\nabla f(x^T)\|^2 = 0$.

(b) Non-asymptotic: For finite T ,

$$\frac{1}{T} \sum_{k=0}^{T-1} \|\nabla f(x^k)\|^2 \leq \frac{2M(f(x^0) - \bar{f})}{T},$$

Since, minimum of non-negative sum term is upper-bounded by average sum, hence

$$\min_{0 \leq k \leq T-1} \|\nabla f(x^k)\|^2 \leq \frac{2M(f(x^0) - \bar{f})}{T}.$$

Remarks:

- Convergence rate: $\mathcal{O}(\frac{1}{\sqrt{T}})$.
- In practice, choose a large T and apply early stopping.

Case 2 :GD for Convex & M-Smooth





Case 2 :GD for Convex & M-Smooth

Theorem (GD For Convex and M- Smooth)

f is M -smooth and convex. and x^ be a global minimizer of f . Running GD*

$$x^{k+1} = x^k - \frac{1}{M} \nabla f(x^k) \quad (k = 0, 1, \dots, T-1)$$

yields



Case 2 :GD for Convex & M-Smooth

Theorem (GD For Convex and M- Smooth)

f is M -smooth and convex. and x^ be a global minimizer of f . Running GD*

$$x^{k+1} = x^k - \frac{1}{M} \nabla f(x^k) \quad (k = 0, 1, \dots, T-1)$$

yields

$$f(x^T) - f(x^*) \leq \frac{M}{2T} \|x^0 - x^*\|^2.$$

Remarks:

- Convergence rate: $O(1/T)$.



Case 2 :GD for Convex & M-Smooth

Theorem (GD For Convex and M- Smooth)

f is M -smooth and convex. and x^ be a global minimizer of f . Running GD*

$$x^{k+1} = x^k - \frac{1}{M} \nabla f(x^k) \quad (k = 0, 1, \dots, T-1)$$

yields

$$f(x^T) - f(x^*) \leq \frac{M}{2T} \|x^0 - x^*\|^2.$$

Remarks:

- Convergence rate: $O(1/T)$.
- Ensures global convergence to the minimizer.



Case 2 :GD for Convex & M-Smooth

Theorem (GD For Convex and M- Smooth)

f is M -smooth and convex. and x^* be a global minimizer of f . Running GD

$$x^{k+1} = x^k - \frac{1}{M} \nabla f(x^k) \quad (k = 0, 1, \dots, T-1)$$

yields

$$f(x^T) - f(x^*) \leq \frac{M}{2T} \|x^0 - x^*\|^2.$$

Remarks:

- Convergence rate: $O(1/T)$.
- Ensures global convergence to the minimizer.
- Guarantee on the final iterate, not on the gradient norm.



Case 2 :GD for Convex & M-Smooth

Theorem (GD For Convex and M- Smooth)

f is M -smooth and convex. and x^* be a global minimizer of f . Running GD

$$x^{k+1} = x^k - \frac{1}{M} \nabla f(x^k) \quad (k = 0, 1, \dots, T-1)$$

yields

$$f(x^T) - f(x^*) \leq \frac{M}{2T} \|x^0 - x^*\|^2.$$

Remarks:

- Convergence rate: $O(1/T)$.
- Ensures global convergence to the minimizer.
- Guarantee on the final iterate, not on the gradient norm.
- Requires differentiability; non-smooth cases call for subgradient methods.



$$f(x^*) \geq f(x^k) + \langle \nabla f(x^k), x^* - x^k \rangle \quad (\text{convexity})$$



$$f(x^*) \geq f(x^k) + \langle \nabla f(x^k), x^* - x^k \rangle \quad (\text{convexity})$$

$$f(x^k) \leq f(x^*) - \langle \nabla f(x^k), x^* - x^k \rangle$$

$$f(x^*) \geq f(x^k) + \langle \nabla f(x^k), x^* - x^k \rangle \quad (\text{convexity})$$

$$f(x^k) \leq f(x^*) - \langle \nabla f(x^k), x^* - x^k \rangle$$

$$f(x^{k+1}) \leq f(x^k) - \frac{1}{2M} \|\nabla f(x^k)\|^2 \quad (\text{Descent Lemma})$$



$$f(x^*) \geq f(x^k) + \langle \nabla f(x^k), x^* - x^k \rangle \quad (\text{convexity})$$

$$f(x^k) \leq f(x^*) - \langle \nabla f(x^k), x^* - x^k \rangle$$

$$f(x^{k+1}) \leq f(x^k) - \frac{1}{2M} \|\nabla f(x^k)\|^2 \quad (\text{Descent Lemma})$$

$$f(x^{k+1}) \leq f(x^*) - \langle \nabla f(x^k), x^* - x^k \rangle - \frac{1}{2M} \|\nabla f(x^k)\|^2$$

$$f(x^*) \geq f(x^k) + \langle \nabla f(x^k), x^* - x^k \rangle \quad (\text{convexity})$$

$$f(x^k) \leq f(x^*) - \langle \nabla f(x^k), x^* - x^k \rangle$$

$$f(x^{k+1}) \leq f(x^k) - \frac{1}{2M} \|\nabla f(x^k)\|^2 \quad (\text{Descent Lemma})$$

$$\begin{aligned} f(x^{k+1}) &\leq f(x^*) - \langle \nabla f(x^k), x^* - x^k \rangle - \frac{1}{2M} \|\nabla f(x^k)\|^2 \\ &= f(x^*) + \langle \nabla f(x^k), x^k - x^* \rangle - \frac{1}{2M} \|\nabla f(x^k)\|^2 \end{aligned}$$

$$f(x^*) \geq f(x^k) + \langle \nabla f(x^k), x^* - x^k \rangle \quad (\text{convexity})$$

$$f(x^k) \leq f(x^*) - \langle \nabla f(x^k), x^* - x^k \rangle$$

$$f(x^{k+1}) \leq f(x^k) - \frac{1}{2M} \|\nabla f(x^k)\|^2 \quad (\text{Descent Lemma})$$

$$\begin{aligned} f(x^{k+1}) &\leq f(x^*) - \langle \nabla f(x^k), x^* - x^k \rangle - \frac{1}{2M} \|\nabla f(x^k)\|^2 \\ &= f(x^*) + \langle \nabla f(x^k), x^k - x^* \rangle - \frac{1}{2M} \|\nabla f(x^k)\|^2 \\ &= f(x^*) + \frac{M}{2} \left(\|x^k - x^*\|^2 - \|x^k - x^* - \frac{1}{M} \nabla f(x^k)\|^2 \right) \end{aligned}$$

$$f(x^*) \geq f(x^k) + \langle \nabla f(x^k), x^* - x^k \rangle \quad (\text{convexity})$$

$$f(x^k) \leq f(x^*) - \langle \nabla f(x^k), x^* - x^k \rangle$$

$$f(x^{k+1}) \leq f(x^k) - \frac{1}{2M} \|\nabla f(x^k)\|^2 \quad (\text{Descent Lemma})$$

$$\begin{aligned} f(x^{k+1}) &\leq f(x^*) - \langle \nabla f(x^k), x^* - x^k \rangle - \frac{1}{2M} \|\nabla f(x^k)\|^2 \\ &= f(x^*) + \langle \nabla f(x^k), x^k - x^* \rangle - \frac{1}{2M} \|\nabla f(x^k)\|^2 \\ &= f(x^*) + \frac{M}{2} \left(\|x^k - x^*\|^2 - \|x^k - x^* - \frac{1}{M} \nabla f(x^k)\|^2 \right) \\ &= f(x^*) + \frac{M}{2} (\|x^k - x^*\|^2 - \|x^{k+1} - x^*\|^2). \end{aligned}$$

$$f(x^*) \geq f(x^k) + \langle \nabla f(x^k), x^* - x^k \rangle \quad (\text{convexity})$$

$$f(x^k) \leq f(x^*) - \langle \nabla f(x^k), x^* - x^k \rangle$$

$$f(x^{k+1}) \leq f(x^k) - \frac{1}{2M} \|\nabla f(x^k)\|^2 \quad (\text{Descent Lemma})$$

$$\begin{aligned} f(x^{k+1}) &\leq f(x^*) - \langle \nabla f(x^k), x^* - x^k \rangle - \frac{1}{2M} \|\nabla f(x^k)\|^2 \\ &= f(x^*) + \langle \nabla f(x^k), x^k - x^* \rangle - \frac{1}{2M} \|\nabla f(x^k)\|^2 \\ &= f(x^*) + \frac{M}{2} \left(\|x^k - x^*\|^2 - \|x^k - x^* - \frac{1}{M} \nabla f(x^k)\|^2 \right) \\ &= f(x^*) + \frac{M}{2} (\|x^k - x^*\|^2 - \|x^{k+1} - x^*\|^2). \end{aligned}$$



Summing over $k = 0, \dots, T - 1$ gives

$$\sum_{k=0}^{T-1} (f(x^{k+1}) - f(x^*)) \leq$$



Summing over $k = 0, \dots, T - 1$ gives

$$\sum_{k=0}^{T-1} (f(x^{k+1}) - f(x^*)) \leq \frac{M}{2} (\|x^0 - x^*\|^2 - \|x^T - x^*\|^2) \leq$$



Summing over $k = 0, \dots, T - 1$ gives

$$\sum_{k=0}^{T-1} (f(x^{k+1}) - f(x^*)) \leq \frac{M}{2} (\|x^0 - x^*\|^2 - \|x^T - x^*\|^2) \leq \frac{M}{2} \|x^0 - x^*\|^2.$$



Summing over $k = 0, \dots, T - 1$ gives

$$\sum_{k=0}^{T-1} (f(x^{k+1}) - f(x^*)) \leq \frac{M}{2} (\|x^0 - x^*\|^2 - \|x^T - x^*\|^2) \leq \frac{M}{2} \|x^0 - x^*\|^2.$$

Since $\{f(x^k)\}$ is non-increasing due to Decent lemma and hence any term will be bounded by average quantity,



Summing over $k = 0, \dots, T - 1$ gives

$$\sum_{k=0}^{T-1} (f(x^{k+1}) - f(x^*)) \leq \frac{M}{2} (\|x^0 - x^*\|^2 - \|x^T - x^*\|^2) \leq \frac{M}{2} \|x^0 - x^*\|^2.$$

Since $\{f(x^k)\}$ is non-increasing due to Decent lemma and hence any term will be bounded by average quantity,

$$f(x^T) - f(x^*) \leq \frac{1}{T} \sum_{k=0}^{T-1} (f(x^k) - f(x^*)) \leq \frac{M}{2T} \|x^0 - x^*\|^2.$$



Case 3:M-Smooth and m-Strongly Convex

Lemma

If f is m -strongly convex and let x^ is a minima of f then*

$$\|\nabla f(x)\|^2 \geq 2m(f(x) - f(x^*)) \quad (3)$$



Case 3:M-Smooth and m-Strongly Convex

Lemma

If f is m -strongly convex and let x^ is a minima of f then*

$$\|\nabla f(x)\|^2 \geq 2m(f(x) - f(x^*)) \quad (3)$$

Proof:

- since f is m -strongly convex



Case 3: M -Smooth and m -Strongly Convex

Lemma

If f is m -strongly convex and let x^* is a minima of f then

$$\|\nabla f(x)\|^2 \geq 2m(f(x) - f(x^*)) \quad (3)$$

Proof:

- since f is m -strongly convex

$$f(y) \geq f(x) + \langle \nabla f(x), y - x \rangle + \frac{m}{2} \|y - x\|^2.$$

Case 3: M -Smooth and m -Strongly Convex

Lemma

If f is m -strongly convex and let x^* is a minima of f then

$$\|\nabla f(x)\|^2 \geq 2m(f(x) - f(x^*)) \quad (3)$$

Proof:

- since f is m -strongly convex

$$f(y) \geq f(x) + \langle \nabla f(x), y - x \rangle + \frac{m}{2} \|y - x\|^2.$$

- Fix x and optimize over y then



Case 3:M-Smooth and m-Strongly Convex

Lemma

If f is m -strongly convex and let x^* is a minima of f then

$$\|\nabla f(x)\|^2 \geq 2m(f(x) - f(x^*)) \quad (3)$$

Proof:

- since f is m -strongly convex

$$f(y) \geq f(x) + \langle \nabla f(x), y - x \rangle + \frac{m}{2} \|y - x\|^2.$$

- Fix x and optimize over y then

$$\nabla f(x) + \frac{m}{2}(2(y - x)) = 0$$

Case 3: M -Smooth and m -Strongly Convex

Lemma

If f is m -strongly convex and let x^* is a minima of f then

$$\|\nabla f(x)\|^2 \geq 2m(f(x) - f(x^*)) \quad (3)$$

Proof:

- since f is m -strongly convex

$$f(y) \geq f(x) + \langle \nabla f(x), y - x \rangle + \frac{m}{2} \|y - x\|^2.$$

- Fix x and optimize over y then

$$\begin{aligned} \nabla f(x) + \frac{m}{2}(2(y - x)) &= 0 \\ \Rightarrow y &= x - \frac{1}{m} \nabla f(x). \end{aligned}$$



- Substitute the value of y in RHS of above equation and LHS in $f(x^*)$,



- Substitute the value of y in RHS of above equation and LHS in $f(x^*)$,

$$f(x^*) \geq f(x) + \langle \nabla f(x), -\frac{1}{m} \nabla f(x) \rangle + \frac{m}{2} \left\| -\frac{1}{m} \nabla f(x) \right\|^2$$

- Substitute the value of y in RHS of above equation and LHS in $f(x^*)$,

$$\begin{aligned} f(x^*) &\geq f(x) + \langle \nabla f(x), -\frac{1}{m} \nabla f(x) \rangle + \frac{m}{2} \left\| -\frac{1}{m} \nabla f(x) \right\|^2 \\ &= f(x) - \frac{1}{m} \langle \nabla f(x), \nabla f(x) \rangle + \frac{1}{2m} \|\nabla f(x)\|^2 \end{aligned}$$

- Substitute the value of y in RHS of above equation and LHS in $f(x^*)$,

$$\begin{aligned} f(x^*) &\geq f(x) + \langle \nabla f(x), -\frac{1}{m} \nabla f(x) \rangle + \frac{m}{2} \left\| -\frac{1}{m} \nabla f(x) \right\|^2 \\ &= f(x) - \frac{1}{m} \langle \nabla f(x), \nabla f(x) \rangle + \frac{1}{2m} \|\nabla f(x)\|^2 \\ &= f(x) - \frac{1}{2m} \|\nabla f(x)\|^2 \end{aligned}$$

- Substitute the value of y in RHS of above equation and LHS in $f(x^*)$,

$$\begin{aligned} f(x^*) &\geq f(x) + \langle \nabla f(x), -\frac{1}{m} \nabla f(x) \rangle + \frac{m}{2} \left\| -\frac{1}{m} \nabla f(x) \right\|^2 \\ &= f(x) - \frac{1}{m} \langle \nabla f(x), \nabla f(x) \rangle + \frac{1}{2m} \|\nabla f(x)\|^2 \\ &= f(x) - \frac{1}{2m} \|\nabla f(x)\|^2 \\ \Rightarrow \|\nabla f(x)\|^2 &\geq 2m(f(x) - f(x^*)). \end{aligned}$$



Theorem

Let f be M -smooth and m -strongly convex. We run GD with $\alpha_k = \frac{1}{M}$ then,



GD For M-Smooth and m-strongly convex

Theorem

Let f be M -smooth and m -strongly convex. We run GD with $\alpha_k = \frac{1}{M}$ then,

$$f(x^T) - f(x^*) \leq \left(1 - \frac{m}{M}\right)^T (f(x^0) - f(x^*)). \quad (4)$$

Proof:



GD For M-Smooth and m-strongly convex

Theorem

Let f be M -smooth and m -strongly convex. We run GD with $\alpha_k = \frac{1}{M}$ then,

$$f(x^T) - f(x^*) \leq \left(1 - \frac{m}{M}\right)^T (f(x^0) - f(x^*)). \quad (4)$$

Proof:

- Using Descent lemma:we get

$$f(x^{k+1}) \leq f(x^k) - \frac{1}{2M} \|\nabla f(x^k)\|^2$$



GD For M-Smooth and m-strongly convex

Theorem

Let f be M -smooth and m -strongly convex. We run GD with $\alpha_k = \frac{1}{M}$ then,

$$f(x^T) - f(x^*) \leq \left(1 - \frac{m}{M}\right)^T (f(x^0) - f(x^*)). \quad (4)$$

Proof:

- Using Descent lemma:we get

$$f(x^{k+1}) \leq f(x^k) - \frac{1}{2M} \|\nabla f(x^k)\|^2$$

- Using previous lemma for m-strongly convex function



GD For M-Smooth and m-strongly convex

Theorem

Let f be M -smooth and m -strongly convex. We run GD with $\alpha_k = \frac{1}{M}$ then,

$$f(x^T) - f(x^*) \leq \left(1 - \frac{m}{M}\right)^T (f(x^0) - f(x^*)). \quad (4)$$

Proof:

- Using Descent lemma:we get

$$f(x^{k+1}) \leq f(x^k) - \frac{1}{2M} \|\nabla f(x^k)\|^2$$

- Using previous lemma for m-strongly convex function ,we get

$$f(x^{k+1}) \leq f(x^k) - \frac{m}{M} (f(x^k) - f(x^*))$$



$$f(x^{k+1}) - f(x^*) \leq f(x^k) - \frac{m}{M}(f(x^k) - f(x^*)) - f(x^*)$$



$$\begin{aligned} f(x^{k+1}) - f(x^*) &\leq f(x^k) - \frac{m}{M}(f(x^k) - f(x^*)) - f(x^*) \\ &= \left(1 - \frac{m}{M}\right)[f(x^k) - f(x^*)] \end{aligned}$$



$$\begin{aligned}f(x^{k+1}) - f(x^*) &\leq f(x^k) - \frac{m}{M}(f(x^k) - f(x^*)) - f(x^*) \\&= \left(1 - \frac{m}{M}\right)[f(x^k) - f(x^*)] \\&\leq \left(1 - \frac{m}{M}\right)^2[f(x^{k-1}) - f(x^*)]\end{aligned}$$



$$\begin{aligned}f(x^{k+1}) - f(x^*) &\leq f(x^k) - \frac{m}{M}(f(x^k) - f(x^*)) - f(x^*) \\&= \left(1 - \frac{m}{M}\right)[f(x^k) - f(x^*)] \\&\leq \left(1 - \frac{m}{M}\right)^2[f(x^{k-1}) - f(x^*)] \\&\vdots\end{aligned}$$



$$\begin{aligned}f(x^{k+1}) - f(x^*) &\leq f(x^k) - \frac{m}{M}(f(x^k) - f(x^*)) - f(x^*) \\&= \left(1 - \frac{m}{M}\right)[f(x^k) - f(x^*)] \\&\leq \left(1 - \frac{m}{M}\right)^2[f(x^{k-1}) - f(x^*)] \\&\vdots \\&\leq \left(1 - \frac{m}{M}\right)^k[f(x^0) - f(x^*)]\end{aligned}$$



$$\begin{aligned}
 f(x^{k+1}) - f(x^*) &\leq f(x^k) - \frac{m}{M}(f(x^k) - f(x^*)) - f(x^*) \\
 &= \left(1 - \frac{m}{M}\right)[f(x^k) - f(x^*)] \\
 &\leq \left(1 - \frac{m}{M}\right)^2 [f(x^{k-1}) - f(x^*)] \\
 &\vdots \\
 &\leq \left(1 - \frac{m}{M}\right)^k [f(x^0) - f(x^*)]
 \end{aligned}$$

- ① In this case, we can give a guarantee on $\|x^T - x^*\|$. The choice of step-size in that case would be $\alpha_k = \frac{2}{M+m}$
- ② Left as exercise!

Convergence Rates of Gradient Descent



Assumptions	Rate	Guarantee



Convergence Rates of Gradient Descent

Assumptions	Rate	Guarantee
f is M -smooth	$\min_{0 \leq k < T} \ \nabla f(x_k)\ ^2 \leq \frac{2M(f(x_0) - f^*)}{T}$	Best-iterate gradient norm



Convergence Rates of Gradient Descent

Assumptions	Rate	Guarantee
f is M -smooth	$\frac{\min_{0 \leq k < T} \ \nabla f(x_k)\ ^2}{2M(f(x_0) - f^*)} \leq$	Best-iterate gradient norm
f is M -smooth and convex	$f(x_T) - f^* \leq \frac{M}{2T} \ x_0 - x^*\ ^2$	Final-iterate function gap

Convergence Rates of Gradient Descent

Assumptions	Rate	Guarantee
f is M -smooth	$\min_{0 \leq k < T} \ \nabla f(x_k)\ ^2 \leq \frac{2M(f(x_0) - f^*)}{T}$	Best-iterate gradient norm
f is M -smooth and convex	$f(x_T) - f^* \leq \frac{M}{2T} \ x_0 - x^*\ ^2$	Final-iterate function gap
f is M -smooth, m -strongly convex	$f(x_T) - f^* \leq \left(1 - \frac{m}{M}\right)^T (f(x_0) - f^*)$	Exponential (linear) convergence

Gradient Descent: Rates under Various Assumptions



Let Here, $\kappa = M/m$, $R = \|x_0 - x^*\|$, and $\Delta_0 = f(x_0) - f^*$.

Assumptions	Criterion	Iterations

Gradient Descent: Rates under Various Assumptions



Let Here, $\kappa = M/m$, $R = \|x_0 - x^*\|$, and $\Delta_0 = f(x_0) - f^*$.

Assumptions	Criterion	Iterations
f is m -strongly convex, M -smooth	$\ x_T - x^*\ \leq \varepsilon$	$O(\kappa \ln(R/\varepsilon))$

Gradient Descent: Rates under Various Assumptions



Let Here, $\kappa = M/m$, $R = \|x_0 - x^*\|$, and $\Delta_0 = f(x_0) - f^*$.

Assumptions	Criterion	Iterations
f is m -strongly convex, M -smooth	$\ x_T - x^*\ \leq \varepsilon$	$O(\kappa \ln(R/\varepsilon))$
f is convex, M -smooth	$f(x_T) - f^* \leq \varepsilon$	$O\left(\frac{MR^2}{\varepsilon}\right)$

Gradient Descent: Rates under Various Assumptions



Let Here, $\kappa = M/m$, $R = \|x_0 - x^*\|$, and $\Delta_0 = f(x_0) - f^*$.

Assumptions	Criterion	Iterations
f is m -strongly convex, M -smooth	$\ x_T - x^*\ \leq \varepsilon$	$O(\kappa \ln(R/\varepsilon))$
f is convex, M -smooth	$f(x_T) - f^* \leq \varepsilon$	$O(\frac{M R^2}{\varepsilon})$
f is nonconvex, M -smooth	$\min_{0 \leq k < T} \ \nabla f(x_k)\ \leq \varepsilon$	$O(\frac{M \Delta_0}{\varepsilon^2})$



Consider the following constrained optimization problem:

$$\min_{x \in \Omega} f(x) \quad (5)$$



Consider the following constrained optimization problem:

$$\min_{x \in \Omega} f(x) \quad (5)$$

where $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is a continuously differentiable and M -smooth function,



Consider the following constrained optimization problem:

$$\min_{x \in \Omega} f(x) \quad (5)$$

where $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is a continuously differentiable and M -smooth function, and $\Omega \subseteq \mathbb{R}^d$ is a closed convex set.



Consider the following constrained optimization problem:

$$\min_{x \in \Omega} f(x) \quad (5)$$

where $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is a continuously differentiable and M -smooth function, and $\Omega \subseteq \mathbb{R}^d$ is a closed convex set.

Euclidean projection





Let $\Omega \subseteq \mathbb{R}^d$ be a closed and convex set.



Let $\Omega \subseteq \mathbb{R}^d$ be a closed and convex set. The projection of a point $x \in \mathbb{R}^d$ onto Ω is defined as:



Let $\Omega \subseteq \mathbb{R}^d$ be a closed and convex set. The projection of a point $x \in \mathbb{R}^d$ onto Ω is defined as:

$$P_{\Omega}(x)$$



Let $\Omega \subseteq \mathbb{R}^d$ be a closed and convex set. The projection of a point $x \in \mathbb{R}^d$ onto Ω is defined as:

$$P_{\Omega}(x) := \arg \min_{z \in \Omega} \frac{1}{2} \|z - x\|^2 \quad (6)$$

Example 1: Non-Negative Orthant





Example 1: Non-Negative Orthant

Consider the set $\Omega \subseteq \mathbb{R}^d$ defined as the **non-negative orthant**:



Example 1: Non-Negative Orthant

Consider the set $\Omega \subseteq \mathbb{R}^d$ defined as the **non-negative orthant**:

$$\Omega := \left\{ x \in \mathbb{R}^d ; x_i \geq 0, \quad \forall i = 1, \dots, d \right\} \quad (7)$$



Example 1: Non-Negative Orthant

Consider the set $\Omega \subseteq \mathbb{R}^d$ defined as the **non-negative orthant**:

$$\Omega := \left\{ x \in \mathbb{R}^d ; x_i \geq 0, \quad \forall i = 1, \dots, d \right\} \quad (7)$$

The projection of a vector $y \in \mathbb{R}^d$ onto Ω is given component-wise by:



Example 1: Non-Negative Orthant

Consider the set $\Omega \subseteq \mathbb{R}^d$ defined as the **non-negative orthant**:

$$\Omega := \left\{ x \in \mathbb{R}^d ; x_i \geq 0, \quad \forall i = 1, \dots, d \right\} \quad (7)$$

The projection of a vector $y \in \mathbb{R}^d$ onto Ω is given component-wise by:

$$[P_{\Omega}(y)]_i = \max\{0, y_i\}, \quad \text{for each } i = 1, \dots, d \quad (8)$$

Example 2: Box



Example 2: Box



Consider the set $\Omega \subseteq \mathbb{R}^d$ defined as:

$$\Omega := \left\{ x \in \mathbb{R}^d ; \|x\|_{\infty} \leq 1 \right\} \quad (9)$$



Example 2: Box

Consider the set $\Omega \subseteq \mathbb{R}^d$ defined as:

$$\Omega := \left\{ x \in \mathbb{R}^d ; \|x\|_{\infty} \leq 1 \right\} \quad (9)$$

Exercise: Given a point $y \in \mathbb{R}^d$, determine the projection $P_{\Omega}(y)$ onto Ω .





Consider the set $\Omega \subseteq \mathbb{R}^d$ defined as:

$$\Omega := \left\{ x \in \mathbb{R}^d ; \|x\|_2 \leq 1 \right\} \quad (10)$$



L_2 Norm Ball

Consider the set $\Omega \subseteq \mathbb{R}^d$ defined as:

$$\Omega := \left\{ x \in \mathbb{R}^d ; \|x\|_2 \leq 1 \right\} \quad (10)$$

The projection $P_\Omega(y)$ onto Ω is given by:



L_2 Norm Ball

Consider the set $\Omega \subseteq \mathbb{R}^d$ defined as:

$$\Omega := \left\{ x \in \mathbb{R}^d ; \|x\|_2 \leq 1 \right\} \quad (10)$$

The projection $P_\Omega(y)$ onto Ω is given by:

- If $\|y\|_2 \leq 1$,



L_2 Norm Ball

Consider the set $\Omega \subseteq \mathbb{R}^d$ defined as:

$$\Omega := \left\{ x \in \mathbb{R}^d ; \|x\|_2 \leq 1 \right\} \quad (10)$$

The projection $P_\Omega(y)$ onto Ω is given by:

- If $\|y\|_2 \leq 1$, then $P_\Omega(y) = y$



L_2 Norm Ball

Consider the set $\Omega \subseteq \mathbb{R}^d$ defined as:

$$\Omega := \left\{ x \in \mathbb{R}^d ; \|x\|_2 \leq 1 \right\} \quad (10)$$

The projection $P_\Omega(y)$ onto Ω is given by:

- If $\|y\|_2 \leq 1$, then $P_\Omega(y) = y$
- If $\|y\|_2 > 1$,



L_2 Norm Ball

Consider the set $\Omega \subseteq \mathbb{R}^d$ defined as:

$$\Omega := \left\{ x \in \mathbb{R}^d ; \|x\|_2 \leq 1 \right\} \quad (10)$$

The projection $P_\Omega(y)$ onto Ω is given by:

- If $\|y\|_2 \leq 1$, then $P_\Omega(y) = y$
- If $\|y\|_2 > 1$, then $P_\Omega(y) = \frac{y}{\|y\|_2}$



L_2 Norm Ball

Consider the set $\Omega \subseteq \mathbb{R}^d$ defined as:

$$\Omega := \left\{ x \in \mathbb{R}^d ; \|x\|_2 \leq 1 \right\} \quad (10)$$

The projection $P_\Omega(y)$ onto Ω is given by:

- If $\|y\|_2 \leq 1$, then $P_\Omega(y) = y$
- If $\|y\|_2 > 1$, then $P_\Omega(y) = \frac{y}{\|y\|_2}$

Contraction/Non-expansive Property of Projection:



L_2 Norm Ball

Consider the set $\Omega \subseteq \mathbb{R}^d$ defined as:

$$\Omega := \left\{ x \in \mathbb{R}^d ; \|x\|_2 \leq 1 \right\} \quad (10)$$


The projection $P_\Omega(y)$ onto Ω is given by:

- If $\|y\|_2 \leq 1$, then $P_\Omega(y) = y$
- If $\|y\|_2 > 1$, then $P_\Omega(y) = \frac{y}{\|y\|_2}$

Contraction/Non-expansive Property of Projection:

$$\|P_\Omega(x) - P_\Omega(y)\| \leq \|x - y\|, \quad \forall x, y \in \mathbb{R}^n \quad (11)$$

Figure: Illustration of Contraction Property





Projected Gradient Descent:

$$x^{k+1} = P_{\Omega}[x^k - \alpha_k \nabla f(x_k)]$$

- ① Thanks to the Contraction property of Projection, the distance gets smaller after projection
- ② The GD algorithm remains the same with the projection step
- ③ The convergence rates remain the same



- [1] Benjamin Recht and Stephen J. Wright.
Optimization for Data Analysis. 2022.
- [2] Stephen Boyd and Lieven Vandenberghe.
Convex Optimization.
Cambridge University Press, 2004.
- [3] Jorge Nocedal and Stephen J. Wright.
Numerical Optimization, 2nd Edition.
Springer, 2006.

Thank You!