

# Convexity, Gradient Descent, Convergence Guarantees

Avishek Ghosh

Assistant Professor, Dept. of CSE, IIT Bombay



August 3, 2025



1. Convexity and Smoothness
2. Strong Convexity and Examples
3. Gradient Descent and Decent lemma for Smooth functions





## Definition (Convex Set)

A set  $\mathcal{X} \subseteq \mathbb{R}^n$  is *convex* if for any points  $x, y \in \mathcal{X}$  and any scalar  $\lambda \in [0, 1]$ , the combination


$$\lambda x + (1 - \lambda) y \in \mathcal{X}.$$

# Convex set


## Definition (Convex Set)

A set  $\mathcal{X} \subseteq \mathbb{R}^n$  is *convex* if for any points  $x, y \in \mathcal{X}$  and any scalar  $\lambda \in [0, 1]$ , the combination

$$\lambda x + (1 - \lambda) y \in \mathcal{X}.$$



*Convex Set*



*Non-convex Set*

Figure: Convex and Non-convex set.






## Definition (Convex Function)

Let  $\mathcal{X} \subseteq \mathbb{R}^n$  be a convex set. A function  $f : \mathcal{X} \rightarrow \mathbb{R}$  is called *convex* if, for all  $x, y \in \mathcal{X}$  and every  $\lambda \in [0, 1]$ ,

$$f(\lambda x + (1 - \lambda) y) \leq \lambda f(x) + (1 - \lambda) f(y).$$

# Illustration of Convex function





# First order Characterization of Convex Functions





## Theorem

*Let  $\mathcal{X} \subseteq \mathbb{R}^d$  be a convex set and  $f: \mathcal{X} \rightarrow \mathbb{R}$  be differentiable. Then  $f$  is convex if and only if for all  $x, y \in \mathcal{X}$ ,*




## Theorem

Let  $\mathcal{X} \subseteq \mathbb{R}^d$  be a convex set and  $f: \mathcal{X} \rightarrow \mathbb{R}$  be differentiable. Then  $f$  is convex if and only if for all  $x, y \in \mathcal{X}$ ,

$$f(y) \geq f(x) + \nabla f(x)^\top (y - x).$$

**Proof:** Exercise!

# Figure of First order characterization of Convexity



# Second order Characterization of Convex Functions





## Theorem

*Let  $\mathcal{X} \subseteq \mathbb{R}^d$  be a convex set and  $f: \mathcal{X} \rightarrow \mathbb{R}$  be differentiable. Then  $f$  is convex if and only if for all  $x, y \in \mathcal{X}$ ,*



## Theorem

Let  $\mathcal{X} \subseteq \mathbb{R}^d$  be a convex set and  $f: \mathcal{X} \rightarrow \mathbb{R}$  be differentiable. Then  $f$  is convex if and only if for all  $x, y \in \mathcal{X}$ ,

$$\nabla^2 f(x) \succeq 0$$

This means the Hessian matrix  $\nabla^2 f(x)$  is Positive Semi definite (meaning that the eigen values are non-negative) **Examples?**

If  $d = 1$ , implies  $f''(x) \geq 0$  for all  $x$



## Theorem

Let  $\mathcal{X} \subseteq \mathbb{R}^d$  be a convex set and  $f: \mathcal{X} \rightarrow \mathbb{R}$  be differentiable. Then  $f$  is convex if and only if for all  $x, y \in \mathcal{X}$ ,

$$\nabla^2 f(x) \succeq 0$$

This means the Hessian matrix  $\nabla^2 f(x)$  is Positive Semi definite (meaning that the eigen values are non-negative) **Examples?**

If  $d = 1$ , implies  $f''(x) \geq 0$  for all  $x$

**Proof of Theorem:** Exercise!





## Definition

Let  $f : \mathcal{X} \rightarrow \mathbb{R}$ . A point  $x^* \in \mathcal{X}$  is

- **Local minimum** if there exists an open neighborhood  $U$  of  $x^*$  such that

$$f(x^*) \leq f(y) \quad \forall y \in U.$$



## Definition

Let  $f : \mathcal{X} \rightarrow \mathbb{R}$ . A point  $x^* \in \mathcal{X}$  is

- **Local minimum** if there exists an open neighborhood  $U$  of  $x^*$  such that

$$f(x^*) \leq f(y) \quad \forall y \in U.$$

- **Global minimum** if

$$f(x^*) \leq f(y) \quad \forall y \in \mathcal{X}.$$



## Definition

Let  $f : \mathcal{X} \rightarrow \mathbb{R}$ . A point  $x^* \in \mathcal{X}$  is

- **Local minimum** if there exists an open neighborhood  $U$  of  $x^*$  such that

$$f(x^*) \leq f(y) \quad \forall y \in U.$$

- **Global minimum** if

$$f(x^*) \leq f(y) \quad \forall y \in \mathcal{X}.$$

**Question:** Prove that, for convex functions, all local minima are global minima.





## Definition (Smooth Function)

Suppose  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  satisfies

$$\|\nabla f(x) - \nabla f(y)\|_2 \leq M\|x - y\|_2$$

Where  $M > 0$ , then we call  $f$  as  $M$ -smooth/  $M$ -Gradient Lipschitz.



## Definition (Smooth Function)

Suppose  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  satisfies

$$\|\nabla f(x) - \nabla f(y)\|_2 \leq M\|x - y\|_2$$

Where  $M > 0$ , then we call  $f$  as  $M$ -smooth/  $M$ -Gradient Lipschitz.

**Example:** Quadratic  $f(x) = \frac{1}{2}x^T Qx - b^T x$ ;  $Q$  symmetric



## Definition (Smooth Function)

Suppose  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  satisfies

$$\|\nabla f(x) - \nabla f(y)\|_2 \leq M\|x - y\|_2$$

Where  $M > 0$ , then we call  $f$  as  $M$ -smooth/  $M$ -Gradient Lipschitz.

**Example:** Quadratic  $f(x) = \frac{1}{2}x^T Qx - b^T x$ ;  $Q$  symmetric

$$\begin{aligned}\|\nabla f(x) - \nabla f(y)\| &= \|(Qx - b) - (Qy - b)\| \\ &= \|Q(x - y)\| \leq M\|x - y\|\end{aligned}$$



## Definition (Smooth Function)

Suppose  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  satisfies

$$\|\nabla f(x) - \nabla f(y)\|_2 \leq M\|x - y\|_2$$

Where  $M > 0$ , then we call  $f$  as  $M$ -smooth/  $M$ -Gradient Lipschitz.

**Example:** Quadratic  $f(x) = \frac{1}{2}x^T Qx - b^T x$ ;  $Q$  symmetric

$$\begin{aligned}\|\nabla f(x) - \nabla f(y)\| &= \|(Qx - b) - (Qy - b)\| \\ &= \|Q(x - y)\| \leq M\|x - y\|\end{aligned}$$

Hence Quadratic objective function is  $M$ -smooth.



# Equivalent Characterizations of Smoothness





## Theorem

*The following statements are equivalent (  $\forall x, y$  ):*

①  $\|\nabla f(x) - \nabla f(y)\| \leq M\|x - y\|$



## Theorem

*The following statements are equivalent (  $\forall x, y$  ):*

- ①  $\|\nabla f(x) - \nabla f(y)\| \leq M\|x - y\|$
- ②  $\langle \nabla f(x) - \nabla f(y), x - y \rangle \leq M\|x - y\|^2$

# Equivalent Characterizations of Smoothness

## Theorem

*The following statements are equivalent (  $\forall x, y$  ):*

- ①  $\|\nabla f(x) - \nabla f(y)\| \leq M\|x - y\|$
- ②  $\langle \nabla f(x) - \nabla f(y), x - y \rangle \leq M\|x - y\|^2$
- ③  $f(y) - \{f(x) + \langle \nabla f(x), y - x \rangle\} \leq \frac{M}{2}\|x - y\|^2$

# Equivalent Characterizations of Smoothness

## Theorem

*The following statements are equivalent (  $\forall x, y$  ):*

- ①  $\|\nabla f(x) - \nabla f(y)\| \leq M\|x - y\|$
- ②  $\langle \nabla f(x) - \nabla f(y), x - y \rangle \leq M\|x - y\|^2$
- ③  $f(y) - \{f(x) + \langle \nabla f(x), y - x \rangle\} \leq \frac{M}{2}\|x - y\|^2$
- ④  $\langle \nabla f(x) - \nabla f(y), x - y \rangle \geq \frac{1}{M}\|\nabla f(x) - \nabla f(y)\|^2$

# Equivalent Characterizations of Smoothness

## Theorem

*The following statements are equivalent (  $\forall x, y$  ):*

- ①  $\|\nabla f(x) - \nabla f(y)\| \leq M\|x - y\|$
- ②  $\langle \nabla f(x) - \nabla f(y), x - y \rangle \leq M\|x - y\|^2$
- ③  $f(y) - \{f(x) + \langle \nabla f(x), y - x \rangle\} \leq \frac{M}{2}\|x - y\|^2$
- ④  $\langle \nabla f(x) - \nabla f(y), x - y \rangle \geq \frac{1}{M}\|\nabla f(x) - \nabla f(y)\|^2$
- ⑤  $f$  is twice differentiable then:

$$-M\mathbb{I} \preceq \nabla^2 f \preceq M\mathbb{I}$$

where  $M\mathbb{I} - \nabla^2 f$  is P.S.D. matrix.



For  $(1) \Rightarrow (2)$ , Use Cauchy-Schwartz Inequality



For (1)  $\Rightarrow$  (2), Use Cauchy-Schwartz Inequality

$$\langle \nabla f(x) - \nabla f(y), x - y \rangle$$





For (1)  $\Rightarrow$  (2), Use Cauchy-Schwartz Inequality

$$\langle \nabla f(x) - \nabla f(y), x - y \rangle \leq \|\nabla f(x) - \nabla f(y)\| \|x - y\|$$



For (1)  $\Rightarrow$  (2), Use Cauchy-Schwartz Inequality

$$\begin{aligned}\langle \nabla f(x) - \nabla f(y), x - y \rangle &\leq \|\nabla f(x) - \nabla f(y)\| \|x - y\| \\ &\leq M \|x - y\|^2\end{aligned}$$

For (2)  $\Rightarrow$  (3), we use the Interpolation method where  $G : [0, 1] \rightarrow \mathbb{R}$ ,



For (1)  $\Rightarrow$  (2), Use Cauchy-Schwartz Inequality

$$\begin{aligned}\langle \nabla f(x) - \nabla f(y), x - y \rangle &\leq \|\nabla f(x) - \nabla f(y)\| \|x - y\| \\ &\leq M \|x - y\|^2\end{aligned}$$

For (2)  $\Rightarrow$  (3), we use the Interpolation method where  $G : [0, 1] \rightarrow \mathbb{R}$ ,

$$G(t) = f(x + t(y - x)) - f(x) - \langle \nabla f(x), t(y - x) \rangle$$



For (1)  $\Rightarrow$  (2), Use Cauchy-Schwartz Inequality

$$\begin{aligned}\langle \nabla f(x) - \nabla f(y), x - y \rangle &\leq \|\nabla f(x) - \nabla f(y)\| \|x - y\| \\ &\leq M \|x - y\|^2\end{aligned}$$

For (2)  $\Rightarrow$  (3), we use the Interpolation method where  $G : [0, 1] \rightarrow \mathbb{R}$ ,

$$G(t) = f(x + t(y - x)) - f(x) - \langle \nabla f(x), t(y - x) \rangle$$

Observe that  $G(1) = \text{LHS of (3)}$  and  $G(0) = 0$ .



For (1)  $\Rightarrow$  (2), Use Cauchy-Schwartz Inequality

$$\begin{aligned}\langle \nabla f(x) - \nabla f(y), x - y \rangle &\leq \|\nabla f(x) - \nabla f(y)\| \|x - y\| \\ &\leq M \|x - y\|^2\end{aligned}$$

For (2)  $\Rightarrow$  (3), we use the Interpolation method where  $G : [0, 1] \rightarrow \mathbb{R}$ ,

$$G(t) = f(x + t(y - x)) - f(x) - \langle \nabla f(x), t(y - x) \rangle$$

Observe that  $G(1) = \text{LHS of (3)}$  and  $G(0) = 0$ .

By the fundamental theorem of calculus,



For (1)  $\Rightarrow$  (2), Use Cauchy-Schwartz Inequality

$$\begin{aligned}\langle \nabla f(x) - \nabla f(y), x - y \rangle &\leq \|\nabla f(x) - \nabla f(y)\| \|x - y\| \\ &\leq M \|x - y\|^2\end{aligned}$$

For (2)  $\Rightarrow$  (3), we use the Interpolation method where  $G : [0, 1] \rightarrow \mathbb{R}$ ,

$$G(t) = f(x + t(y - x)) - f(x) - \langle \nabla f(x), t(y - x) \rangle$$

Observe that  $G(1) = \text{LHS of (3)}$  and  $G(0) = 0$ .

By the fundamental theorem of calculus,

$$G(1) - G(0) = \int_0^1 G'(t) dt$$



$$\text{LHS of (3)} = \int_0^1 \langle \nabla f(x + t(y - x)), (y - x) \rangle - 0 - \langle \nabla f(x), y - x \rangle dt$$



$$\begin{aligned}\text{LHS of (3)} &= \int_0^1 \langle \nabla f(x + t(y - x)), (y - x) \rangle - 0 - \langle \nabla f(x), y - x \rangle dt \\ &= \int_0^1 \langle \nabla f(x + t(y - x)) - \nabla f(x), y - x \rangle dt\end{aligned}$$





$$\begin{aligned}
 \text{LHS of (3)} &= \int_0^1 \langle \nabla f(x + t(y - x)), (y - x) \rangle - 0 - \langle \nabla f(x), y - x \rangle dt \\
 &= \int_0^1 \langle \nabla f(x + t(y - x)) - \nabla f(x), y - x \rangle dt \\
 &= \int_0^1 \langle \nabla f(x + t(y - x)) - \nabla f(x), t(y - x) \rangle \frac{dt}{t}
 \end{aligned}$$



$$\begin{aligned}
 \text{LHS of (3)} &= \int_0^1 \langle \nabla f(x + t(y - x)), (y - x) \rangle - 0 - \langle \nabla f(x), y - x \rangle dt \\
 &= \int_0^1 \langle \nabla f(x + t(y - x)) - \nabla f(x), y - x \rangle dt \\
 &= \int_0^1 \langle \nabla f(x + t(y - x)) - \nabla f(x), t(y - x) \rangle \frac{dt}{t}
 \end{aligned}$$

Using (2),



$$\begin{aligned}
 \text{LHS of (3)} &= \int_0^1 \langle \nabla f(x + t(y - x)), (y - x) \rangle - 0 - \langle \nabla f(x), y - x \rangle dt \\
 &= \int_0^1 \langle \nabla f(x + t(y - x)) - \nabla f(x), y - x \rangle dt \\
 &= \int_0^1 \langle \nabla f(x + t(y - x)) - \nabla f(x), t(y - x) \rangle \frac{dt}{t}
 \end{aligned}$$

Using (2),

$$G(1) - G(0) \leq \int_0^1 Mt \|x - y\|^2 dt$$



$$\begin{aligned}\text{LHS of (3)} &= \int_0^1 \langle \nabla f(x + t(y - x)), (y - x) \rangle - 0 - \langle \nabla f(x), y - x \rangle dt \\ &= \int_0^1 \langle \nabla f(x + t(y - x)) - \nabla f(x), y - x \rangle dt \\ &= \int_0^1 \langle \nabla f(x + t(y - x)) - \nabla f(x), t(y - x) \rangle \frac{dt}{t}\end{aligned}$$

Using (2),

$$\begin{aligned}G(1) - G(0) &\leq \int_0^1 Mt \|x - y\|^2 dt \\ &= M \|x - y\|^2 \int_0^1 t dt = \frac{M}{2} \|x - y\|^2\end{aligned}$$



$$\begin{aligned}
 \text{LHS of (3)} &= \int_0^1 \langle \nabla f(x + t(y - x)), (y - x) \rangle - 0 - \langle \nabla f(x), y - x \rangle dt \\
 &= \int_0^1 \langle \nabla f(x + t(y - x)) - \nabla f(x), y - x \rangle dt \\
 &= \int_0^1 \langle \nabla f(x + t(y - x)) - \nabla f(x), t(y - x) \rangle \frac{dt}{t}
 \end{aligned}$$

Using (2),

$$\begin{aligned}
 G(1) - G(0) &\leq \int_0^1 M t \|x - y\|^2 dt \\
 &= M \|x - y\|^2 \int_0^1 t dt = \frac{M}{2} \|x - y\|^2
 \end{aligned}$$

For (3)  $\Rightarrow$  (5), we utilize the Taylor Series Expansion.



$$\begin{aligned}
 \text{LHS of (3)} &= \int_0^1 \langle \nabla f(x + t(y - x)), (y - x) \rangle - 0 - \langle \nabla f(x), y - x \rangle dt \\
 &= \int_0^1 \langle \nabla f(x + t(y - x)) - \nabla f(x), y - x \rangle dt \\
 &= \int_0^1 \langle \nabla f(x + t(y - x)) - \nabla f(x), t(y - x) \rangle \frac{dt}{t}
 \end{aligned}$$

Using (2),

$$\begin{aligned}
 G(1) - G(0) &\leq \int_0^1 M t \|x - y\|^2 dt \\
 &= M \|x - y\|^2 \int_0^1 t dt = \frac{M}{2} \|x - y\|^2
 \end{aligned}$$

For (3)  $\Rightarrow$  (5), we utilize the Taylor Series Expansion.

For (4)  $\Rightarrow$  (1), we utilize the Cauchy Schwartz Inequality.



## Definition (Srong Convexity)



## Definition (Strong Convexity)

Let  $\mathcal{C} \subseteq \mathbb{R}^d$  be a nonempty convex set. A function  $f: \mathcal{C} \rightarrow \mathbb{R}$  is called  *$m$ -strongly convex* if there exists  $m > 0$  such that for all  $x, y \in \mathcal{C}$  and every  $\lambda \in [0, 1]$ ,

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda) f(y) - \frac{m}{2} \lambda(1 - \lambda) \|x - y\|^2.$$

## Remark





## Definition (Srong Convexity)

Let  $\mathcal{C} \subseteq \mathbb{R}^d$  be a nonempty convex set. A function  $f: \mathcal{C} \rightarrow \mathbb{R}$  is called *m-strongly convex* if there exists  $m > 0$  such that for all  $x, y \in \mathcal{C}$  and every  $\lambda \in [0, 1]$ ,

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda) f(y) - \frac{m}{2} \lambda(1 - \lambda) \|x - y\|^2.$$

## Remark

- **Convexity:** Ensures for any  $x, y \in \mathcal{C}$  and  $\lambda \in [0, 1]$ ,

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda) f(y).$$

The graph lies below the chord connecting  $(x, f(x))$  and  $(y, f(y))$ .



- **$m$ -Strong Convexity:** Strengthens convexity by quantifying the gap with a quadratic term:



- **$m$ -Strong Convexity:** Strengthens convexity by quantifying the gap with a quadratic term:

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda) f(y) - \frac{m}{2} \lambda(1 - \lambda) \|x - y\|^2.$$



- **$m$ -Strong Convexity:** Strengthens convexity by quantifying the gap with a quadratic term:

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda) f(y) - \frac{m}{2} \lambda(1 - \lambda) \|x - y\|^2.$$

- **Interpretation:** Strong convexity not only bounds  $f$  below its secant, but guarantees a minimum separation (proportional to  $\|x - y\|^2$ ) between the function's graph and that chord (see below Figure).

# Figure: Interpretation of $m$ -strongly convex Function




Figure:  $m$ -strongly convex function  $f$

# Equivalent Characterizations of Strong Convexity





## Theorem

*Let  $\mathcal{C} \subseteq \mathbb{R}^d$  be a nonempty convex set and  $f: \mathcal{C} \rightarrow \mathbb{R}$  be continuously differentiable. Then the following are equivalent:*

- Ⓐ  $f$  is  $m$ -strongly convex.*



## Theorem

Let  $\mathcal{C} \subseteq \mathbb{R}^d$  be a nonempty convex set and  $f: \mathcal{C} \rightarrow \mathbb{R}$  be continuously differentiable. Then the following are equivalent:

- a  $f$  is  $m$ -strongly convex.
- b  $\|\nabla f(x) - \nabla f(y)\| \geq m \|x - y\|$  for all  $x, y \in \mathcal{C}$ .





## Theorem

Let  $\mathcal{C} \subseteq \mathbb{R}^d$  be a nonempty convex set and  $f: \mathcal{C} \rightarrow \mathbb{R}$  be continuously differentiable. Then the following are equivalent:

- a  $f$  is  $m$ -strongly convex.
- b  $\|\nabla f(x) - \nabla f(y)\| \geq m \|x - y\|$  for all  $x, y \in \mathcal{C}$ .
- c  $(\nabla f(x) - \nabla f(y))^\top (x - y) \geq m \|x - y\|^2$  for all  $x, y \in \mathcal{C}$ .



## Theorem

Let  $\mathcal{C} \subseteq \mathbb{R}^d$  be a nonempty convex set and  $f: \mathcal{C} \rightarrow \mathbb{R}$  be continuously differentiable. Then the following are equivalent:

- a  $f$  is  $m$ -strongly convex.
- b  $\|\nabla f(x) - \nabla f(y)\| \geq m \|x - y\|$  for all  $x, y \in \mathcal{C}$ .
- c  $(\nabla f(x) - \nabla f(y))^\top (x - y) \geq m \|x - y\|^2$  for all  $x, y \in \mathcal{C}$ .
- d  $f(y) - (f(x) + \nabla f(x)^\top (y - x)) \geq \frac{m}{2} \|x - y\|^2$  for all  $x, y \in \mathcal{C}$ .



## Theorem

Let  $\mathcal{C} \subseteq \mathbb{R}^d$  be a nonempty convex set and  $f: \mathcal{C} \rightarrow \mathbb{R}$  be continuously differentiable. Then the following are equivalent:

- a  $f$  is  $m$ -strongly convex.
- b  $\|\nabla f(x) - \nabla f(y)\| \geq m \|x - y\|$  for all  $x, y \in \mathcal{C}$ .
- c  $(\nabla f(x) - \nabla f(y))^\top (x - y) \geq m \|x - y\|^2$  for all  $x, y \in \mathcal{C}$ .
- d  $f(y) - (f(x) + \nabla f(x)^\top (y - x)) \geq \frac{m}{2} \|x - y\|^2$  for all  $x, y \in \mathcal{C}$ .
- e  $(\nabla f(x) - \nabla f(y))^\top (x - y) \leq \frac{1}{m} \|\nabla f(x) - \nabla f(y)\|^2$  for all  $x, y \in \mathcal{C}$ .



## Theorem

Let  $\mathcal{C} \subseteq \mathbb{R}^d$  be a nonempty convex set and  $f: \mathcal{C} \rightarrow \mathbb{R}$  be continuously differentiable. Then the following are equivalent:

- a  $f$  is  $m$ -strongly convex.
- b  $\|\nabla f(x) - \nabla f(y)\| \geq m \|x - y\|$  for all  $x, y \in \mathcal{C}$ .
- c  $(\nabla f(x) - \nabla f(y))^\top (x - y) \geq m \|x - y\|^2$  for all  $x, y \in \mathcal{C}$ .
- d  $f(y) - (f(x) + \nabla f(x)^\top (y - x)) \geq \frac{m}{2} \|x - y\|^2$  for all  $x, y \in \mathcal{C}$ .
- e  $(\nabla f(x) - \nabla f(y))^\top (x - y) \leq \frac{1}{m} \|\nabla f(x) - \nabla f(y)\|^2$  for all  $x, y \in \mathcal{C}$ .
- f If  $f$  is twice differentiable, then  $\nabla^2 f(x) \succeq mI$  for all  $x \in \mathcal{C}$ .



## Example 1: Quadratic Function

Let  $f: \mathbb{R}^d \rightarrow \mathbb{R}$  be defined by

$$f(x) = \frac{1}{2} x^\top Q x - b^\top x,$$

where  $Q \in \mathbb{R}^{d \times d}$  is symmetric positive definite and  $b \in \mathbb{R}^d$ .



## Example 1: Quadratic Function

Let  $f: \mathbb{R}^d \rightarrow \mathbb{R}$  be defined by

$$f(x) = \frac{1}{2} x^\top Q x - b^\top x,$$

where  $Q \in \mathbb{R}^{d \times d}$  is symmetric positive definite and  $b \in \mathbb{R}^d$ .

- $\nabla f(x) - \nabla f(y) = Q(x - y),$



## Example 1: Quadratic Function

Let  $f: \mathbb{R}^d \rightarrow \mathbb{R}$  be defined by

$$f(x) = \frac{1}{2} x^\top Q x - b^\top x,$$

where  $Q \in \mathbb{R}^{d \times d}$  is symmetric positive definite and  $b \in \mathbb{R}^d$ .

- $\nabla f(x) - \nabla f(y) = Q(x - y),$
- Then

$$\|\nabla f(x) - \nabla f(y)\| = \|Q(x - y)\|$$



## Example 1: Quadratic Function

Let  $f: \mathbb{R}^d \rightarrow \mathbb{R}$  be defined by

$$f(x) = \frac{1}{2} x^\top Q x - b^\top x,$$

where  $Q \in \mathbb{R}^{d \times d}$  is symmetric positive definite and  $b \in \mathbb{R}^d$ .

- $\nabla f(x) - \nabla f(y) = Q(x - y),$
- Then

$$\|\nabla f(x) - \nabla f(y)\| = \|Q(x - y)\|$$

- Since  $Q \succeq \lambda_{\min}(Q) I$ , It follows





## Example 1: Quadratic Function

Let  $f: \mathbb{R}^d \rightarrow \mathbb{R}$  be defined by

$$f(x) = \frac{1}{2} x^\top Q x - b^\top x,$$

where  $Q \in \mathbb{R}^{d \times d}$  is symmetric positive definite and  $b \in \mathbb{R}^d$ .

- $\nabla f(x) - \nabla f(y) = Q(x - y)$ ,
- Then

$$\|\nabla f(x) - \nabla f(y)\| = \|Q(x - y)\|$$

- Since  $Q \succeq \lambda_{\min}(Q) I$ , It follows

$$\|\nabla f(x) - \nabla f(y)\| \geq \lambda_{\min}(Q) \|x - y\|.$$



## Example 1: Quadratic Function

Let  $f: \mathbb{R}^d \rightarrow \mathbb{R}$  be defined by

$$f(x) = \frac{1}{2} x^\top Q x - b^\top x,$$

where  $Q \in \mathbb{R}^{d \times d}$  is symmetric positive definite and  $b \in \mathbb{R}^d$ .

- $\nabla f(x) - \nabla f(y) = Q(x - y)$ ,
- Then

$$\|\nabla f(x) - \nabla f(y)\| = \|Q(x - y)\|$$

- Since  $Q \succeq \lambda_{\min}(Q) I$ , It follows

$$\|\nabla f(x) - \nabla f(y)\| \geq \lambda_{\min}(Q) \|x - y\|.$$

Thus By Above Theorem ,  $f$  is  $\lambda_{\min}(Q)$ -strongly convex.



## Example 2: Log–Sum–Exp Function

Consider

$$f: [0, \infty) \rightarrow \mathbb{R}, \quad f(x) = \log(1 + e^x).$$

Then:

- **Gradient:**



## Example 2: Log-Sum-Exp Function

Consider

$$f: [0, \infty) \rightarrow \mathbb{R}, \quad f(x) = \log(1 + e^x).$$

Then:

- **Gradient:**

$$\nabla f(x) = \frac{e^x}{1 + e^x}.$$

- **Hessian:**



## Example 2: Log-Sum-Exp Function

Consider

$$f: [0, \infty) \rightarrow \mathbb{R}, \quad f(x) = \log(1 + e^x).$$

Then:

- **Gradient:**

$$\nabla f(x) = \frac{e^x}{1 + e^x}.$$

- **Hessian:**

$$\nabla^2 f(x) = \frac{e^x}{(1 + e^x)^2} \geq 0,$$

so  $f$  is convex on  $[0, \infty)$ .

## Example 2: Log-Sum-Exp Function

Consider

$$f: [0, \infty) \rightarrow \mathbb{R}, \quad f(x) = \log(1 + e^x).$$

Then:

- **Gradient:**

$$\nabla f(x) = \frac{e^x}{1 + e^x}.$$

- **Hessian:**

$$\nabla^2 f(x) = \frac{e^x}{(1 + e^x)^2} \geq 0,$$

so  $f$  is convex on  $[0, \infty)$ .

- **Convex but not strongly convex on  $[0, \infty)$ :**

## Example 2: Log-Sum-Exp Function

Consider

$$f: [0, \infty) \rightarrow \mathbb{R}, \quad f(x) = \log(1 + e^x).$$

Then:

- **Gradient:**

$$\nabla f(x) = \frac{e^x}{1 + e^x}.$$

- **Hessian:**

$$\nabla^2 f(x) = \frac{e^x}{(1 + e^x)^2} \geq 0,$$

so  $f$  is convex on  $[0, \infty)$ .

- **Convex but not strongly convex on  $[0, \infty)$ :**  $\inf_{x \geq 0} \nabla^2 f(x) = 0$ , so no  $m > 0$  exists with  $\nabla^2 f(x) \geq m$  for all  $x \geq 0$ .



- **Strong convexity on  $[0, \beta]$ :**



## Example 2: Contd..

- **Strong convexity on  $[0, \beta]$ :** For any  $\beta > 0$ , set

$$m = \min_{x \in [0, \beta]} \nabla^2 f(x) = \frac{e^\beta}{(1 + e^\beta)^2} > 0.$$

## Example 2: Contd..

- **Strong convexity on  $[0, \beta]$ :** For any  $\beta > 0$ , set

$$m = \min_{x \in [0, \beta]} \nabla^2 f(x) = \frac{e^\beta}{(1 + e^\beta)^2} > 0.$$

Then  $\nabla^2 f(x) \geq m$  on  $[0, \beta]$ , so  $f$  is  $m$ -strongly convex there.

- **Smoothness:**



## Example 2: Contd..

- **Strong convexity on  $[0, \beta]$ :** For any  $\beta > 0$ , set

$$m = \min_{x \in [0, \beta]} \nabla^2 f(x) = \frac{e^\beta}{(1 + e^\beta)^2} > 0.$$

Then  $\nabla^2 f(x) \geq m$  on  $[0, \beta]$ , so  $f$  is  $m$ -strongly convex there.

- **Smoothness:**  $\nabla^2 f(x) \leq \nabla^2 f(0) = \frac{1}{4}$  for all  $x \geq 0$ , hence  $f$  is  $M$ -smooth with

$$M = \frac{1}{4}.$$



## Example 3: Quartic Function on $[-1, 1]$

- Let  $f: [-1, 1] \rightarrow \mathbb{R}$  be defined by  $f(x) = x^4$ .



## Example 3: Quartic Function on $[-1, 1]$

- Let  $f: [-1, 1] \rightarrow \mathbb{R}$  be defined by  $f(x) = x^4$ .
- Its second derivative is

$$f''(x) = 12x^2,$$



## Example 3: Quartic Function on $[-1, 1]$

- Let  $f: [-1, 1] \rightarrow \mathbb{R}$  be defined by  $f(x) = x^4$ .
- Its second derivative is

$$f''(x) = 12x^2,$$

so  $0 \leq f''(x) \leq 12$  for all  $x \in [-1, 1]$ .

- Hence  $f$  is convex and  $M$ -smooth with  $M = 12$ .



## Example 3: Quartic Function on $[-1, 1]$

- Let  $f: [-1, 1] \rightarrow \mathbb{R}$  be defined by  $f(x) = x^4$ .
- Its second derivative is

$$f''(x) = 12x^2,$$

so  $0 \leq f''(x) \leq 12$  for all  $x \in [-1, 1]$ .

- Hence  $f$  is convex and  $M$ -smooth with  $M = 12$ .
- However, since  $f''(0) = 0$ , there is no  $m > 0$  such that  $f''(x) \geq m$  on  $[-1, 1]$ ; thus  $f$  is not strongly convex.

## Example 4: Invex Function (Non-Convex but Smooth)



- Define

$$f: \left[-\frac{1}{2}, \infty\right) \times [1, \infty) \rightarrow \mathbb{R}, \quad f(x, y) = y(x^2 - 1)^2.$$



## Example 4: Invex Function (Non-Convex but Smooth)



- Define

$$f: \left[-\frac{1}{2}, \infty\right) \times [1, \infty) \rightarrow \mathbb{R}, \quad f(x, y) = y(x^2 - 1)^2.$$

- Hessian matrix

## Example 4: Invex Function (Non-Convex but Smooth)



- Define

$$f: [-\frac{1}{2}, \infty) \times [1, \infty) \rightarrow \mathbb{R}, \quad f(x, y) = y(x^2 - 1)^2.$$

- Hessian matrix

$$\nabla^2 f(x, y) = \begin{pmatrix} 4y(3x^2 - 1) & 4x(x^2 - 1) \\ 4x(x^2 - 1) & 0 \end{pmatrix}.$$

## Example 4: Invex Function (Non-Convex but Smooth)



- Define

$$f: [-\frac{1}{2}, \infty) \times [1, \infty) \rightarrow \mathbb{R}, \quad f(x, y) = y(x^2 - 1)^2.$$

- Hessian matrix

$$\nabla^2 f(x, y) = \begin{pmatrix} 4y(3x^2 - 1) & 4x(x^2 - 1) \\ 4x(x^2 - 1) & 0 \end{pmatrix}.$$

- Since  $\nabla^2 f(x, y)$  is not positive definite on the entire domain,  $f$  is *not convex*.

## Example 4: Invex Function (Non-Convex but Smooth)



- Define

$$f: [-\tfrac{1}{2}, \infty) \times [1, \infty) \rightarrow \mathbb{R}, \quad f(x, y) = y(x^2 - 1)^2.$$

- Hessian matrix

$$\nabla^2 f(x, y) = \begin{pmatrix} 4y(3x^2 - 1) & 4x(x^2 - 1) \\ 4x(x^2 - 1) & 0 \end{pmatrix}.$$

- Since  $\nabla^2 f(x, y)$  is not positive definite on the entire domain,  $f$  is *not convex*.
- $f \in C^2$  with continuous Hessian  $\Rightarrow$  on any bounded subset of  $[-\frac{1}{2}, \infty) \times [1, \infty)$ ,  $f$  is  $M$ -smooth.



# Gradient Descent(GD) Formulation

**Problem:**

$$\min_{x \in \mathbb{R}^d} f(x)$$

.



# Gradient Descent(GD) Formulation

**Problem:**

$$\min_{x \in \mathbb{R}^d} f(x)$$

**Gradient Descent Update:**



# Gradient Descent(GD) Formulation

**Problem:**

$$\min_{x \in \mathbb{R}^d} f(x)$$

**Gradient Descent Update:**

$$x_{k+1} = x_k - \alpha_k \nabla f(x_k), \quad \alpha_k > 0 \text{ (step size).}$$



# Gradient Descent(GD) Formulation

**Problem:**

$$\min_{x \in \mathbb{R}^d} f(x)$$

**Gradient Descent Update:**

$$x_{k+1} = x_k - \alpha_k \nabla f(x_k), \quad \alpha_k > 0 \text{ (step size).}$$

- Choose  $\alpha_k$  appropriately (for example, through line search methods)





# Gradient Descent(GD) Formulation

## Problem:

$$\min_{x \in \mathbb{R}^d} f(x)$$

## Gradient Descent Update:

$$x_{k+1} = x_k - \alpha_k \nabla f(x_k), \quad \alpha_k > 0 \text{ (step size).}$$

- Choose  $\alpha_k$  appropriately (for example, through line search methods)

Now from next slide ,we'll see about GD convergence results in the following cases:

- Smooth functions



# Gradient Descent(GD) Formulation

## Problem:

$$\min_{x \in \mathbb{R}^d} f(x)$$

## Gradient Descent Update:

$$x_{k+1} = x_k - \alpha_k \nabla f(x_k), \quad \alpha_k > 0 \text{ (step size).}$$

- Choose  $\alpha_k$  appropriately (for example, through line search methods)

Now from next slide ,we'll see about GD convergence results in the following cases:

- Smooth functions
- Smooth + Convex functions



# Gradient Descent(GD) Formulation

## Problem:

$$\min_{x \in \mathbb{R}^d} f(x)$$

## Gradient Descent Update:

$$x_{k+1} = x_k - \alpha_k \nabla f(x_k), \quad \alpha_k > 0 \text{ (step size).}$$

- Choose  $\alpha_k$  appropriately (for example, through line search methods)

Now from next slide ,we'll see about GD convergence results in the following cases:

- Smooth functions
- Smooth + Convex functions
- Smooth + Strongly Convex functions



- [1] Benjamin Recht and Stephen J. Wright.  
*Optimization for Data Analysis*. 2022.
- [2] Stephen Boyd and Lieven Vandenberghe.  
*Convex Optimization*.  
Cambridge University Press, 2004.
- [3] Jorge Nocedal and Stephen J. Wright.  
*Numerical Optimization*, 2nd Edition.  
Springer, 2006.

**Thank You!**