

# Linear Algebra and Calculus refresher

★ Star 2,633

## General notations

### Definitions

**Vector** — We note  $x \in \mathbb{R}^n$  a vector with  $n$  entries, where  $x_i \in \mathbb{R}$  is the  $i^{th}$  entry:

$$x = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \in \mathbb{R}^n$$

**Matrix** — We note  $A \in \mathbb{R}^{m \times n}$  a matrix with  $n$  rows and  $m$ , where  $A_{i,j} \in \mathbb{R}$  is the entry located in the  $i^{th}$  row and  $j^{th}$  column:

$$A = \begin{pmatrix} A_{1,1} & \cdots & A_{1,n} \\ \vdots & & \vdots \\ A_{m,1} & \cdots & A_{m,n} \end{pmatrix} \in \mathbb{R}^{m \times n}$$

*Remark: the vector  $x$  defined above can be viewed as a  $n \times 1$  matrix and is more particularly called a column-vector.*

### Main matrices

**Identity matrix** — The identity matrix  $I \in \mathbb{R}^{n \times n}$  is a square matrix with ones in its diagonal and zero everywhere else:

$$I = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & 1 \end{pmatrix}$$

*Remark: for all matrices  $A \in \mathbb{R}^{n \times n}$ , we have  $A \times I = I \times A = A$ .*

**Diagonal matrix** — A diagonal matrix  $D \in \mathbb{R}^{n \times n}$  is a square matrix with nonzero values in its diagonal and zero everywhere else:

$$D = \begin{pmatrix} d_1 & 0 & \cdots & 0 \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & d_n \end{pmatrix}$$

*Remark: we also note  $D$  as  $\text{diag}(d_1, \dots, d_n)$ .*

## Matrix operations

---

### Multiplication

**Vector-vector** — There are two types of vector-vector products:

- inner product: for  $x, y \in \mathbb{R}^n$ , we have:

$$x^T y = \sum_{i=1}^n x_i y_i \in \mathbb{R}$$

- outer product: for  $x \in \mathbb{R}^m, y \in \mathbb{R}^n$ , we have:

$$xy^T = \begin{pmatrix} x_1 y_1 & \cdots & x_1 y_n \\ \vdots & & \vdots \\ x_m y_1 & \cdots & x_m y_n \end{pmatrix} \in \mathbb{R}^{m \times n}$$

**Matrix-vector** — The product of matrix  $A \in \mathbb{R}^{m \times n}$  and vector  $x \in \mathbb{R}^n$  is a vector of size  $\mathbb{R}^m$ , such that:

$$Ax = \begin{pmatrix} a_{r,1}^T x \\ \vdots \\ a_{r,m}^T x \end{pmatrix} = \sum_{i=1}^n a_{c,i} x_i \in \mathbb{R}^m$$

where  $a_{r,i}^T$  are the vector rows and  $a_{c,j}$  are the vector columns of  $A$ , and  $x_i$  are the entries of  $x$ .

**Matrix-matrix** — The product of matrices  $A \in \mathbb{R}^{m \times n}$  and  $B \in \mathbb{R}^{n \times p}$  is a matrix of size  $\mathbb{R}^{m \times p}$ , such that:

$$AB = \begin{pmatrix} a_{r,1}^T b_{c,1} & \cdots & a_{r,1}^T b_{c,p} \\ \vdots & & \vdots \\ a_{r,m}^T b_{c,1} & \cdots & a_{r,m}^T b_{c,p} \end{pmatrix} = \sum_{i=1}^n a_{c,i} b_{r,i}^T \in \mathbb{R}^{m \times p}$$

where  $a_{r,i}^T, b_{r,i}^T$  are the vector rows and  $a_{c,j}, b_{c,j}$  are the vector columns of  $A$  and  $B$  respectively.

## Other operations

**Transpose** — The transpose of a matrix  $A \in \mathbb{R}^{m \times n}$ , noted  $A^T$ , is such that its entries are flipped:

$$\forall i, j, \quad A_{i,j}^T = A_{j,i}$$

*Remark: for matrices  $A, B$ , we have  $(AB)^T = B^T A^T$ .*

**Inverse** — The inverse of an invertible square matrix  $A$  is noted  $A^{-1}$  and is the only matrix such that:

$$AA^{-1} = A^{-1}A = I$$

*Remark: not all square matrices are invertible. Also, for matrices  $A, B$ , we have  $(AB)^{-1} = B^{-1}A^{-1}$*

**Trace** — The trace of a square matrix  $A$ , noted  $\text{tr}(A)$ , is the sum of its diagonal entries:

$$\text{tr}(A) = \sum_{i=1}^n A_{i,i}$$

*Remark: for matrices  $A, B$ , we have  $\text{tr}(A^T) = \text{tr}(A)$  and  $\text{tr}(AB) = \text{tr}(BA)$*

**Determinant** — The determinant of a square matrix  $A \in \mathbb{R}^{n \times n}$ , noted  $|A|$  or  $\det(A)$  is expressed recursively in terms of  $A_{\setminus i, \setminus j}$ , which is the matrix  $A$  without its  $i^{\text{th}}$  row and  $j^{\text{th}}$  column, as follows:

$$\det(A) = |A| = \sum_{j=1}^n (-1)^{i+j} A_{i,j} |A_{\setminus i, \setminus j}|$$

*Remark:  $A$  is invertible if and only if  $|A| \neq 0$ . Also,  $|AB| = |A||B|$  and  $|A^T| = |A|$ .*

## Matrix properties

---

### Definitions

**Symmetric decomposition** — A given matrix  $A$  can be expressed in terms of its symmetric and antisymmetric parts as follows:

$$A = \underbrace{\frac{A + A^T}{2}}_{\text{Symmetric}} + \underbrace{\frac{A - A^T}{2}}_{\text{Antisymmetric}}$$

**Norm** — A norm is a function  $N : V \longrightarrow [0, +\infty[$  where  $V$  is a vector space, and such that for all  $x, y \in V$ , we have:

- $N(x + y) \leq N(x) + N(y)$
- $N(ax) = |a|N(x)$  for  $a$  scalar
- if  $N(x) = 0$ , then  $x = 0$

For  $x \in V$ , the most commonly used norms are summed up in the table below:

Norm	Notation	Definition	Use case
Manhattan, $L^1$	$\ x\ _1$	$\sum_{i=1}^n  x_i $	LASSO regularization
Euclidean, $L^2$	$\ x\ _2$	$\sqrt{\sum_{i=1}^n x_i^2}$	Ridge regularization
$p$ -norm, $L^p$	$\ x\ _p$	$\left(\sum_{i=1}^n x_i^p\right)^{\frac{1}{p}}$	Hölder inequality
Infinity, $L^\infty$	$\ x\ _\infty$	$\max_i  x_i $	Uniform convergence

**Linearly dependence** — A set of vectors is said to be linearly dependent if one of the vectors in the set can be defined as a linear combination of the others.

*Remark: if no vector can be written this way, then the vectors are said to be linearly independent.*

**Matrix rank** — The rank of a given matrix  $A$  is noted  $\text{rank}(A)$  and is the dimension of the vector space generated by its columns. This is equivalent to the maximum number of linearly independent columns of  $A$ .

**Positive semi-definite matrix** — A matrix  $A \in \mathbb{R}^{n \times n}$  is positive semi-definite (PSD) and is noted  $A \succeq 0$  if we have:

$$\boxed{A = A^T} \quad \text{and} \quad \boxed{\forall x \in \mathbb{R}^n, \quad x^T A x \geq 0}$$

*Remark: similarly, a matrix  $A$  is said to be positive definite, and is noted  $A \succ 0$ , if it is a PSD matrix which satisfies for all non-zero vector  $x$ ,  $x^T A x > 0$ .*

**Eigenvalue, eigenvector** — Given a matrix  $A \in \mathbb{R}^{n \times n}$ ,  $\lambda$  is said to be an eigenvalue of  $A$  if there exists a vector  $z \in \mathbb{R}^n \setminus \{0\}$ , called eigenvector, such that we have:

$$\boxed{Az = \lambda z}$$

**Spectral theorem** — Let  $A \in \mathbb{R}^{n \times n}$ . If  $A$  is symmetric, then  $A$  is diagonalizable by a real orthogonal matrix  $U \in \mathbb{R}^{n \times n}$ . By noting  $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$ , we have:

$$\boxed{\exists \Lambda \text{ diagonal, } A = U\Lambda U^T}$$

**Singular-value decomposition** — For a given matrix  $A$  of dimensions  $m \times n$ , the singular-value decomposition (SVD) is a factorization technique that guarantees the existence of  $U$   $m \times m$  unitary,  $\Sigma$   $m \times n$  diagonal and  $V$   $n \times n$  unitary matrices, such that:

$$\boxed{A = U\Sigma V^T}$$

## Matrix calculus

---

**Gradient** — Let  $f : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$  be a function and  $A \in \mathbb{R}^{m \times n}$  be a matrix. The gradient of  $f$  with respect to  $A$  is a  $m \times n$  matrix, noted  $\nabla_A f(A)$ , such that:

$$\boxed{\left(\nabla_A f(A)\right)_{i,j} = \frac{\partial f(A)}{\partial A_{i,j}}}$$

*Remark: the gradient of  $f$  is only defined when  $f$  is a function that returns a scalar.*

**Hessian** — Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function and  $x \in \mathbb{R}^n$  be a vector. The hessian of  $f$  with respect to  $x$  is a  $n \times n$  symmetric matrix, noted  $\nabla_x^2 f(x)$ , such that:

$$\boxed{\left(\nabla_x^2 f(x)\right)_{i,j} = \frac{\partial^2 f(x)}{\partial x_i \partial x_j}}$$

*Remark: the hessian of  $f$  is only defined when  $f$  is a function that returns a scalar.*

**Gradient operations** — For matrices  $A, B, C$ , the following gradient properties are worth having in mind:

$$\nabla_A \text{tr}(AB) = B^T$$

$$\nabla_{A^T} f(A) = (\nabla_A f(A))^T$$

$$\nabla_A \text{tr}(ABA^T C) = CAB + C^T AB^T$$

$$\nabla_A |A| = |A|(A^{-1})^T$$



(<https://twitter.com/shervinea>)



(<https://linkedin.com/in/shervineamidi>)



(<https://github.com/shervinea>)



(<https://scholar.google.com/citations?user=nMnMTm8AAAAJ>)

