CLASS 9 · PART TEST-1 · STUDENT PERFORMANCE REPORT

Masarapu Kartikeya

Pragya G9 (EXT) Achievers · Adm No **246416594**

Campus HYD/Nallakunta DD Colony
JEE Mains 08 Oct 2026
JEE Advanced 09 Oct 2026

JEE Mains

08 Oct 2026 · 74 questions

207 / 296 — **69.9%**

31	22	4/79	4
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **194.8 / 300** (65.0%) → this test **69.9%** ▲ **5.0**

89 marks lost: 45 on 9 wrong answers (incl. 9 negative) · 44 on 11 skipped

JEE Advanced

09 Oct 2026 · 50 questions

70 / 176 — **39.8%**

230	165	19/78	30†
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **97.9 / 180** (54.4%) → this test **39.8%** ▼ **14.6**

106 marks lost: 89 on 23 wrong answers (incl. 7 negative) · 11 on 3 skipped · 6 on partly-correct answers

1 Subject scorecard

Your marks in each subject, against the 79 students of this program on campus and against your own weekly-test average.

SUBJECT	MARKS	SCORE VS CAMPUS	SUBJECT AIR	CAMPUS AVG / BEST	WEEKLY AVG	CHANGE	RIGHT / WRONG / SKIP
JEE Mains 207 / 296 · 69.9%							
Maths	**80**/100	80%	**44**	43.1 / 95	**76.0**/100 76.0%	▲ **4.0**	21 / 4 / 0
Physics	**52**/100	52%	**120**	21.9 / 90	**64.2**/100 64.2%	▼ **12.2**	14 / 4 / 7
Chemistry	**75**/96	78%	**18**	41.4 / 81	**54.5**/100 54.5%	▲ **23.7**	19 / 1 / 4
JEE Advanced 70 / 176 · 39.8%							
Maths	**27**/60	45%	**389**	24.5 / 60	**35.0**/60 58.4%	▼ **13.4**	9 / 7 / 1
Physics	**14**/60	23%	**423**	12.3 / 56	**32.7**/60 54.5%	▼ **31.2**	5 / 10 / 2
Chemistry	**29**/56	52%	**93**	15.8 / 37	**30.3**/60 50.5%	▲ **1.3**	10 / 6 / 0

Bar = your score % | campus average | best on campus Change = this test % − weekly-test average %

What this report says

1. **Maths** is your strongest JEE Mains subject: 80/100 (80%), All-India subject rank 44.
2. **Physics** needs the most work in JEE Mains: 52/100 (52%), subject rank 120; you lost 11 of its 25 questions.
3. **Chemistry** is your strongest JEE Advanced subject: 29/56 (52%), All-India subject rank 93.
4. **Physics** needs the most work in JEE Advanced: 14/60 (23%), subject rank 423; you lost 12 of its 17 questions.
5. Biggest fall against your weekly-test average: **JEE Advanced Physics** – 23% here against 54% on average.
6. Biggest improvement: **JEE Mains Chemistry** – 78% here against 54% on average.
7. **1 question** you lost were answered correctly by 70%+ of the batch's top 100 – about **6 marks** within easy reach (Advanced Q5).
8. Negative marking cost you **9 in JEE Mains and 7 in JEE Advanced**.
9. You left **11 in JEE Mains and 3 in JEE Advanced** questions blank, and answered 9 in JEE Mains and 23 in JEE Advanced wrong – both are marks to recover.

JEE Mains – question by question

207 / 296 · 69.9%
08 Oct 2026

2 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	2	3	4	5	6	**7**	8	9	10	11	**12**	13	14	15	16	17	18	19	20	21	22	23	**24**	**25**
Physics	**26**	27	28	29	**30**	**31**	**32**	33	34	35	36	37	**38**	39	40	**41**	**42**	**43**	44	**45**	46	**47**	48	49	**50**
Chemistry	51	52	53	54	55	56	57	~~58~~	59	60	61	62	**63**	64	65	66	**67**	68	69	70	**71**	**72**	73	74	**75**

3 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 79 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40–69% · **Tough one** below 40%

Maths

4 of 25 questions lost · **20** marks · ● 2 within reach · ● 2 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q25	Within reach	**Trigonometric Equations**	Numerical · +4 / −1	**Wrong**	5	**67%**	57%
Q12	Within reach	**Inverse Trigonometric Functions**	Single correct · +4 / −1	**Wrong**	5	**64%**	52%
Q7	Tough one	**Trigonometric Equations** – $\sqrt{3}\sin \pi x + \cos \pi x$ = quadratic – number of solutions	Single correct · +4 / −1	**Wrong**	5	**37%**	16%
Q24	Tough one	**Functions** – Even function with GIF, fractional part and signum	Numerical · +4 / −1	**Wrong**	5	**20%**	16%

Physics

11 of 25 questions lost · **48** marks · ● 3 within reach · ● 8 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q50	Within reach	**Momentum & Collisions**	Numerical · +4 / −1	**Wrong**	5	**60%**	41%
Q41	Within reach	**Laws of Motion**	Single correct · +4 / −1	**Skipped**	4	**50%**	39%
Q30	Within reach	**Circular Motion**	Single correct · +4 / −1	**Skipped**	4	**43%**	27%
Q47	Tough one	**Laws of Motion**	Numerical · +4 / −1	**Skipped**	4	**36%**	24%
Q32	Tough one	**Circular Motion** – Blocks on a rotating frame – ω when sliding starts	Single correct · +4 / −1	**Wrong**	5	**33%**	28%
Q31	Tough one	**Work, Energy & Power** – KE vs height for a stone thrown up from a cliff	Single correct · +4 / −1	**Wrong**	5	**32%**	10%
Q42	Tough one	**Work, Energy & Power** – Change in PE of a ball thrown from a tower	Single correct · +4 / −1	**Skipped**	4	**32%**	15%
Q26	Tough one	**Momentum & Collisions** – KE loss in head-on collision from relative velocities	Single correct · +4 / −1	**Skipped**	4	**31%**	22%
Q38	Tough one	**Work, Energy & Power** – Work along different paths; conservative or not	Single correct · +4 / −1	**Skipped**	4	**25%**	13%
Q43	Tough one	**Laws of Motion** – Tension in a spring pulled by 5 N at each end	Single correct · +4 / −1	**Skipped**	4	**24%**	10%
Q45	Tough one	**Work, Energy & Power** – Speed on hitting the ground – energy conservation	Single correct · +4 / −1	**Wrong**	5	**18%**	6%

Chemistry

5 of 24 questions lost · 21 marks · ● 5 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q72	Tough one	**Hydrocarbons**	Numerical · +4 / −1	Skipped	4	28%	23%
Q67	Tough one	**General Organic Chemistry** – Structure and hybridisation of the methyl carbocation	Single correct · +4 / −1	Skipped	4	26%	28%
Q63	Tough one	**Hydrocarbons** – Monochlorination products incl. stereoisomers	Single correct · +4 / −1	Wrong	5	16%	20%
Q75	Tough one	**General Organic Chemistry** – Bases stronger than aniline	Numerical · +4 / −1	Skipped	4	11%	9%
Q71	Tough one	**Isomerism** – Count of optically active compounds	Numerical · +4 / −1	Skipped	4	6%	10%

4 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Trigonometric Equations	7	18/28		44%	Q7, Q25
Functions	7	23/28		35%	Q24
Inverse Trigonometric Functions	10	35/40		48%	Q12
Sequences & Series	1	4/4		45%	– none
Physics					
Work, Energy & Power	5	2/20		9%	Q31, Q38, Q42, Q45
Momentum & Collisions	3	3/12		25%	Q26, Q50
Laws of Motion	8	20/32		23%	Q41, Q43, Q47
Circular Motion	7	19/28		30%	Q30, Q32
Friction	2	8/8		17%	– none
Chemistry					
Hydrocarbons	4	7/16		42%	Q63, Q72
General Organic Chemistry	10	32/40		32%	Q67, Q75
Isomerism	10	36/40		55%	Q71

Bar = your % of the chapter's marks | campus average

5 JEE Mains · Maths – questions to revisit

4 questions · easy misses first
cropped from the question paper

Q25 — Trigonometric Equations

JEE Mains · Maths · Numerical · marking +4 / −1 · marks lost 5
Top 100: **67%** got it right · Campus: **57%** (45 of 79)

Within reach
You: Wrong

25. Thanos has created a destruction field along $y = \sin x$. Thor is running along $y = \frac{x}{10}$.

To protect him (Thor) Doctor Strange creates a portal at every point where thor's path meets the destruction field. How many portals must be created?

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q12 — Inverse Trigonometric Functions

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5
Top 100: **64%** got it right · Campus: **52%** (41 of 79)

Within reach
You: Wrong

12. Let 'S' be the complete set of solutions of the inequation $\left(\sec^{-1}x\right)^2 - 5\left(\sec^{-1}x\right) + 6 > 0$

Which of the following intervals is not contained in the solution set?

A) $(\sec 2, \sec 3)$ B) $(-\infty, \sec 2)$ C) $(\sec 3, -1]$ D) $[1, \infty)$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q7 — Trigonometric Equations – √3 sin πx + cos πx = quadratic – number of solutions

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5
Top 100: **37%** got it right · Campus: **16%** (13 of 79)

Tough one
You: Wrong

7. Number of solutions of $\sqrt{3}\sin(\pi x) + \cos(\pi x) = x^2 - \frac{2}{3}x + \frac{19}{9}$ is .....

A) 0 B) 1 C) 2 D) 3

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q24

Functions – Even function with GIF, fractional part and signum

JEE Mains · Maths · Numerical · marking +4 / −1 · marks lost 5

Top 100: **20%** got it right · Campus: **16%** (13 of 79)

Tough one

You: Wrong

24. Let $f(x)=\left([a]^2-5[a]+4\right)x^3-\left(6\{a\}^2-5\{a\}+1\right)x-(\tan x)(\operatorname{sgn}(x))$ (where [.] = GIF, $\{.\}=F.P.F, \operatorname{sgn}(x)=signum(x)$). If $f(x)$ is an even function of 'x', then the sum of possible values of 'a' is .....?

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Mains · Physics – questions to revisit

11 questions · easy misses first
cropped from the question paper

Q50 — Momentum & Collisions

JEE Mains · Physics · Numerical · marking +4 / −1 · marks lost 5
Top 100: **60%** got it right · Campus: **41%** (32 of 79)

Within reach
You: Wrong

50. When kinetic energy of a body becomes 64 times of its original value, the percentage increase in momentum of body will be ……%

Why did I lose this question? (write the reason and the correct approach)


☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q41 — Laws of Motion

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4
Top 100: **50%** got it right · Campus: **39%** (31 of 79)

Within reach
You: Skipped

41. For the arrangement of pulley shown in fig, the effort (P) required to raise the given load (w). assume efficiency of the system as 80%



A) 200N B) 250N C) 500N D) 1000N

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q30

Circular Motion

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4

Top 100: **43%** got it right · Campus: **27%** (21 of 79)

Within reach

You: Skipped

30. When a bucket containing water is rotated fast in a vertical circle of radius R, the water in the bucket doesn't spill provided that bucket is

A) whirled around with maximum speed of $\sqrt{2gR}$

B) whirled around with minimum speed of $\sqrt{\frac{gR}{2}}$

C) having a rpm of $n=\sqrt{900g/\pi^2R}$

D) having a rpm of $n=\sqrt{3600g/\pi^2R}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q47

Laws of Motion

JEE Mains · Physics · Numerical · marking +4 / −1 · marks lost 4

Top 100: **36%** got it right · Campus: **24%** (19 of 79)

Tough one

You: Skipped

47. A body with mass 5kg is acted upon by a force $\vec{F}=\left(-3\hat{i}+4\hat{j}\right)N$. If its initial velocity at $t=0\text{sec}$ is $u=\left(6\hat{i}-12\hat{j}\right)m/s$. The time at which it will just have a velocity along y-axis .....sec

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q32 **Circular Motion – Blocks on a rotating frame – ω when sliding starts**


JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **33%** got it right · Campus: **28%** (22 of 79)

Tough one

You: Wrong

32. Two blocks of masses m and M are connected by a chord passing around a frictionless pulley which is attached to a rotating frame, which rotates about a vertical axis with angular velocity ω. If co-efficient of friction between two masses and the surface be μ_1 and μ_2 respectively. The value of ω at which block starts sliding radially $(M > m)$



A) $\omega = \left[g \left[\frac{\mu_1 M + \mu_2 m}{m r_1 - M r_1} \right] \right]^{1/2}$

B) $\omega = \left[g \left[\frac{\mu_1 m + \mu_2 M}{M r_2 - m r_1} \right] \right]^{1/2}$

C) $\omega = \left[g \left[\frac{\mu_1 m - \mu_2 M}{M r_2 - m r_1} \right] \right]^{1/2}$

D) $\omega = \left[g \left[\frac{\mu_1 M - \mu_2 m}{M r_2 - m r_1} \right] \right]^{1/2}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q31

Work, Energy & Power – KE vs height for a stone thrown up from a cliff

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5


Top 100: **32%** got it right · Campus: **10%** (8 of 79)

Tough one


You: Wrong

31. A stone of mass 50g is thrown vertically upwards with a velocity of 12m/s from edge of a cliff 44m high. Identify kinetic energy of stone versus height from bottom of cliff $\left(g = 10m/s^2\right)$ (ignore air resistance)


A)




B)



C)



D)



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q42

Work, Energy & Power – Change in PE of a ball thrown from a tower

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4

Top 100: **32%** got it right · Campus: **15%** (12 of 79)

Tough one

You: Skipped

42. At t =0 sec a 1kg ball is thrown from top to a tall tower with velocity $\vec{V} = \left(8\hat{i} + 9\hat{j}\right)$ m/s. The change in potential energy of ball – earth system between $t = 0$ *to* $t = 5$sec is

$\left(g = 10m/s^2\right)$

A) 800J B) 1536J C) 2100J D) -800J

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q26 **Momentum & Collisions** – **KE loss in head-on collision from relative velocities**
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4
Top 100: **31%** got it right · Campus: **22%** (17 of 79)

Tough one
You: Skipped


26. Two bodies of masses 3 kg and 2 kg collide head on. Their relative velocities before and after collision are 15m/s and 5 m/s respectively. The loss of kinetic energy of system is
A) 120J B) 100J C) 80J D) 240J

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q38 **Work, Energy & Power** – **Work along different paths; conservative or not**
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4
Top 100: **25%** got it right · Campus: **13%** (10 of 79)

Tough one
You: Skipped

38. A particle is moved along the different paths OAC, OBC and ODC as shown in the fig. path ODC is parabola, $y = 4x^2$. The work done by a force $\vec{F} = xy\hat{i} + x^2 y\hat{j}$ on the particle along



Column – I	**Column – II**
a)Path OAC	P) 2J
b) Path OBC	Q) $\frac{19}{3}$J
c) Path ODC	R) Conservative
d) Force is	S) 8J
	T) Non- conservative

A) $a \to Q, b \to S, c \to P, d \to T$ B) $a \to S, b \to P, c \to Q, d \to T$

C) $a \to S, b \to Q, c \to P, d \to T$ D) $a \to S, b \to Q, c \to S, d \to T$

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q43

Laws of Motion – Tension in a spring pulled by 5 N at each end

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4

Top 100: **24%** got it right · Campus: **10%** (8 of 79)

Tough one

You: Skipped

43. The tension in the massless spring is



A) zero B) 10N C) 2.5N D) 5N

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q45

Work, Energy & Power – Speed on hitting the ground – energy conservation

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **18%** got it right · Campus: **6%** (5 of 79)

Tough one

You: Wrong

45. A cricket bowler releases the ball in two different ways

i) giving it only horizontal velocity and

ii) giving it horizontal velocity and small downward velocity

The speeds at the time release is same. Both are released at a height H from ground then which one will have greater speed when ball hits the ground?(Neglect air resistance)

A) Ist way B) IInd way C) same in two ways D) None of the above

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Mains · Chemistry — questions to revisit


5 questions · easy misses first
cropped from the question paper

Q72

Hydrocarbons
JEE Mains · Chemistry · Numerical · marking +4 / −1 · marks lost 4
Top 100: **28%** got it right · Campus: **23%** (18 of 79)

Tough one
You: Skipped

72. How many products will be formed for given reaction, including stereo isomers



Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q67

General Organic Chemistry — **Structure and hybridisation of the methyl carbocation**
JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 4
Top 100: **26%** got it right · Campus: **28%** (22 of 79)

Tough one
You: Skipped

67. Wrong statement regarding methyl carbonium ion $\overset{\oplus}{C}H_3$

A) It is sp^2 hybridized

B) Vacant orbital is sp^2 hybridized

C) vacant orbital is perpendicular to molecular planar and is pure p-orbital

D) It is electrophilic with sextet of electron

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q63

Hydrocarbons – **Monochlorination products incl. stereoisomers**


JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5

Top 100: **16%** got it right · Campus: **20%** (16 of 79)

Tough one

You: Wrong

63. Number of monochlorinated product when following compound undergo reaction with $Cl_2 / h\nu$ (including stereo isomers)



A) 5 B) 9 C) 10 D) 6

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q75

General Organic Chemistry – **Bases stronger than aniline**








JEE Mains · Chemistry · Numerical · marking +4 / −1 · marks lost 4

Top 100: **11%** got it right · Campus: **9%** (7 of 79)

Tough one

You: Skipped

75. How many of the following are more basic then Aniline?

NH_3, $PhCH_2NH_2$,  ,  ,  ,  ,  ,  , $CH_3—CH_2—NH_2$, 

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q71

Isomerism – Count of optically active compounds


JEE Mains · Chemistry · Numerical · marking +4 / −1 · marks lost 4

Top 100: **6%** got it right · Campus: **10%** (8 of 79)

Tough one

You: Skipped

71. Total no of compounds which are optically active are _____



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced – question by question

70 / 176 · 39.8%
09 Oct 2026

6 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	2	**3**	4	**5**	6	**7**	**8**	**9**	10	11	**12**	**13**	14	15	16	**17**
Physics	**18**	**19**	**20**	21	22	**23**	**24**	25	**26**	**27**	28	**29**	**30**	31	**32**	**33**	**34**
Chemistry	35	36	37	38	39	40	41	42	**43**	44	~~45~~	**46**	**47**	48	**49**	**50**	**51**

7 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 78 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40–69% · **Tough one** below 40%

Maths

8 of 17 questions lost · **33** marks · ● 1 easy miss · ● 4 within reach · ● 3 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q5	Easy miss	**Inverse Trigonometric Functions**	Multiple correct · +4 / −2	**Wrong**	6	**74%**	74%
Q12	Within reach	**Functions**	Integer · +4 / 0	**Wrong**	4	**55%**	40%
Q9	Within reach	**Inverse Trigonometric Functions**	Integer · +4 / 0	**Wrong**	4	**54%**	58%
Q13	Within reach	**Inverse Trigonometric Functions**	Integer · +4 / 0	**Wrong**	4	**54%**	45%
Q7	Within reach	**Functions**	Multiple correct · +4 / −2	**Skipped**	4	**49%**	44%
Q3	Tough one	**Trigonometric Equations**	Single correct · +3 / −1	**Wrong**	4	**36%**	46%
Q8	Tough one	**Functions** – Period from two lines of symmetry	Integer · +4 / 0	**Wrong**	4	**12%**	13%
Q17	Tough one	**Trigonometric Equations** – Sum of solutions of $\cos(n\sqrt{(x-100)})\cdot\cos(n\sqrt{x}) = 1$	Paragraph · +3 / 0	**Wrong**	3	**6%**	6%

Physics

12 of 17 questions lost · **46** marks · ● 2 within reach · ● 10 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q23	Within reach	**Laws of Motion**	Multiple correct · +4 / −2	**Skipped**	4	56%	51%
Q19	Within reach	**Circular Motion**	Single correct · +3 / −1	**Skipped**	3	40%	38%
Q24	Tough one	**Circular Motion**	Multiple correct · +4 / −2	**Wrong**	6	34%	27%
Q20	Tough one	**Work, Energy & Power** – Maximum spring extension – energy method	Single correct · +3 / −1	**Wrong**	4	29%	27%
Q26	Tough one	**Circular Motion**	Integer · +4 / 0	**Wrong**	4	24%	17%
Q29	Tough one	**Circular Motion** – Net acceleration in a vertical circle at 37°	Integer · +4 / 0	**Wrong**	4	19%	14%
Q30	Tough one	**Friction** – Work by friction seen from an accelerating observer	Integer · +4 / 0	**Wrong**	4	19%	12%
Q18	Tough one	**Friction** – Static vs kinetic friction under a time-varying force	Single correct · +3 / −1	**Wrong**	4	16%	9%
Q34	Tough one	**Work, Energy & Power** – Work by conservative forces from the U–x graph	Paragraph · +3 / 0	**Wrong**	3	12%	9%
Q27	Tough one	**Friction** – Net work done by friction between sliding blocks	Integer · +4 / 0	**Wrong**	4	8%	10%
Q32	Tough one	**Friction** – Total force exerted by a block on an accelerating cube	Paragraph · +3 / 0	**Wrong**	3	8%	6%
Q33	Tough one	**Work, Energy & Power** – Work-energy theorem from a–t and U–x graphs	Paragraph · +3 / 0	**Wrong**	3	2%	1%






Chemistry

6 of 16 questions lost · **21** marks (+6 on partly-correct answers) · ● 2 within reach · ● 4 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q49	Within reach	**General Organic Chemistry**	Paragraph · +3 / 0	**Wrong**	3	53%	46%
Q43	Within reach	**Hydrocarbons** – Alkane from O_2 needed; count secondary carbons	Integer · +4 / 0	**Wrong**	4	47%	28%
Q46	Tough one	**Isomerism** – Count of optically active compounds	Integer · +4 / 0	**Wrong**	4	15%	15%
Q51	Tough one	**Hydrocarbons** – Chiral products of free-radical monobromination	Paragraph · +3 / 0	**Wrong**	3	15%	4%
Q47	Tough one	**Hydrocarbons** – Dichlorination isomers of C_5H_{10} (VD 35)	Integer · +4 / 0	**Wrong**	4	7%	5%
Q50	Tough one	**Hydrocarbons** – Preparation of alkanes – count correct statements	Paragraph · +3 / 0	**Wrong**	3	4%	3%

8 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject. Chapter marks are approximate (≈): partly-correct answers cannot be traced to a question.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Inverse Trigonometric Functions	6	≈ **7**/21		55%	**Q5**, **Q9**, **Q13**
Functions	6	≈ **10**/22		40%	**Q7**, **Q8**, **Q12**
Trigonometric Equations	5	≈ **10**/17		35%	**Q3**, **Q17**
Physics					
Circular Motion	4	≈ **-2**/15		17%	**Q19**, **Q24**, **Q26**, **Q29**
Friction	5	≈ **2**/17		12%	**Q18**, **Q27**, **Q30**, **Q32**
Work, Energy & Power	4	≈ **2**/12		22%	**Q20**, **Q33**, **Q34**
Laws of Motion	3	≈ **8**/12		39%	**Q23**
Momentum & Collisions	1	≈ **4**/4		33%	– none
Chemistry					
Hydrocarbons	5	≈ **4**/18		22%	**Q43**, **Q47**, **Q50**, **Q51**
Isomerism	2	≈ **4**/8		16%	**Q46**
General Organic Chemistry	9	≈ **27**/30		43%	**Q49**

Bar = your % of the chapter's marks | campus average

9 JEE Advanced · Maths – questions to revisit

8 questions · easy misses first
cropped from the question paper

Q5 Inverse Trigonometric Functions

JEE Advanced · Maths · Multiple correct · marking +4 / −2 · marks lost 6
Top 100: **74%** got it right · Campus: **74%** (58 of 78)

Easy miss
You: Wrong

5. Consider the equation : $\sin^{-1}(x)+\cos^{-1}(x)+\tan^{-1}(x)=k$. For which of the following values of 'k' does the equation have at least one solution?

A) 0.1 B) 1 C) 2 D) 3

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q12 Functions

JEE Advanced · Maths · Integer · marking +4 / 0 · marks lost 4
Top 100: **55%** got it right · Campus: **40%** (31 of 78)

Within reach
You: Wrong

12. Consider the function $f(x)=\log_{([x-1])}\left(\frac{|x|}{x}\right),\ ([.]=GIF)$

Let 'n' be the smallest positive integer in the domain of f. Let 'S' be the sum of all integers in the range of f. Then $n+S=$.....

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q9 Inverse Trigonometric Functions

JEE Advanced · Maths · Integer · marking +4 / 0 · marks lost 4
Top 100: **54%** got it right · Campus: **58%** (45 of 78)

Within reach
You: Wrong

9. $\sum_{n=1}^{5}\sin^{-1}\left(\sin\left(2n-1\right)\right)=$.....

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q13 Inverse Trigonometric Functions

JEE Advanced · Maths · Integer · marking +4 / 0 · marks lost 4
Top 100: **54%** got it right · Campus: **45%** (35 of 78)

Within reach
You: Wrong

13. If $\cos^{-1}(x_1)+\cos^{-1}(x_2)+\cos^{-1}(x_3)+.....+\cos^{-1}(x_{2026})=0$, then $x_2+x_4+x_6+.....+x_{2026}=.....$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q7 Functions

JEE Advanced · Maths · Multiple correct · marking +4 / −2 · marks lost 4
Top 100: **49%** got it right · Campus: **44%** (34 of 78)

Within reach
You: Skipped

7. Let $f:N\to N$ be such that $f(1)=1$ and $f(n+1)=2f(n)+1$. Then $\forall n\in N$, there exists a geometric progression of integral terms such that :

A) $f(n)$ is its nth term

B) $(f(n)+1)$ is its nth term

C) $f(n)$ is the sum of its first 'n' terms

D) $(f(n)+1)$ is the sum of its first 'n' terms

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q3 Trigonometric Equations

JEE Advanced · Maths · Single correct · marking +3 / −1 · marks lost 4
Top 100: **36%** got it right · Campus: **46%** (36 of 78)

Tough one
You: Wrong

3. If the equation $1+\sin^2(ax)=\cos x$ has a unique solution in 'x' then 'a' must be

A) rational B) irrational C) integer D) whole number

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q8

Functions – Period from two lines of symmetry

JEE Advanced · Maths · Integer · marking +4 / 0 · marks lost 4

Top 100: **12%** got it right · Campus: **13%** (10 of 78)

Tough one

You: Wrong

8. Let $a \in R$. If $f(a-x) = f(a+x) \forall x \in R$, then $x = a$ is a line of symmetry in the graph of $y = f(x)$. Consider a non-constant function $g(x)$ such that $g(1+x) = g(1-x) \forall x \in R$ and $g(2026+x) = g(2026-x) \forall x \in R$. Then all such $g(x)$ must be periodic. The smallest positive number that could be a period for all such $g(x)$ is .....

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q17

Trigonometric Equations – Sum of solutions of cos(π√(x−100))·cos(π√x) = 1

JEE Advanced · Maths · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **6%** got it right · Campus: **6%** (5 of 78)

Tough one

You: Wrong

PARAGRAPH-II

Given positive integer 'k' , if we want to find positive integer 'n' and 'm' such that $n^2 - m^2 = k$ then we use $(n+m)(n-m) = k$ where we equate $n+m$ & $n-m$ to factors of k to find all possible values of (n,m).

Eg: $n^2 - m^2 = 6 \Rightarrow (n+m)(n-m) = (\pm 6)(\pm 1)$ or $(\pm 3)(\pm 2)$

17. The sum of real solution(s) of $\cos\left(\pi\sqrt{x-100}\right)\cos\left(\pi\sqrt{x}\right) = 1$ is .....

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced · Physics – questions to revisit

12 questions · easy misses first
cropped from the question paper

Q23 Laws of Motion

JEE Advanced · Physics · Multiple correct · marking +4 / −2 · marks lost 4
Top 100: **56%** got it right · Campus: **51%** (40 of 78)

Within reach
You: Skipped

23. An object of mass m is at rest in equilibrium at the origin. At $t = 0$sec a force $\vec{F}(t)$ is applied which has the components $F_x(y) = k_1 + k_2 y$ $F_y(t) = k_3 t$ where k_1, k_2, k_3 are constants . Then

A) X-component of velocity at any time 't' is $V_x = \frac{k_1 t}{m} + \frac{k_2 k_3}{24 m^2} t^4$

B) X-component of acceleration at anytime 't' is $a_x = \left(\frac{k_1}{m} + \frac{k_2 k_3}{6 m^2} t^3 \right)$

C) Y- coordinate of object at any time 't' is $\frac{k_3 t^2}{3m}$

D) Y- coordinate of object at any time 't' is $\frac{k_3 t^3}{6m}$

Why did I lose this question? (write the reason and the correct approach)


☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q19 Circular Motion

JEE Advanced · Physics · Single correct · marking +3 / −1 · marks lost 3
Top 100: **40%** got it right · Campus: **38%** (30 of 78)

Within reach
You: Skipped

19. If a particle starts from 'A' moving along the curved semi circular path as shown in figure with tangential acceleration 'a' then acceleration at B in magnitude is



A) $2a\left[\sqrt{1+\pi^2}\right]$ B) $a\left[\sqrt{1+\pi^2}\right]$ C) $a\left[\sqrt{\pi^2-1}\right]$ D) $a\pi\left[\sqrt{1+\pi^2}\right]$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q24

Circular Motion

JEE Advanced · Physics · Multiple correct · marking +4 / −2 · marks lost 6

Top 100: **34%** got it right · Campus: **27%** (21 of 78)

Tough one

You: Wrong

24. A particle is revolving in a circle of radius R with initial speed u. it starts retarding with constant retardation $\frac{u^2}{4\pi R}$. The number of revolutions it makes in a time $\frac{8\pi R}{u}$ is

A) 3 B) 2 C) 4 D) 6

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q20

Work, Energy & Power – Maximum spring extension – energy method


JEE Advanced · Physics · Single correct · marking +3 / −1 · marks lost 4

Top 100: **29%** got it right · Campus: **27%** (21 of 78)

Tough one

You: Wrong

20. All the surfaces are smooth system is released from rest with spring unstretched. Then maximum extension in spring will be $(m_A = m; m_B = 2m)$



A) $\frac{mg}{K}$ B) $\frac{2mg}{K}$ C) $\frac{3mg}{K}$ D) $\frac{4mg}{K}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q26 — Circular Motion

JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **24%** got it right · Campus: **17%** (13 of 78)

Tough one

You: Wrong

26. A particle of mass 2.5kg is moving in gravity free space with velocity $3\hat{i}m/s$. At $t=0\text{sec}$ a force of magnitude 67.5N starts acting on the particle such that it is always perpendicular to instantaneous velocity. The minimum time after which particle has same velocity as the initial is $\frac{2\pi}{x}$ sec. The value of x ……..

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q29 — Circular Motion – Net acceleration in a vertical circle at 37°

JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **19%** got it right · Campus: **14%** (11 of 78)

Tough one

You: Wrong

29. A small ball of mass m is attached to the end of string of length $l=1m$ whose other end is fixed. From its lowest position, the ball is given a kinetic energy $\frac{mgl}{5}$. Net acceleration $\left(in\ m/s^2\right)$ of ball at the instant when string makes an angle $\theta=37^{o}$ with vertical is ……$\left(g=10m/s^2\right)$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q30

Friction – Work by friction seen from an accelerating observer


JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **19%** got it right · Campus: **12%** (9 of 78)

Tough one

You: Wrong

30. An observer and a vehicle, both starts moving together from rest with acceleration $5m/s^2$ and $2m/s^2$ respectively. There is a 2 kg block on the floor of the vehicle and $\mu = 0.3$ between their surfaces. The work done by the frictional force on 2kg block as observed by running observer , during first 2seconds of motion is $\underline{-x}$ (J) $\left(g = 10m/s^2\right)$ find 'x'



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q18

Friction – Static vs kinetic friction under a time-varying force


JEE Advanced · Physics · Single correct · marking +3 / −1 · marks lost 4

Top 100: **16%** got it right · Campus: **9%** (7 of 78)

Tough one

You: Wrong

18. The force acting on the block of mass 1 kg is given by $F = 5 - 2t$. The frictional force acting on the block after time $t = 2\text{sec}$ will be $\left(\mu = 0.2, g = 10m/s^2\right)$



A) 2N B) 3N C) 1N D) zero

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q34

Work, Energy & Power – **Work by conservative forces from the U-x graph**

JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **12%** got it right · Campus: **9%** (7 of 78)


Tough one

You: Wrong

PARAGRAPH-II

A particle of mass 1kg is moving along x-axis its velocity is 6m/s at x=0. Acceleration versus time and P.E versus displacement of particle are shown $\left(g = 10 m/s^2\right)$

Acceleration a is in $m/\sec^2$ b



34. Work done by conservative forces _____ J

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q27

Friction – **Net work done by friction between sliding blocks**


JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **8%** got it right · Campus: **10%** (8 of 78)

Tough one

You: Wrong

27. Block A of mass 1kg is placed on rough surface of block B of mass 3kg. Block B is placed on smooth horizontal surface. Blocks are given velocities as shown. If net workdone by frictional force is "-n" J. value of n is ……..



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q32

Friction – Total force exerted by a block on an accelerating cube

JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3


Top 100: **8%** got it right · Campus: **6%** (5 of 78)

Tough one

You: Wrong

PARAGRAPH-I

A solid block of mass 2 kg is resting inside a cube as shown in figure. The cube is moving with a velocity $\vec{V} = \left(5t\hat{i} + 2t\hat{j}\right) m/s$. Here 't' is second. The block is in rest w.r.t cube and $\mu = 0.5$ between block and surface of cube. $\left(g = 10 m/s^2\right)$



32. Total force exerted by block on cube is _____ N

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q33

Work, Energy & Power – Work-energy theorem from a-t and U-x graphs

JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **2%** got it right · Campus: **1%** (1 of 78)


Tough one

You: Wrong

PARAGRAPH-II

A particle of mass 1kg is moving along x-axis its velocity is 6m/s at x=0. Acceleration versus time and P.E versus displacement of particle are shown $\left(g = 10 m/s^2\right)$

Acceleration a is in $m/\sec^2$ b



33. Work done by all forces _____ J

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced · Chemistry – questions to revisit

6 questions · easy misses first
cropped from the question paper

Q49 General Organic Chemistry

JEE Advanced · Chemistry · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **53%** got it right · Campus: **46%** (36 of 78)

Within reach
You: Wrong

PARAGRAPH-I

Electrophiles and nucleophiles are fundamental concepts of organic chemistry playing crucial roles in various reactions. Electron loving species that accept electron pairs to form covalent bonds are electrophiles, Nucleus loving species that donate electron pair to form covalent bonds.

49. How many of the following are nucleophiles

A) OR^- B) CH_3^+ C) NH_3 D) $AlCl_3$

E) NO_2^- F) NO_2^+ G) Br^+

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q43 Hydrocarbons – Alkane from O_2 needed; count secondary carbons

JEE Advanced · Chemistry · Integer · marking +4 / 0 · marks lost 4
Top 100: **47%** got it right · Campus: **28%** (22 of 78)

Within reach
You: Wrong

43. one mole of alkane (Y) requires 8 moles of oxygen for complete combustion on chlorination with $Cl_2 / h\gamma$, gives four monochlorinated products (including stereo). The total number of secondary carbon atoms per each (Y) is ……..

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q46

Isomerism – Count of optically active compounds


JEE Advanced · Chemistry · Integer · marking +4 / 0 · marks lost 4

Top 100: **15%** got it right · Campus: **15%** (12 of 78)

Tough one

You: Wrong

46. How many given compounds are optically active



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q51

Hydrocarbons – Chiral products of free-radical monobromination

JEE Advanced · Chemistry · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **15%** got it right · Campus: **4%** (3 of 78)


Tough one

You: Wrong

PARAGRAPH-II

Alkanes are aliphatic saturated hydrocarbons and have the general molecular formula C_nH_{2n+2}. The simplest one is methane (CH_4), is also the most abundant. Large amounts are present in atmosphere, in the ground, and in the oceans. Alkanes are most reduced forms of organic compounds. Hence, they can be prepared by reduction of any organic compound. Alkanes are prepared by the catalytic hydrogenation of alkenes or alkynes.

51. In the following monobromination reaction, the number of possible chiral products is



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q47 **Hydrocarbons – Dichlorination isomers of C_5H_{10} (VD 35)**

JEE Advanced · Chemistry · Integer · marking +4 / 0 · marks lost 4

Top 100: **7%** got it right · Campus: **5%** (4 of 78)

Tough one

You: Wrong

47. A hydrocarbon [X] is having vapour density 35 and it forms only 1 monochloro product under sunlight. How many isomeric products (including stereo) are possible when [X] undergoes dichlorination?

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed


PARAGRAPH-II

Alkanes are aliphatic saturated hydrocarbons and have the general molecular formula C_nH_{2n+2}. The simplest one is methane (CH_4), is also the most abundant. Large amounts are present in atmosphere, in the ground, and in the oceans. Alkanes are most reduced forms of organic compounds. Hence, they can be prepared by reduction of any organic compound. Alkanes are prepared by the catalytic hydrogenation of alkenes or alkynes.

50. Find the No of correct statements out of the following

A) Alkanes are prepared by catalytic hydrogenation of Alkenes where both hydrogen atoms add to the same face of the alkene (syn-addition).

B) Alkyl halides when reacts with Lithium Aluminium Hydride form corresponding alkane: $R-X \xrightarrow{LiAlH_4} R-H$ for R : methyl , $1^0, 2^0$




C)In case of Tertiary Alkyl halide is observed.


D) Grignard's reagent when treated with a polar protic solvent gives alkane:

$R-Mg+HA \rightarrow R-H+MgX(A)$

E) The different compounds obtained by the following reaction are 7 (only organic compounds)



F) Rate of sodalime decarboxylation for the following acids follow:



G) The side products of potassium butanoate during Kolbe's Electrolysis are propane, propene and propyl butanoate

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

10 How to read this report

Where the numbers come from. Marks, percentages, All-India / State / Campus ranks, subject marks, subject All-India ranks and the wrong / skipped questions are the official results in the institute's *All India Marks Analysis* (graded with the final answer key). Maximum marks leave out deleted questions — JEE Mains: Q58 — JEE Advanced: Q45. **Program rank** = rank in Pragya G9 (EXT) Achievers across all campuses, from the batch's question-wise sheet († JEE Advanced: that sheet was graded before the final key, so this rank may differ slightly). **Top 100** = the batch's top 100 across all campuses, from the same question-wise sheet; * where it has no data for a question, the campus figure decides the priority. **Campus** = all students of this program on campus. **Priority:** Easy miss = 70%+ of the top 100 got it right; Within reach = 40–69%; Tough one = below 40%. **Marks lost** on a question = its marks plus the negative mark if answered wrong, its marks if skipped. In JEE Advanced, 6 marks went on multiple-correct answers that were only partly correct. **Weekly-test average** = your average over the weekly Mains tests (WTM) or Advanced tests (WTA) before this Part Test, from the campus averages sheet. The weekly Advanced tests are marked out of 240; their averages are scaled to 180 to match this paper (× 180/240, so 80 per subject becomes 60). Percentages are unchanged.