CLASS 10 · PART TEST-1 · STUDENT PERFORMANCE REPORT

Dharshan Raj S

Pragya G10 (EXT) Achievers · Adm No **246551298**

Campus HYD/Nallakunta DD Colony
JEE Mains 08 Oct 2026
JEE Advanced 09 Oct 2026

JEE Mains

08 Oct 2026 · 74 questions

219 / 296 — **74.0%**

20	15	7/65	7
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **236.4 / 300** (78.8%), 5 of 6 tests → this test **74.0%** ▼ **4.8**

77 marks lost: 65 on 13 wrong answers (incl. 13 negative) · 12 on 3 skipped

JEE Advanced

09 Oct 2026 · 50 questions

136 / 176 — **77.3%**

17	10	2/61	2
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **125.2 / 180** (69.5%), 5 of 6 tests → this test **77.3%** ▲ **7.7**

40 marks lost: 32 on 8 wrong answers (incl. 3 negative) · 6 on 2 skipped · 2 on partly-correct answers

1 Subject scorecard

Your marks in each subject, against the 65 students of this program on campus and against your own weekly-test average.

SUBJECT	MARKS	SCORE VS CAMPUS	SUBJECT AIR	CAMPUS AVG / BEST	WEEKLY AVG	CHANGE	RIGHT / WRONG / SKIP
JEE Mains 219 / 296 · 74.0%							
Maths	66/100	66%	85	58.2 / 100	78.3/100 78.3%	▼ 12.3	18 / 6 / 1
Physics	82/100	82%	35	53.8 / 100	85.9/100 85.9%	▼ 3.9	21 / 2 / 2
Chemistry	71/96	74%	22	55.9 / 91	72.4/100 72.4%	▲ 1.6	19 / 5 / 0
JEE Advanced 136 / 176 · 77.3%							
Maths	56/60	93%	4	30.5 / 57	48.8/60 81.3%	▲ 12.0	16 / 1 / 0
Physics	42/56	75%	30	24.5 / 56	42.8/60 71.3%	▲ 3.7	12 / 2 / 2
Chemistry	38/60	63%	158	38.2 / 60	33.6/60 56.1%	▲ 7.3	12 / 5 / 0

Bar = your score % | campus average | best on campus Change = this test % − weekly-test average %

What this report says

1. **Physics** is your strongest JEE Mains subject: 82/100 (82%), All-India subject rank 35.
2. **Maths** needs the most work in JEE Mains: 66/100 (66%), subject rank 85; you lost 7 of its 25 questions.
3. **Maths** is your strongest JEE Advanced subject: 56/60 (93%), All-India subject rank 4.
4. **Chemistry** needs the most work in JEE Advanced: 38/60 (63%), subject rank 158; you lost 5 of its 17 questions.
5. Biggest fall against your weekly-test average: **JEE Mains Maths** – 66% here against 78% on average.
6. Biggest improvement: **JEE Advanced Maths** – 93% here against 81% on average.
7. **2 questions** you lost were answered correctly by 70%+ of the batch's top 100 – about **9 marks** within easy reach (Mains Q15 · Advanced Q42).
8. Negative marking cost you **13 in JEE Mains and 3 in JEE Advanced**.
9. You attempted almost everything (only 5 questions skipped across both papers), so the marks to gain are in **accuracy**: wrong answers – 13 in JEE Mains and 8 in JEE Advanced.

JEE Mains – question by question

219 / 296 · 74.0%
08 Oct 2026

2 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	2	3	4	5	**6**	7	**8**	9	10	11	12	13	**14**	**15**	**16**	17	18	**19**	**20**	21	22	23	24	25
Physics	26	27	28	29	**30**	31	32	33	**34**	35	36	**37**	38	39	40	41	**42**	43	44	45	46	47	48	49	50
Chemistry	51	**52**	53	**54**	55	~~56~~	57	58	59	60	61	62	**63**	64	**65**	**66**	67	68	69	70	71	72	73	74	75

3 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 65 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40-69% · **Tough one** below 40%

Maths

7 of 25 questions lost · **34** marks · 1 easy miss · 2 within reach · 4 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q15	Easy miss	**Indefinite Integration**	Single correct · +4 / −1	Wrong	5	74%	77%
Q19	Within reach	**Differential Equations**	Single correct · +4 / −1	Wrong	5	68%	77%
Q8	Within reach	**Definite Integration**	Single correct · +4 / −1	Wrong	5	56%	38%
Q14	Tough one	**Definite Integration**	Single correct · +4 / −1	Wrong	5	39%	34%
Q6	Tough one	**Indefinite Integration** – $\int \sqrt{(x^2+1)}\,[\ln(x^2+1) - 2\ln x] / x^4\,dx$	Single correct · +4 / −1	Wrong	5	33%	31%
Q20	Tough one	**Definite Integration** – King's property with f(x) + f(y) = f(xy)	Single correct · +4 / −1	Skipped	4	29%	32%
Q16	Tough one	**Definite Integration** – $\int_0^{100} ([\tan^{-1}x] + [\cot^{-1}x])\,dx$ with GIF	Single correct · +4 / −1	Wrong	5	16%	11%

Physics

4 of 25 questions lost · **18** marks · 3 within reach · 1 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q37	Within reach	**Current Electricity** – Infinite ladder – voltage across the 2nd shunt resistor	Single correct · +4 / −1	Skipped	4	52%	46%
Q34	Within reach	**Current Electricity**	Single correct · +4 / −1	Skipped	4	45%	37%
Q42	Within reach	**Kinematics** – Distance (not displacement) in 3 s for a ball thrown up	Single correct · +4 / −1	Wrong	5	44%	48%
Q30	Tough one	**Electrostatics** – Field energy inside vs outside a charged sphere	Single correct · +4 / −1	Wrong	5	17%	11%





Chemistry

5 of 24 questions lost · **25** marks · 3 within reach · 2 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q66	Within reach	**Aldehydes & Ketones**	Single correct · +4 / −1	Wrong	5	65%	74%
Q65	Within reach	**Aldehydes & Ketones**	Single correct · +4 / −1	Wrong	5	60%	55%
Q54	Within reach	**Alcohols, Phenols & Ethers**	Single correct · +4 / −1	Wrong	5	47%	49%
Q63	Tough one	**Aldehydes & Ketones** – Enol-ether hydrolysis; iodoform-positive products	Single correct · +4 / −1	Wrong	5	32%	45%
Q52	Tough one	**Alcohols, Phenols & Ethers** – Ether vs butanol miscibility – assertion / reason	Single correct · +4 / −1	Wrong	5	24%	22%

4 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Definite Integration	9	**17**/36		29%	**Q8**, **Q14**, **Q16**, **Q20**
Indefinite Integration	7	**18**/28		61%	**Q6**, **Q15**
Differential Equations	4	**11**/16		84%	**Q19**
Area Under Curves	5	**20**/20		86%	– none
Physics					
Current Electricity	8	**24**/32		57%	**Q34**, **Q37**
Electrostatics	8	**27**/32		27%	**Q30**
Kinematics	9	**31**/36		75%	**Q42**
Chemistry					
Aldehydes & Ketones	14	**41**/56		65%	**Q63**, **Q65**, **Q66**
Alcohols, Phenols & Ethers	10	**30**/40		48%	**Q52**, **Q54**

Bar = your % of the chapter's marks | campus average

5 JEE Mains · Maths – questions to revisit

7 questions · easy misses first
cropped from the question paper

Q15 — Indefinite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5
Top 100: **74%** got it right · Campus: **77%** (50 of 65)

Easy miss
You: Wrong

15. If $\int \frac{4e^{x}+6e^{-x}}{9e^{x}-4e^{-x}}dx = ax + b\ln\left(9e^{2x}+c\right)+d$ then descending order of a,b,c is (where 'd' is constant of integration)

A) a,b,c B) b,c,a C) b,a,c D) c,a,b

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q19 — Differential Equations

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5
Top 100: **68%** got it right · Campus: **77%** (50 of 65)

Within reach
You: Wrong

19. The equation of the curve whose subnormal is equal to constant 'a' is

A) $y = ax + c$ B) $y^2 = 2ax + 2c$ C) $ay^2 = x^3 + ac$ D) $y^2 = ax + c$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q8 — Definite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5
Top 100: **56%** got it right · Campus: **38%** (25 of 65)

Within reach
You: Wrong

8. If $\int_{0}^{\frac{\pi}{2}} \frac{\tan x}{\tan x+\sec x}dx = m(\pi+n)$, then $m\times n$ is equal to

A) $-\frac{1}{2}$ B) 1 C) -1 D) $\frac{1}{2}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q14 — Definite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5

Top 100: **39%** got it right · Campus: **34%** (22 of 65)

Tough one

You: Wrong

14. Let $I_1=\int_0^{\frac{\pi}{2}}\ln(\sin x)\ln(\sin 2x)\,dx$ and $I_2=\int_0^{\frac{\pi}{2}}(\ln(\sin 2x))^2\,dx$ then $\frac{\pi(\ln 2)^2}{2I_1-I_2}=$

A) $\frac{1}{2}$ B) 2 C) −2 D) $-\frac{1}{2}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q6 — Indefinite Integration – ∫ √(x²+1)[ln(x²+1) − 2 ln x] / x⁴ dx

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5

Top 100: **33%** got it right · Campus: **31%** (20 of 65)

Tough one

You: Wrong

6. $\int\frac{\sqrt{x^2+1}\left[\ln(x^2+1)-2\ln x\right]}{x^4}\,dx$ is equal to (c is arbitrary constant)

A) $\frac{(x^2+1)^{3/2}}{x^3}\left[\frac{2}{3}-\ln\left(\frac{x^2+1}{x^2}\right)\right]+c$ B) $\frac{(x^2+1)^{3/2}}{3x^3}\left[\ln\left(\frac{x^2+1}{x^2}\right)-\frac{2}{3}\right]+c$

C) $\frac{2(x^2+1)^{3/2}}{3x^3}\left[\frac{1}{3}-\frac{1}{2}\ln\left(\frac{x^2+1}{x^2}\right)\right]+c$ D) $\frac{(x^2+1)^{3/2}}{3x^3}\left[\ln\left(\frac{x^2+1}{x^2}\right)+\frac{2}{3}\right]+c$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q20 — Definite Integration – King's property with f(x) + f(y) = f(xy)

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **29%** got it right · Campus: **32%** (21 of 65)

Tough one

You: Skipped

20. Let $f:(0,\infty)\to R$ be a continuous function, satisfying $f(x)+f(y)=f(xy)$. Then the value of $\int_{\sqrt{f(e)}}^{\sqrt{f(e^2)}}\frac{x\tan x^2}{\tan x^2+\tan\left(f(e^3)-x^2\right)}\,dx$ is equal to

A) $\frac{1}{2}f(e^2)$ B) $\frac{1}{4}f(e^2)$ C) $\frac{1}{2}f(e)$ D) $\frac{f(e)}{4}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q16 **Definite Integration – $\int_0^{100}([\tan^{-1}x]+[\cot^{-1}x])\,dx$ with GIF**

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5

Top 100: **16%** got it right · Campus: **11%** (7 of 65)

Tough one

You: Wrong

16. If $\int_0^{100}\left(\left[\tan^{-1}x\right]+\left[\cot^{-1}x\right]\right)dx=\ldots$ [.] denotes the greatest integer function

A) $100-\cot 1$ B) $100+2\cot 1$ C) $100-2\cot 2$ D) $100+2\cot 2$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Mains · Physics – questions to revisit

4 questions · easy misses first
cropped from the question paper

Q37 — Current Electricity – Infinite ladder – voltage across the 2nd shunt resistor

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4
Top 100: **52%** got it right · Campus: **46%** (30 of 65)

Within reach
You: Skipped

37. A 24 V battery of negligible internal resistance is connected to an infinite ladder network. In each section, both the top rail and the bottom rail contain a 1 Ω resistor, and a 4 Ω resistor is connected across the rails. The potential difference across the second 4 Ω resistor (counting from the battery) is

A) 12 V B) 8 V C) 6 V D) 3 V

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q34 — Current Electricity

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4
Top 100: **45%** got it right · Campus: **37%** (24 of 65)

Within reach
You: Skipped

34. In a Wheatstone bridge ABCD, a 12 V battery of negligible internal resistance is connected between A and C. The arms are AB = 2 Ω, BC = 3 Ω, AD = 3 Ω and DC = 3 Ω. An ammeter of resistance 0.3 Ω is connected between B and D. The current through the ammeter is

A) 0.2 A B) 0.6 A C) 0.4 A D) 1.2 A

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q42 — Kinematics – Distance (not displacement) in 3 s for a ball thrown up

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **44%** got it right · Campus: **48%** (31 of 65)

Within reach
You: Wrong

42. A ball is thrown vertically upward with a speed of 20 m/s (take g = 10 m/s^2). The total distance travelled by the ball in the first 3 s is

A) 15 m B) 25 m C) 20 m D) 45 m

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q30 **Electrostatics – Field energy inside vs outside a charged sphere**

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **17%** got it right · Campus: **11%** (7 of 65)

Tough one

You: Wrong

30. A solid sphere of radius R carries a total charge Q distributed uniformly throughout its volume. The ratio of the electrostatic field energy stored inside the sphere to that stored outside the sphere is

A) 3 : 5 B) 1 : 5 C) 1 : 3 D) 1 : 2

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Mains · Chemistry – questions to revisit


5 questions · easy misses first
cropped from the question paper

Q66 Aldehydes & Ketones


JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5
Top 100: **65%** got it right · Campus: **74%** (48 of 65)


Within reach
You: Wrong


66.




Product (D) will be

A) 

B) 

C) 

D) 

Why did I lose this question? (write the reason and the correct approach)


☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q65 Aldehydes & Ketones

JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5
Top 100: **60%** got it right · Campus: **55%** (36 of 65)

Within reach
You: Wrong

65. Which sequence of steps describes the best synthesis of 2-methyl 3-pentanone?



A) $1)1-propanol+(CH_3)_2CHMgBr, diethyl\ ether\ 2)H_3O^+\ 3)PCC.CH_2Cl_2$

B) $1)1-propanol+Na_2Cr_2O_7, H_2SO_4, H_2, heat\ 2)SOCl_2\ 3)(CH_3)_2CHCl, AlCl_3$

C) $1)1-propanol+PCC, CH_2Cl_2\ 2)(CH_3)_2CHLi, diethyl\ ether\ 3)H_3O^+\ 4)Na_2Cr_2O_7, H_2SO_4, H_2O\ heat$

D) $1)2-propanol+Na_2Cr_2O_7, H_2SO_4, H_2O\ heat\ 2)CH_3CH_2CH_2Li, diethyl\ ether\ 3)H_3O^+\ 4)PCC, CH_2Cl_2$


Why did I lose this question? (write the reason and the correct approach)


☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed


Q54 **Alcohols, Phenols & Ethers**
JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5
Top 100: **47%** got it right · Campus: **49%** (32 of 65)


Within reach
You: Wrong

54. Identify the correct acidic strength order of following pairs

A) 

B) 

C) 

D) 

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q63

Aldehydes & Ketones – Enol-ether hydrolysis; iodoform-positive products

JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5

Top 100: **32%** got it right · Campus: **45%** (29 of 65)


Tough one


You: Wrong

63. $\underset{C_5H_{10}O}{(A)} \xrightarrow{H_3O^{\oplus}} B + C; (B) and (C)$ both give +ve Iodoferm test.

Compound (A) is

A) $CH_3-CH=CH-O-CH_2-CH_3$

B) 

C) 

D) Both (B) and (C)

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q52

Alcohols, Phenols & Ethers – Ether vs butanol miscibility – assertion / reason

JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5

Top 100: **24%** got it right · Campus: **22%** (14 of 65)

Tough one

You: Wrong

52. **Assertion(A) :** Ethoxy ethane and butanol-1 have almost same miscibility with water.
Reason(R) : Ethoxy ethane and butanol-1 have form Hydrogen bond with water.

A) Both (A) and (R) are correct and (R) is the correct explanation of (A)

B) (A) is correct but (R) is not correct

C) Both (A) and (R) are not correct

D) Both (A) and (R) are correct and (R) is not the correct explanation of (A)

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced – question by question

136 / 176 · 77.3%
09 Oct 2026

6 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
Physics	18	19	20	21	22	23	24	~~25~~	26	27	28	29	30	31	32	33	34
Chemistry	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51

7 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 61 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40–69% · **Tough one** below 40%

Maths

1 of 17 questions lost · 4 marks · ● 1 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q12	Tough one	**Differential Equations** – Solve a first-order DE and match its constants	Integer · +4 / 0	Wrong	4	35%	26%

Physics

4 of 16 questions lost · 14 marks · ● 4 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q20	Tough one	**Kinematics** – Shell fired from a plateau – minimum landing distance	Single correct · +3 / −1	Wrong	4	29%	23%
Q27	Tough one	**Kinematics** – Projection angle that maximises area under the trajectory	Integer · +4 / 0	Wrong	4	27%	23%
Q31	Tough one	**Kinematics** – Hexagon projectile – maximum height	Paragraph · +3 / 0	Skipped	3	5%	5%
Q32	Tough one	**Kinematics** – Hexagon projectile – speed at the top	Paragraph · +3 / 0	Skipped	3	5%	5%




Chemistry

5 of 17 questions lost · 20 marks (+2 on partly-correct answers) · ● 1 easy miss · ● 1 within reach · ● 3 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q42	Easy miss	**Aldehydes & Ketones**	Integer · +4 / 0	Wrong	4	74%	87%
Q40	Within reach	**Alcohols, Phenols & Ethers**	Multiple correct · +4 / −2	Wrong	6	51%	44%
Q50	Tough one	**Carboxylic Acids** – Molar mass of an acid from NaOH titration	Paragraph · +3 / 0	Wrong	3	36%	52%
Q46	Tough one	**Alcohols, Phenols & Ethers** – Identifying polar aprotic solvents	Integer · +4 / 0	Wrong	4	24%	8%
Q51	Tough one	**Carboxylic Acids** – Moles of $LiAlH_4$ to reduce RCOOH to RCH_2OH	Paragraph · +3 / 0	Wrong	3	17%	23%

8 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject. Chapter marks are approximate (≈): partly-correct answers cannot be traced to a question.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Differential Equations	5	≈ **14**/18		30%	**Q12**
Area Under Curves	3	≈ **10**/10		60%	– none
Definite Integration	5	≈ **17**/17		54%	– none
Indefinite Integration	4	≈ **15**/15		71%	– none
Physics					
Kinematics	8	≈ **14**/28		41%	**Q20**, **Q27**, **Q31**, **Q32**
Current Electricity	2	≈ **7**/7		68%	– none
Electrostatics	6	≈ **21**/21		45%	– none
Chemistry					
Carboxylic Acids	4	≈ **8**/14		48%	**Q50**, **Q51**
Alcohols, Phenols & Ethers	7	≈ **16**/26		62%	**Q40**, **Q46**
Aldehydes & Ketones	6	≈ **16**/20		83%	**Q42**

Bar = your % of the chapter's marks | campus average

9 JEE Advanced · Maths – questions to revisit

1 question · easy misses first
cropped from the question paper

Q12 **Differential Equations** – Solve a first-order DE and match its constants

JEE Advanced · Maths · Integer · marking +4 / 0 · marks lost 4
Top 100: **35%** got it right · Campus: **26%** (16 of 61)

Tough one
You: Wrong

12. The differential equation $\left(xy^3\left(1+\cos x\right)-y\right)dx+xdy=0$ represents the curve $\frac{x^2}{ay^2}=\frac{x^3}{b}+x^2\sin x+cx\cos x-d\sin x+k$. Then a+b+c+d is equal to (where 'k' is arbitrary constant).

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced · Physics – questions to revisit


4 questions · easy misses first
cropped from the question paper

Q20

Kinematics – Shell fired from a plateau – minimum landing distance
JEE Advanced · Physics · Single correct · marking +3 / −1 · marks lost 4
Top 100: **29%** got it right · Campus: **23%** (14 of 61)

Tough one
You: Wrong

20. A gun is mounted on a plateau 960m away from its edge as shown. Height of plateau is 960m. The gun can fire shells with a velocity 100 m/s at any angle of the following choices. What is the minimum distance OP(=x) from the edge of plateau where shell of



gun can reach? (given g=10m/s^2, $\sin 74^0 = \sin 106^0 \simeq 0.96$)

A) 480 m B) 720 m C) 360 m D) 960 m

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q27

Kinematics – Projection angle that maximises area under the trajectory
JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4
Top 100: **27%** got it right · Campus: **23%** (14 of 61)

Tough one
You: Wrong

27. A projectile is thrown with a speed 10 m/s, at an angle θ with the horizontal. For a certain value of angle of projection $\theta = \frac{(n+1)\pi}{12}$ the area bound by the trajectory with horizontal is maximum value of n is (g=10 m/s^2). Assume no airdrag.


Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q31 **Kinematics – Hexagon projectile – maximum height**
JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **5%** got it right · Campus: **5%** (3 of 61)

Tough one
You: Skipped

PARAGRAPH-I



A regular hexagon of side a=$10\sqrt{3}$ m is shown in figure. From any point a particle is projected such that it touches B,C,D and E. The velocity of particle is v m/s. O is the centre of the hexagon. x-axis is parallel to horizontal and y-axis is perpendicular to AE and CD (g=10 m/s^2) $\left(\sqrt{3}=1.73\right)$.

31. The maximum height from the horizontal surface X' attained by the projectile is ___ m.


Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q32 **Kinematics – Hexagon projectile – speed at the top**
JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **5%** got it right · Campus: **5%** (3 of 61)

Tough one
You: Skipped

PARAGRAPH-I



A regular hexagon of side a=$10\sqrt{3}$ m is shown in figure. From any point a particle is projected such that it touches B,C,D and E. The velocity of particle is v m/s. O is the centre of the hexagon. x-axis is parallel to horizontal and y-axis is perpendicular to AE and CD (g=10 m/s^2) $\left(\sqrt{3}=1.73\right)$.

32. The velocity of projectile at maximum height is _____ m/s

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced · Chemistry – questions to revisit

5 questions · easy misses first
cropped from the question paper

Q42

Aldehydes & Ketones
JEE Advanced · Chemistry · Integer · marking +4 / 0 · marks lost 4
Top 100: **74%** got it right · Campus: **87%** (53 of 61)

Easy miss
You: Wrong

42. 

How many among the following compounds will give the above result?
i.Cyclohexanone
ii.Acetone
iii.Propionaldehyde
iv.Acetophenone
v.Acetaldehyde
vi.Benzophenone
vii.Benzaldehyde

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed


Q40 **Alcohols, Phenols & Ethers**

JEE Advanced · Chemistry · Multiple correct · marking +4 / −2 · marks lost 6


Top 100: **51%** got it right · Campus: **44%** (27 of 61)

Within reach


You: Wrong


40. 

major product is

A) 

B)$C_2H_5O\text{-}CH_3$

C) 

D) 

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q50 **Carboxylic Acids** – **Molar mass of an acid from NaOH titration**
JEE Advanced · Chemistry · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **36%** got it right · Campus: **52%** (32 of 61)

Tough one
You: Wrong

PARAGRAPH-II

Carboxylic acids are organic compounds containing the carbonyl(>C=O) and hydroxyl (-OH) functional groups combined as a carboxyl group(-COOH). Due to resonance stabilization of the carboxylate anion, carboxylic acids exhibit higher acidity compared to alcohols and phenols. They participate in several characteristic reactions, including nucleophilic acyl substitution (esterification, amide formation, acid halide synthesis), decarboxylation, and reduction to primary alcohols. Furthermore, the presence of the –COOH group strongly influences physical properties, enabling extensive intermolecular hydrogen bonding that results in higher boiling points and stable dimeric structures in non-polar solvents.

50. A 0.600 g sample of a pure monoprotic carboxylic acid (R-COOH) requires 25.00 mL of 0.200 M NaOH for complete neutralization. Calculate the molar mass of the carboxylic acid?


Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q46 **Alcohols, Phenols & Ethers** – **Identifying polar aprotic solvents**
JEE Advanced · Chemistry · Integer · marking +4 / 0 · marks lost 4
Top 100: **24%** got it right · Campus: **8%** (5 of 61)

Tough one
You: Wrong

46. How many of the following are polar aprotic solvents?



[P]

Diethyl ether,


[Q]

Benzene,

[R]

Ethyl alcohol,

[S]



[T]

Petrol

[U]

Acetone

[Q]

DMSO (dimethyl sulphoxide

[W]

DMF (dimethyl formamide

[X]

formic acid

[Y]

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q51 **Carboxylic Acids – Moles of $LiAlH_4$ to reduce RCOOH to RCH_2OH**

JEE Advanced · Chemistry · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **17%** got it right · Campus: **23%** (14 of 61)

Tough one

You: Wrong

PARAGRAPH-II

Carboxylic acids are organic compounds containing the carbonyl(>C=O) and hydroxyl (-OH) functional groups combined as a carboxyl group(-COOH). Due to resonance stabilization of the carboxylate anion, carboxylic acids exhibit higher acidity compared to alcohols and phenols. They participate in several characteristic reactions, including nucleophilic acyl substitution (esterification, amide formation, acid halide synthesis), decarboxylation, and reduction to primary alcohols. Furthermore, the presence of the –COOH group strongly influences physical properties, enabling extensive intermolecular hydrogen bonding that results in higher boiling points and stable dimeric structures in non-polar solvents.

51. The number of moles of $LiAlH_4$ required theoretically to reduce eight moles of carboxylic acid to primary alcohol

$$RCOOH \xrightarrow[2.H_3O^+]{1.LiAiH_4} RCH_2OH$$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

10 How to read this report

Where the numbers come from. Marks, percentages, All-India / State / Campus ranks, subject marks, subject All-India ranks and the wrong / skipped questions are the official results in the institute's *All India Marks Analysis* (graded with the final answer key). Maximum marks leave out deleted questions — JEE Mains: Q56 — JEE Advanced: Q25. **Program rank** = rank in Pragya G10 (EXT) Achievers across all campuses, from the batch's question-wise sheet. **Top 100** = the batch's top 100 across all campuses, from the same question-wise sheet; * where it has no data for a question, the campus figure decides the priority. **Campus** = all students of this program on campus. **Priority:** Easy miss = 70%+ of the top 100 got it right; Within reach = 40–69%; Tough one = below 40%. **Marks lost** on a question = its marks plus the negative mark if answered wrong, its marks if skipped. In JEE Advanced, 2 marks went on multiple-correct answers that were only partly correct. **Weekly-test average** = your average over the weekly Mains tests (WTM) or Advanced tests (WTA) before this Part Test, from the campus averages sheet. The weekly Advanced tests are marked out of 240; their averages are scaled to 180 to match this paper (× 180/240, so 80 per subject becomes 60). Percentages are unchanged.