CLASS 10 · PART TEST-1 · STUDENT PERFORMANCE REPORT

Gajavali Anjaneyulu

Pragya G10 (EXT) Achievers · Adm No **235416384**

Campus HYD/Nallakunta DD Colony
JEE Mains 08 Oct 2026
JEE Advanced 09 Oct 2026

JEE Mains

08 Oct 2026 · 74 questions

169 / 296 — **57.1%**

117	83	33/65	43
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **190.0 / 300** (63.3%) → this test **57.1%** ▼ **6.2**

127 marks lost: 115 on 23 wrong answers (incl. 23 negative) · 12 on 3 skipped

JEE Advanced

09 Oct 2026 · 50 questions

118 / 176 — **67.0%**

52	36	9/61	13
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **92.6 / 180** (51.4%), 5 of 6 tests → this test **67.0%** ▲ **15.6**

58 marks lost: 51 on 14 wrong answers (incl. 4 negative) · 7 on 2 skipped

1 Subject scorecard

Your marks in each subject, against the 65 students of this program on campus and against your own weekly-test average.

SUBJECT	MARKS	SCORE VS CAMPUS	SUBJECT AIR	CAMPUS AVG / BEST	WEEKLY AVG	CHANGE	RIGHT / WRONG / SKIP
JEE Mains 169 / 296 · 57.1%							
Maths	62/100	62%	125	58.2 / 100	69.1/100 69.1%	▼ 7.1	17 / 6 / 2
Physics	55/100	55%	227	53.8 / 100	61.9/100 61.9%	▼ 6.9	16 / 9 / 0
Chemistry	52/96	54%	151	55.9 / 91	59.0/100 59.0%	▼ 4.8	15 / 8 / 1
JEE Advanced 118 / 176 · 67.0%							
Maths	43/60	72%	68	30.5 / 57	38.3/60 63.8%	▲ 7.9	13 / 4 / 0
Physics	25/56	45%	273	24.5 / 56	29.1/60 48.5%	▼ 3.8	7 / 8 / 1
Chemistry	50/60	83%	25	38.2 / 60	25.2/60 42.1%	▲ 41.3	14 / 2 / 1

Bar = your score % | campus average | best on campus Change = this test % − weekly-test average %

What this report says

1. **Maths** is your strongest JEE Mains subject: 62/100 (62%), All-India subject rank 125.
2. **Chemistry** needs the most work in JEE Mains: 52/96 (54%), subject rank 151; you lost 9 of its 24 questions.
3. **Chemistry** is your strongest JEE Advanced subject: 50/60 (83%), All-India subject rank 25.
4. **Physics** needs the most work in JEE Advanced: 25/56 (45%), subject rank 273; you lost 9 of its 16 questions.
5. Biggest fall against your weekly-test average: **JEE Mains Maths** – 62% here against 69% on average.
6. Biggest improvement: **JEE Advanced Chemistry** – 83% here against 42% on average.
7. **4 questions** you lost were answered correctly by 70%+ of the batch's top 100 – about **20 marks** within easy reach (Mains Q17, Q25, Q38, Q57).
8. Negative marking cost you **23 in JEE Mains and 4 in JEE Advanced**.
9. You attempted almost everything (only 5 questions skipped across both papers), so the marks to gain are in **accuracy**: wrong answers – 23 in JEE Mains and 14 in JEE Advanced.

JEE Mains – question by question

169 / 296 · 57.1%
08 Oct 2026

2 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	2	3	4	**5**	**6**	7	8	**9**	10	**11**	12	13	14	15	16	**17**	**18**	19	20	21	22	**23**	24	**25**
Physics	26	**27**	**28**	**29**	**30**	31	32	33	**34**	**35**	36	**37**	**38**	39	40	41	42	43	44	45	**46**	47	48	49	50
Chemistry	51	**52**	53	**54**	**55**	~~56~~	**57**	58	**59**	60	61	62	63	64	65	66	67	68	**69**	70	71	72	**73**	**74**	**75**

3 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 65 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40-69% · **Tough one** below 40%

Maths

8 of 25 questions lost · **38** marks · 2 easy miss · 3 within reach · 3 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q25	Easy miss	**Differential Equations**	Numerical · +4 / −1	Wrong	5	78%	89%
Q17	Easy miss	**Indefinite Integration**	Single correct · +4 / −1	Wrong	5	74%	66%
Q9	Within reach	**Indefinite Integration**	Single correct · +4 / −1	Wrong	5	66%	63%
Q23	Within reach	**Area Under Curves**	Numerical · +4 / −1	Skipped	4	66%	72%
Q5	Within reach	**Definite Integration**	Single correct · +4 / −1	Wrong	5	40%	32%
Q18	Tough one	**Definite Integration** – ∫ of min{|x − 3k|} – periodic zig-zag	Single correct · +4 / −1	Wrong	5	37%	40%
Q6	Tough one	**Indefinite Integration** – ∫ √(x²+1) [ln(x²+1) − 2 ln x] / x⁴ dx	Single correct · +4 / −1	Skipped	4	33%	31%
Q11	Tough one	**Definite Integration**	Single correct · +4 / −1	Wrong	5	29%	26%

Physics

9 of 25 questions lost · **45** marks · 1 easy miss · 6 within reach · 2 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q38	Easy miss	**Current Electricity**	Single correct · +4 / −1	Wrong	5	88%	91%
Q29	Within reach	**Electrostatics**	Single correct · +4 / −1	Wrong	5	61%	48%
Q28	Within reach	**Electrostatics**	Single correct · +4 / −1	Wrong	5	54%	45%
Q37	Within reach	**Current Electricity** – Infinite ladder – voltage across the 2nd shunt resistor	Single correct · +4 / −1	Wrong	5	52%	46%
Q35	Within reach	**Current Electricity**	Single correct · +4 / −1	Wrong	5	50%	48%
Q34	Within reach	**Current Electricity**	Single correct · +4 / −1	Wrong	5	45%	37%
Q27	Within reach	**Electrostatics** – Potential at the centre of a thick charged shell	Single correct · +4 / −1	Wrong	5	42%	32%
Q46	Tough one	**Electrostatics**	Numerical · +4 / −1	Wrong	5	25%	25%
Q30	Tough one	**Electrostatics** – Field energy inside vs outside a charged sphere	Single correct · +4 / −1	Wrong	5	17%	11%

Chemistry

9 of 24 questions lost · 44 marks · 1 easy miss · 5 within reach · 3 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q57	Easy miss	Aldehydes & Ketones	Single correct · +4 / −1	Wrong	5	74%	82%
Q59	Within reach	Alcohols, Phenols & Ethers	Single correct · +4 / −1	Wrong	5	65%	52%
Q55	Within reach	Alcohols, Phenols & Ethers	Single correct · +4 / −1	Wrong	5	54%	71%
Q69	Within reach	Aldehydes & Ketones – Order of hydration constants of carbonyls	Single correct · +4 / −1	Skipped	4	52%	49%
Q54	Within reach	Alcohols, Phenols & Ethers	Single correct · +4 / −1	Wrong	5	47%	49%
Q73	Within reach	Alcohols, Phenols & Ethers	Numerical · +4 / −1	Wrong	5	43%	49%
Q75	Tough one	Aldehydes & Ketones	Numerical · +4 / −1	Wrong	5	27%	38%
Q52	Tough one	Alcohols, Phenols & Ethers – Ether vs butanol miscibility – assertion / reason	Single correct · +4 / −1	Wrong	5	24%	22%
Q74	Tough one	Alcohols, Phenols & Ethers – Site of benzylation of 2-naphthoxide	Numerical · +4 / −1	Wrong	5	12%	15%

4 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Indefinite Integration	7	14/28		61%	Q6, Q9, Q17
Definite Integration	9	21/36		29%	Q5, Q11, Q18
Differential Equations	4	11/16		84%	Q25
Area Under Curves	5	16/20		86%	Q23
Physics					
Electrostatics	8	7/32		27%	Q27, Q28, Q29, Q30, Q46
Current Electricity	8	12/32		57%	Q34, Q35, Q37, Q38
Kinematics	9	36/36		75%	– none
Chemistry					
Alcohols, Phenols & Ethers	10	10/40		48%	Q52, Q54, Q55, Q59, Q73, Q74
Aldehydes & Ketones	14	42/56		65%	Q57, Q69, Q75

Bar = your % of the chapter's marks | campus average

5 JEE Mains · Maths – questions to revisit

8 questions · easy misses first
cropped from the question paper

Q25 — Differential Equations

JEE Mains · Maths · Numerical · marking +4 / −1 · marks lost 5
Top 100: **78%** got it right · Campus: **89%** (58 of 65)

Easy miss
You: Wrong

25. If the solution $y(x)$ of the given differential equation $(e^y+1)\cos x dx + e^y \sin x dy = 0$ passes through the point $\left(\frac{\pi}{2}, 0\right)$, then the value of $e^{y\left(\frac{\pi}{6}\right)}$ is equal to

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q17 — Indefinite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5
Top 100: **74%** got it right · Campus: **66%** (43 of 65)

Easy miss
You: Wrong

17. $\int \left(1+x-x^{-1}\right)e^{x+x^{-1}}dx =$ _____ (c is arbitrary constant)

A) $(1+x)e^{x+x^{-1}}+c$ B) $(x-1)e^{x+x^{-1}}+c$ C) $-xe^{x+x^{-1}}+c$ D) $xe^{x+x^{-1}}+c$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q9

Indefinite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5

Top 100: **66%** got it right · Campus: **63%** (41 of 65)

Within reach

You: Wrong

9. $\int \frac{dx}{\sin x + \sec x} =$ (c is arbitrary constant)

A) $\frac{1}{2\sqrt{3}} \ln \left| \frac{\sin x - \cos x + \sqrt{3}}{\cos x - \sin x + \sqrt{3}} \right| + \tan^{-1}(\sin x + \cos x) + c$

B) $\frac{1}{2\sqrt{3}} \ln \left| \frac{\sin x - \cos x - \sqrt{3}}{\sin x - \cos x + \sqrt{3}} \right| + c$

C) $\frac{1}{2\sqrt{3}} \ln \left| (\sin x - \cos x + \sqrt{3})(\sin x - \cos x - \sqrt{3}) \right| + c$

D) $\frac{1}{2\sqrt{3}} \ln \left| (\sin x - \cos x + \sqrt{3})(\sin x - \cos x - \sqrt{3}) \right| + \tan^{-1}(\sin x + \cos x) + c$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q23

Area Under Curves

JEE Mains · Maths · Numerical · marking +4 / −1 · marks lost 4

Top 100: **66%** got it right · Campus: **72%** (47 of 65)

Within reach

You: Skipped

23. The area bounded by the lines $y = ||x-1|-2|$ with x-axis is _____ (in sq.units)

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q5

Definite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5

Top 100: **40%** got it right · Campus: **32%** (21 of 65)

Within reach

You: Wrong

5. $\int_{-1}^{1} \sin x \ln\left(1^x + 2^x + 3^x + 6^x\right) dx =$

A) $(\ln 6)\left(\cos\left(\frac{\pi+4}{4}\right)\right)$

B) $(\ln 6)\left(\sin\left(\frac{\pi+4}{4}\right)\right)$

C) $\sqrt{2}(\ln 6)\left(\sin\left(\frac{4-\pi}{4}\right)\right)$

D) $\sqrt{2}(\ln 6)\left(\sin\left(\frac{\pi+4}{4}\right)\right)$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q18

Definite Integration – ∫ of min{|x − 3k|} – periodic zig-zag

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5

Top 100: **37%** got it right · Campus: **40%** (26 of 65)

Tough one

You: Wrong

18. Let $f(x) = \min\{\ldots, |x+6|, |x+3|, |x|, |x-3|, |x-6|, \ldots\}$ then $\int_{-4}^{3} f(x)\,dx =$ _____

A) 4

B) $\frac{9}{2}$

C) 5

D) $\frac{11}{2}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q6

Indefinite Integration – ∫ √(x²+1)[ln(x²+1) − 2 ln x] / x⁴ dx

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **33%** got it right · Campus: **31%** (20 of 65)

Tough one

You: Skipped

6. $\int \frac{\sqrt{x^2+1}\left[\ln\left(x^2+1\right) - 2\ln x\right]}{x^4} dx$ is equal to (c is arbitrary constant)

A) $\frac{\left(x^2+1\right)^{3/2}}{x^3}\left[\frac{2}{3} - \ln\left(\frac{x^2+1}{x^2}\right)\right] + c$

B) $\frac{\left(x^2+1\right)^{3/2}}{3x^3}\left[\ln\left(\frac{x^2+1}{x^2}\right) - \frac{2}{3}\right] + c$

C) $\frac{2\left(x^2+1\right)^{3/2}}{3x^3}\left[\frac{1}{3} - \frac{1}{2}\ln\left(\frac{x^2+1}{x^2}\right)\right] + c$

D) $\frac{\left(x^2+1\right)^{3/2}}{3x^3}\left[\ln\left(\frac{x^2+1}{x^2}\right) + \frac{2}{3}\right] + c$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q11

Definite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5

Top 100: **29%** got it right · Campus: **26%** (17 of 65)

Tough one

You: Wrong

11. Let $f(x)=\sin x+\sum_{r=0}^{2}\cos\left(x+\frac{2r\pi}{3}\right)$, $g(x)=\cos x+\sum_{r=0}^{2}\sin\left(x+\frac{2r\pi}{3}\right)$ then $\int_{0}^{\frac{\pi}{2}} f(g(x))f(x)\,dx=$

A) $1+\sin 1$ B) $1-\sin 1$ C) $1+\cos 1$ D) $1-\cos 1$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Mains · Physics – questions to revisit

9 questions · easy misses first
cropped from the question paper

Q38

Current Electricity
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **88%** got it right · Campus: **91%** (59 of 65)

Easy miss
You: Wrong

38. Four wires each of resistance R, form a square ABCD. The equivalent resistance between two adjacent corners A and B is

A) $\frac{R}{2}$ B) $\frac{3R}{4}$ C) R D) $\frac{4R}{3}$

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q29

Electrostatics
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **61%** got it right · Campus: **48%** (31 of 65)

Within reach
You: Wrong

29. A point charge q is placed on the axis of a circular disc of radius R at a distance $\frac{R}{\sqrt{3}}$ from its centre. The electric flux through the disc is

A) $\frac{q\left(2-\sqrt{3}\right)}{4\varepsilon_0}$ B) $\frac{q}{6\varepsilon_0}$ C) $\frac{q}{2\varepsilon_0}$ D) $\frac{q}{4\varepsilon_0}$

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q28 Electrostatics

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **54%** got it right · Campus: **45%** (29 of 65)

Within reach

You: Wrong

28. Two fixed point charges +Q each are separated by '2a'. A particle of mass 'm' and charge −q is placed on the perpendicular bisector of the line joining them, at a small distance y (≪ a) from the midpoint, and released. Assuming only electrostatic forces act, the time period of its oscillation is

A) $2\pi\sqrt{\frac{\pi\varepsilon_0 ma^3}{Qq}}$ B) $2\pi\sqrt{\frac{2\pi\varepsilon_0 ma^3}{Qq}}$ C) $2\pi\sqrt{\frac{4\pi\varepsilon_0 ma^3}{Qq}}$ D) $\pi\sqrt{\frac{\pi\varepsilon_0 ma^3}{Qq}}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q37 Current Electricity – Infinite ladder – voltage across the 2nd shunt resistor

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **52%** got it right · Campus: **46%** (30 of 65)

Within reach

You: Wrong

37. A 24 V battery of negligible internal resistance is connected to an infinite ladder network. In each section, both the top rail and the bottom rail contain a 1 Ω resistor, and a 4 Ω resistor is connected across the rails. The potential difference across the second 4 Ω resistor (counting from the battery) is

A) 12 V B) 8 V C) 6 V D) 3 V

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q35 **Current Electricity**
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **50%** got it right · Campus: **48%** (31 of 65)

Within reach
You: Wrong

35. A 12 V battery of negligible internal resistance is connected to the input terminals of an infinite ladder network. Each section of the ladder has a 1 Ω resistor in the top (series) arm and a 2 Ω resistor connected across the two rails. The current through the first 2 Ω shunt resistor is

A) 6 A B) 3 A C) 2 A D) 4 A

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q34 **Current Electricity**
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **45%** got it right · Campus: **37%** (24 of 65)

Within reach
You: Wrong

34. In a Wheatstone bridge ABCD, a 12 V battery of negligible internal resistance is connected between A and C. The arms are AB = 2 Ω, BC = 3 Ω, AD = 3 Ω and DC = 3 Ω. An ammeter of resistance 0.3 Ω is connected between B and D. The current through the ammeter is

A) 0.2 A B) 0.6 A C) 0.4 A D) 1.2 A

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q27 **Electrostatics – Potential at the centre of a thick charged shell**
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **42%** got it right · Campus: **32%** (21 of 65)

Within reach
You: Wrong

27. A thick spherical shell of inner radius ‘a’ and outer radius ‘b’ carries a uniform volume charge density ρ. The electric potential at the centre of the shell (taking V = 0 at infinity) is

A) $\rho\frac{(b^2-a^2)}{12\varepsilon_0}$ B) $\rho\frac{(b^3-a^3)}{3\varepsilon_0 b}$ C) $\rho\frac{(b^2-a^2)}{2\varepsilon_0}$ D) $\rho\frac{(b^3-a^3)}{3\varepsilon_0 a}$

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q46 **Electrostatics**
JEE Mains · Physics · Numerical · marking +4 / −1 · marks lost 5
Top 100: **25%** got it right · Campus: **25%** (16 of 65)

Tough one
You: Wrong

46. In a region the electric field is $E=\left(5x\hat{i}+3y\hat{j}+4\hat{k}\right)N/C$, with x, y in meters. A cube of side 1m has one vertex at the origin and its edges along the positive x, y and z axis. The net electric flux through the cube (in Nm^2/C) is _____.

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q30 **Electrostatics – Field energy inside vs outside a charged sphere**
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **17%** got it right · Campus: **11%** (7 of 65)

Tough one
You: Wrong

30. A solid sphere of radius R carries a total charge Q distributed uniformly throughout its volume. The ratio of the electrostatic field energy stored inside the sphere to that stored outside the sphere is

A) 3 : 5 B) 1 : 5 C) 1 : 3 D) 1 : 2

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Mains · Chemistry – questions to revisit


9 questions · easy misses first
cropped from the question paper

Q57


Aldehydes & Ketones
JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5
Top 100: **74%** got it right · Campus: **82%** (53 of 65)

Easy miss
You: Wrong


57. Rank of the following in order of increasing value of the equilibrium constant for hydration K_{hyd} (smallest value first)



(1)



(2)



(3)

A) 1<2<3 B) 3<1<2 C) 2<1<3 D) 2<3<1

Why did I lose this question? (write the reason and the correct approach)


☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q59

Alcohols, Phenols & Ethers
JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5
Top 100: **65%** got it right · Campus: **52%** (34 of 65)

Within reach
You: Wrong

59. Total number of stereoisomers of the product (B) will be



A) 2 B) 3 C) 4 D) 5

Why did I lose this question? (write the reason and the correct approach)


☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q55

Alcohols, Phenols & Ethers
JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5
Top 100: **54%** got it right · Campus: **71%** (46 of 65)

Within reach
You: Wrong

55. Given below are two statements for the



$$C_2H_5O^- + H_2O \rightleftharpoons C_2H_5OH + OH^-; K_2$$

Statement I : The value of K_2 less than K_1

Statement II : Phenoxide is a weaker base than ethoxide

In the light of the above statement, choose the most appropriate answer from the options given below.

A) Both Statement I and Statement II are correct

B) Both Statement I and Statement II are incorrect

C) Statement I is correct but Statement II is incorrect

D) Statement II is correct but Statement I is incorrect

Why did I lose this question? (write the reason and the correct approach)


☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q69

Aldehydes & Ketones – Order of hydration constants of carbonyls
JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 4
Top 100: **52%** got it right · Campus: **49%** (32 of 65)

Within reach
You: Skipped

69. Arrange in their increasing order of equilibrium constants for hydration?



A) A < B < C < D < E

B) A < C < B < E < D

C) A < C < E < B < D

D) C < A < B < E < D

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q54

Alcohols, Phenols & Ethers


JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5


Top 100: **47%** got it right · Campus: **49%** (32 of 65)


Within reach


You: Wrong

54. Identify the correct acidic strength order of following pairs

A) 

B) 

C) 

D) 

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q73

Alcohols, Phenols & Ethers

JEE Mains · Chemistry · Numerical · marking +4 / −1 · marks lost 5

Top 100: **43%** got it right · Campus: **49%** (32 of 65)

Within reach

You: Wrong

73. How many chiral alcohols, including stereoisomers can give 2-chloro-2-methylbutane as the major product on reaction with $NaCl / H_2SO_4$ by hydride ion shift during its path?

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q75


Aldehydes & Ketones

JEE Mains · Chemistry · Numerical · marking +4 / −1 · marks lost 5

Top 100: **27%** got it right · Campus: **38%** (25 of 65)

Tough one

You: Wrong

75. 

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q52

Alcohols, Phenols & Ethers – Ether vs butanol miscibility – assertion / reason

JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5

Top 100: **24%** got it right · Campus: **22%** (14 of 65)

Tough one

You: Wrong

52. **Assertion(A) :** Ethoxy ethane and butanol-1 have almost same miscibility with water.
Reason(R) : Ethoxy ethane and butanol-1 have form Hydrogen bond with water.
A) Both (A) and (R) are correct and (R) is the correct explanation of (A)
B) (A) is correct but (R) is not correct
C) Both (A) and (R) are not correct
D) Both (A) and (R) are correct and (R) is not the correct explanation of (A)

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q74

Alcohols, Phenols & Ethers – Site of benzylation of 2-naphthoxide


JEE Mains · Chemistry · Numerical · marking +4 / −1 · marks lost 5

Top 100: **12%** got it right · Campus: **15%** (10 of 65)

Tough one

You: Wrong

74. Which site is predominately benzylated in the given reaction?



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced – question by question

118 / 176 · 67.0%
09 Oct 2026

6 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
Physics	18	19	20	21	22	23	24	~~25~~	26	27	28	29	30	31	32	33	34
Chemistry	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51

7 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 61 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40–69% · **Tough one** below 40%

Maths

4 of 17 questions lost · 17 marks · 2 within reach · 2 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q14	Within reach	**Definite Integration**	Paragraph · +3 / 0	Wrong	3	60%	56%
Q7	Within reach	**Differential Equations** – Curve from tangent-intercept ratio BP : AP = 3 : 1	Multiple correct · +4 / −2	Wrong	6	43%	26%
Q12	Tough one	**Differential Equations** – Solve a first-order DE and match its constants	Integer · +4 / 0	Wrong	4	35%	26%
Q11	Tough one	**Differential Equations**	Integer · +4 / 0	Wrong	4	25%	21%

Physics

9 of 16 questions lost · 31 marks · 4 within reach · 5 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q21	Within reach	**Kinematics**	Single correct · +3 / −1	Wrong	4	64%	51%
Q18	Within reach	**Current Electricity**	Single correct · +3 / −1	Wrong	4	57%	49%
Q28	Within reach	**Electrostatics** – Flux through a cube with charges at its vertices	Integer · +4 / 0	Wrong	4	54%	44%
Q33	Within reach	**Electrostatics** – Flux through a ring – solid angle	Paragraph · +3 / 0	Wrong	3	42%	38%
Q34	Tough one	**Electrostatics** – Flux through a cube from a wire along one edge	Paragraph · +3 / 0	Wrong	3	33%	23%
Q20	Tough one	**Kinematics** – Shell fired from a plateau – minimum landing distance	Single correct · +3 / −1	Skipped	3	29%	23%
Q27	Tough one	**Kinematics** – Projection angle that maximises area under the trajectory	Integer · +4 / 0	Wrong	4	27%	23%
Q31	Tough one	**Kinematics** – Hexagon projectile – maximum height	Paragraph · +3 / 0	Wrong	3	5%	5%
Q32	Tough one	**Kinematics** – Hexagon projectile – speed at the top	Paragraph · +3 / 0	Wrong	3	5%	5%

Chemistry

3 of 17 questions lost · 10 marks · ● 3 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q50	Tough one	**Carboxylic Acids** – Molar mass of an acid from NaOH titration	Paragraph · +3 / 0	Wrong	3	36%	52%
Q46	Tough one	**Alcohols, Phenols & Ethers** – Identifying polar aprotic solvents	Integer · +4 / 0	Skipped	4	24%	8%
Q51	Tough one	**Carboxylic Acids** – Moles of $LiAlH_4$ to reduce RCOOH to RCH_2OH	Paragraph · +3 / 0	Wrong	3	17%	23%

8 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject. Chapter marks are approximate (≈): partly-correct answers cannot be traced to a question.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Differential Equations	5	≈ 4/18		30%	Q7, Q11, Q12
Definite Integration	5	≈ 14/17		54%	Q14
Area Under Curves	3	≈ 10/10		60%	– none
Indefinite Integration	4	≈ 15/15		71%	– none
Physics					
Kinematics	8	≈ 11/28		41%	Q20, Q21, Q27, Q31, Q32
Current Electricity	2	≈ 3/7		68%	Q18
Electrostatics	6	≈ 11/21		45%	Q28, Q33, Q34
Chemistry					
Carboxylic Acids	4	≈ 8/14		48%	Q50, Q51
Alcohols, Phenols & Ethers	7	≈ 22/26		62%	Q46
Aldehydes & Ketones	6	≈ 20/20		83%	– none

Bar = your % of the chapter's marks | campus average

9 JEE Advanced · Maths – questions to revisit

4 questions · easy misses first
cropped from the question paper

Q14

Definite Integration
JEE Advanced · Maths · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **60%** got it right · Campus: **56%** (34 of 61)

Within reach
You: Wrong

PARAGRAPH-I

$$\int_0^a f(x)dx=\int_0^a f(a-x)dx\,;\quad \int_a^b f(x)dx=\int_a^b f(a+b-x)dx\,;\quad \int_0^{2a} f(x)dx=\int_0^a\left(f(x)+f(2a-x)\right)dx$$

14. If $\int_0^{\frac{\pi}{2}}\ln(\sin x)dx=\frac{-\pi}{q}\ln(p),(p,q\in N)$ then find the least possible value of p+q

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q7

Differential Equations – Curve from tangent-intercept ratio BP : AP = 3 : 1
JEE Advanced · Maths · Multiple correct · marking +4 / −2 · marks lost 6
Top 100: **43%** got it right · Campus: **26%** (16 of 61)

Within reach
You: Wrong

7. A curve $y=f(x)$ passes through the point (1,1) and tangent at P(x,y) cuts the x-axis and y-axis at A and B, respectively such that BP:AP=3:1, then

A) Equation of the curve is $xy^3=1$

B) Equation of normal at the point (1,1) is $x-3y+2=0$

C) Curve passes through the point $\left(\frac{1}{8},2\right)$

D) Equation of the curve is $x\frac{dy}{dx}+3y=0$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q12 Differential Equations – Solve a first-order DE and match its constants

JEE Advanced · Maths · Integer · marking +4 / 0 · marks lost 4

Top 100: **35%** got it right · Campus: **26%** (16 of 61)

Tough one

You: Wrong

12. The differential equation $\left(xy^3\left(1+\cos x\right)-y\right)dx+xdy=0$ represents the curve $\frac{x^2}{ay^2}=\frac{x^3}{b}+x^2\sin x+cx\cos x-d\sin x+k$. Then a+b+c+d is equal to (where 'k' is arbitrary constant).

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q11 Differential Equations

JEE Advanced · Maths · Integer · marking +4 / 0 · marks lost 4

Top 100: **25%** got it right · Campus: **21%** (13 of 61)

Tough one

You: Wrong

11. Let $e^{-x}\frac{d^2y}{dx^2}-\frac{dy}{dx}-y=0$ such that at the point $(0,1)$, $\frac{dy}{dx}=1$. Then the number of point of intersection of the given curve with $y=e^{e^x}$ is

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced · Physics – questions to revisit

9 questions · easy misses first
cropped from the question paper

Q21

Kinematics
JEE Advanced · Physics · Single correct · marking +3 / −1 · marks lost 4
Top 100: **64%** got it right · Campus: **51%** (31 of 61)

Within reach
You: Wrong

21. Two boats were going down stream with different velocities. When one overtook the other a plastic ball was dropped from one of the boats. Sometime later both boats turned back simultaneously and went at the same speeds as before (relative to water) towards the spot where the ball had been dropped. Which boat will reach the ball first?
A) the boat with greater velocity (relative to water)
B) the boat with lesser velocity (relative to water)
C) both will reach the ball simultaneously
D) data insufficient


Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q18

Current Electricity
JEE Advanced · Physics · Single correct · marking +3 / −1 · marks lost 4
Top 100: **57%** got it right · Campus: **49%** (30 of 61)

Within reach
You: Wrong

18. A hemi-spherical network of radius ‘a’ is made by using a conducting wire of resistance



per unit length ‘r’. The equivalent resistance across OP is:

A) $\frac{ra(\pi+3)}{8}$ B) $\frac{ra(\pi+2)}{8}$ C) $\frac{ra(\pi+4)}{8}$ D) $\frac{ra(\pi+1)}{8}$

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q28 **Electrostatics – Flux through a cube with charges at its vertices**
JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4
Top 100: **54%** got it right · Campus: **44%** (27 of 61)

Within reach
You: Wrong

28. A cube has charges each of value +Q placed at the vertices. The flux associated with the cube is $(X-1)\frac{Q}{\varepsilon_0}$, value of x is:

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q33 **Electrostatics – Flux through a ring – solid angle**
JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **42%** got it right · Campus: **38%** (23 of 61)

Within reach
You: Wrong


PARAGRAPH-II

The electric flux linked with a surface is the number of electric field lines associated with that area. Gauss law helps determine the flux

$$\oint \vec{E}.\vec{ds} = \frac{Q_{inc}}{\varepsilon_0}$$

For an open surface the flux will change depending on the solid angle. Considering this answer the following questions.

33.



The flux associated with the ring due to the charge q is given by Xq/ε_0. The value of X is approx.

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q34 **Electrostatics – Flux through a cube from a wire along one edge**
JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **33%** got it right · Campus: **23%** (14 of 61)

Tough one
You: Wrong


PARAGRAPH-II

The electric flux linked with a surface is the number of electric field lines associated with that area. Gauss law helps determine the flux

$$\oint \vec{E}.\vec{ds} = \frac{Q_{inc}}{\varepsilon_0}$$

For an open surface the flux will change depending on the solid angle. Considering this answer the following questions.

34.



A cube of edge length a contains an infinite wire of linear charge density λ at one of edges the flux associated with cube is given by $P\lambda a/\varepsilon_0$.The value of P upto two decimal places is :


Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q20 **Kinematics – Shell fired from a plateau – minimum landing distance**
JEE Advanced · Physics · Single correct · marking +3 / −1 · marks lost 3
Top 100: **29%** got it right · Campus: **23%** (14 of 61)

Tough one
You: Skipped

20. A gun is mounted on a plateau 960m away from its edge as shown. Height of plateau is 960m. The gun can fire shells with a velocity 100 m/s at any angle of the following choices. What is the minimum distance OP(=x) from the edge of plateau where shell of



gun can reach? (given $g=10m/s^2$, $\sin 74^0 = \sin 106^0 \simeq 0.96$)

A) 480 m B) 720 m C) 360 m D) 960 m

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q27

Kinematics – Projection angle that maximises area under the trajectory

JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **27%** got it right · Campus: **23%** (14 of 61)

Tough one

You: Wrong

27. A projectile is thrown with a speed 10 m/s, at an angle θ with the horizontal. For a certain value of angle of projection $\theta = \frac{(n+1)\pi}{12}$ the area bound by the trajectory with horizontal is maximum value of n is (g=10 m/s^2). Assume no airdrag.

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q31

Kinematics – Hexagon projectile – maximum height


JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **5%** got it right · Campus: **5%** (3 of 61)

Tough one

You: Wrong

PARAGRAPH-I



A regular hexagon of side a=$10\sqrt{3}$ m is shown in figure. From any point a particle is projected such that it touches B,C,D and E. The velocity of particle is v m/s. O is the centre of the hexagon. x-axis is parallel to horizontal and y-axis is perpendicular to AE and CD (g=10 m/s^2) $\left(\sqrt{3}=1.73\right)$.

31. The maximum height from the horizontal surface X' attained by the projectile is ___ m.


Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q32 **Kinematics – Hexagon projectile – speed at the top**
JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **5%** got it right · Campus: **5%** (3 of 61)

Tough one
You: Wrong

PARAGRAPH-I



A regular hexagon of side a=$10\sqrt{3}$ m is shown in figure. From any point a particle is projected such that it touches B,C,D and E. The velocity of particle is v m/s. O is the centre of the hexagon. x-axis is parallel to horizontal and y-axis is perpendicular to AE and CD (g=10 m/s^2) $\left(\sqrt{3}=1.73\right)$.

32. The velocity of projectile at maximum height is _____ m/s

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced · Chemistry – questions to revisit

3 questions · easy misses first
cropped from the question paper

Q50 **Carboxylic Acids – Molar mass of an acid from NaOH titration**
JEE Advanced · Chemistry · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **36%** got it right · Campus: **52%** (32 of 61)

Tough one
You: Wrong

PARAGRAPH-II

Carboxylic acids are organic compounds containing the carbonyl(>C=O) and hydroxyl (-OH) functional groups combined as a carboxyl group(-COOH). Due to resonance stabilization of the carboxylate anion, carboxylic acids exhibit higher acidity compared to alcohols and phenols. They participate in several characteristic reactions, including nucleophilic acyl substitution (esterification, amide formation, acid halide synthesis), decarboxylation, and reduction to primary alcohols. Furthermore, the presence of the –COOH group strongly influences physical properties, enabling extensive intermolecular hydrogen bonding that results in higher boiling points and stable dimeric structures in non-polar solvents.

50. A 0.600 g sample of a pure monoprotic carboxylic acid (R-COOH) requires 25.00 mL of 0.200 M NaOH for complete neutralization. Calculate the molar mass of the carboxylic acid?


Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q46 **Alcohols, Phenols & Ethers** – **Identifying polar aprotic solvents**
JEE Advanced · Chemistry · Integer · marking +4 / 0 · marks lost 4
Top 100: **24%** got it right · Campus: **8%** (5 of 61)

Tough one
You: Skipped

46. How many of the following are polar aprotic solvents?



[P]

Diethyl ether,


[Q]

Benzene,

[R]

Ethyl alcohol,

[S]



[T]

Petrol

[U]

Acetone

[Q]

DMSO (dimethyl sulphoxide

[W]

DMF (dimethyl formamide

[X]

formic acid

[Y]

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q51 **Carboxylic Acids – Moles of $LiAlH_4$ to reduce RCOOH to RCH_2OH**

JEE Advanced · Chemistry · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **17%** got it right · Campus: **23%** (14 of 61)

Tough one

You: Wrong

PARAGRAPH-II

Carboxylic acids are organic compounds containing the carbonyl(>C=O) and hydroxyl (-OH) functional groups combined as a carboxyl group(-COOH). Due to resonance stabilization of the carboxylate anion, carboxylic acids exhibit higher acidity compared to alcohols and phenols. They participate in several characteristic reactions, including nucleophilic acyl substitution (esterification, amide formation, acid halide synthesis), decarboxylation, and reduction to primary alcohols. Furthermore, the presence of the –COOH group strongly influences physical properties, enabling extensive intermolecular hydrogen bonding that results in higher boiling points and stable dimeric structures in non-polar solvents.

51. The number of moles of $LiAlH_4$ required theoretically to reduce eight moles of carboxylic acid to primary alcohol

$$RCOOH \xrightarrow[2.H_3O^+]{1.LiAiH_4} RCH_2OH$$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

10 How to read this report

Where the numbers come from. Marks, percentages, All-India / State / Campus ranks, subject marks, subject All-India ranks and the wrong / skipped questions are the official results in the institute's *All India Marks Analysis* (graded with the final answer key). Maximum marks leave out deleted questions — JEE Mains: Q56 — JEE Advanced: Q25. **Program rank** = rank in Pragya G10 (EXT) Achievers across all campuses, from the batch's question-wise sheet. **Top 100** = the batch's top 100 across all campuses, from the same question-wise sheet; * where it has no data for a question, the campus figure decides the priority. **Campus** = all students of this program on campus. **Priority:** Easy miss = 70%+ of the top 100 got it right; Within reach = 40–69%; Tough one = below 40%. **Marks lost** on a question = its marks plus the negative mark if answered wrong, its marks if skipped. **Weekly-test average** = your average over the weekly Mains tests (WTM) or Advanced tests (WTA) before this Part Test, from the campus averages sheet. The weekly Advanced tests are marked out of 240; their averages are scaled to 180 to match this paper (× 180/240, so 80 per subject becomes 60). Percentages are unchanged.