CLASS 10 · PART TEST-1 · STUDENT PERFORMANCE REPORT

Mancha Sagar Babu

Pragya G10 (EXT) Achievers · Adm No **235416675**

Campus HYD/Nallakunta DD Colony
JEE Mains 08 Oct 2026
JEE Advanced 09 Oct 2026

JEE Mains

08 Oct 2026 · 74 questions

213 / 296 — **72.0%**

27	20	9/65	9
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **182.2 / 300** (60.8%) → this test **72.0%** ▲ **11.2**

83 marks lost: 75 on 15 wrong answers (incl. 15 negative) · 8 on 2 skipped

JEE Advanced

09 Oct 2026 · 50 questions

107 / 176 — **60.8%**

99	68	17/61	25
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **101.0 / 180** (56.1%) → this test **60.8%** ▲ **4.7**

69 marks lost: 49 on 14 wrong answers (incl. 2 negative) · 17 on 5 skipped · 3 on partly-correct answers

1 Subject scorecard

Your marks in each subject, against the 65 students of this program on campus and against your own weekly-test average.

SUBJECT	MARKS	SCORE VS CAMPUS	SUBJECT AIR	CAMPUS AVG / BEST	WEEKLY AVG	CHANGE	RIGHT / WRONG / SKIP
JEE Mains 213 / 296 · 72.0%							
Maths	75/100	75%	34	58.2 / 100	64.8/100 64.8%	▲ 10.2	20 / 5 / 0
Physics	66/100	66%	122	53.8 / 100	59.4/100 59.4%	▲ 6.6	18 / 6 / 1
Chemistry	72/96	75%	16	55.9 / 91	58.0/100 58.0%	▲ 17.0	19 / 4 / 1
JEE Advanced 107 / 176 · 60.8%							
Maths	57/60	95% Best on campus	2	30.5 / 57	39.3/60 65.5%	▲ 29.5	16 / 1 / 0
Physics	24/56	43%	310	24.5 / 56	30.8/60 51.3%	▼ 8.4	7 / 7 / 2
Chemistry	26/60	43%	425	38.2 / 60	30.9/60 51.5%	▼ 8.1	8 / 6 / 3

Bar = your score % | campus average | best on campus Change = this test % − weekly-test average %

What this report says

1. **Maths** is your strongest JEE Mains subject: 75/100 (75%), All-India subject rank 34.
2. **Physics** needs the most work in JEE Mains: 66/100 (66%), subject rank 122; you lost 7 of its 25 questions.
3. **Maths** is your strongest JEE Advanced subject: 57/60 (95%), All-India subject rank 2 – the best score on campus.
4. **Physics** needs the most work in JEE Advanced: 24/56 (43%), subject rank 310; you lost 9 of its 16 questions.
5. Biggest fall against your weekly-test average: **JEE Advanced Physics** – 43% here against 51% on average.
6. Biggest improvement: **JEE Advanced Maths** – 95% here against 66% on average.
7. **2 questions** you lost were answered correctly by 70%+ of the batch's top 100 – about **10 marks** within easy reach (Mains Q10, Q17).
8. Negative marking cost you **15 in JEE Mains and 2 in JEE Advanced**.
9. You attempted almost everything (only 7 questions skipped across both papers), so the marks to gain are in **accuracy**: wrong answers – 15 in JEE Mains and 14 in JEE Advanced.

JEE Mains – question by question

213 / 296 · 72.0%
08 Oct 2026

2 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	2	3	4	5	6	7	8	9	**10**	11	12	13	14	15	16	**17**	**18**	19	20	21	**22**	**23**	24	25
Physics	26	**27**	28	**29**	**30**	31	32	33	34	35	36	**37**	38	39	40	41	42	43	44	45	**46**	**47**	**48**	49	50
Chemistry	51	**52**	53	54	55	~~56~~	57	58	59	60	61	**62**	**63**	64	65	66	67	68	**69**	70	71	72	73	**74**	75

3 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 65 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40-69% · **Tough one** below 40%

Maths

5 of 25 questions lost · 25 marks · 2 easy miss · 1 within reach · 2 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q10	Easy miss	**Differential Equations**	Single correct · +4 / −1	Wrong	5	76%	85%
Q17	Easy miss	**Indefinite Integration**	Single correct · +4 / −1	Wrong	5	74%	66%
Q23	Within reach	**Area Under Curves**	Numerical · +4 / −1	Wrong	5	66%	72%
Q18	Tough one	**Definite Integration** – ∫ of min{|x − 3k|} – periodic zig-zag	Single correct · +4 / −1	Wrong	5	37%	40%
Q22	Tough one	**Definite Integration** – Definite integral of a product × sum of reciprocals	Numerical · +4 / −1	Wrong	5	20%	20%

Physics

7 of 25 questions lost · 34 marks · 5 within reach · 2 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q29	Within reach	**Electrostatics**	Single correct · +4 / −1	Wrong	5	61%	48%
Q37	Within reach	**Current Electricity** – Infinite ladder – voltage across the 2nd shunt resistor	Single correct · +4 / −1	Wrong	5	52%	46%
Q48	Within reach	**Current Electricity**	Numerical · +4 / −1	Wrong	5	47%	54%
Q27	Within reach	**Electrostatics** – Potential at the centre of a thick charged shell	Single correct · +4 / −1	Wrong	5	42%	32%
Q47	Within reach	**Electrostatics**	Numerical · +4 / −1	Wrong	5	42%	34%
Q46	Tough one	**Electrostatics**	Numerical · +4 / −1	Skipped	4	25%	25%
Q30	Tough one	**Electrostatics** – Field energy inside vs outside a charged sphere	Single correct · +4 / −1	Wrong	5	17%	11%

Chemistry

5 of 24 questions lost · **24** marks · ● 2 within reach · ● 3 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q62	Within reach	**Aldehydes & Ketones**	Single correct · +4 / −1	**Skipped**	4	68%	65%
Q69	Within reach	**Aldehydes & Ketones** – Order of hydration constants of carbonyls	Single correct · +4 / −1	**Wrong**	5	52%	49%
Q63	Tough one	**Aldehydes & Ketones** – Enol-ether hydrolysis; iodoform-positive products	Single correct · +4 / −1	**Wrong**	5	32%	45%
Q52	Tough one	**Alcohols, Phenols & Ethers** – Ether vs butanol miscibility – assertion / reason	Single correct · +4 / −1	**Wrong**	5	24%	22%
Q74	Tough one	**Alcohols, Phenols & Ethers** – Site of benzylation of 2-naphthoxide	Numerical · +4 / −1	**Wrong**	5	12%	15%

4 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Differential Equations	4	11/16		84%	Q10
Definite Integration	9	26/36		29%	Q18, Q22
Area Under Curves	5	15/20		86%	Q23
Indefinite Integration	7	23/28		61%	Q17
Physics					
Electrostatics	8	8/32		27%	Q27, Q29, Q30, Q46, Q47
Current Electricity	8	22/32		57%	Q37, Q48
Kinematics	9	36/36		75%	– none
Chemistry					
Alcohols, Phenols & Ethers	10	30/40		48%	Q52, Q74
Aldehydes & Ketones	14	42/56		65%	Q62, Q63, Q69

Bar = your % of the chapter's marks | campus average

5 JEE Mains · Maths – questions to revisit

5 questions · easy misses first
cropped from the question paper

Q10 Differential Equations

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5
Top 100: **76%** got it right · Campus: **85%** (55 of 65)

Easy miss
You: Wrong

10. The solution of $\frac{xdy}{x^2+y^2}=\left(\frac{y}{x^2+y^2}-1\right)dx$ is_____ (c is arbitrary constant)

A) $y=x\cot(c-x)$ B) $y=x\tan(c-x)$ C) $y=x\cos(c-x)$ D) $y=x\sin(c-x)$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q17 Indefinite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5
Top 100: **74%** got it right · Campus: **66%** (43 of 65)

Easy miss
You: Wrong

17. $\int\left(1+x-x^{-1}\right)e^{x+x^{-1}}dx=$_____ (c is arbitrary constant)

A) $(1+x)e^{x+x^{-1}}+c$ B) $(x-1)e^{x+x^{-1}}+c$ C) $-xe^{x+x^{-1}}+c$ D) $xe^{x+x^{-1}}+c$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q23 Area Under Curves

JEE Mains · Maths · Numerical · marking +4 / −1 · marks lost 5
Top 100: **66%** got it right · Campus: **72%** (47 of 65)

Within reach
You: Wrong

23. The area bounded by the lines $y=\left||x-1|-2\right|$ with x-axis is _____ (in sq.units)

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q18

Definite Integration – ∫ of min{|x − 3k|} – periodic zig-zag

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5

Top 100: **37%** got it right · Campus: **40%** (26 of 65)

Tough one

You: Wrong

18. Let $f(x)=\min\{...,|x+6|,|x+3|,|x|,|x-3|,|x-6|,...\}$ then $\int_{-4}^{3} f(x)\,dx=$ _____

A) 4 B) $\frac{9}{2}$ C) 5 D) $\frac{11}{2}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q22

Definite Integration – Definite integral of a product × sum of reciprocals

JEE Mains · Maths · Numerical · marking +4 / −1 · marks lost 5

Top 100: **20%** got it right · Campus: **20%** (13 of 65)

Tough one

You: Wrong

22. $\int_{0}^{102}(x-1)(x-2)....(x-100)\left(\frac{1}{x-1}+\frac{1}{x-2}+...+\frac{1}{x-100}\right)dx = p(p!)$ then the number of positive integral divisors of p is equal to_____ (where, $p\in N$, $p!$ means the product of first p natural numbers).

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Mains · Physics – questions to revisit

7 questions · easy misses first
cropped from the question paper

Q29

Electrostatics
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **61%** got it right · Campus: **48%** (31 of 65)

Within reach
You: Wrong

29. A point charge q is placed on the axis of a circular disc of radius R at a distance $\frac{R}{\sqrt{3}}$ from its centre. The electric flux through the disc is

A) $\frac{q\left(2-\sqrt{3}\right)}{4\varepsilon_0}$ B) $\frac{q}{6\varepsilon_0}$ C) $\frac{q}{2\varepsilon_0}$ D) $\frac{q}{4\varepsilon_0}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q37

Current Electricity – Infinite ladder – voltage across the 2nd shunt resistor
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **52%** got it right · Campus: **46%** (30 of 65)

Within reach
You: Wrong

37. A 24 V battery of negligible internal resistance is connected to an infinite ladder network. In each section, both the top rail and the bottom rail contain a 1 Ω resistor, and a 4 Ω resistor is connected across the rails. The potential difference across the second 4 Ω resistor (counting from the battery) is

A) 12 V B) 8 V C) 6 V D) 3 V

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q48

Current Electricity


JEE Mains · Physics · Numerical · marking +4 / −1 · marks lost 5

Top 100: **47%** got it right · Campus: **54%** (35 of 65)

Within reach

You: Wrong

48. Three branches are connected between points P and Q. **Branch 1:** A 12 V battery in series



with 2Ω. **Branch 2:** A 2 V battery in series with 1 Ω. **Branch 3:** A single 2 Ω resistor. Both batteries have their positive terminals towards P. The current (in A) through the 2 Ω resistor of Branch 3 is _____.

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q27

Electrostatics – Potential at the centre of a thick charged shell

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **42%** got it right · Campus: **32%** (21 of 65)

Within reach

You: Wrong

27. A thick spherical shell of inner radius 'a' and outer radius 'b' carries a uniform volume charge density ρ. The electric potential at the centre of the shell (taking V = 0 at infinity) is

A) $\rho\frac{\left(b^2-a^2\right)}{12\varepsilon_0}$ B) $\rho\frac{\left(b^3-a^3\right)}{3\varepsilon_0 b}$ C) $\rho\frac{\left(b^2-a^2\right)}{2\varepsilon_0}$ D) $\rho\frac{\left(b^3-a^3\right)}{3\varepsilon_0 a}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q47 **Electrostatics**
JEE Mains · Physics · Numerical · marking +4 / −1 · marks lost 5
Top 100: **42%** got it right · Campus: **34%** (22 of 65)

Within reach
You: Wrong

47. A non-conducting sphere of radius R has a volume charge density $\rho = \rho_0 \left(\frac{r}{R}\right)^2$, where 'r' is the distance from the centre. The electric field at its surface is $\rho_0 \frac{R}{n\varepsilon_0}$. The value of n is _____.

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q46 **Electrostatics**
JEE Mains · Physics · Numerical · marking +4 / −1 · marks lost 4
Top 100: **25%** got it right · Campus: **25%** (16 of 65)

Tough one
You: Skipped

46. In a region the electric field is $E = \left(5x\hat{i} + 3y\hat{j} + 4\hat{k}\right) N/C$, with x, y in meters. A cube of side 1m has one vertex at the origin and its edges along the positive x, y and z axis. The net electric flux through the cube (in Nm^2/C) is _____.

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q30 **Electrostatics – Field energy inside vs outside a charged sphere**
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **17%** got it right · Campus: **11%** (7 of 65)

Tough one
You: Wrong

30. A solid sphere of radius R carries a total charge Q distributed uniformly throughout its volume. The ratio of the electrostatic field energy stored inside the sphere to that stored outside the sphere is

A) 3 : 5 B) 1 : 5 C) 1 : 3 D) 1 : 2

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed


JEE Mains · Chemistry – questions to revisit

5 questions · easy misses first
cropped from the question paper

Q62

Aldehydes & Ketones
JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 4
Top 100: **68%** got it right · Campus: **65%** (42 of 65)

Within reach
You: Skipped

62. 

Product of the reaction will be

A) Recemic mixture
B) Diastereomeric pair
C) Mesomer
D) Optically pure isomer


Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q69

Aldehydes & Ketones – Order of hydration constants of carbonyls
JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5
Top 100: **52%** got it right · Campus: **49%** (32 of 65)

Within reach
You: Wrong

69. Arrange in their increasing order of equilibrium constants for hydration?



A) A < B < C < D < E
B) A < C < B < E < D
C) A < C < E < B < D
D) C < A < B < E < D

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q63 **Aldehydes & Ketones** – Enol-ether hydrolysis; iodoform-positive products

JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5

Top 100: **32%** got it right · Campus: **45%** (29 of 65)


Tough one


You: Wrong

63. $\underset{C_5H_{10}O}{(A)} \xrightarrow{H_3O^{\oplus}} B + C; (B) and (C)$ both give +ve Iodoferm test.

Compound (A) is

A) $CH_3-CH=CH-O-CH_2-CH_3$

B) 

C) 

D) Both (B) and (C)

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q52 **Alcohols, Phenols & Ethers** – Ether vs butanol miscibility – assertion / reason

JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5

Top 100: **24%** got it right · Campus: **22%** (14 of 65)

Tough one

You: Wrong

52. **Assertion(A) :** Ethoxy ethane and butanol-1 have almost same miscibility with water.
Reason(R) : Ethoxy ethane and butanol-1 have form Hydrogen bond with water.

A) Both (A) and (R) are correct and (R) is the correct explanation of (A)

B) (A) is correct but (R) is not correct

C) Both (A) and (R) are not correct

D) Both (A) and (R) are correct and (R) is not the correct explanation of (A)

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q74 **Alcohols, Phenols & Ethers – Site of benzylation of 2-naphthoxide**


JEE Mains · Chemistry · Numerical · marking +4 / −1 · marks lost 5

Top 100: **12%** got it right · Campus: **15%** (10 of 65)

Tough one

You: Wrong

74. Which site is predominately benzylated in the given reaction?



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced – question by question

107 / 176 · 60.8%
09 Oct 2026

6 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
Physics	18	19	20	21	22	23	24	~~25~~	26	27	28	29	30	31	32	33	34
Chemistry	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51

7 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 61 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40–69% · **Tough one** below 40%

Maths

1 of 17 questions lost · **3** marks · 1 within reach

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q15	Within reach	Definite Integration	Paragraph · +3 / 0	Wrong	3	41%	39%

Physics

9 of 16 questions lost · **32** marks · 5 within reach · 4 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q21	Within reach	Kinematics	Single correct · +3 / −1	Wrong	4	64%	51%
Q18	Within reach	Current Electricity	Single correct · +3 / −1	Wrong	4	57%	49%
Q28	Within reach	**Electrostatics** – Flux through a cube with charges at its vertices	Integer · +4 / 0	Wrong	4	54%	44%
Q29	Within reach	Electrostatics	Integer · +4 / 0	Wrong	4	54%	34%
Q33	Within reach	**Electrostatics** – Flux through a ring – solid angle	Paragraph · +3 / 0	Skipped	3	42%	38%
Q34	Tough one	**Electrostatics** – Flux through a cube from a wire along one edge	Paragraph · +3 / 0	Skipped	3	33%	23%
Q27	Tough one	**Kinematics** – Projection angle that maximises area under the trajectory	Integer · +4 / 0	Wrong	4	27%	23%
Q31	Tough one	**Kinematics** – Hexagon projectile – maximum height	Paragraph · +3 / 0	Wrong	3	5%	5%
Q32	Tough one	**Kinematics** – Hexagon projectile – speed at the top	Paragraph · +3 / 0	Wrong	3	5%	5%

Chemistry

9 of 17 questions lost · 31 marks (+3 on partly-correct answers) · 5 within reach · 4 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q37	Within reach	Alcohols, Phenols & Ethers	Single correct · +3 / −1	Skipped	3	68%	75%
Q44	Within reach	Carboxylic Acids	Integer · +4 / 0	Skipped	4	68%	80%
Q39	Within reach	Alcohols, Phenols & Ethers	Multiple correct · +4 / −2	Skipped	4	66%	77%
Q48	Within reach	Aldehydes & Ketones	Paragraph · +3 / 0	Wrong	3	66%	77%
Q49	Within reach	Aldehydes & Ketones	Paragraph · +3 / 0	Wrong	3	59%	70%
Q50	Tough one	Carboxylic Acids – Molar mass of an acid from NaOH titration	Paragraph · +3 / 0	Wrong	3	36%	52%
Q47	Tough one	Carboxylic Acids	Integer · +4 / 0	Wrong	4	32%	33%
Q46	Tough one	Alcohols, Phenols & Ethers – Identifying polar aprotic solvents	Integer · +4 / 0	Wrong	4	24%	8%
Q51	Tough one	Carboxylic Acids – Moles of $LiAlH_4$ to reduce RCOOH to RCH_2OH	Paragraph · +3 / 0	Wrong	3	17%	23%

8 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject. Chapter marks are approximate (≈): partly-correct answers cannot be traced to a question.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Definite Integration	5	≈ 14/17		54%	Q15
Area Under Curves	3	≈ 10/10		60%	– none
Differential Equations	5	≈ 18/18		30%	– none
Indefinite Integration	4	≈ 15/15		71%	– none
Physics					
Electrostatics	6	≈ 7/21		45%	Q28, Q29, Q33, Q34
Current Electricity	2	≈ 3/7		68%	Q18
Kinematics	8	≈ 14/28		41%	Q21, Q27, Q31, Q32
Chemistry					
Carboxylic Acids	4	≈ 0/14		48%	Q44, Q47, Q50, Q51
Alcohols, Phenols & Ethers	7	≈ 15/26		62%	Q37, Q39, Q46
Aldehydes & Ketones	6	≈ 14/20		83%	Q48, Q49

Bar = your % of the chapter's marks | campus average

9 JEE Advanced · Maths – questions to revisit

1 question · easy misses first
cropped from the question paper

Q15 Definite Integration

JEE Advanced · Maths · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **41%** got it right · Campus: **39%** (24 of 61)

Within reach
You: Wrong

PARAGRAPH-I

$$\int_0^a f(x)dx = \int_0^a f(a-x)dx\,; \quad \int_a^b f(x)dx = \int_a^b f(a+b-x)dx\,; \quad \int_0^{2a} f(x)dx = \int_0^a \left(f(x)+f(2a-x)\right)dx$$

15. If $\left|\int_0^{\frac{\pi}{2}} \ln(1+\cos x)\,dx\right| = \frac{\pi}{q}\ln(p), (p,q \in N)$ then find the least possible value p+q

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced · Physics – questions to revisit

9 questions · easy misses first
cropped from the question paper

Q21

Kinematics
JEE Advanced · Physics · Single correct · marking +3 / −1 · marks lost 4
Top 100: **64%** got it right · Campus: **51%** (31 of 61)

Within reach
You: Wrong

21. Two boats were going down stream with different velocities. When one overtook the other a plastic ball was dropped from one of the boats. Sometime later both boats turned back simultaneously and went at the same speeds as before (relative to water) towards the spot where the ball had been dropped. Which boat will reach the ball first?
A) the boat with greater velocity (relative to water)
B) the boat with lesser velocity (relative to water)
C) both will reach the ball simultaneously
D) data insufficient


Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q18

Current Electricity
JEE Advanced · Physics · Single correct · marking +3 / −1 · marks lost 4
Top 100: **57%** got it right · Campus: **49%** (30 of 61)

Within reach
You: Wrong

18. A hemi-spherical network of radius ‘a’ is made by using a conducting wire of resistance



per unit length ‘r’. The equivalent resistance across OP is:

A) $\frac{ra(\pi+3)}{8}$ B) $\frac{ra(\pi+2)}{8}$ C) $\frac{ra(\pi+4)}{8}$ D) $\frac{ra(\pi+1)}{8}$

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q28

Electrostatics – Flux through a cube with charges at its vertices

JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **54%** got it right · Campus: **44%** (27 of 61)

Within reach

You: Wrong

28. A cube has charges each of value +Q placed at the vertices. The flux associated with the cube is $(X-1)\frac{Q}{\varepsilon_0}$, value of x is:

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q29

Electrostatics


JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **54%** got it right · Campus: **34%** (21 of 61)

Within reach

You: Wrong

29. An electron is projected from a distance d and with initial velocity u parallel to a uniformly charge flat conducting plate as shown. It strikes the plate after traveling a distance l along the direction of projection. The surface charge density of the conducting plate is $\sigma = \frac{nd\varepsilon_0 mu^2}{el^2}$, The value of n is:



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q33 **Electrostatics – Flux through a ring – solid angle**
JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **42%** got it right · Campus: **38%** (23 of 61)

Within reach
You: Skipped


PARAGRAPH-II

The electric flux linked with a surface is the number of electric field lines associated with that area. Gauss law helps determine the flux

$$\oint \vec{E}.\vec{ds} = \frac{Q_{inc}}{\varepsilon_0}$$

For an open surface the flux will change depending on the solid angle. Considering this answer the following questions.

33.



The flux associated with the ring due to the charge q is given by Xq/ε_0. The value of X is approx.

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q34

Electrostatics – Flux through a cube from a wire along one edge

JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **33%** got it right · Campus: **23%** (14 of 61)

Tough one

You: Skipped


PARAGRAPH-II

The electric flux linked with a surface is the number of electric field lines associated with that area. Gauss law helps determine the flux

$$\oint \vec{E}.\vec{ds} = \frac{Q_{inc}}{\varepsilon_0}$$

For an open surface the flux will change depending on the solid angle. Considering this answer the following questions.

34.



A cube of edge length a contains an infinite wire of linear charge density λ at one of edges the flux associated with cube is given by $P\lambda a/\varepsilon_0$.The value of P upto two decimal places is :

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q27

Kinematics – Projection angle that maximises area under the trajectory

JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **27%** got it right · Campus: **23%** (14 of 61)

Tough one

You: Wrong

27. A projectile is thrown with a speed 10 m/s, at an angle θ with the horizontal. For a certain value of angle of projection $\theta = \frac{(n+1)\pi}{12}$ the area bound by the trajectory with horizontal is maximum value of n is (g=10 m/s^2). Assume no airdrag.


Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q31 **Kinematics – Hexagon projectile – maximum height**
JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **5%** got it right · Campus: **5%** (3 of 61)

Tough one
You: Wrong

PARAGRAPH-I



A regular hexagon of side a=$10\sqrt{3}$ m is shown in figure. From any point a particle is projected such that it touches B,C,D and E. The velocity of particle is v m/s. O is the centre of the hexagon. x-axis is parallel to horizontal and y-axis is perpendicular to AE and CD (g=10 m/s^2) $\left(\sqrt{3}=1.73\right)$.

31. The maximum height from the horizontal surface X' attained by the projectile is ___ m.


Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q32 **Kinematics – Hexagon projectile – speed at the top**
JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **5%** got it right · Campus: **5%** (3 of 61)

Tough one
You: Wrong

PARAGRAPH-I



A regular hexagon of side a=$10\sqrt{3}$ m is shown in figure. From any point a particle is projected such that it touches B,C,D and E. The velocity of particle is v m/s. O is the centre of the hexagon. x-axis is parallel to horizontal and y-axis is perpendicular to AE and CD (g=10 m/s^2) $\left(\sqrt{3}=1.73\right)$.

32. The velocity of projectile at maximum height is _____ m/s

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed


JEE Advanced · Chemistry – questions to revisit

9 questions · easy misses first
cropped from the question paper

Q37 Alcohols, Phenols & Ethers

JEE Advanced · Chemistry · Single correct · marking +3 / −1 · marks lost 3
Top 100: **68%** got it right · Campus: **75%** (46 of 61)

Within reach
You: Skipped



37. How many litres of gas 'P' is formed in above reaction in NTP. (molar volume of gas at NTP is 22.4 L)

A)22.4 L B)33.6 L C)44.8 L D)66 L

Why did I lose this question? (write the reason and the correct approach)


☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q44 Carboxylic Acids

JEE Advanced · Chemistry · Integer · marking +4 / 0 · marks lost 4
Top 100: **68%** got it right · Campus: **80%** (49 of 61)

Within reach
You: Skipped

44. How many of the following acids will show higher reactivity towards esterification reaction as compared to acetic acid?



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q39 Alcohols, Phenols & Ethers


JEE Advanced · Chemistry · Multiple correct · marking +4 / −2 · marks lost 4

Top 100: **66%** got it right · Campus: **77%** (47 of 61)

Within reach

You: Skipped

39.



Major product (Y) Major product (X)

Select the correct statement(s) regarding above reactions?

A) 'X' is O-alkylated product and 'Y' is C-alkylated product

B) Both X and Y are C-alkylated products only.

C) Mixture of X and Y can be separated dil NaOH

D) Product 'Y' can react with sodium metal.

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q48 Aldehydes & Ketones

JEE Advanced · Chemistry · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **66%** got it right · Campus: **77%** (47 of 61)

Within reach

You: Wrong

PARAGRAPH-I

When a carbonyl compound reacts with hydroxylamine(NH_2OH), it undergoes nucleophilic addition followed by dehydration to form an oxime (R_2C=N-OH) and water. The rate of oxime formation is maximum at an optimum pH≈4.5. At lower pH, the hydroxylamine gets protonated into NH_3^+OH, losing its nucleophilicity, whereas at higher pH, protonation of the carbonyl oxygen is insufficient to activate the carbonyl group.

48. If 5.8g of acetone (M=58 g mol^{-1}) reacts completely with an excess of hydroxylamine, Calculate the mass of acetoxime (M=73 g mol^{-1}) formed.

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q49 **Aldehydes & Ketones**
JEE Advanced · Chemistry · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **59%** got it right · Campus: **70%** (43 of 61)

Within reach
You: Wrong

PARAGRAPH-I

When a carbonyl compound reacts with hydroxylamine(NH_2OH), it undergoes nucleophilic addition followed by dehydration to form an oxime ($R_2C=N\text{-}OH$) and water. The rate of oxime formation is maximum at an optimum pH≈4.5. At lower pH, the hydroxylamine gets protonated into NH_3^+OH, losing its nucleophilicity, whereas at higher pH, protonation of the carbonyl oxygen is insufficient to activate the carbonyl group.

49. In a reaction vessel, 0.1 mol of benzaldehyde reacts completely with 0.1 mol of hydroxylamine to form benzaldoxime. Calculate the total volume of water (H_2O) vapor produced in liters at STP (1 mol of gas at STP = 22.4 L).

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q50

Carboxylic Acids – Molar mass of an acid from NaOH titration

JEE Advanced · Chemistry · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **36%** got it right · Campus: **52%** (32 of 61)

Tough one

You: Wrong

PARAGRAPH-II

Carboxylic acids are organic compounds containing the carbonyl(>C=O) and hydroxyl (-OH) functional groups combined as a carboxyl group(-COOH). Due to resonance stabilization of the carboxylate anion, carboxylic acids exhibit higher acidity compared to alcohols and phenols. They participate in several characteristic reactions, including nucleophilic acyl substitution (esterification, amide formation, acid halide synthesis), decarboxylation, and reduction to primary alcohols. Furthermore, the presence of the –COOH group strongly influences physical properties, enabling extensive intermolecular hydrogen bonding that results in higher boiling points and stable dimeric structures in non-polar solvents.

50. A 0.600 g sample of a pure monoprotic carboxylic acid (R-COOH) requires 25.00 mL of 0.200 M NaOH for complete neutralization. Calculate the molar mass of the carboxylic acid?

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q47

Carboxylic Acids


JEE Advanced · Chemistry · Integer · marking +4 / 0 · marks lost 4


Top 100: **32%** got it right · Campus: **33%** (20 of 61)


Tough one


You: Wrong

47. At room temperature how many of the following compounds can decompose $NaHCO_3$ to liberate CO_2?

a) 

b) 

c) 

d) 


Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q46 **Alcohols, Phenols & Ethers** – **Identifying polar aprotic solvents**
JEE Advanced · Chemistry · Integer · marking +4 / 0 · marks lost 4
Top 100: **24%** got it right · Campus: **8%** (5 of 61)

Tough one
You: Wrong

46. How many of the following are polar aprotic solvents?



[P]

Diethyl ether,


[Q]

Benzene,

[R]

Ethyl alcohol,

[S]



[T]

Petrol

[U]

Acetone

[Q]

DMSO (dimethyl sulphoxide

[W]

DMF (dimethyl formamide

[X]

formic acid

[Y]

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q51 **Carboxylic Acids – Moles of $LiAlH_4$ to reduce RCOOH to RCH_2OH**

JEE Advanced · Chemistry · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **17%** got it right · Campus: **23%** (14 of 61)

Tough one

You: Wrong

PARAGRAPH-II

Carboxylic acids are organic compounds containing the carbonyl(>C=O) and hydroxyl (-OH) functional groups combined as a carboxyl group(-COOH). Due to resonance stabilization of the carboxylate anion, carboxylic acids exhibit higher acidity compared to alcohols and phenols. They participate in several characteristic reactions, including nucleophilic acyl substitution (esterification, amide formation, acid halide synthesis), decarboxylation, and reduction to primary alcohols. Furthermore, the presence of the –COOH group strongly influences physical properties, enabling extensive intermolecular hydrogen bonding that results in higher boiling points and stable dimeric structures in non-polar solvents.

51. The number of moles of $LiAlH_4$ required theoretically to reduce eight moles of carboxylic acid to primary alcohol

$$RCOOH \xrightarrow[2.H_3O^+]{1.LiAiH_4} RCH_2OH$$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

10 How to read this report

Where the numbers come from. Marks, percentages, All-India / State / Campus ranks, subject marks, subject All-India ranks and the wrong / skipped questions are the official results in the institute's *All India Marks Analysis* (graded with the final answer key). Maximum marks leave out deleted questions — JEE Mains: Q56 — JEE Advanced: Q25. **Program rank** = rank in Pragya G10 (EXT) Achievers across all campuses, from the batch's question-wise sheet. **Top 100** = the batch's top 100 across all campuses, from the same question-wise sheet; * where it has no data for a question, the campus figure decides the priority. **Campus** = all students of this program on campus. **Priority:** Easy miss = 70%+ of the top 100 got it right; Within reach = 40–69%; Tough one = below 40%. **Marks lost** on a question = its marks plus the negative mark if answered wrong, its marks if skipped. In JEE Advanced, 3 marks went on multiple-correct answers that were only partly correct. **Weekly-test average** = your average over the weekly Mains tests (WTM) or Advanced tests (WTA) before this Part Test, from the campus averages sheet. The weekly Advanced tests are marked out of 240; their averages are scaled to 180 to match this paper (× 180/240, so 80 per subject becomes 60). Percentages are unchanged.