CLASS 10 · PART TEST-1 · STUDENT PERFORMANCE REPORT

Ragi Tanmay Sai

Pragya G10 (EXT) Achievers · Adm No **235416398**

Campus HYD/Nallakunta DD Colony
JEE Mains 08 Oct 2026
JEE Advanced 09 Oct 2026

JEE Mains

08 Oct 2026 · 74 questions

248 / 296 — **83.8%**

7	5	2/65	2
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **228.8 / 300** (76.2%) → this test **83.8%** ▲ **7.5**

48 marks lost: 20 on 4 wrong answers (incl. 4 negative) · 28 on 7 skipped

JEE Advanced

09 Oct 2026 · 50 questions

111 / 176 — **63.1%**

82	56	13/61	20
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **107.7 / 180** (59.9%) → this test **63.1%** ▲ **3.2**

65 marks lost: 42 on 11 wrong answers (incl. 3 negative) · 21 on 6 skipped · 2 on partly-correct answers

1 Subject scorecard

Your marks in each subject, against the 65 students of this program on campus and against your own weekly-test average.

SUBJECT	MARKS	SCORE VS CAMPUS	SUBJECT AIR	CAMPUS AVG / BEST	WEEKLY AVG	CHANGE	RIGHT / WRONG / SKIP
JEE Mains 248 / 296 · 83.8%							
Maths	88/100	88%	7	58.2 / 100	80.2/100 80.2%	▲ 7.8	22 / 0 / 3
Physics	87/100	87%	16	53.8 / 100	76.0/100 76.0%	▲ 11.0	22 / 1 / 2
Chemistry	73/96	76%	13	55.9 / 91	72.7/100 72.7%	▲ 3.4	19 / 3 / 2
JEE Advanced 111 / 176 · 63.1%							
Maths	42/60	70%	78	30.5 / 57	41.2/60 68.7%	▲ 1.3	12 / 4 / 1
Physics	22/56	39%	373	24.5 / 56	33.5/60 55.9%	▼ 16.6	7 / 5 / 4
Chemistry	47/60	78%	51	38.2 / 60	33.0/60 54.9%	▲ 23.4	14 / 2 / 1

Bar = your score % | campus average | best on campus Change = this test % − weekly-test average %

What this report says

1. **Maths** is your strongest JEE Mains subject: 88/100 (88%), All-India subject rank 7.
2. **Chemistry** needs the most work in JEE Mains: 73/96 (76%), subject rank 13; you lost 5 of its 24 questions.
3. **Chemistry** is your strongest JEE Advanced subject: 47/60 (78%), All-India subject rank 51.
4. **Physics** needs the most work in JEE Advanced: 22/56 (39%), subject rank 373; you lost 9 of its 16 questions.
5. Biggest fall against your weekly-test average: **JEE Advanced Physics** – 39% here against 56% on average.
6. Biggest improvement: **JEE Advanced Chemistry** – 78% here against 55% on average.
7. **3 questions** you lost were answered correctly by 70%+ of the batch's top 100 – about **14 marks** within easy reach (Mains Q64 · Advanced Q9, Q23).
8. Negative marking cost you **4 in JEE Mains and 3 in JEE Advanced**.
9. You left **7 in JEE Mains and 6 in JEE Advanced** questions blank, and answered 4 in JEE Mains and 11 in JEE Advanced wrong – both are marks to recover.

JEE Mains – question by question

248 / 296 · 83.8%
08 Oct 2026

2 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25
Physics	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50
Chemistry	51	52	53	54	55	~~56~~	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75

3 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 65 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40-69% · **Tough one** below 40%

Maths

3 of 25 questions lost · **12** marks · ● 3 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q11	Tough one	**Definite Integration**	Single correct · +4 / −1	Skipped	4	29%	26%
Q20	Tough one	**Definite Integration** – King's property with f(x) + f(y) = f(xy)	Single correct · +4 / −1	Skipped	4	29%	32%
Q16	Tough one	**Definite Integration** – $\int_0^{100}$ ([tan⁻¹x] + [cot⁻¹x]) dx with GIF	Single correct · +4 / −1	Skipped	4	16%	11%

Physics

3 of 25 questions lost · **13** marks · ● 1 within reach · ● 2 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q34	Within reach	**Current Electricity**	Single correct · +4 / −1	Skipped	4	45%	37%
Q46	Tough one	**Electrostatics**	Numerical · +4 / −1	Wrong	5	25%	25%
Q30	Tough one	**Electrostatics** – Field energy inside vs outside a charged sphere	Single correct · +4 / −1	Skipped	4	17%	11%



Chemistry

5 of 24 questions lost · **23** marks · ● 1 easy miss · ● 1 within reach · ● 3 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q64	Easy miss	**Aldehydes & Ketones**	Single correct · +4 / −1	Skipped	4	77%	74%
Q53	Within reach	**Alcohols, Phenols & Ethers**	Single correct · +4 / −1	Wrong	5	68%	69%
Q75	Tough one	**Aldehydes & Ketones**	Numerical · +4 / −1	Wrong	5	27%	38%
Q52	Tough one	**Alcohols, Phenols & Ethers** – Ether vs butanol miscibility – assertion / reason	Single correct · +4 / −1	Wrong	5	24%	22%
Q74	Tough one	**Alcohols, Phenols & Ethers** – Site of benzylation of 2-naphthoxide	Numerical · +4 / −1	Skipped	4	12%	15%

4 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Definite Integration	9	**24**/36		29%	**Q11**, **Q16**, **Q20**
Area Under Curves	5	**20**/20		86%	– none
Differential Equations	4	**16**/16		84%	– none
Indefinite Integration	7	**28**/28		61%	– none
Physics					
Electrostatics	8	**23**/32		27%	**Q30**, **Q46**
Current Electricity	8	**28**/32		57%	**Q34**
Kinematics	9	**36**/36		75%	– none
Chemistry					
Alcohols, Phenols & Ethers	10	**26**/40		48%	**Q52**, **Q53**, **Q74**
Aldehydes & Ketones	14	**47**/56		65%	**Q64**, **Q75**

Bar = your % of the chapter's marks | campus average

5 JEE Mains · Maths – questions to revisit

3 questions · easy misses first
cropped from the question paper

Q11 — Definite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4
Top 100: **29%** got it right · Campus: **26%** (17 of 65)

Tough one
You: Skipped

11. Let $f(x)=\sin x+\sum_{r=0}^{2}\cos\left(x+\frac{2r\pi}{3}\right)$, $g(x)=\cos x+\sum_{r=0}^{2}\sin\left(x+\frac{2r\pi}{3}\right)$ then $\int_{0}^{\frac{\pi}{2}} f(g(x))f(x)\,dx=$

A) $1+\sin 1$ B) $1-\sin 1$ C) $1+\cos 1$ D) $1-\cos 1$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q20 — Definite Integration – King's property with f(x) + f(y) = f(xy)

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4
Top 100: **29%** got it right · Campus: **32%** (21 of 65)

Tough one
You: Skipped

20. Let $f:(0,\infty)\to R$ be a continuous function, satisfying $f(x)+f(y)=f(xy)$. Then the value of $\int_{\sqrt{f(e)}}^{\sqrt{f(e^2)}}\frac{x\tan x^2}{\tan x^2+\tan\left(f(e^3)-x^2\right)}dx$ is equal to

A) $\frac{1}{2}f(e^2)$ B) $\frac{1}{4}f(e^2)$ C) $\frac{1}{2}f(e)$ D) $\frac{f(e)}{4}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q16 — Definite Integration – $\int_0^{100}([\tan^{-1}x]+[\cot^{-1}x])\,dx$ with GIF

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4
Top 100: **16%** got it right · Campus: **11%** (7 of 65)

Tough one
You: Skipped

16. If $\int_{0}^{100}\left(\left[\tan^{-1}x\right]+\left[\cot^{-1}x\right]\right)dx=\ldots$ [.] denotes the greatest integer function

A) $100-\cot 1$ B) $100+2\cot 1$ C) $100-2\cot 2$ D) $100+2\cot 2$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Mains · Physics – questions to revisit

3 questions · easy misses first
cropped from the question paper

Q34 — Current Electricity

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4
Top 100: **45%** got it right · Campus: **37%** (24 of 65)

Within reach
You: Skipped

34. In a Wheatstone bridge ABCD, a 12 V battery of negligible internal resistance is connected between A and C. The arms are AB = 2 Ω, BC = 3 Ω, AD = 3 Ω and DC = 3 Ω. An ammeter of resistance 0.3 Ω is connected between B and D. The current through the ammeter is

A) 0.2 A B) 0.6 A C) 0.4 A D) 1.2 A

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q46 — Electrostatics

JEE Mains · Physics · Numerical · marking +4 / −1 · marks lost 5
Top 100: **25%** got it right · Campus: **25%** (16 of 65)

Tough one
You: Wrong

46. In a region the electric field is $E = \left(5x\hat{i} + 3y\hat{j} + 4\hat{k}\right) N/C$, with x, y in meters. A cube of side 1m has one vertex at the origin and its edges along the positive x, y and z axis. The net electric flux through the cube (in Nm^2/C) is _____.

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q30 — Electrostatics – Field energy inside vs outside a charged sphere

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4
Top 100: **17%** got it right · Campus: **11%** (7 of 65)

Tough one
You: Skipped

30. A solid sphere of radius R carries a total charge Q distributed uniformly throughout its volume. The ratio of the electrostatic field energy stored inside the sphere to that stored outside the sphere is

A) 3 : 5 B) 1 : 5 C) 1 : 3 D) 1 : 2

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed


JEE Mains · Chemistry – questions to revisit

5 questions · easy misses first
cropped from the question paper

Q64

Aldehydes & Ketones
JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 4
Top 100: **77%** got it right · Campus: **74%** (48 of 65)

Easy miss
You: Skipped

64. 

CH_3MgBr / H^+ (1) $KMnO_4(colddil)$ (2) CrO_3 (3) H^+ / Δ (4)

For the above conversion the correct order of reagents used is :

A) $1 \rightarrow 2 \rightarrow 3 \rightarrow 4$ B) $1 \rightarrow 4 \rightarrow 3 \rightarrow 2$ C) $1 \rightarrow 4 \rightarrow 2 \rightarrow 3$ D) $2 \rightarrow 3 \rightarrow 4 \rightarrow 1$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q53

Alcohols, Phenols & Ethers
JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5
Top 100: **68%** got it right · Campus: **69%** (45 of 65)

Within reach
You: Wrong

53. Phenol industrially prepared from

A) Benzene sulphonic acid B) Cumene
C) Benzene diazonium chloride D) Chlorobenzene


Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q75

Aldehydes & Ketones
JEE Mains · Chemistry · Numerical · marking +4 / −1 · marks lost 5
Top 100: **27%** got it right · Campus: **38%** (25 of 65)

Tough one
You: Wrong

75. 

Number of possible product(S) including stereoisomerism

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q52

Alcohols, Phenols & Ethers – Ether vs butanol miscibility – assertion / reason
JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5
Top 100: **24%** got it right · Campus: **22%** (14 of 65)

Tough one
You: Wrong

52. **Assertion(A) :** Ethoxy ethane and butanol-1 have almost same miscibility with water.
Reason(R) : Ethoxy ethane and butanol-1 have form Hydrogen bond with water.
A) Both (A) and (R) are correct and (R) is the correct explanation of (A)
B) (A) is correct but (R) is not correct
C) Both (A) and (R) are not correct
D) Both (A) and (R) are correct and (R) is not the correct explanation of (A)


Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q74

Alcohols, Phenols & Ethers – Site of benzylation of 2-naphthoxide
JEE Mains · Chemistry · Numerical · marking +4 / −1 · marks lost 4
Top 100: **12%** got it right · Campus: **15%** (10 of 65)

Tough one
You: Skipped

74. Which site is predominately benzylated in the given reaction?



Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced – question by question

111 / 176 · 63.1%
09 Oct 2026

6 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
Physics	18	19	20	21	22	23	24	~~25~~	26	27	28	29	30	31	32	33	34
Chemistry	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51

7 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 61 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40–69% · **Tough one** below 40%

Maths

5 of 17 questions lost · 18 marks · 1 easy miss · 4 within reach

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q9	Easy miss	Indefinite Integration	Integer · +4 / 0	Wrong	4	71%	67%
Q1	Within reach	Indefinite Integration	Single correct · +3 / −1	Wrong	4	66%	70%
Q13	Within reach	Definite Integration	Integer · +4 / 0	Skipped	4	64%	70%
Q14	Within reach	Definite Integration	Paragraph · +3 / 0	Wrong	3	60%	56%
Q15	Within reach	Definite Integration	Paragraph · +3 / 0	Wrong	3	41%	39%

Physics

9 of 16 questions lost · 34 marks · 1 easy miss · 4 within reach · 4 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q23	Easy miss	Electrostatics	Multiple correct · +4 / −2	Wrong	6	88%	84%
Q21	Within reach	Kinematics	Single correct · +3 / −1	Skipped	3	64%	51%
Q26	Within reach	Kinematics	Integer · +4 / 0	Skipped	4	59%	67%
Q28	Within reach	**Electrostatics** – Flux through a cube with charges at its vertices	Integer · +4 / 0	Skipped	4	54%	44%
Q29	Within reach	Electrostatics	Integer · +4 / 0	Wrong	4	54%	34%
Q20	Tough one	**Kinematics** – Shell fired from a plateau – minimum landing distance	Single correct · +3 / −1	Skipped	3	29%	23%
Q27	Tough one	**Kinematics** – Projection angle that maximises area under the trajectory	Integer · +4 / 0	Wrong	4	27%	23%
Q31	Tough one	**Kinematics** – Hexagon projectile – maximum height	Paragraph · +3 / 0	Wrong	3	5%	5%
Q32	Tough one	**Kinematics** – Hexagon projectile – speed at the top	Paragraph · +3 / 0	Wrong	3	5%	5%



Chemistry

3 of 17 questions lost · 11 marks (+2 on partly-correct answers) · 3 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q47	Tough one	Carboxylic Acids	Integer · +4 / 0	Wrong	4	32%	33%
Q46	Tough one	**Alcohols, Phenols & Ethers** – Identifying polar aprotic solvents	Integer · +4 / 0	Wrong	4	24%	8%
Q51	Tough one	**Carboxylic Acids** – Moles of $LiAlH_4$ to reduce RCOOH to RCH_2OH	Paragraph · +3 / 0	Skipped	3	17%	23%

8 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject. Chapter marks are approximate (≈): partly-correct answers cannot be traced to a question.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Definite Integration	5	≈ **7**/17		54%	**Q13**, **Q14**, **Q15**
Indefinite Integration	4	≈ **7**/15		71%	**Q1**, **Q9**
Area Under Curves	3	≈ **10**/10		60%	– none
Differential Equations	5	≈ **18**/18		30%	– none
Physics					
Kinematics	8	≈ **8**/28		41%	**Q20**, **Q21**, **Q26**, **Q27**, **Q31**, **Q32**
Electrostatics	6	≈ **7**/21		45%	**Q23**, **Q28**, **Q29**
Current Electricity	2	≈ **7**/7		68%	– none
Chemistry					
Carboxylic Acids	4	≈ **7**/14		48%	**Q47**, **Q51**
Alcohols, Phenols & Ethers	7	≈ **22**/26		62%	**Q46**
Aldehydes & Ketones	6	≈ **20**/20		83%	– none

Bar = your % of the chapter's marks | campus average

9 JEE Advanced · Maths – questions to revisit

5 questions · easy misses first
cropped from the question paper

Q9 Indefinite Integration

JEE Advanced · Maths · Integer · marking +4 / 0 · marks lost 4
Top 100: **71%** got it right · Campus: **67%** (41 of 61)

Easy miss
You: Wrong

9. If the primitive of the function $f(x)=\frac{\sin^4 x}{\cos^2 x}$ with respect to x is tan(x)+g(x)+C where g(0)=0 and C is indefinite integration constant, then the value of $8\left(\frac{\pi}{8}+g\left(\frac{\pi}{12}\right)\right)$ is equal to

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q1 Indefinite Integration

JEE Advanced · Maths · Single correct · marking +3 / −1 · marks lost 4
Top 100: **66%** got it right · Campus: **70%** (43 of 61)

Within reach
You: Wrong

1. $\int \frac{2e^{2x}+e^x}{3e^{2x}-2e^x-1}dx$ is equal to

A) $\frac{1}{4}\ln\left|\frac{(e^x-1)^{1/3}}{(3e^x+1)^3}\right|+C$

B) $\frac{1}{4}\ln\left|\frac{(e^x-1)^3}{(3e^x+1)^{1/3}}\right|+C$

C) $\frac{1}{3}\ln\left|\frac{(e^x-1)^4}{(3e^x+1)^{1/4}}\right|+C$

D) $\frac{1}{3}\ln\left|\frac{(e^x-1)^{1/4}}{(3e^x+1)^4}\right|+C$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q13 Definite Integration

JEE Advanced · Maths · Integer · marking +4 / 0 · marks lost 4
Top 100: **64%** got it right · Campus: **70%** (43 of 61)

Within reach
You: Skipped

13. Let $t(x)=\sqrt{1+x\sqrt{1+(x+1)\sqrt{1+(x+2)(x+4)}}}$ then $\int_0^{100} t(x)dx$ is K. Find $\frac{K}{1700}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q14 Definite Integration

JEE Advanced · Maths · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **60%** got it right · Campus: **56%** (34 of 61)

Within reach

You: Wrong

PARAGRAPH-I

$$\int_0^a f(x)dx = \int_0^a f(a-x)dx\,; \quad \int_a^b f(x)dx = \int_a^b f(a+b-x)dx\,; \quad \int_0^{2a} f(x)dx = \int_0^a \left(f(x)+f(2a-x)\right)dx$$

14. If $\int_0^{\frac{\pi}{2}} \ln(\sin x)dx = \frac{-\pi}{q}\ln(p), (p,q \in N)$ then find the least possible value of p+q

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q15 Definite Integration

JEE Advanced · Maths · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **41%** got it right · Campus: **39%** (24 of 61)

Within reach

You: Wrong

PARAGRAPH-I

$$\int_0^a f(x)dx = \int_0^a f(a-x)dx\,; \quad \int_a^b f(x)dx = \int_a^b f(a+b-x)dx\,; \quad \int_0^{2a} f(x)dx = \int_0^a \left(f(x)+f(2a-x)\right)dx$$

15. If $\left|\int_0^{\frac{\pi}{\pi}} \ln(1+\cos x)dx\right| = \frac{\pi}{q}\ln(p), (p,q \in N)$ then find the least possible value p+q

Why did I lose this question? (write the reason and the correct approach)


☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q23 Electrostatics

JEE Advanced · Physics · Multiple correct · marking +4 / −2 · marks lost 6
Top 100: **88%** got it right · Campus: **84%** (51 of 61)

Easy miss
You: Wrong

23. Two large thin conducting plates with small gap in between are placed in a uniform electric field 'E' (perpendicular to plates). Area of each plate is A and charges +Q and –Q are given to these plates as shown in figure. If points R,S and T as shown in the figure are



three points in space, then the

A) field point R is E

B) field at point S is E

C) field at point T is $E+\frac{Q}{\varepsilon_0 A}$

D) field at point S is $E+\frac{Q}{\varepsilon_0 A}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q21 Kinematics

JEE Advanced · Physics · Single correct · marking +3 / −1 · marks lost 3
Top 100: **64%** got it right · Campus: **51%** (31 of 61)

Within reach
You: Skipped

21. Two boats were going down stream with different velocities. When one overtook the other a plastic ball was dropped from one of the boats. Sometime later both boats turned back simultaneously and went at the same speeds as before (relative to water) towards the spot where the ball had been dropped. Which boat will reach the ball first?
A) the boat with greater velocity (relative to water)
B) the boat with lesser velocity (relative to water)
C) both will reach the ball simultaneously
D) data insufficient

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q26

Kinematics


JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **59%** got it right · Campus: **67%** (41 of 61)

Within reach

You: Skipped

26. Rain is falling with speed $12\sqrt{2}$ m/s at an angle 45^0 with vertical line. A man in a glider going at a speed of v at an angle of 37^0 with respect to ground. The speed of the glider is 5P in (m/s) so that rain appears to him falling vertically. Find the value of P, Consider motion of glider and rain drops in same vertical plane.



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q28

Electrostatics – Flux through a cube with charges at its vertices

JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **54%** got it right · Campus: **44%** (27 of 61)

Within reach

You: Skipped

28. A cube has charges each of value +Q placed at the vertices. The flux associated with the cube is $(X-1)\frac{Q}{\varepsilon_0}$, value of x is:

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q29

Electrostatics


JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **54%** got it right · Campus: **34%** (21 of 61)

Within reach

You: Wrong

29. An electron is projected from a distance d and with initial velocity u parallel to a uniformly charge flat conducting plate as shown. It strikes the plate after traveling a distance l along the direction of projection. The surface charge density of the conducting plate is $\sigma = \frac{nd\varepsilon_0 mu^2}{el^2}$, The value of n is:



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q20

Kinematics – Shell fired from a plateau – minimum landing distance


JEE Advanced · Physics · Single correct · marking +3 / −1 · marks lost 3

Top 100: **29%** got it right · Campus: **23%** (14 of 61)

Tough one

You: Skipped

20. A gun is mounted on a plateau 960m away from its edge as shown. Height of plateau is 960m. The gun can fire shells with a velocity 100 m/s at any angle of the following choices. What is the minimum distance OP(=x) from the edge of plateau where shell of



gun can reach? (given $g=10m/s^2$, $\sin 74^0 = \sin 106^0 \simeq 0.96$)

A) 480 m B) 720 m C) 360 m D) 960 m

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q27

Kinematics – Projection angle that maximises area under the trajectory

JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **27%** got it right · Campus: **23%** (14 of 61)

Tough one

You: Wrong

27. A projectile is thrown with a speed 10 m/s, at an angle θ with the horizontal. For a certain value of angle of projection $\theta = \frac{(n+1)\pi}{12}$ the area bound by the trajectory with horizontal is maximum value of n is (g=10 m/s^2). Assume no airdrag.

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q31

Kinematics – Hexagon projectile – maximum height


JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **5%** got it right · Campus: **5%** (3 of 61)

Tough one

You: Wrong

PARAGRAPH-I



A regular hexagon of side a=$10\sqrt{3}$ m is shown in figure. From any point a particle is projected such that it touches B,C,D and E. The velocity of particle is v m/s. O is the centre of the hexagon. x-axis is parallel to horizontal and y-axis is perpendicular to AE and CD (g=10 m/s^2) $\left(\sqrt{3}=1.73\right)$.

31. The maximum height from the horizontal surface X' attained by the projectile is ___ m.

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q32 **Kinematics – Hexagon projectile – speed at the top**


JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **5%** got it right · Campus: **5%** (3 of 61)

Tough one

You: Wrong

PARAGRAPH-I



A regular hexagon of side $a=10\sqrt{3}$ m is shown in figure. From any point a particle is projected such that it touches B,C,D and E. The velocity of particle is v m/s. O is the centre of the hexagon. x-axis is parallel to horizontal and y-axis is perpendicular to AE and CD ($g=10$ m/s^2) $\left(\sqrt{3}=1.73\right)$.

32. The velocity of projectile at maximum height is _____ m/s

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced · Chemistry — questions to revisit


3 questions · easy misses first
cropped from the question paper


Q47


Carboxylic Acids
JEE Advanced · Chemistry · Integer · marking +4 / 0 · marks lost 4
Top 100: **32%** got it right · Campus: **33%** (20 of 61)


Tough one
You: Wrong

47. At room temperature how many of the following compounds can decompose $NaHCO_3$ to liberate CO_2?

a) 

b) 

c) 

d) 

Why did I lose this question? (write the reason and the correct approach)


☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q46

Alcohols, Phenols & Ethers – **Identifying polar aprotic solvents**
JEE Advanced · Chemistry · Integer · marking +4 / 0 · marks lost 4
Top 100: **24%** got it right · Campus: **8%** (5 of 61)

Tough one
You: Wrong

46. How many of the following are polar aprotic solvents?



[P]

Diethyl ether,


[Q]

Benzene,

[R]

Ethyl alcohol,

[S]



[T]

Petrol

[U]

Acetone

[Q]

DMSO (dimethyl sulphoxide

[W]

DMF (dimethyl formamide

[X]

formic acid

[Y]

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q51 **Carboxylic Acids – Moles of $LiAlH_4$ to reduce RCOOH to RCH_2OH**

JEE Advanced · Chemistry · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **17%** got it right · Campus: **23%** (14 of 61)

Tough one

You: Skipped

PARAGRAPH-II

Carboxylic acids are organic compounds containing the carbonyl(>C=O) and hydroxyl (-OH) functional groups combined as a carboxyl group(-COOH). Due to resonance stabilization of the carboxylate anion, carboxylic acids exhibit higher acidity compared to alcohols and phenols. They participate in several characteristic reactions, including nucleophilic acyl substitution (esterification, amide formation, acid halide synthesis), decarboxylation, and reduction to primary alcohols. Furthermore, the presence of the –COOH group strongly influences physical properties, enabling extensive intermolecular hydrogen bonding that results in higher boiling points and stable dimeric structures in non-polar solvents.

51. The number of moles of $LiAlH_4$ required theoretically to reduce eight moles of carboxylic acid to primary alcohol

$$RCOOH \xrightarrow[2.H_3O^+]{1.LiAiH_4} RCH_2OH$$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

10 How to read this report

Where the numbers come from. Marks, percentages, All-India / State / Campus ranks, subject marks, subject All-India ranks and the wrong / skipped questions are the official results in the institute's *All India Marks Analysis* (graded with the final answer key). Maximum marks leave out deleted questions — JEE Mains: Q56 — JEE Advanced: Q25. **Program rank** = rank in Pragya G10 (EXT) Achievers across all campuses, from the batch's question-wise sheet. **Top 100** = the batch's top 100 across all campuses, from the same question-wise sheet; * where it has no data for a question, the campus figure decides the priority. **Campus** = all students of this program on campus. **Priority:** Easy miss = 70%+ of the top 100 got it right; Within reach = 40–69%; Tough one = below 40%. **Marks lost** on a question = its marks plus the negative mark if answered wrong, its marks if skipped. In JEE Advanced, 2 marks went on multiple-correct answers that were only partly correct. **Weekly-test average** = your average over the weekly Mains tests (WTM) or Advanced tests (WTA) before this Part Test, from the campus averages sheet. The weekly Advanced tests are marked out of 240; their averages are scaled to 180 to match this paper (× 180/240, so 80 per subject becomes 60). Percentages are unchanged.