CLASS 8 · PART TEST-1 · STUDENT PERFORMANCE REPORT

Aleti Dhanu Sai Sagar

Pragya G8 (EXI) Achievers · Adm No **257423306**

Campus HYD/Nallakunta DD Colony
JEE Mains 08 Oct 2026
JEE Advanced 09 Oct 2026

JEE Mains

08 Oct 2026 · 74 questions

148 / 296 — **50.0%**

180	124	9/63	56
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **174.0 / 300** (58.0%), 3 of 6 tests → this test **50.0% ▼ 8.0**

148 marks lost: 80 on 16 wrong answers (incl. 16 negative) · 68 on 17 skipped

JEE Advanced

09 Oct 2026 · 48 questions

82 / 171 — **48.0%**

278	193	12/63	94†
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **108.0 / 180** (60.0%), 3 of 6 tests → this test **48.0% ▼ 12.0**

89 marks lost: 46 on 12 wrong answers (incl. 4 negative) · 37 on 11 skipped · 6 on partly-correct answers

1 Subject scorecard

Your marks in each subject, against the 63 students of this program on campus and against your own weekly-test average.

SUBJECT	MARKS	SCORE VS CAMPUS	SUBJECT AIR	CAMPUS AVG / BEST	WEEKLY AVG	CHANGE	RIGHT / WRONG / SKIP
JEE Mains 148 / 296 · 50.0%							
Maths	**48**/96	50%	**157**	37.0 / 77	**54.7**/100 54.7%	▼ **4.7**	13 / 4 / 7
Physics	**20**/100	20%	**639**	18.3 / 60	**55.0**/100 55.0%	▼ **35.0**	7 / 8 / 10
Chemistry	**80**/100	80%	**116**	37.4 / 86	**64.3**/100 64.3%	▲ **15.7**	21 / 4 / 0
JEE Advanced 82 / 171 · 48.0%							
Maths	**29**/54	54%	**202**	24.2 / 41	**34.7**/60 57.8%	▼ **4.1**	9 / 3 / 3
Physics	**19**/60	32%	**389**	9.4 / 40	**35.7**/60 59.5%	▼ **27.8**	6 / 5 / 6
Chemistry	**34**/57	60%	**435**	23.9 / 50	**37.7**/60 62.8%	▼ **3.1**	10 / 4 / 2

Bar = your score % | campus average | best on campus Change = this test % − weekly-test average %

What this report says

1. **Chemistry** is your strongest JEE Mains subject: 80/100 (80%), All-India subject rank 116.
2. **Physics** needs the most work in JEE Mains: 20/100 (20%), subject rank 639; you lost 18 of its 25 questions.
3. **Chemistry** is your strongest JEE Advanced subject: 34/57 (60%), All-India subject rank 435.
4. **Physics** needs the most work in JEE Advanced: 19/60 (32%), subject rank 389; you lost 11 of its 17 questions.
5. Biggest fall against your weekly-test average: **JEE Mains Physics** – 20% here against 55% on average.
6. Biggest improvement: **JEE Mains Chemistry** – 80% here against 64% on average.
7. **11 questions** you lost were answered correctly by 70%+ of the batch's top 100 – about **50 marks** within easy reach (Mains Q1, Q6, Q26, Q46, Q62, Q74 · Advanced Q9, Q24, Q38, Q40, Q47).
8. Negative marking cost you **16 in JEE Mains and 4 in JEE Advanced**.
9. You left **17 in JEE Mains and 11 in JEE Advanced** questions blank, and answered 16 in JEE Mains and 12 in JEE Advanced wrong – both are marks to recover.

JEE Mains – question by question

148 / 296 · 50.0%
08 Oct 2026

2 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	~~20~~	21	22	23	24	25
Physics	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50
Chemistry	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75

3 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 63 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40–69% · **Tough one** below 40%

Maths

11 of 24 questions lost · **48** marks · 2 easy miss · 6 within reach · 3 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q1	Easy miss	**Surds & Indices**	Single correct · +4 / −1	**Skipped**	4	**89%**	86%
Q6	Easy miss	**Quadratic Equations**	Single correct · +4 / −1	**Wrong**	5	**76%**	38%
Q7	Within reach	**Logarithms & Exponents**	Single correct · +4 / −1	**Skipped**	4	**65%**	35%
Q12	Within reach	**Sequences & Series**	Single correct · +4 / −1	**Wrong**	5	**59%**	19%
Q14	Within reach	**Logarithms & Exponents** – Log-exponential equation; counting solutions	Single correct · +4 / −1	**Skipped**	4	**53%**	19%
Q17	Within reach	**Surds & Indices** – Cube-root surd identity (matrix match)	Single correct · +4 / −1	**Skipped**	4	**53%**	41%
Q11	Within reach	**Quadratic Equations**	Single correct · +4 / −1	**Skipped**	4	**46%**	21%
Q4	Within reach	**Sequences & Series** – Identify AP / GP / HP from a surd expression	Single correct · +4 / −1	**Skipped**	4	**42%**	19%
Q22	Tough one	**Surds & Indices**	Numerical · +4 / −1	**Skipped**	4	**25%**	11%
Q5	Tough one	**Quadratic Equations** – Range of a rational quadratic expression	Single correct · +4 / −1	**Wrong**	5	**21%**	6%
Q16	Tough one	**Quadratic Equations** – Roots in GP, condition on discriminant	Single correct · +4 / −1	**Wrong**	5	**19%**	3%

Physics

18 of 25 questions lost · **80** marks · ● 2 easy miss · ● 7 within reach · ● 9 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q26	Easy miss	**Vectors**	Single correct · +4 / −1	**Skipped**	4	**91%**	86%
Q46	Easy miss	**Vectors**	Numerical · +4 / −1	**Wrong**	5	**75%**	43%
Q29	Within reach	**Vectors** – Angles between three vectors summing to zero	Single correct · +4 / −1	**Skipped**	4	**62%**	24%
Q40	Within reach	**Laws of Motion** – Velocity over fixed distance vs mass ($v \propto 1/\sqrt{m}$)	Single correct · +4 / −1	**Wrong**	5	**62%**	30%
Q44	Within reach	**Laws of Motion** – Normal reactions on a cylinder resting on edges	Single correct · +4 / −1	**Wrong**	5	**53%**	29%
Q50	Within reach	**Laws of Motion**	Numerical · +4 / −1	**Wrong**	5	**44%**	17%
Q27	Within reach	**Vectors** – Vector components under axis rotation	Single correct · +4 / −1	**Skipped**	4	**43%**	6%
Q30	Within reach	**Laws of Motion** – Tension / normal reaction in two-block systems (matrix match)	Single correct · +4 / −1	**Wrong**	5	**42%**	30%
Q35	Within reach	**Kinematics** – $a = kv^2$ – integrate to get v(s)	Single correct · +4 / −1	**Skipped**	4	**41%**	5%
Q41	Tough one	**Laws of Motion** – Newton's 3rd law – fan blowing on own sail	Single correct · +4 / −1	**Wrong**	5	**31%**	11%
Q42	Tough one	**Laws of Motion** – Extra tension when bird leaves the wire	Single correct · +4 / −1	**Skipped**	4	**31%**	14%
Q38	Tough one	**Kinematics** – Resistance $\propto v^2$ – time inside plank	Single correct · +4 / −1	**Skipped**	4	**27%**	11%
Q45	Tough one	**Laws of Motion** – Contact force between pushed blocks (statement type)	Single correct · +4 / −1	**Skipped**	4	**27%**	10%
Q36	Tough one	**Kinematics** – a(t) with modulus – area under a-t	Single correct · +4 / −1	**Wrong**	5	**24%**	5%
Q43	Tough one	**Laws of Motion** – Normal reaction across an inclined contact	Single correct · +4 / −1	**Wrong**	5	**20%**	10%
Q47	Tough one	**Vectors** – Net displacement from multi-leg journey	Numerical · +4 / −1	**Skipped**	4	**15%**	3%
Q48	Tough one	**Kinematics** – Distance from a-t graph	Numerical · +4 / −1	**Skipped**	4	**15%**	3%
Q49	Tough one	**Kinematics** – Free fall then deceleration – find height	Numerical · +4 / −1	**Skipped**	4	**15%**	2%




Chemistry

4 of 25 questions lost · **20** marks · ● 2 easy miss · ● 1 within reach · ● 1 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q62	Easy miss	**Atomic Structure**	Single correct · +4 / −1	**Wrong**	5	**94%**	76%
Q74	Easy miss	**Periodic Classification**	Numerical · +4 / −1	**Wrong**	5	**70%**	46%
Q59	Within reach	**Chemical Bonding** – VBT, orbital overlap, hybridisation statements	Single correct · +4 / −1	**Wrong**	5	**50%**	29%
Q67	Tough one	**Chemical Bonding** – Strongest ionic bond (charge and size)	Single correct · +4 / −1	**Wrong**	5	**32%**	13%

4 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Quadratic Equations	5	**1**/20		11%	**Q5**, **Q6**, **Q11**, **Q16**
Logarithms & Exponents	3	**4**/12		40%	**Q7**, **Q14**
Sequences & Series	5	**11**/20		33%	**Q4**, **Q12**
Surds & Indices	11	**32**/44		53%	**Q1**, **Q17**, **Q22**
Physics					
Laws of Motion	8	**-6**/32		7%	**Q30**, **Q40**, **Q41**, **Q42**, **Q43**, **Q44**, **Q45**, **Q50**
Vectors	7	**7**/28		27%	**Q26**, **Q27**, **Q29**, **Q46**, **Q47**
Kinematics	10	**19**/40		21%	**Q35**, **Q36**, **Q38**, **Q48**, **Q49**
Chemistry					
Chemical Bonding	6	**14**/24		34%	**Q59**, **Q67**
Atomic Structure	5	**15**/20		31%	**Q62**
Periodic Classification	5	**15**/20		58%	**Q74**
Mole Concept & Basics	9	**36**/36		32%	– none

Bar = your % of the chapter's marks | campus average

5 JEE Mains · Maths – questions to revisit

11 questions · easy misses first
cropped from the question paper

Q1 Surds & Indices

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4
Top 100: **89%** got it right · Campus: **86%** (54 of 63)

Easy miss
You: Skipped

1. The values of x which satisfy the equation
$\sqrt{5x^2-8x+3}-\sqrt{5x^2-9x+4}=\sqrt{2x^2-2x}-\sqrt{2x^2-3x+1}$ are
(A)3 (B)2 (C)1 (D)0

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q6 Quadratic Equations

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5
Top 100: **76%** got it right · Campus: **38%** (24 of 63)

Easy miss
You: Wrong

6. If the sum of the roots of the quadratic equation $ax^2+bx+c=0$ is equal to the sum of the squares of the reciprocals, then
(A) ab^2, ca^2, bc^2 are in A.P. (B) a^2b, c^2b, b^2c are in A.P.
(C) ab^2, ca^2, bc^2 are in G.P. (D) a^2b, c^2b, b^2c are in G.P.

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q7 Logarithms & Exponents

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4
Top 100: **65%** got it right · Campus: **35%** (22 of 63)

Within reach
You: Skipped

7. The roots of the equation $2^{x+2}27^{x/(x-1)}=9$ are given by
(A)1 − $\log_2 3$, 2 (B)−2, 1 − $\log_3 2$
(C)−2, 1 − $\log_2 3$ (D) 2, −1 $\log_2 3$

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q12

Sequences & Series
JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5
Top 100: **59%** got it right · Campus: **19%** (12 of 63)

Within reach
You: Wrong

12. If $4a^2 + 9b^2 + 16c^2 = 2(3ab + 6bc + 4ca)$, where a, b, c are nonzero numbers, then a, b, c are in
(A)A.P. (B)G.P. (C)H.P. (D)Neither A.P nor G.P nor H.P

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q14

Logarithms & Exponents – Log-exponential equation; counting solutions
JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4
Top 100: **53%** got it right · Campus: **19%** (12 of 63)

Within reach
You: Skipped

14. The equation $x^{(3/4)(\log_2^x)^2 + \log_2^x - (5/4)} = \sqrt{2}$ has
(A)at least one real solution (B)exactly four real solutions
(C)exactly two irrational solution (D)exactly four non-real solutions

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q17 Surds & Indices – Cube-root surd identity (matrix match)

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **53%** got it right · Campus: **41%** (26 of 63)

Within reach

You: Skipped

17. Let $x=\left(a+\sqrt{a^2+b^3}\right)^{\frac{1}{3}}+\left(a-\sqrt{a^2+b^3}\right)^{\frac{1}{3}}$ and $x^3+Abx-B.a=0$, Then match COLUMN-I with respect to COLUMN-II

	COLUMN-I		COLUMN-II
I)	A + B	P)	8
II)	A – B	Q)	9
III)	A^B	R)	1
IV)	B^A	S)	5
		T)	3

(A) I – S; II – R; III – Q; IV – P (B) I – S; II – T; III – P; IV – Q

(C) I – Q; II – P; III – T; IV – S (D) I – Q; II – P; III – S; IV – R

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q11 Quadratic Equations

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **46%** got it right · Campus: **21%** (13 of 63)

Within reach

You: Skipped

11. The range of values of a for which all the roots of the equation $(a-1)\left(1+x+x^2\right)^2=(a+1)\left(1+x^2+x^4\right)$ are imaginary is

(A) $(-\infty,-2]$ (B) $(2,\infty)$ (C) $-2<a<2$ (D) $(-\infty,2]$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q4

Sequences & Series – Identify AP / GP / HP from a surd expression

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **42%** got it right · Campus: **19%** (12 of 63)

Within reach

You: Skipped

4. If a, b, c are in A.P., then $\frac{1}{\sqrt{a}+\sqrt{b}}, \frac{1}{\sqrt{a}+\sqrt{c}}, \frac{1}{\sqrt{b}+\sqrt{c}}$ are in

(A)A.P. (B)G.P. (C)H.P. (D)Neither A.P nor G.P nor H.P

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q22

Surds & Indices

JEE Mains · Maths · Numerical · marking +4 / −1 · marks lost 4

Top 100: **25%** got it right · Campus: **11%** (7 of 63)

Tough one

You: Skipped

22. If $y=\frac{(p+1)^{\frac{-}{3}}+(p-1)^{\frac{-}{3}}}{(p+1)^{\frac{1}{3}}-(p-1)^{\frac{1}{3}}}$ then the value of the expression $y^3-3py^2+3y-p+3$ is_____?

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q5

Quadratic Equations – Range of a rational quadratic expression

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5

Top 100: **21%** got it right · Campus: **6%** (4 of 63)

Tough one

You: Wrong

5. If x is real, then $\frac{x^2+2x+c}{x^2+4x+3c}$ can take all real values if

(A)$0 < c < 2$ (B)$0 \le c \le 1$

(C)$-1 < c < 1$ (D)$0 < c < 1$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q16 **Quadratic Equations – Roots in GP, condition on discriminant**

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5

Top 100: **19%** got it right · Campus: **3%** (2 of 63)

Tough one

You: Wrong

16. Let α, β be the roots of the equation $ax^2 + bx + c = 0$ and $\Delta = b^2 - 4ac$. If $\alpha + \beta, \alpha^2 + \beta^2, \alpha^3 + \beta^3$ are in G.P., then

(A) $\Delta = 0$ (B) $\Delta \neq 0$ (C) $b\Delta = 0$ (D) $c\Delta = 0$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Mains · Physics – questions to revisit

18 questions · easy misses first
cropped from the question paper

Q26 — Vectors

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4
Top 100: **91%** got it right · Campus: **86%** (54 of 63)

Easy miss
You: Skipped

26. ABCDEF is regular hexagon with point O as centre. The value of $\overrightarrow{AB}+\overrightarrow{AC}+\overrightarrow{AD}+\overrightarrow{AE}+\overrightarrow{AF}$ is
(A) $2\overrightarrow{AB}$ (B) $4\overrightarrow{AO}$ (C) $6\overrightarrow{AO}$ (D) 0

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q46 — Vectors

JEE Mains · Physics · Numerical · marking +4 / −1 · marks lost 5
Top 100: **75%** got it right · Campus: **43%** (27 of 63)

Easy miss
You: Wrong

46. If the resultant of two vectors having magnitudes of 6 and 3 is 9, then the dot product of two vectors is:

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q29 — Vectors – Angles between three vectors summing to zero

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4
Top 100: **62%** got it right · Campus: **24%** (15 of 63)

Within reach
You: Skipped

29. Given: $\vec{a}+\vec{b}+\vec{c}=0$. Out of the three vectors $\vec{a},\vec{b}$ and $\vec{c}$, two are equal in magnitude. The magnitude of the third vector is $\sqrt{2}$ times that of either of the two having equal magnitude. The angles between the vector are
(A) $90^0,135^0,135^0$ (B) $30^0,60^0,90^0$
(C) $45^0,45^0,90^0$ (D) $45^0,60^0,90^0$

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q40

Laws of Motion – Velocity over fixed distance vs mass (v ∝ 1/√m)

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **62%** got it right · Campus: **30%** (19 of 63)

Within reach

You: Wrong

40. A constant force acts on a mass m at rest. Velocity acquired in travelling a fixed distance is directly proportional to

(A) $\sqrt{m}$ (B) m (C) $1/\sqrt{m}$ (D) $1/m$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q44

Laws of Motion – Normal reactions on a cylinder resting on edges


JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **53%** got it right · Campus: **29%** (18 of 63)

Within reach

You: Wrong

44. A cylinder of mass M and radius R is resting on two corner edges A and B as shown in the figure. The normal reaction at the edges A and B are (neglect friction)



(A) $N_A = \sqrt{2}N_B$ (B) $N_B = \sqrt{3}N_A$ (C) $N_A = \sqrt{3}N_B$ (D) $N_B = \frac{2\sqrt{3}Mg}{5}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q50 **Laws of Motion**
JEE Mains · Physics · Numerical · marking +4 / −1 · marks lost 5
Top 100: **44%** got it right · Campus: **17%** (11 of 63)

Within reach
You: Wrong

50. A constant force acts on a body of mass 1kg for 5 seconds, so that the body begins to move with zero initial velocity on a smooth horizontal plane. After 5 seconds the force ceases to act. In the next 5 second, the body describes 100m. What is force (in Newton) that acted on the body

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q27 **Vectors – Vector components under axis rotation**
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4
Top 100: **43%** got it right · Campus: **6%** (4 of 63)

Within reach
You: Skipped

27. The components of a vector along x and y directions are (n+1) and 1 respectively . If the co-ordinate system is rotated by an angle $\theta = 60^0$, then the components change to n and 3. The value of n is

(A) 2 (B) $\cos 60^0$ (C) $\sin 60^0$ (D) 3.5

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q30

Laws of Motion – Tension / normal reaction in two-block systems (matrix match)

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **42%** got it right · Campus: **30%** (19 of 63)

Within reach

You: Wrong

30. Column-I gives four different situations involving two blocks of mass m_1 and m_2 placed in different ways on a smooth horizontal surface as shown. In each of the situations horizontal forces F_1 and F_2 are applied on blocks of mass m_1 and m_2 respectively and also $m_2 F_1 < m_1 F_2$. Match the statements in column I with corresponding results in column-II

Column-I	Column-II
(A)  . Both the blocks are connected by massless inelastic string. The magnitude of tension in the string is	(p) $\frac{m_1 m_2}{m_1+m_2}\left(\frac{F_1}{m_1}-\frac{F_2}{m_2}\right)$
(B)  . Both the blocks are connected by massless inelastic string. The magnitude of tension in the string is	(q) $\frac{m_1 m_2}{m_1+m_2}\left(\frac{F_1}{m_1}+\frac{F_2}{m_2}\right)$
(C)  . The magnitude of normal reaction between the blocks is	(r) $\frac{m_1 m_2}{m_1+m_2}\left(\frac{F_2}{m_2}-\frac{F_1}{m_1}\right)$
(D)  . The magnitude of normal reaction between the blocks is	(s) $m_1 m_2\left(\frac{F_1+F_2}{m_1+m_2}\right)$

(A) (A) – r ; (B) – q ; (C) – q ; (D) – r

(B) (A) – q ; (B) – r ; (C) – r ; (D) – q

(C) (A) – q ; (B) – r ; (C) – q ; (D) – r

(D) (A) – q ; (B) – p ; (C) – r ; (D) – r

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q35

Kinematics – a = kv² – integrate to get v(s)

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4

Top 100: **41%** got it right · Campus: **5%** (3 of 63)

Within reach

You: Skipped

35. If a particle accelerates with $a = kv^2$ and initial velocity = u, then find velocity after displacement s

(A) $v = ue^{ks}$ (B) $v = u^{ks}e^{s}$ (C) $v = u^{k}e^{s}$ (D) $v = eu^{ks}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q41

Laws of Motion – Newton's 3rd law – fan blowing on own sail

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **31%** got it right · Campus: **11%** (7 of 63)

Tough one

You: Wrong

41. On a stationary sail boat, air is blown at the sails from a fan attached to the boat. The boat will:

(A) Move forward (B) move backward

(C) spin around (D) remain stationary

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q42

Laws of Motion – Extra tension when bird leaves the wire

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4

Top 100: **31%** got it right · Campus: **14%** (9 of 63)

Tough one

You: Skipped

42. A bird dismounts on a stretched telegraph wire. The additional tension produced in the wire is

(A) Zero (B) greater than the weight of the bird

(C) Less than the weight of the bird (D) equal to the weight of the bird

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q38

Kinematics – Resistance ∝ v² – time inside plank

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4

Top 100: **27%** got it right · Campus: **11%** (7 of 63)

Tough one

You: Skipped

38. Having gone through a plank of thickness h, a bullet changed its velocity from v_0 to v. Find the time of motion of the bullet in the plank assuming resistance force to be proportional to the square of the velocity

(A) $\left[\frac{h(v_0-v)}{vv_0 l\,n\frac{v_0}{v}}\right]$ (B) $\left[\frac{(v_0-v)}{hvv_0 l\,n\left(\frac{v_0}{v}\right)}\right]$ (C) $\left[\frac{vv_0 \ln\left(\frac{v_0}{v}\right)}{h(v_0-v)}\right]$ (D) $\left[\frac{\ln\frac{v_0}{v}}{vv_0(v_0-v)}\right]$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q45

Laws of Motion – Contact force between pushed blocks (statement type)


JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4

Top 100: **27%** got it right · Campus: **10%** (6 of 63)

Tough one

You: Skipped

45. Statement-1: Block A is moving on a horizontal surface towards right under action of force F. all surfaces are smooth. At the instant shown the force exerted by block A on block B is equal to net force on block B.



Statement-2: From Newton's third law, the force exerted by block A on B is equal in magnitude to force exerted by block B on A.

(A) Statement 1 is true, statement 2 is true, statement 2 is correct explanation for statement 1

(B) Statement 1 is true, statement 2 is true, statement 2 is not correct explanation for statement 1

(C) Statement 1 is true, statement 2 is false

(D) Statement 1 is false, statement 2 is true

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q36

Kinematics – a(t) with modulus – area under a-t

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **24%** got it right · Campus: **5%** (3 of 63)

Tough one

You: Wrong

36. Starting from rest a particle moves in a straight line with acceleration $a=\{2+|t-2|\}\,m/s^2$. Velocity of the particle at the end of 4s will be

(A) 16 m/s (B) 20 m/s (C) 8 m/s (D) 12 m/s

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q43 Laws of Motion – Normal reaction across an inclined contact


JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **20%** got it right · Campus: **10%** (6 of 63)

Tough one

You: Wrong

43. Two blocks 'A" and 'B' each of mass 'm' are placed on a smooth horizontal surface. Two horizontal forces F and 2F are applied on both the blocks 'A' and 'B' respectively as shown in figure. The block A does not slide on block B. Then the normal reaction acting between the two blocks is



(A) F (B) $F/2$ (C) $F/\sqrt{3}$ (D) 3F

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q47 Vectors – Net displacement from multi-leg journey

JEE Mains · Physics · Numerical · marking +4 / −1 · marks lost 4

Top 100: **15%** got it right · Campus: **3%** (2 of 63)

Tough one

You: Skipped

47. In going from one city to another , a car travels 75 km north, 60 km north –west and 20 km east. The magnitude(in km) of the vector drawn between the two cities is $\left(Take 1/\sqrt{2}=0.7\right)$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q48

Kinematics – Distance from a-t graph


JEE Mains · Physics · Numerical · marking +4 / −1 · marks lost 4

Top 100: **15%** got it right · Campus: **3%** (2 of 63)

Tough one

You: Skipped

48. An experiment on the take off performance of an aeroplane shows that the acceleration varies as shown in the fig and it takes 12 sec to take off from a rest position.The distance along the runway covered by the aeroplane is(in m).



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q49

Kinematics – Free fall then deceleration – find height

JEE Mains · Physics · Numerical · marking +4 / −1 · marks lost 4

Top 100: **15%** got it right · Campus: **2%** (1 of 63)

Tough one

You: Skipped

49. A parachutist after bailing out falls 50m without friction. When parachute opens, it decelerates at $2m/\sec^2$. He reaches the ground with a speed of 3m/s. The height he bail out is nearly (in m) (Take g=10m/sec^2) .

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Mains · Chemistry – questions to revisit

4 questions · easy misses first
cropped from the question paper

Q62 Atomic Structure

JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5
Top 100: **94%** got it right · Campus: **76%** (48 of 63)

Easy miss
You: Wrong

62. Match the column for the energy and radius of Hydrogen.

Column I	**Column II**
(A) Radius of 1st orbit	(P) – 13.6 eV/atom
(B) Radius of 2nd orbit	(Q) – 3.4 eV/atom
(C) Energy of 1st orbit	(R) 0.529 Å
(D) Energy of 2nd orbit	(S) 2.116 Å

A) [A → R; B → S; C →P; D → Q]
B) [A → S; B → R; C →P; D → Q]
C) [A → Q; B → S; C →R; D →P]
D) [A → P; B → R; C →S; D → Q]

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q74 Periodic Classification

JEE Mains · Chemistry · Numerical · marking +4 / −1 · marks lost 5
Top 100: **70%** got it right · Campus: **46%** (29 of 63)

Easy miss
You: Wrong

74. An element X belongs to the 3rd period and has 5 valence electrons. Another element Y belongs to the same period and has 2 valence electrons.
Find the sum of the atomic numbers of X and Y.

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q59

Chemical Bonding – VBT, orbital overlap, hybridisation statements

JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5

Top 100: **50%** got it right · Campus: **29%** (18 of 63)

Within reach

You: Wrong

59. Statement I: According to VBT, a covalent bond is formed by the overlap of half-filled atomic orbitals containing electrons with opposite spins.
Statement II: Greater the extent of effective orbital overlap, stronger is the covalent bond formed.
Statement III: Hybridisation involves the mixing of atomic orbitals of comparable energy to form equivalent hybrid orbitals.
Which of the following is correct?
(A) Only Statement I is correct
(B) Statements I and II are correct
(C) Statements I, II and III are correct
(D) Statements II and III are correct

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q67

Chemical Bonding – Strongest ionic bond (charge and size)

JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5

Top 100: **32%** got it right · Campus: **13%** (8 of 63)

Tough one

You: Wrong

67. Which combination will give strongest ionic bond?
(A) K^+ and Cl^- (B) K^+ and O^{2-}
(C) Ca^{2+} and Cl^- (D) Ca^{2+} and O^{2-}

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced – question by question

82 / 171 · 48.0%
09 Oct 2026

6 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · ● Easy miss · ● Within reach · ● Tough one

Maths	1	2	3	**4**	5	6	7	8	**9**	**10**	11	12	**13**	~~14~~	**15**	**16**	~~17~~
Physics	18	19	**20**	**21**	22	**23**	**24**	25	**26**	**27**	28	29	**30**	**31**	**32**	**33**	**34**
Chemistry	**35**	36	37	**38**	39	**40**	41	42	43	44	45	46	**47**	48	**49**	~~50~~	**51**

7 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 63 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40–69% · **Tough one** below 40%

Maths

6 of 15 questions lost · **22** marks (+3 on partly-correct answers) · ● 1 easy miss · ● 3 within reach · ● 2 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q9	Easy miss	**Quadratic Equations**	Integer · +4 / 0	Wrong	4	73%	56%
Q4	Within reach	**Logarithms** – Logs in AP – solve for x	Single correct · +3 / −1	Wrong	4	68%	35%
Q10	Within reach	**Quadratic Equations** – Roots of two linked quadratics	Integer · +4 / 0	Skipped	4	50%	25%
Q15	Within reach	**Sequences & Series** – AP: ratio of sums of 14 terms	Paragraph · +3 / 0	Wrong	3	42%	8%
Q16	Tough one	**Logarithms** – Log equations – A + B + C	Paragraph · +3 / 0	Skipped	3	18%	3%
Q13	Tough one	**Quadratic Equations** – Spot the polynomial identity, then sum P(1) … P(2023)	Integer · +4 / 0	Skipped	4	– no data*	2%

Physics

11 of 17 questions lost · **38** marks (+3 on partly-correct answers) · ● 1 easy miss · ● 5 within reach · ● 5 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q24	Easy miss	**Laws of Motion** – Tensions in two inclined strings	Multiple correct · +4 / −2	Skipped	4	70%	44%
Q21	Within reach	**Kinematics** – Retardation from the t–x relation	Single correct · +3 / −1	Skipped	3	62%	24%
Q26	Within reach	**Vectors**	Integer · +4 / 0	Wrong	4	56%	32%
Q20	Within reach	**Kinematics**	Single correct · +3 / −1	Skipped	3	52%	22%
Q23	Within reach	**Laws of Motion** – Block on block, no friction – motion of m	Multiple correct · +4 / −2	Skipped	4	51%	22%
Q30	Within reach	**Kinematics** – a from v(t) at a given position	Integer · +4 / 0	Wrong	4	43%	8%
Q33	Tough one	**Kinematics** – Train passage – retardation to avoid collision	Paragraph · +3 / 0	Skipped	3	15%	8%
Q34	Tough one	**Kinematics** – Train passage – time of retardation	Paragraph · +3 / 0	Skipped	3	15%	10%
Q27	Tough one	**Laws of Motion** – Normal reaction between pushed blocks	Integer · +4 / 0	Wrong	4	11%	2%
Q31	Tough one	**Vectors** – Carrom passage – displacement centre → edge	Paragraph · +3 / 0	Wrong	3	7%	2%
Q32	Tough one	**Vectors** – Carrom passage – displacement edge → hole	Paragraph · +3 / 0	Wrong	3	6%	2%

Chemistry

6 of 16 questions lost · **23** marks · ● 3 easy miss · ● 2 within reach · ● 1 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q38	Easy miss	**Chemical Bonding** – Pair of species with the same shape (VSEPR)	Single correct · +3 / −1	**Wrong**	**4**	**92%**	49%
Q40	Easy miss	**Atomic Structure**	Multiple correct · +4 / −2	**Wrong**	**6**	**82%**	44%
Q47	Easy miss	**Mole Concept & Basics** – Mass of water from 8 g H_2 + 32 g O_2 (limiting reagent)	Integer · +4 / 0	**Wrong**	**4**	**72%**	32%
Q49	Within reach	**Mole Concept & Basics** – Mass of oxygen in 24 g of the same copper oxide	Paragraph · +3 / 0	**Wrong**	**3**	**60%**	49%
Q35	Within reach	**Mole Concept & Basics** – Limiting reagent over two-step reaction	Single correct · +3 / −1	**Skipped**	**3**	**51%**	24%
Q51	Tough one	**Atomic Structure** – Hydrogen-like ion – find Z from photon energies	Paragraph · +3 / 0	**Skipped**	**3**	**0%**	5%

8 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject. Chapter marks are approximate (≈): partly-correct answers cannot be traced to a question.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Logarithms	4	≈ **6**/13		50%	**Q4**, **Q16**
Quadratic Equations	7	≈ **14**/26		43%	**Q9**, **Q10**, **Q13**
Sequences & Series	4	≈ **12**/15		56%	**Q15**
Physics					
Laws of Motion	4	≈ **4**/16		6%	**Q23**, **Q24**, **Q27**
Kinematics	8	≈ **11**/27		24%	**Q20**, **Q21**, **Q30**, **Q33**, **Q34**
Vectors	5	≈ **7**/17		23%	**Q26**, **Q31**, **Q32**
Chemistry					
Chemical Bonding	1	≈ **-1**/3		41%	**Q38**
Atomic Structure	4	≈ **5**/14		33%	**Q40**, **Q51**
Mole Concept & Basics	9	≈ **23**/33		44%	**Q35**, **Q47**, **Q49**
Periodic Classification	2	≈ **7**/7		68%	– none

Bar = your % of the chapter's marks | campus average

9 JEE Advanced · Maths – questions to revisit

6 questions · easy misses first
cropped from the question paper

Q9 Quadratic Equations

JEE Advanced · Maths · Integer · marking +4 / 0 · marks lost 4
Top 100: **73%** got it right · Campus: **56%** (35 of 63)

Easy miss
You: Wrong

9. If α, β are roots of the equation $\lambda\left(x^2 - x\right) + x + 5 = 0$ and if λ_1 & λ_2 are two values of λ for which α, β are related by $\frac{\alpha}{\beta} + \frac{\beta}{\alpha} = \frac{4}{5}$ and the value of $\frac{\lambda_1}{\lambda_2} + \frac{\lambda_2}{\lambda_1} = 255 - k^2$;then absolute value of k is____?

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q4 Logarithms – Logs in AP – solve for x

JEE Advanced · Maths · Single correct · marking +3 / −1 · marks lost 4
Top 100: **68%** got it right · Campus: **35%** (22 of 63)

Within reach
You: Wrong

4. If $\log_3 2, \log_3\left(2^x - 5\right)$ & $\log_3\left(2^x - \frac{7}{2}\right)$ are in AP then the value of x is?

A)2 B)3 C)4 D)2&3

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q10 Quadratic Equations – Roots of two linked quadratics

JEE Advanced · Maths · Integer · marking +4 / 0 · marks lost 4
Top 100: **50%** got it right · Campus: **25%** (16 of 63)

Within reach
You: Skipped

10. Let a and b be roots of the equation $x^2 - 10cx - 11d = 0$ and the roots of $x^2 - 10ax - 11b = 0$ are c and d; then the value of $\frac{a+b+c+d}{605}$ is____?(Given $a \neq b \neq c \neq d \neq 0$)

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q15

Sequences & Series – AP: ratio of sums of 14 terms

JEE Advanced · Maths · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **42%** got it right · Campus: **8%** (5 of 63)

Within reach

You: Wrong

Sum of n terms of two different Arithmetic progressions are in the ratio 5n+4:9n+6 then

15. The ratio of the sum of their first 14 terms is Q, then [100Q] is _____(where [.] is G.I.F)

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q16

Logarithms – Log equations – A + B + C

JEE Advanced · Maths · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **18%** got it right · Campus: **3%** (2 of 63)

Tough one

You: Skipped

PARAGRAPH-II

Let **A** denote the product xyz where x, y, z are solutions of the system of equations

$\log_3 x = \log_3 5 - \log_3 7, \log_5 y = \log_5 7 - \log_5 3, \log_7 z = \log_7 3 - \log_7 5$

Let **B** denote the sum of square of the solutions of the equation

$\log_2\left(\log_2 x^6 - 3\right) - \log_2\left(\log_2 x^4 - 5\right) = \log_2 3$

Let **C** denote the characteristic of the number K, where

$K = \log_2\left(\log_2 3\right) - \log_2\left(\log_4 3\right) + \log_2\left(\log_4 5\right) - \log_2\left(\log_6 5\right) + \log_2\left(\log_6 7\right) - \log_2\left(\log_8 7\right)$

16. The value of A+B+C is____?

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q13

Quadratic Equations – Spot the polynomial identity, then sum P(1) ... P(2023)

JEE Advanced · Maths · Integer · marking +4 / 0 · marks lost 4

Top 100: **no data** got it right · Campus: **2%** (1 of 63)

Tough one

You: Skipped

13. Let $P(x)=\left(1+\sqrt{2}\right)\left(1+\sqrt{3}\right)\left(x-\sqrt{2}\right)\left(x-\sqrt{3}\right)+\left(1+\sqrt{3}\right)\left(\sqrt{3}+\sqrt{2}\right)(x-1)\left(x-\sqrt{2}\right)$ $-2\left(1+\sqrt{2}\right)\left(\sqrt{3}+\sqrt{2}\right)(x-1)\left(x-\sqrt{3}\right)\ \forall x\in R$. Then the value of $P(1)+P(2)+P(3)+...P(2023)$ is_____?

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced · Physics – questions to revisit


11 questions · easy misses first
cropped from the question paper

Q24

Laws of Motion – Tensions in two inclined strings
JEE Advanced · Physics · Multiple correct · marking +4 / −2 · marks lost 4
Top 100: **70%** got it right · Campus: **44%** (28 of 63)

Easy miss
You: Skipped

24. A body of mass 5 kg is suspended by the strings making angles 60^0 and 30^0 with the horizontal as shown in the figure($g = 10ms^{-2}$). Then



A) $T_1 = 25N$ B)$T_2 = 25N$ C) $T_1 = 25\sqrt{3}N$ D) $T_2 = 25\sqrt{3}N$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q21

Kinematics – Retardation from the t-x relation
JEE Advanced · Physics · Single correct · marking +3 / −1 · marks lost 3
Top 100: **62%** got it right · Campus: **24%** (15 of 63)

Within reach
You: Skipped

21. What is the retardation of a moving particle if the relation between time (t) and position (x) is $t = Ax^2 + Bx$, where A and B are appropriate constants.

A) $2A/(2Ax+B)^3$ B) $2B/(2Ax+B)^3$

C) $-2A/(2Ax+B)^3$ D) $-2B/(2Ax+B)^3$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q26

Vectors

JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **56%** got it right · Campus: **32%** (20 of 63)

Within reach

You: Wrong

26. Two forces 5N and 10N are acting with an inclination of 120^0 between them. The angle which the resultant makes with the 10N is $\frac{180^0}{n}$ Find the value of n.

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q20

Kinematics

JEE Advanced · Physics · Single correct · marking +3 / −1 · marks lost 3

Top 100: **52%** got it right · Campus: **22%** (14 of 63)

Within reach

You: Skipped

20. A particle is projected vertically upwards from the ground at time $t = 0$ and reaches a height h at $t = T$. Greatest height of the particle is

A) $\left(gT^2 + h\right)^2 / 8gT^2$

B) $2\left(gT^2 + h\right)^2 / 8gT^2$

C) $\left(gT^2 + 2h\right)^2 / 8gT^2$

D) $2h$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q23

Laws of Motion – Block on block, no friction – motion of m


JEE Advanced · Physics · Multiple correct · marking +4 / −2 · marks lost 4

Top 100: **51%** got it right · Campus: **22%** (14 of 63)

Within reach

You: Skipped

23. In the figure small block is kept on M, then



A) the acceleration of m w.r.t. ground is $\frac{F}{m}$

B) the acceleration of m w.r.t. ground is zero

C) the time taken by m to separate from M is $\sqrt{\frac{2\ell m}{F}}$

D) the time taken by m to separate from M is $\sqrt{\frac{2\ell M}{F}}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q30

Kinematics – a from v(t) at a given position

JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **43%** got it right · Campus: **8%** (5 of 63)

Within reach

You: Wrong

30. A body starts from origin and moves along x axis such that at any instant velocity is $v_t=4t^3-2t$ where t is in second and v_t is in ms^{-1}. The acceleration of the particle when it is 2m from the origin is $(20+K)\,ms^{-2}$, where K is

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q33

Kinematics – Train passage – retardation to avoid collision

JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **15%** got it right · Campus: **8%** (5 of 63)

Tough one

You: Skipped

PASSAGE-II:

The driver of a train A moving with a uniform speed of 144 km/h sights another train B, 1km ahead of him. The train B is moving with a uniform speed of 108 km/h. the driver of the train A immediately applies brakes producing a constant retardation and just manages to avoid a collision.

33. What is the retardation of the train A(in ms^{-2})_____

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q34

Kinematics – Train passage – time of retardation

JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **15%** got it right · Campus: **10%** (6 of 63)

Tough one

You: Skipped

PASSAGE-II:

The driver of a train A moving with a uniform speed of 144 km/h sights another train B, 1km ahead of him. The train B is moving with a uniform speed of 108 km/h. the driver of the train A immediately applies brakes producing a constant retardation and just manages to avoid a collision.

34. For how long is this retardation produced(in sec)_____

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q27

Laws of Motion – Normal reaction between pushed blocks


JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **11%** got it right · Campus: **2%** (1 of 63)

Tough one

You: Wrong

27. The normal reaction between 2 kg and 3 kg block is



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q31

Vectors – Carrom passage – displacement centre → edge

JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **7%** got it right · Campus: **2%** (1 of 63)

Tough one

You: Wrong

PASSAGE-1:

A Corrom board $(4\,ft \times 4\,ft\ square)$ has queen at the centre. The queen , hit by the striker moves to the front edge, rebounds and goes in the hole behind the striking line.

Answer the following question.

31. The magnitude of the displacement of the queen from the centre to the front edge is $\frac{2}{3}\sqrt{a}\,ft$, the value of a is _____

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q32

Vectors – Carrom passage – displacement edge → hole

JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **6%** got it right · Campus: **2%** (1 of 63)

Tough one

You: Wrong

PASSAGE-1:

A Corrom board $(4\,ft \times 4\,ft\ square)$ has queen at the centre. The queen , hit by the striker moves to the front edge, rebounds and goes in the hole behind the striking line.

Answer the following question.

32. The magnitude of the displacement of the queen from the front edge to the hole is $\frac{4}{3}\sqrt{b}\,ft$, the value of b is_____

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced · Chemistry – questions to revisit

6 questions · easy misses first
cropped from the question paper

Q38 — Chemical Bonding – Pair of species with the same shape (VSEPR)

JEE Advanced · Chemistry · Single correct · marking +3 / −1 · marks lost 4
Top 100: **92%** got it right · Campus: **49%** (31 of 63)

Easy miss — You: Wrong

38. The pair of species with similar shape is

(A) PCl_3, NH_3 (B) CF_4, SF_4 (C) $PbCl_2$, CO_2 (D) PF_5, IF_5

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q40 — Atomic Structure

JEE Advanced · Chemistry · Multiple correct · marking +4 / −2 · marks lost 6
Top 100: **82%** got it right · Campus: **44%** (28 of 63)

Easy miss — You: Wrong

40. Which of the following properties is/are proportional to the energy of the electromagnetic radiation ?

(A) Frequency (B) Wave number
(C) Wavelength (D) Number of photons

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q47 — Mole Concept & Basics – Mass of water from 8 g H_2 + 32 g O_2 (limiting reagent)

JEE Advanced · Chemistry · Integer · marking +4 / 0 · marks lost 4
Top 100: **72%** got it right · Campus: **32%** (20 of 63)

Easy miss — You: Wrong

47. For $2H_2 + O_2 \rightarrow 2H_2O$, 8 g H_2 reacts with 32 g O_2. Find the mass of water formed.

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q49 **Mole Concept & Basics** – Mass of oxygen in 24 g of the same copper oxide
JEE Advanced · Chemistry · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **60%** got it right · Campus: **49%** (31 of 63)

Within reach
You: Wrong

PARAGRAPH-I

An unknown copper oxide is analysed. A 12.0 g sample contains 9.6 g Cu, so the remaining 2.4 g is oxygen. Taking Cu = 64 and O = 16, the moles of Cu and O are equal

49. If the same oxide has the same composition, what mass of oxygen is present in a 24 g sample?

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q35 **Mole Concept & Basics** – Limiting reagent over two-step reaction
JEE Advanced · Chemistry · Single correct · marking +3 / −1 · marks lost 3
Top 100: **51%** got it right · Campus: **24%** (15 of 63)

Within reach
You: Skipped

35. 5 moles of A, 6 moles of Z and mixed with sufficient amount of C to produce final product F. How many maximum moles of 'F' can be produced as per the given sequence of reaction ?

$A + 2Z \rightarrow B$; $B + C \rightarrow Z + F$

(A) 3 (B) 2 (C) 4 (D) 5

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q51 **Atomic Structure** – Hydrogen-like ion – find Z from photon energies
JEE Advanced · Chemistry · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **0%** got it right · Campus: **5%** (3 of 63)

Tough one
You: Skipped

PARAGRAPH-II

When electron jumps from lower orbit to higher orbit energy is gained and when electron jumps from higher orbit to lower orbit energy is released.

51. A hydrogen like species (atomic number Z) is present in a higher excited state of quantum number n. This excited atom can make a transition to the first excited state by successive emission of two photons of energies 10.20 eV and 17.0 eV respectively. Alternatively the atom from the same excited state can make a transition to the second excited state by successive emission of two photons of energy 4.25 eV and 5.95 eV respectively. Determine the value of Z.

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

10 How to read this report

Where the numbers come from. Marks, percentages, All-India / State / Campus ranks, subject marks, subject All-India ranks and the wrong / skipped questions are the official results in the institute's *All India Marks Analysis* (graded with the final answer key). Maximum marks leave out deleted questions — JEE Mains: Q20 — JEE Advanced: Q14, Q17, Q50. **Program rank** = rank in Pragya G8 (EXI) Achievers across all campuses, from the batch's question-wise sheet († JEE Advanced: that sheet was graded before the final key, so this rank may differ slightly). **Top 100** = the batch's top 100 across all campuses, from the same question-wise sheet; * where it has no data for a question, the campus figure decides the priority. In JEE Advanced that sheet predates the final key, so its figures for the revised questions Q25, Q48, Q51 use the old answers. **Campus** = all students of this program on campus. **Priority:** Easy miss = 70%+ of the top 100 got it right; Within reach = 40–69%; Tough one = below 40%. **Marks lost** on a question = its marks plus the negative mark if answered wrong, its marks if skipped. In JEE Advanced, 6 marks went on multiple-correct answers that were only partly correct. **Weekly-test average** = your average over the weekly Mains tests (WTM) or Advanced tests (WTA) before this Part Test, from the campus averages sheet. The weekly Advanced tests are marked out of 240; their averages are scaled to 180 to match this paper (× 180/240, so 80 per subject becomes 60). Percentages are unchanged.