CLASS 8 · PART TEST-1 · STUDENT PERFORMANCE REPORT

Fareeha Shaik

Pragya G8 (EXI) Achievers · Adm No **246562182**

Campus HYD/Nallakunta DD Colony
JEE Mains 08 Oct 2026
JEE Advanced 09 Oct 2026

JEE Mains

08 Oct 2026 · 74 questions

168 / 296 — **56.8%**

112	76	5/63	40
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **204.1 / 300** (68.0%), 5 of 6 tests → this test **56.8% ▼ 11.3**

128 marks lost: 100 on 20 wrong answers (incl. 20 negative) · 28 on 7 skipped

JEE Advanced

09 Oct 2026 · 48 questions

106 / 171 — **62.0%**

97	66	4/63	29†
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **90.8 / 180** (50.5%), 5 of 6 tests → this test **62.0% ▲ 11.5**

65 marks lost: 56 on 15 wrong answers (incl. 5 negative) · 4 on 1 skipped · 5 on partly-correct answers

1 Subject scorecard

Your marks in each subject, against the 63 students of this program on campus and against your own weekly-test average.

SUBJECT	MARKS	SCORE VS CAMPUS	SUBJECT AIR	CAMPUS AVG / BEST	WEEKLY AVG	CHANGE	RIGHT / WRONG / SKIP
JEE Mains 168 / 296 · 56.8%							
Maths	63/96	66%	36	37.0 / 77	71.2/100 71.2%	▼ 5.5	17 / 5 / 2
Physics	38/100	38%	191	18.3 / 60	70.2/100 70.2%	▼ 32.2	12 / 10 / 3
Chemistry	67/100	67%	294	37.4 / 86	62.5/100 62.5%	▲ 4.5	18 / 5 / 2
JEE Advanced 106 / 171 · 62.0%							
Maths	34/54	63%	109	24.2 / 41	32.8/60 54.6%	▲ 8.4	11 / 3 / 1
Physics	26/60	43%	173	9.4 / 40	36.0/60 60.0%	▼ 16.6	8 / 9 / 0
Chemistry	46/57	81%	126	23.9 / 50	22.3/60 37.2%	▲ 43.5	13 / 3 / 0

Bar = your score % | campus average | best on campus Change = this test % − weekly-test average %

What this report says

1. **Chemistry** is your strongest JEE Mains subject: 67/100 (67%), All-India subject rank 294.
2. **Physics** needs the most work in JEE Mains: 38/100 (38%), subject rank 191; you lost 13 of its 25 questions.
3. **Chemistry** is your strongest JEE Advanced subject: 46/57 (81%), All-India subject rank 126.
4. **Physics** needs the most work in JEE Advanced: 26/60 (43%), subject rank 173; you lost 9 of its 17 questions.
5. Biggest fall against your weekly-test average: **JEE Mains Physics** – 38% here against 70% on average.
6. Biggest improvement: **JEE Advanced Chemistry** – 81% here against 37% on average.
7. **8 questions** you lost were answered correctly by 70%+ of the batch's top 100 – about **37 marks** within easy reach (Mains Q21, Q28, Q32, Q46, Q69 · Advanced Q18, Q38, Q42).
8. Negative marking cost you **20 in JEE Mains and 5 in JEE Advanced**.
9. You attempted almost everything (only 8 questions skipped across both papers), so the marks to gain are in **accuracy**: wrong answers – 20 in JEE Mains and 15 in JEE Advanced.

JEE Mains – question by question

168 / 296 · 56.8%
08 Oct 2026

2 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · ● Easy miss · ● Within reach · ● Tough one

Maths	1	2	3	**4**	5	6	7	8	9	10	11	12	13	14	15	**16**	**17**	18	**19**	~~20~~	**21**	22	**23**	24	**25**
Physics	26	**27**	**28**	29	30	31	**32**	33	34	**35**	**36**	37	38	39	**40**	**41**	42	**43**	44	**45**	**46**	**47**	**48**	**49**	50
Chemistry	51	52	**53**	**54**	**55**	56	57	**58**	59	60	61	62	63	64	**65**	66	**67**	68	**69**	70	71	72	73	74	75

3 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 63 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40–69% · **Tough one** below 40%

Maths

7 of 24 questions lost · **33** marks · ● 1 easy miss · ● 4 within reach · ● 2 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q21	Easy miss	**Surds & Indices**	Numerical · +4 / −1	Wrong	5	74%	54%
Q19	Within reach	**Sequences & Series**	Single correct · +4 / −1	Wrong	5	68%	44%
Q17	Within reach	**Surds & Indices** – Cube-root surd identity (matrix match)	Single correct · +4 / −1	Wrong	5	53%	41%
Q25	Within reach	**Surds & Indices**	Numerical · +4 / −1	Skipped	4	44%	21%
Q4	Within reach	**Sequences & Series** – Identify AP / GP / HP from a surd expression	Single correct · +4 / −1	Wrong	5	42%	19%
Q23	Tough one	**Surds & Indices** – Cube root of a surd expression	Numerical · +4 / −1	Skipped	4	24%	13%
Q16	Tough one	**Quadratic Equations** – Roots in GP, condition on discriminant	Single correct · +4 / −1	Wrong	5	19%	3%

Physics

13 of 25 questions lost · **62** marks · ● 3 easy miss · ● 3 within reach · ● 7 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q28	Easy miss	**Vectors** – Resultant ⊥ smaller vector; find magnitudes	Single correct · +4 / −1	**Wrong**	5	**77%**	30%
Q32	Easy miss	**Kinematics**	Single correct · +4 / −1	**Wrong**	5	**75%**	29%
Q46	Easy miss	**Vectors**	Numerical · +4 / −1	**Wrong**	5	**75%**	43%
Q40	Within reach	**Laws of Motion** – Velocity over fixed distance vs mass ($v \propto 1/\sqrt{m}$)	Single correct · +4 / −1	**Wrong**	5	**62%**	30%
Q27	Within reach	**Vectors** – Vector components under axis rotation	Single correct · +4 / −1	**Skipped**	4	**43%**	6%
Q35	Within reach	**Kinematics** – $a = kv^2$ – integrate to get v(s)	Single correct · +4 / −1	**Skipped**	4	**41%**	5%
Q41	Tough one	**Laws of Motion** – Newton's 3rd law – fan blowing on own sail	Single correct · +4 / −1	**Wrong**	5	**31%**	11%
Q45	Tough one	**Laws of Motion** – Contact force between pushed blocks (statement type)	Single correct · +4 / −1	**Wrong**	5	**27%**	10%
Q36	Tough one	**Kinematics** – a(t) with modulus – area under a-t	Single correct · +4 / −1	**Wrong**	5	**24%**	5%
Q43	Tough one	**Laws of Motion** – Normal reaction across an inclined contact	Single correct · +4 / −1	**Wrong**	5	**20%**	10%
Q47	Tough one	**Vectors** – Net displacement from multi-leg journey	Numerical · +4 / −1	**Skipped**	4	**15%**	3%
Q48	Tough one	**Kinematics** – Distance from a-t graph	Numerical · +4 / −1	**Wrong**	5	**15%**	3%
Q49	Tough one	**Kinematics** – Free fall then deceleration – find height	Numerical · +4 / −1	**Wrong**	5	**15%**	2%




Chemistry

7 of 25 questions lost · **33** marks · ● 1 easy miss · ● 5 within reach · ● 1 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q69	Easy miss	**Chemical Bonding**	Single correct · +4 / −1	**Wrong**	5	**75%**	44%
Q54	Within reach	**Mole Concept & Basics** – Minimum molecular mass of insulin from 3.4% S	Single correct · +4 / −1	**Skipped**	4	**67%**	25%
Q55	Within reach	**Mole Concept & Basics** – Laws of chemical combination	Single correct · +4 / −1	**Wrong**	5	**66%**	17%
Q53	Within reach	**Mole Concept & Basics** – Valence electrons in N^{3-} (mole → electrons)	Single correct · +4 / −1	**Skipped**	4	**65%**	25%
Q65	Within reach	**Periodic Classification** – Largest radius among isoelectronic ions	Single correct · +4 / −1	**Wrong**	5	**64%**	33%
Q58	Within reach	**Atomic Structure**	Single correct · +4 / −1	**Wrong**	5	**57%**	16%
Q67	Tough one	**Chemical Bonding** – Strongest ionic bond (charge and size)	Single correct · +4 / −1	**Wrong**	5	**32%**	13%

4 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Sequences & Series	5	**10**/20		33%	**Q4**, **Q19**
Surds & Indices	11	**26**/44		53%	**Q17**, **Q21**, **Q23**, **Q25**
Quadratic Equations	5	**15**/20		11%	**Q16**
Logarithms & Exponents	3	**12**/12		40%	– none
Physics					
Vectors	7	**10**/28		27%	**Q27**, **Q28**, **Q46**, **Q47**
Laws of Motion	8	**12**/32		7%	**Q40**, **Q41**, **Q43**, **Q45**
Kinematics	10	**16**/40		21%	**Q32**, **Q35**, **Q36**, **Q48**, **Q49**
Chemistry					
Chemical Bonding	6	**14**/24		34%	**Q67**, **Q69**
Mole Concept & Basics	9	**23**/36		32%	**Q53**, **Q54**, **Q55**
Atomic Structure	5	**15**/20		31%	**Q58**
Periodic Classification	5	**15**/20		58%	**Q65**

Bar = your % of the chapter's marks | campus average

5 JEE Mains · Maths – questions to revisit

7 questions · easy misses first
cropped from the question paper

Q21 — Surds & Indices

JEE Mains · Maths · Numerical · marking +4 / −1 · marks lost 5
Top 100: **74%** got it right · Campus: **54%** (34 of 63)

Easy miss
You: Wrong

21. If $x = \sqrt[3]{7+5\sqrt{2}} - \dfrac{1}{\sqrt[3]{7+5\sqrt{2}}}$ the value of the expression $x^3 + 3x - 14$ is ?

1 1

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q19 — Sequences & Series

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5
Top 100: **68%** got it right · Campus: **44%** (28 of 63)

Within reach
You: Wrong

19. If T_k denotes the kth term of an H.P from the beginning and $\dfrac{T_2}{T_6} = 9$, then $\dfrac{T_{10}}{T_4}$ is equal to

(A) $\dfrac{17}{5}$ (B) $\dfrac{5}{17}$ (C) $\dfrac{7}{19}$ (D) $\dfrac{9}{7}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q17

Surds & Indices – Cube-root surd identity (matrix match)

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5

Top 100: **53%** got it right · Campus: **41%** (26 of 63)

Within reach

You: Wrong

17. Let $x=\left(a+\sqrt{a^2+b^3}\right)^{\frac{1}{3}}+\left(a-\sqrt{a^2+b^3}\right)^{\frac{1}{3}}$ and $x^3+Abx-B.a=0$, Then match COLUMN-I with respect to COLUMN-II

	COLUMN-I		COLUMN-II
I)	A + B	P)	8
II)	A – B	Q)	9
III)	A^B	R)	1
IV)	B^A	S)	5
		T)	3

(A) I – S; II – R; III – Q; IV – P (B) I – S; II – T; III – P; IV – Q

(C) I – Q; II – P; III – T; IV – S (D) I – Q; II – P; III – S; IV – R

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q25

Surds & Indices

JEE Mains · Maths · Numerical · marking +4 / −1 · marks lost 4

Top 100: **44%** got it right · Campus: **21%** (13 of 63)

Within reach

You: Skipped

25. If $\left(2+\sqrt{3}\right)^{x^2-7x+1}+\left(2-\sqrt{3}\right)^{x^2-7x-1}=\frac{2}{2-\sqrt{3}}$, then the sum of values of 'x' satisfying this equation is equal to_____ ?

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q4

Sequences & Series – Identify AP / GP / HP from a surd expression

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5

Top 100: **42%** got it right · Campus: **19%** (12 of 63)

Within reach

You: Wrong

4. If a, b, c are in A.P., then $\frac{1}{\sqrt{a}+\sqrt{b}}, \frac{1}{\sqrt{a}+\sqrt{c}}, \frac{1}{\sqrt{b}+\sqrt{c}}$ are in

(A)A.P. (B)G.P. (C)H.P. (D)Neither A.P nor G.P nor H.P

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q23

Surds & Indices – Cube root of a surd expression

JEE Mains · Maths · Numerical · marking +4 / −1 · marks lost 4

Top 100: **24%** got it right · Campus: **13%** (8 of 63)

Tough one

You: Skipped

23. If $\sqrt[3]{38\sqrt{14}-100\sqrt{2}} = \sqrt{x}-\sqrt{y}$, then the value of $x-y$ is__?

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q16

Quadratic Equations – Roots in GP, condition on discriminant

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5

Top 100: **19%** got it right · Campus: **3%** (2 of 63)

Tough one

You: Wrong

16. Let α, β be the roots of the equation $ax^2+bx+c=0$ and $\Delta = b^2-4ac$. If $\alpha+\beta, \alpha^2+\beta^2, \alpha^3+\beta^3$ are in G.P., then

(A)$\Delta = 0$ (B)$\Delta \neq 0$ (C)$b\Delta = 0$ (D)$c\Delta = 0$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Mains · Physics – questions to revisit

13 questions · easy misses first
cropped from the question paper

Q28

Vectors – Resultant ⊥ smaller vector; find magnitudes
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **77%** got it right · Campus: **30%** (19 of 63)

Easy miss
You: Wrong

28. The simple sum of two co-initial vectors is 16 units . Their vector sum is 8 units. The resultant of the vectors is perpendicular to the smaller vector . The magnitudes of the two vectors are

(A) 22units and 14 units
(B) 4 units and 12 units
(C) 6 units and 10 units
(D) 8 units and 12 units

Why did I lose this question? (write the reason and the correct approach)


☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q32

Kinematics
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **75%** got it right · Campus: **29%** (18 of 63)

Easy miss
You: Wrong

32. The velocity versus time curve of a moving points is given below. The maximum acceleration is



(A) 1 cm/s^2 (B) 2 cm/s^2 (C) 3 cm/s^2 (D) 4 cm/s^2

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q46 — Vectors

JEE Mains · Physics · Numerical · marking +4 / −1 · marks lost 5
Top 100: **75%** got it right · Campus: **43%** (27 of 63)

Easy miss
You: Wrong

46. If the resultant of two vectors having magnitudes of 6 and 3 is 9, then the dot product of two vectors is:

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q40 — Laws of Motion – Velocity over fixed distance vs mass (v ∝ 1/√m)

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **62%** got it right · Campus: **30%** (19 of 63)

Within reach
You: Wrong

40. A constant force acts on a mass m at rest. Velocity acquired in travelling a fixed distance is directly proportional to

(A) $\sqrt{m}$ (B) m (C) $1/\sqrt{m}$ (D) $1/m$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q27 — Vectors – Vector components under axis rotation

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4
Top 100: **43%** got it right · Campus: **6%** (4 of 63)

Within reach
You: Skipped

27. The components of a vector along x and y directions are (n+1) and 1 respectively . If the co-ordinate system is rotated by an angle $\theta = 60^0$, then the components change to n and 3. The value of n is

(A) 2 (B) $\cos 60^0$ (C) $\sin 60^0$ (D) 3.5

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q35

Kinematics – a = kv² – integrate to get v(s)

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4

Top 100: **41%** got it right · Campus: **5%** (3 of 63)

Within reach

You: Skipped

35. If a particle accelerates with $a = kv^2$ and initial velocity = u, then find velocity after displacement s

(A) $v = ue^{ks}$ (B) $v = u^{ks}e^{s}$ (C) $v = u^{k}e^{s}$ (D) $v = eu^{ks}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q41

Laws of Motion – Newton's 3rd law – fan blowing on own sail

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **31%** got it right · Campus: **11%** (7 of 63)

Tough one

You: Wrong

41. On a stationary sail boat, air is blown at the sails from a fan attached to the boat. The boat will:

(A) Move forward (B) move backward

(C) spin around (D) remain stationary

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q45

Laws of Motion – Contact force between pushed blocks (statement type)


JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **27%** got it right · Campus: **10%** (6 of 63)

Tough one

You: Wrong

45. Statement-1: Block A is moving on a horizontal surface towards right under action of force F. all surfaces are smooth. At the instant shown the force exerted by block A on block B is equal to net force on block B.



Statement-2: From Newton's third law, the force exerted by block A on B is equal in magnitude to force exerted by block B on A.

(A) Statement 1 is true, statement 2 is true, statement 2 is correct explanation for statement 1

(B) Statement 1 is true, statement 2 is true, statement 2 is not correct explanation for statement 1

(C) Statement 1 is true, statement 2 is false

(D) Statement 1 is false, statement 2 is true

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q36

Kinematics – a(t) with modulus – area under a-t

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **24%** got it right · Campus: **5%** (3 of 63)

Tough one

You: Wrong

36. Starting from rest a particle moves in a straight line with acceleration $a = \{2 + |t-2|\}\, m/s^2$. Velocity of the particle at the end of 4s will be

(A) 16 m/s (B) 20 m/s (C) 8 m/s (D) 12 m/s

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q43

Laws of Motion – Normal reaction across an inclined contact


JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **20%** got it right · Campus: **10%** (6 of 63)

Tough one

You: Wrong

43. Two blocks 'A" and 'B' each of mass 'm' are placed on a smooth horizontal surface. Two horizontal forces F and 2F are applied on both the blocks 'A' and 'B' respectively as shown in figure. The block A does not slide on block B. Then the normal reaction acting between the two blocks is



(A) F (B) $F/2$ (C) $F/\sqrt{3}$ (D) 3F

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q47

Vectors – Net displacement from multi-leg journey

JEE Mains · Physics · Numerical · marking +4 / −1 · marks lost 4

Top 100: **15%** got it right · Campus: **3%** (2 of 63)

Tough one

You: Skipped

47. In going from one city to another , a car travels 75 km north, 60 km north –west and 20 km east. The magnitude(in km) of the vector drawn between the two cities is $\left(Take 1/\sqrt{2} = 0.7\right)$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q48

Kinematics – Distance from a-t graph


JEE Mains · Physics · Numerical · marking +4 / −1 · marks lost 5

Top 100: **15%** got it right · Campus: **3%** (2 of 63)

Tough one

You: Wrong

48. An experiment on the take off performance of an aeroplane shows that the acceleration varies as shown in the fig and it takes 12 sec to take off from a rest position.The distance along the runway covered by the aeroplane is(in m).



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q49

Kinematics – Free fall then deceleration – find height

JEE Mains · Physics · Numerical · marking +4 / −1 · marks lost 5

Top 100: **15%** got it right · Campus: **2%** (1 of 63)

Tough one

You: Wrong

49. A parachutist after bailing out falls 50m without friction. When parachute opens, it decelerates at $2m/\sec^2$. He reaches the ground with a speed of 3m/s. The height he bail out is nearly (in m) (Take g=10m/sec^2) .

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Mains · Chemistry – questions to revisit

7 questions · easy misses first
cropped from the question paper

Q69 Chemical Bonding

JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5
Top 100: **75%** got it right · Campus: **44%** (28 of 63)

Easy miss
You: Wrong

69. In which of the following species the bonds are non-directional ?
(A) NCl_3 (B) RbCl
(C) $BeCl_2$ (D) BCl_3

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q54 Mole Concept & Basics – Minimum molecular mass of insulin from 3.4% S

JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 4
Top 100: **67%** got it right · Campus: **25%** (16 of 63)

Within reach
You: Skipped

54. Insulin contains 3.4% sulphur. What will be the minimum molecular weight of insulin ?
(A) 94.176 (B) 1884 (C) 941.176 (D) 976

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q55 Mole Concept & Basics – Laws of chemical combination

JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5
Top 100: **66%** got it right · Campus: **17%** (11 of 63)

Within reach
You: Wrong

55. 12 g carbon combines with 64 g sulphur to form CS_2. 12 g carbon also combines with 32 g oxygen to form CO_2. 10 g sulphur combines with 10 g oxygen to form SO_2. These data illustrate the -
(A) Law of multiple proportions
(B) Law of definite proportions
(C) Law of reciprocal proportions
(D) Law of gaseous volumes.

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q53 Mole Concept & Basics – Valence electrons in N^{3-} (mole → electrons)

JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 4

Top 100: **65%** got it right · Campus: **25%** (16 of 63)

Within reach

You: Skipped

53. If N_A is Avogadro number, then the number of valence electrons in 4.2 g of N^{3-} ions is

(A) 2.4 N_A (B) 4.2 N_A (C) 1.6 N_A (D) 3.2 N_A

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q65 Periodic Classification – Largest radius among isoelectronic ions

JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5

Top 100: **64%** got it right · Campus: **33%** (21 of 63)

Within reach

You: Wrong

65. Which of the following has largest radius ?

(A) Mg^{2+} (B) Na^{+} (C) O^{2-} (D) F^{-}

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q58 Atomic Structure

JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5

Top 100: **57%** got it right · Campus: **16%** (10 of 63)

Within reach

You: Wrong

58. The ratio of the radii of first orbits of H, He^{+} and Li^{2+} is -

(A) 1 : 2 : 3 (B) 6 : 3 : 2 (C) 1 : 4 : 9 (D) 9 : 4 : 1

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q67 **Chemical Bonding – Strongest ionic bond (charge and size)**
JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5
Top 100: **32%** got it right · Campus: **13%** (8 of 63)

Tough one
You: Wrong

67. Which combination will give strongest ionic bond?

(A) K^+ and Cl^- (B) K^+ and O^{2-}

(C) Ca^{2+} and Cl^- (D) Ca^{2+} and O^{2-}

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced – question by question

106 / 171 · 62.0%
09 Oct 2026

6 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	2	3	4	**5**	6	7	8	9	10	11	**12**	**13**	~~14~~	15	**16**	~~17~~
Physics	**18**	19	**20**	21	22	23	24	25	26	**27**	28	**29**	**30**	**31**	**32**	**33**	**34**
Chemistry	35	36	37	**38**	39	40	41	**42**	43	44	45	46	47	**48**	49	~~50~~	51

7 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 63 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40–69% · **Tough one** below 40%

Maths

4 of 15 questions lost · **17** marks (+3 on partly-correct answers) · ● 2 within reach · ● 2 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q5	Within reach	**Quadratic Equations** – Quadratic with AP coefficients and integer roots (multiple correct)	Multiple correct · +4 / −2	Wrong	6	69%	62%
Q12	Within reach	**Sequences & Series**	Integer · +4 / 0	Wrong	4	68%	38%
Q16	Tough one	**Logarithms** – Log equations – A + B + C	Paragraph · +3 / 0	Wrong	3	18%	3%
Q13	Tough one	**Quadratic Equations** – Spot the polynomial identity, then sum P(1) … P(2023)	Integer · +4 / 0	Skipped	4	– no data*	2%

Physics

9 of 17 questions lost · **32** marks (+2 on partly-correct answers) · ● 1 easy miss · ● 2 within reach · ● 6 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q18	Easy miss	**Vectors**	Single correct · +3 / −1	Wrong	4	78%	46%
Q20	Within reach	**Kinematics**	Single correct · +3 / −1	Wrong	4	52%	22%
Q30	Within reach	**Kinematics** – a from v(t) at a given position	Integer · +4 / 0	Wrong	4	43%	8%
Q33	Tough one	**Kinematics** – Train passage – retardation to avoid collision	Paragraph · +3 / 0	Wrong	3	15%	8%
Q34	Tough one	**Kinematics** – Train passage – time of retardation	Paragraph · +3 / 0	Wrong	3	15%	10%
Q27	Tough one	**Laws of Motion** – Normal reaction between pushed blocks	Integer · +4 / 0	Wrong	4	11%	2%
Q31	Tough one	**Vectors** – Carrom passage – displacement centre → edge	Paragraph · +3 / 0	Wrong	3	7%	2%
Q29	Tough one	**Kinematics**	Integer · +4 / 0	Wrong	4	– no data*	6%
Q32	Tough one	**Vectors** – Carrom passage – displacement edge → hole	Paragraph · +3 / 0	Wrong	3	6%	2%

Chemistry

3 of 16 questions lost · **11** marks · ● 2 easy miss · ● 1 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q38	Easy miss	**Chemical Bonding** – Pair of species with the same shape (VSEPR)	Single correct · +3 / −1	**Wrong**	**4**	**92%**	49%
Q42	Easy miss	**Mole Concept & Basics**	Integer · +4 / 0	**Wrong**	**4**	**91%**	49%
Q48	Tough one	**Mole Concept & Basics** – Empirical formula from % composition	Paragraph · +3 / 0	**Wrong**	**3**	**0%**	59%

8 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject. Chapter marks are approximate (≈): partly-correct answers cannot be traced to a question.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Quadratic Equations	7	≈ **16**/26		43%	**Q5**, **Q13**
Sequences & Series	4	≈ **11**/15		56%	**Q12**
Logarithms	4	≈ **10**/13		50%	**Q16**
Physics					
Kinematics	8	≈ **9**/27		24%	**Q20**, **Q29**, **Q30**, **Q33**, **Q34**
Vectors	5	≈ **7**/17		23%	**Q18**, **Q31**, **Q32**
Laws of Motion	4	≈ **12**/16		6%	**Q27**
Chemistry					
Chemical Bonding	1	≈ **-1**/3		41%	**Q38**
Mole Concept & Basics	9	≈ **26**/33		44%	**Q42**, **Q48**
Atomic Structure	4	≈ **14**/14		33%	– none
Periodic Classification	2	≈ **7**/7		68%	– none

Bar = your % of the chapter's marks | campus average

9 JEE Advanced · Maths – questions to revisit

4 questions · easy misses first
cropped from the question paper

Q5

Quadratic Equations – **Quadratic with AP coefficients and integer roots (multiple correct)**
JEE Advanced · Maths · Multiple correct · marking +4 / −2 · marks lost 6
Top 100: **69%** got it right · Campus: **62%** (39 of 63)

Within reach
You: Wrong

5. Let a, b, c be positive integers in arithmetic progression such that the equation $ax^2 + bx + c = 0$ has only integer solutions. Then which of the following statements is (are) TRUE?

A) $c - b$ is an integer multiple of a

B) Both the roots of the equation $ax^2 + bx + c = 0$ are odd integers

C) If $c = 15$, then $ab = 8$

D) If $b = 8$, then $x = 3$ is a root of the equation $ax^2 + bx + c = 0$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q12

Sequences & Series
JEE Advanced · Maths · Integer · marking +4 / 0 · marks lost 4
Top 100: **68%** got it right · Campus: **38%** (24 of 63)

Within reach
You: Wrong

12. Consider a sequence whose sum to n terms is given by the quadratic function $S_n = 3n^2 + 5n$. For the given sequence, the number 5456 is m^{th} term then value of m is___?

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q16

Logarithms – Log equations – A + B + C

JEE Advanced · Maths · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **18%** got it right · Campus: **3%** (2 of 63)

Tough one

You: Wrong

PARAGRAPH-II

Let **A** denote the product xyz where x, y, z are solutions of the system of equations

$\log_3 x = \log_3 5 - \log_3 7, \log_5 y = \log_5 7 - \log_5 3, \log_7 z = \log_7 3 - \log_7 5$

Let **B** denote the sum of square of the solutions of the equation

$\log_2\left(\log_2 x^6 - 3\right) - \log_2\left(\log_2 x^4 - 5\right) = \log_2 3$

Let **C** denote the characteristic of the number K, where

$K = \log_2\left(\log_2 3\right) - \log_2\left(\log_4 3\right) + \log_2\left(\log_4 5\right) - \log_2\left(\log_6 5\right) + \log_2\left(\log_6 7\right) - \log_2\left(\log_8 7\right)$

16. The value of A+B+C is____?

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q13

Quadratic Equations – Spot the polynomial identity, then sum P(1) ... P(2023)

JEE Advanced · Maths · Integer · marking +4 / 0 · marks lost 4

Top 100: **no data** got it right · Campus: **2%** (1 of 63)

Tough one

You: Skipped

13. Let $P(x) = \left(1+\sqrt{2}\right)\left(1+\sqrt{3}\right)\left(x-\sqrt{2}\right)\left(x-\sqrt{3}\right) + \left(1+\sqrt{3}\right)\left(\sqrt{3}+\sqrt{2}\right)(x-1)\left(x-\sqrt{2}\right)$ $-2\left(1+\sqrt{2}\right)\left(\sqrt{3}+\sqrt{2}\right)(x-1)\left(x-\sqrt{3}\right)\ \forall x \in R$. Then the value of $P(1)+P(2)+P(3)+\ldots P(2023)$ is_____?

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced · Physics – questions to revisit

9 questions · easy misses first
cropped from the question paper

Q18 Vectors

JEE Advanced · Physics · Single correct · marking +3 / −1 · marks lost 4
Top 100: **78%** got it right · Campus: **46%** (29 of 63)

Easy miss
You: Wrong

18. The resultant of two forces, one double the other in magnitude, is perpendicular to the smaller of the two forces. The angle between the two forces is :

A) 120^0 B) 60^0 C) 90^0 D) 150^0

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q20 Kinematics

JEE Advanced · Physics · Single correct · marking +3 / −1 · marks lost 4
Top 100: **52%** got it right · Campus: **22%** (14 of 63)

Within reach
You: Wrong

20. A particle is projected vertically upwards from the ground at time $t = 0$ and reaches a height h at $t = T$. Greatest height of the particle is

A) $\left(gT^2+h\right)^2/8gT^2$ B) $2\left(gT^2+h\right)^2/8gT^2$

C) $\left(gT^2+2h\right)^2/8gT^2$ D) $2h$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q30 Kinematics – a from v(t) at a given position

JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4
Top 100: **43%** got it right · Campus: **8%** (5 of 63)

Within reach
You: Wrong

30. A body starts from origin and moves along x axis such that at any instant velocity is $v_t=4t^3-2t$ where t is in second and v_t is in ms^{-1}. The acceleration of the particle when it is 2m from the origin is $(20+K)\,ms^{-2}$, where K is

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q33

Kinematics – Train passage – retardation to avoid collision

JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **15%** got it right · Campus: **8%** (5 of 63)

Tough one

You: Wrong

PASSAGE-II:

The driver of a train A moving with a uniform speed of 144 km/h sights another train B, 1km ahead of him. The train B is moving with a uniform speed of 108 km/h. the driver of the train A immediately applies brakes producing a constant retardation and just manages to avoid a collision.

33. What is the retardation of the train A(in ms^{-2})_____

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q34

Kinematics – Train passage – time of retardation

JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **15%** got it right · Campus: **10%** (6 of 63)

Tough one

You: Wrong

PASSAGE-II:

The driver of a train A moving with a uniform speed of 144 km/h sights another train B, 1km ahead of him. The train B is moving with a uniform speed of 108 km/h. the driver of the train A immediately applies brakes producing a constant retardation and just manages to avoid a collision.

34. For how long is this retardation produced(in sec)_____

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q27

Laws of Motion – Normal reaction between pushed blocks


JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **11%** got it right · Campus: **2%** (1 of 63)

Tough one

You: Wrong

27. The normal reaction between 2 kg and 3 kg block is



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q31

Vectors – Carrom passage – displacement centre → edge

JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **7%** got it right · Campus: **2%** (1 of 63)

Tough one

You: Wrong

PASSAGE-1:

A Corrom board $(4\,ft \times 4\,ft\ square)$ has queen at the centre. The queen , hit by the striker moves to the front edge, rebounds and goes in the hole behind the striking line.

Answer the following question.

31. The magnitude of the displacement of the queen from the centre to the front edge is $\frac{2}{3}\sqrt{a}\,ft$, the value of a is _____

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q29

Kinematics

JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **no data** got it right · Campus: **6%** (4 of 63)

Tough one

You: Wrong

29. A lead ball is dropped freely into a lake from a diving board 5 m above the water. It strikes the water with a certain velocity and then reaches the bottom with the same constant velocity in 5 second after it is dropped.

Take g=10 ms^{-2} calculate the average velocity of the ball in ms^{-1} just before reaches to the bottom from the freely dropped point .

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q32 **Vectors – Carrom passage – displacement edge → hole**

JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **6%** got it right · Campus: **2%** (1 of 63)

Tough one

You: Wrong

PASSAGE-1:

A Corrom board $(4\,ft \times 4\,ft\ square)$ has queen at the centre. The queen , hit by the striker moves to the front edge, rebounds and goes in the hole behind the striking line.
Answer the following question.

32. The magnitude of the displacement of the queen from the front edge to the hole is $\frac{4}{3}\sqrt{b}\ ft$, the value of b is_____

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced · Chemistry — questions to revisit

3 questions · easy misses first
cropped from the question paper

Q38 — Chemical Bonding — Pair of species with the same shape (VSEPR)

JEE Advanced · Chemistry · Single correct · marking +3 / −1 · marks lost 4
Top 100: **92%** got it right · Campus: **49%** (31 of 63)

Easy miss — You: Wrong

38. The pair of species with similar shape is

(A) PCl_3, NH_3 (B) CF_4, SF_4 (C) $PbCl_2$, CO_2 (D) PF_5, IF_5

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q42 — Mole Concept & Basics

JEE Advanced · Chemistry · Integer · marking +4 / 0 · marks lost 4
Top 100: **91%** got it right · Campus: **49%** (31 of 63)

Easy miss — You: Wrong

42. How many moles of electrons present in 5.1 g NH_3 ?

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q48 — Mole Concept & Basics — Empirical formula from % composition

JEE Advanced · Chemistry · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **0%** got it right · Campus: **59%** (37 of 63)

Tough one — You: Wrong

PARAGRAPH-I

An unknown copper oxide is analysed. A 12.0 g sample contains 9.6 g Cu, so the remaining 2.4 g is oxygen. Taking Cu = 64 and O = 16, the moles of Cu and O are equal

48. What is the sum of the simplest whole-number ratio of Cu:O?

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

10 How to read this report

Where the numbers come from. Marks, percentages, All-India / State / Campus ranks, subject marks, subject All-India ranks and the wrong / skipped questions are the official results in the institute's *All India Marks Analysis* (graded with the final answer key). Maximum marks leave out deleted questions — JEE Mains: Q20 — JEE Advanced: Q14, Q17, Q50. **Program rank** = rank in Pragya G8 (EXI) Achievers across all campuses, from the batch's question-wise sheet († JEE Advanced: that sheet was graded before the final key, so this rank may differ slightly). **Top 100** = the batch's top 100 across all campuses, from the same question-wise sheet; * where it has no data for a question, the campus figure decides the priority. In JEE Advanced that sheet predates the final key, so its figures for the revised questions Q25, Q48, Q51 use the old answers. **Campus** = all students of this program on campus. **Priority:** Easy miss = 70%+ of the top 100 got it right; Within reach = 40–69%; Tough one = below 40%. **Marks lost** on a question = its marks plus the negative mark if answered wrong, its marks if skipped. In JEE Advanced, 5 marks went on multiple-correct answers that were only partly correct. **Weekly-test average** = your average over the weekly Mains tests (WTM) or Advanced tests (WTA) before this Part Test, from the campus averages sheet. The weekly Advanced tests are marked out of 240; their averages are scaled to 180 to match this paper (× 180/240, so 80 per subject becomes 60). Percentages are unchanged.