CLASS 8 · PART TEST-1 · STUDENT PERFORMANCE REPORT

Kolukula Hema Shiva Naga Hariharan

Pragya G8 (EXI) Achievers · Adm No **246562156**

Campus HYD/Nallakunta DD Colony
JEE Mains 08 Oct 2026
JEE Advanced 09 Oct 2026

JEE Mains

08 Oct 2026 · 74 questions

204 / 296 — **68.9%**

26	18	1/63	11
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **236.8 / 300** (78.9%) → this test **68.9%** ▼ **10.0**

92 marks lost: 80 on 16 wrong answers (incl. 16 negative) · 12 on 3 skipped

JEE Advanced

09 Oct 2026 · 48 questions

121 / 171 — **70.8%**

44	29	2/63	17†
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **132.7 / 180** (73.7%) → this test **70.8%** ▼ **2.9**

50 marks lost: 44 on 11 wrong answers (incl. 6 negative) · 4 on 1 skipped · 2 on partly-correct answers

1 Subject scorecard

Your marks in each subject, against the 63 students of this program on campus and against your own weekly-test average.

SUBJECT	MARKS	SCORE VS CAMPUS	SUBJECT AIR	CAMPUS AVG / BEST	WEEKLY AVG	CHANGE	RIGHT / WRONG / SKIP
JEE Mains 204 / 296 · 68.9%							
Maths	77/96	80% Best on campus	5	37.0 / 77	81.2/100 81.2%	▼ 1.0	20 / 3 / 1
Physics	47/100	47%	103	18.3 / 60	80.8/100 80.8%	▼ 33.8	14 / 9 / 2
Chemistry	80/100	80%	116	37.4 / 86	74.7/100 74.7%	▲ 5.3	21 / 4 / 0
JEE Advanced 121 / 171 · 70.8%							
Maths	34/54	63%	109	24.2 / 41	47.0/60 78.4%	▼ 15.4	11 / 4 / 0
Physics	40/60	67% Best on campus	38	9.4 / 40	47.7/60 79.5%	▼ 12.8	12 / 4 / 1
Chemistry	47/57	82%	103	23.9 / 50	38.1/60 63.6%	▲ 18.9	13 / 3 / 0

Bar = your score % | campus average | best on campus Change = this test % − weekly-test average %

What this report says

1. **Maths** is your strongest JEE Mains subject: 77/96 (80%), All-India subject rank 5 — the best score on campus.
2. **Physics** needs the most work in JEE Mains: 47/100 (47%), subject rank 103; you lost 11 of its 25 questions.
3. **Chemistry** is your strongest JEE Advanced subject: 47/57 (82%), All-India subject rank 103.
4. Your JEE Advanced **Physics** score (40/60) is the best on campus — All-India subject rank 38.
5. **Maths** needs the most work in JEE Advanced: 34/54 (63%), subject rank 109; you lost 4 of its 15 questions.
6. Biggest fall against your weekly-test average: **JEE Mains Physics** — 47% here against 81% on average.
7. Biggest improvement: **JEE Advanced Chemistry** — 82% here against 64% on average.
8. **5 questions** you lost were answered correctly by 70%+ of the batch's top 100 — about **25 marks** within easy reach (Mains Q33, Q37, Q39 · Advanced Q6, Q38).
9. Negative marking cost you **16 in JEE Mains and 6 in JEE Advanced**.
10. You attempted almost everything (only 4 questions skipped across both papers), so the marks to gain are in **accuracy**: wrong answers — 16 in JEE Mains and 11 in JEE Advanced.

JEE Mains – question by question

204 / 296 · 68.9%
08 Oct 2026

2 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	~~20~~	21	22	23	24	25
Physics	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50
Chemistry	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75

3 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 63 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40-69% · **Tough one** below 40%

Maths

4 of 24 questions lost · **19** marks · 3 within reach · 1 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q17	Within reach	**Surds & Indices** – Cube-root surd identity (matrix match)	Single correct · +4 / −1	Skipped	4	53%	41%
Q3	Within reach	**Sequences & Series** – Product of GP terms; natural-number GP	Single correct · +4 / −1	Wrong	5	42%	51%
Q4	Within reach	**Sequences & Series** – Identify AP / GP / HP from a surd expression	Single correct · +4 / −1	Wrong	5	42%	19%
Q16	Tough one	**Quadratic Equations** – Roots in GP, condition on discriminant	Single correct · +4 / −1	Wrong	5	19%	3%

Physics

11 of 25 questions lost · **53** marks · 3 easy miss · 3 within reach · 5 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q33	Easy miss	**Kinematics** – Which v–t graph can show real one-dimensional motion	Single correct · +4 / −1	Wrong	5	75%	65%
Q37	Easy miss	**Kinematics** – a–x graph from a given v–x graph ($a = v \cdot dv/dx$)	Single correct · +4 / −1	Wrong	5	71%	54%
Q39	Easy miss	**Kinematics** – a–x graph from a given v–x graph (same question as Q37)	Single correct · +4 / −1	Wrong	5	70%	52%
Q29	Within reach	**Vectors** – Angles between three vectors summing to zero	Single correct · +4 / −1	Wrong	5	62%	24%
Q30	Within reach	**Laws of Motion** – Tension / normal reaction in two-block systems (matrix match)	Single correct · +4 / −1	Wrong	5	42%	30%
Q35	Within reach	**Kinematics** – $a = kv^2$ – integrate to get v(s)	Single correct · +4 / −1	Skipped	4	41%	5%
Q41	Tough one	**Laws of Motion** – Newton's 3rd law – fan blowing on own sail	Single correct · +4 / −1	Wrong	5	31%	11%
Q42	Tough one	**Laws of Motion** – Extra tension when bird leaves the wire	Single correct · +4 / −1	Wrong	5	31%	14%
Q38	Tough one	**Kinematics** – Resistance $\propto v^2$ – time inside plank	Single correct · +4 / −1	Skipped	4	27%	11%
Q36	Tough one	**Kinematics** – a(t) with modulus – area under a-t	Single correct · +4 / −1	Wrong	5	24%	5%
Q43	Tough one	**Laws of Motion** – Normal reaction across an inclined contact	Single correct · +4 / −1	Wrong	5	20%	10%

Chemistry

4 of 25 questions lost · 20 marks · 3 within reach · 1 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q57	Within reach	**Atomic Structure** – Which electron jump gives the red line (smallest energy gap)	Single correct · +4 / −1	Wrong	5	66%	25%
Q65	Within reach	**Periodic Classification** – Largest radius among isoelectronic ions	Single correct · +4 / −1	Wrong	5	64%	33%
Q75	Within reach	**Chemical Bonding** – Hybrid orbitals + lone pairs in NH_3	Numerical · +4 / −1	Wrong	5	60%	30%
Q67	Tough one	**Chemical Bonding** – Strongest ionic bond (charge and size)	Single correct · +4 / −1	Wrong	5	32%	13%

4 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Sequences & Series	5	10/20		33%	Q3, Q4
Quadratic Equations	5	15/20		11%	Q16
Surds & Indices	11	40/44		53%	Q17
Logarithms & Exponents	3	12/12		40%	– none
Physics					
Kinematics	10	12/40		21%	Q33, Q35, Q36, Q37, Q38, Q39
Laws of Motion	8	12/32		7%	Q30, Q41, Q42, Q43
Vectors	7	23/28		27%	Q29
Chemistry					
Chemical Bonding	6	14/24		34%	Q67, Q75
Atomic Structure	5	15/20		31%	Q57
Periodic Classification	5	15/20		58%	Q65
Mole Concept & Basics	9	36/36		32%	– none

Bar = your % of the chapter's marks | campus average

5 JEE Mains · Maths – questions to revisit

4 questions · easy misses first
cropped from the question paper

Q17 — Surds & Indices – Cube-root surd identity (matrix match)

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4
Top 100: **53%** got it right · Campus: **41%** (26 of 63)

Within reach
You: Skipped

17. Let $x=\left(a+\sqrt{a^2+b^3}\right)^{\frac{1}{3}}+\left(a-\sqrt{a^2+b^3}\right)^{\frac{1}{3}}$ and $x^3+Abx-B.a=0$, Then match

COLUMN-I with respect to COLUMN-II

	COLUMN-I		COLUMN-II
I)	A + B	P)	8
II)	A – B	Q)	9
III)	A^B	R)	1
IV)	B^A	S)	5
		T)	3

(A) I – S; II – R; III – Q; IV – P (B) I – S; II – T; III – P; IV – Q
(C) I – Q; II – P; III – T; IV – S (D) I – Q; II – P; III – S; IV – R

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q3 — Sequences & Series – Product of GP terms; natural-number GP

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5
Top 100: **42%** got it right · Campus: **51%** (32 of 63)

Within reach
You: Wrong

3. If $x_1, x_2, x_3, x_4, x_5, \ldots$ is a geometric progression of natural numbers and $x_1x_2x_3x_4 = 64$, then complete set of solutions of x_4 is

(A){1} (B){8} (C){1, 8} (D)cannot be determined

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q4

Sequences & Series – Identify AP / GP / HP from a surd expression

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5

Top 100: **42%** got it right · Campus: **19%** (12 of 63)

Within reach

You: Wrong

4. If a, b, c are in A.P., then $\frac{1}{\sqrt{a}+\sqrt{b}}, \frac{1}{\sqrt{a}+\sqrt{c}}, \frac{1}{\sqrt{b}+\sqrt{c}}$ are in

(A)A.P. (B)G.P. (C)H.P. (D)Neither A.P nor G.P nor H.P

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q16

Quadratic Equations – Roots in GP, condition on discriminant

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5

Top 100: **19%** got it right · Campus: **3%** (2 of 63)

Tough one

You: Wrong

16. Let α, β be the roots of the equation $ax^2+bx+c=0$ and $\Delta = b^2-4ac$. If $\alpha+\beta, \alpha^2+\beta^2, \alpha^3+\beta^3$ are in G.P., then

(A)$\Delta = 0$ (B)$\Delta \neq 0$ (C)$b\Delta = 0$ (D)$c\Delta = 0$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q33 **Kinematics – Which v-t graph can show real one-dimensional motion**


JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5


Top 100: **75%** got it right · Campus: **65%** (41 of 63)


Easy miss


You: Wrong

33. From following graph which of these possibly represents one-dimensional motion of the particle.

(A) 

(B) 

(C) 

(D) 

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q37 **Kinematics – a-x graph from a given v-x graph (a = v·dv/dx)**


JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **71%** got it right · Campus: **54%** (34 of 63)


Easy miss

You: Wrong


37. The v – x graph for a car in a race on a straight road is given. Identify the correct $a-x$ graph.




(A)




(B)



(C)



(D)



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q39

Kinematics – a-x graph from a given v-x graph (same question as Q37)


JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **70%** got it right · Campus: **52%** (33 of 63)


Easy miss

You: Wrong


39. The v – x graph for a car in a race on a straight road is given. Identify the correct $a-x$ graph.




(A)




(B)



(C)



(D)



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q29

Vectors – Angles between three vectors summing to zero

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **62%** got it right · Campus: **24%** (15 of 63)

Within reach

You: Wrong

29. Given: $\vec{a}+\vec{b}+\vec{c}=0$. Out of the three vectors $\vec{a},\vec{b}$ and $\vec{c}$, two are equal in magnitude. The magnitude of the third vector is $\sqrt{2}$ times that of either of the two having equal magnitude. The angles between the vector are

(A) $90^0,135^0,135^0$

(B) $30^0,60^0,90^0$

(C) $45^0,45^0,90^0$

(D) $45^0,60^0,90^0$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q30

Laws of Motion – Tension / normal reaction in two-block systems (matrix match)

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **42%** got it right · Campus: **30%** (19 of 63)

Within reach

You: Wrong

30. Column-I gives four different situations involving two blocks of mass m_1 and m_2 placed in different ways on a smooth horizontal surface as shown. In each of the situations horizontal forces F_1 and F_2 are applied on blocks of mass m_1 and m_2 respectively and also $m_2 F_1 < m_1 F_2$. Match the statements in column I with corresponding results in column-II

Column-I	Column-II
(A)  . Both the blocks are connected by massless inelastic string. The magnitude of tension in the string is	(p) $\frac{m_1 m_2}{m_1+m_2}\left(\frac{F_1}{m_1}-\frac{F_2}{m_2}\right)$
(B)  . Both the blocks are connected by massless inelastic string. The magnitude of tension in the string is	(q) $\frac{m_1 m_2}{m_1+m_2}\left(\frac{F_1}{m_1}+\frac{F_2}{m_2}\right)$
(C)  . The magnitude of normal reaction between the blocks is	(r) $\frac{m_1 m_2}{m_1+m_2}\left(\frac{F_2}{m_2}-\frac{F_1}{m_1}\right)$
(D)  . The magnitude of normal reaction between the blocks is	(s) $m_1 m_2\left(\frac{F_1+F_2}{m_1+m_2}\right)$

(A) (A) – r ; (B) – q ; (C) – q ; (D) – r

(B) (A) – q ; (B) – r ; (C) – r ; (D) – q

(C) (A) – q ; (B) – r ; (C) – q ; (D) – r

(D) (A) – q ; (B) – p ; (C) – r ; (D) – r

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q35

Kinematics – a = kv² – integrate to get v(s)

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4

Top 100: **41%** got it right · Campus: **5%** (3 of 63)

Within reach

You: Skipped

35. If a particle accelerates with $a = kv^2$ and initial velocity = u, then find velocity after displacement s

(A) $v = ue^{ks}$ (B) $v = u^{ks}e^{s}$ (C) $v = u^{k}e^{s}$ (D) $v = eu^{ks}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q41

Laws of Motion – Newton's 3rd law – fan blowing on own sail

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **31%** got it right · Campus: **11%** (7 of 63)

Tough one

You: Wrong

41. On a stationary sail boat, air is blown at the sails from a fan attached to the boat. The boat will:

(A) Move forward (B) move backward

(C) spin around (D) remain stationary

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q42

Laws of Motion – Extra tension when bird leaves the wire

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **31%** got it right · Campus: **14%** (9 of 63)

Tough one

You: Wrong

42. A bird dismounts on a stretched telegraph wire. The additional tension produced in the wire is

(A) Zero (B) greater than the weight of the bird

(C) Less than the weight of the bird (D) equal to the weight of the bird

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q38

Kinematics – Resistance ∝ v² – time inside plank

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4

Top 100: **27%** got it right · Campus: **11%** (7 of 63)

Tough one

You: Skipped

38. Having gone through a plank of thickness h, a bullet changed its velocity from v_0 to v. Find the time of motion of the bullet in the plank assuming resistance force to be proportional to the square of the velocity

(A) $\left[\frac{h(v_0-v)}{vv_0 l\,n\frac{v_0}{v}}\right]$ (B) $\left[\frac{(v_0-v)}{hvv_0 l\,n\left(\frac{v_0}{v}\right)}\right]$ (C) $\left[\frac{vv_0 \ln\left(\frac{v_0}{v}\right)}{h(v_0-v)}\right]$ (D) $\left[\frac{\ln\frac{v_0}{v}}{vv_0(v_0-v)}\right]$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q36

Kinematics – a(t) with modulus – area under a-t

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **24%** got it right · Campus: **5%** (3 of 63)

Tough one

You: Wrong

36. Starting from rest a particle moves in a straight line with acceleration $a = \{2 + |t-2|\}\, m/s^2$. Velocity of the particle at the end of 4s will be

(A) 16 m/s (B) 20 m/s (C) 8 m/s (D) 12 m/s

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q43

Laws of Motion – Normal reaction across an inclined contact


JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **20%** got it right · Campus: **10%** (6 of 63)

Tough one

You: Wrong

43. Two blocks 'A" and 'B' each of mass 'm' are placed on a smooth horizontal surface. Two horizontal forces F and 2F are applied on both the blocks 'A' and 'B' respectively as shown in figure. The block A does not slide on block B. Then the normal reaction acting between the two blocks is



(A) F (B) $F/2$ (C) $F/\sqrt{3}$ (D) 3F

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Mains · Chemistry – questions to revisit

4 questions · easy misses first
cropped from the question paper

Q57

Atomic Structure – Which electron jump gives the red line (smallest energy gap)
JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5
Top 100: **66%** got it right · Campus: **25%** (16 of 63)

Within reach
You: Wrong

57. Suppose that a hypothetical H-like atom gives a red Green, blue & violet line spectrum. Which jump according to figure would give off the red spectral line.



(A) 3 – 1 (B) 2 → 1 (C) 4 – 1 (D) 3 →2

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q65

Periodic Classification – Largest radius among isoelectronic ions
JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5
Top 100: **64%** got it right · Campus: **33%** (21 of 63)

Within reach
You: Wrong

65. Which of the following has largest radius ?
(A) Mg^{2+} (B) Na^{+} (C) O^{2-} (D) F^{-}

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q75

Chemical Bonding – Hybrid orbitals + lone pairs in NH_3
JEE Mains · Chemistry · Numerical · marking +4 / −1 · marks lost 5
Top 100: **60%** got it right · Campus: **30%** (19 of 63)

Within reach
You: Wrong

75. According to VBT, consider the molecule NH_3. The central nitrogen atom undergoes hybridisation before bond formation.
If the number of hybrid orbitals formed is x, and the number of lone pairs present on nitrogen is y, find:
x + y

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q67 **Chemical Bonding – Strongest ionic bond (charge and size)**

JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5

Top 100: **32%** got it right · Campus: **13%** (8 of 63)

Tough one

You: Wrong

67. Which combination will give strongest ionic bond?

(A) K^+ and Cl^- (B) K^+ and O^{2-}

(C) Ca^{2+} and Cl^- (D) Ca^{2+} and O^{2-}

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced – question by question

121 / 171 · 70.8%
09 Oct 2026

6 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	2	3	**4**	**5**	**6**	7	8	9	10	11	12	**13**	~~14~~	15	16	~~17~~
Physics	18	19	20	21	22	23	24	25	26	**27**	**28**	29	**30**	**31**	**32**	33	34
Chemistry	35	36	37	**38**	39	40	41	42	43	44	45	46	47	48	**49**	~~50~~	**51**

7 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 63 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40-69% · **Tough one** below 40%

Maths

4 of 15 questions lost · **20** marks · 1 easy miss · 2 within reach · 1 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q6	Easy miss	**Quadratic Equations** – Other root of $4x^2 + 2x - 1 = 0$ in terms of α	Multiple correct · +4 / −2	Wrong	6	82%	67%
Q5	Within reach	**Quadratic Equations** – Quadratic with AP coefficients and integer roots (multiple correct)	Multiple correct · +4 / −2	Wrong	6	69%	62%
Q4	Within reach	**Logarithms** – Logs in AP – solve for x	Single correct · +3 / −1	Wrong	4	68%	35%
Q13	Tough one	**Quadratic Equations** – Spot the polynomial identity, then sum P(1) … P(2023)	Integer · +4 / 0	Wrong	4	– no data*	2%

Physics

5 of 17 questions lost · **18** marks (+2 on partly-correct answers) · 1 within reach · 4 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q30	Within reach	**Kinematics** – a from v(t) at a given position	Integer · +4 / 0	Wrong	4	43%	8%
Q28	Tough one	**Laws of Motion** – Spring-connected blocks – acceleration	Integer · +4 / 0	Skipped	4	25%	13%
Q27	Tough one	**Laws of Motion** – Normal reaction between pushed blocks	Integer · +4 / 0	Wrong	4	11%	2%
Q31	Tough one	**Vectors** – Carrom passage – displacement centre → edge	Paragraph · +3 / 0	Wrong	3	7%	2%
Q32	Tough one	**Vectors** – Carrom passage – displacement edge → hole	Paragraph · +3 / 0	Wrong	3	6%	2%



Chemistry

3 of 16 questions lost · **10** marks · 1 easy miss · 1 within reach · 1 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q38	Easy miss	**Chemical Bonding** – Pair of species with the same shape (VSEPR)	Single correct · +3 / −1	Wrong	4	92%	49%
Q49	Within reach	**Mole Concept & Basics** – Mass of oxygen in 24 g of the same copper oxide	Paragraph · +3 / 0	Wrong	3	60%	49%
Q51	Tough one	**Atomic Structure** – Hydrogen-like ion – find Z from photon energies	Paragraph · +3 / 0	Wrong	3	0%	5%

8 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject. Chapter marks are approximate (≈): partly-correct answers cannot be traced to a question.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Quadratic Equations	7	≈ **10**/26		43%	**Q5**, **Q6**, **Q13**
Logarithms	4	≈ **9**/13		50%	**Q4**
Sequences & Series	4	≈ **15**/15		56%	– none
Physics					
Laws of Motion	4	≈ **8**/16		6%	**Q27**, **Q28**
Vectors	5	≈ **11**/17		23%	**Q31**, **Q32**
Kinematics	8	≈ **23**/27		24%	**Q30**
Chemistry					
Chemical Bonding	1	≈ **-1**/3		41%	**Q38**
Atomic Structure	4	≈ **11**/14		33%	**Q51**
Mole Concept & Basics	9	≈ **30**/33		44%	**Q49**
Periodic Classification	2	≈ **7**/7		68%	– none

Bar = your % of the chapter's marks | campus average

9 JEE Advanced · Maths – questions to revisit

4 questions · easy misses first
cropped from the question paper

Q6 — Quadratic Equations – Other root of 4x² + 2x – 1 = 0 in terms of α

JEE Advanced · Maths · Multiple correct · marking +4 / –2 · marks lost 6
Top 100: **82%** got it right · Campus: **67%** (42 of 63)

Easy miss
You: Wrong

6. If α is a root of the equation $4x^2+2x-1=0$, then it's other root is given by

A) $4\alpha^3-3\alpha$ B) $4\alpha^3+3\alpha$ C) $\alpha-\frac{1}{2}$ D) $-\alpha-\frac{1}{2}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q5 — Quadratic Equations – Quadratic with AP coefficients and integer roots (multiple correct)

JEE Advanced · Maths · Multiple correct · marking +4 / –2 · marks lost 6
Top 100: **69%** got it right · Campus: **62%** (39 of 63)

Within reach
You: Wrong

5. Let a, b, c be positive integers in arithmetic progression such that the equation $ax^2+bx+c=0$ has only integer solutions. Then which of the following statements is (are) TRUE?

A) $c-b$ is an integer multiple of a

B) Both the roots of the equation $ax^2+bx+c=0$ are odd integers

C) If $c=15$, then $ab=8$

D) If $b=8$, then $x=3$ is a root of the equation $ax^2+bx+c=0$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q4 — Logarithms – Logs in AP – solve for x

JEE Advanced · Maths · Single correct · marking +3 / –1 · marks lost 4
Top 100: **68%** got it right · Campus: **35%** (22 of 63)

Within reach
You: Wrong

4. If $\log_3 2, \log_3(2^x-5)$ & $\log_3\left(2^x-\frac{7}{2}\right)$ are in AP then the value of x is?

A)2 B)3 C)4 D)2&3

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q13 **Quadratic Equations – Spot the polynomial identity, then sum P(1) ... P(2023)**

JEE Advanced · Maths · Integer · marking +4 / 0 · marks lost 4

Top 100: **no data** got it right · Campus: **2%** (1 of 63)

Tough one

You: Wrong

13. Let $P(x)=(1+\sqrt{2})(1+\sqrt{3})(x-\sqrt{2})(x-\sqrt{3})+(1+\sqrt{3})(\sqrt{3}+\sqrt{2})(x-1)(x-\sqrt{2})$ $-2(1+\sqrt{2})(\sqrt{3}+\sqrt{2})(x-1)(x-\sqrt{3})\ \forall x\in R$. Then the value of $P(1)+P(2)+P(3)+...P(2023)$ is_____?

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced · Physics – questions to revisit

5 questions · easy misses first
cropped from the question paper

Q30

Kinematics – a from v(t) at a given position
JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4
Top 100: **43%** got it right · Campus: **8%** (5 of 63)

Within reach
You: Wrong

30. A body starts from origin and moves along x axis such that at any instant velocity is $v_t=4t^3-2t$ where t is in second and v_t is in ms^{-1}. The acceleration of the particle when it is 2m from the origin is $(20+K)\,ms^{-2}$, where K is

Why did I lose this question? (write the reason and the correct approach)


☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q28

Laws of Motion – Spring-connected blocks – acceleration
JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4
Top 100: **25%** got it right · Campus: **13%** (8 of 63)

Tough one
You: Skipped

28. If the acceleration of 3 kg block is 4 m/s^2 to the left. What is the magnitude of acceleration of 8 kg block.



Why did I lose this question? (write the reason and the correct approach)


☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q27

Laws of Motion – Normal reaction between pushed blocks
JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4
Top 100: **11%** got it right · Campus: **2%** (1 of 63)

Tough one
You: Wrong

27. The normal reaction between 2 kg and 3 kg block is



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q31

Vectors – Carrom passage – displacement centre → edge

JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **7%** got it right · Campus: **2%** (1 of 63)

Tough one

You: Wrong

PASSAGE-1:

A Corrom board $(4\,ft \times 4\,ft\ square)$ has queen at the centre. The queen , hit by the striker moves to the front edge, rebounds and goes in the hole behind the striking line.

Answer the following question.

31. The magnitude of the displacement of the queen from the centre to the front edge is $\frac{2}{3}\sqrt{a}\,ft$, the value of a is _____

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q32

Vectors – Carrom passage – displacement edge → hole

JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **6%** got it right · Campus: **2%** (1 of 63)

Tough one

You: Wrong

PASSAGE-1:

A Corrom board $(4\,ft \times 4\,ft\ square)$ has queen at the centre. The queen , hit by the striker moves to the front edge, rebounds and goes in the hole behind the striking line.

Answer the following question.

32. The magnitude of the displacement of the queen from the front edge to the hole is $\frac{4}{3}\sqrt{b}\,ft$, the value of b is_____

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced · Chemistry – questions to revisit

3 questions · easy misses first
cropped from the question paper

Q38 — Chemical Bonding – Pair of species with the same shape (VSEPR)

JEE Advanced · Chemistry · Single correct · marking +3 / −1 · marks lost 4
Top 100: **92%** got it right · Campus: **49%** (31 of 63)

Easy miss
You: Wrong

38. The pair of species with similar shape is

(A) PCl_3, NH_3 (B) CF_4, SF_4 (C) $PbCl_2$, CO_2 (D) PF_5, IF_5

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q49 — Mole Concept & Basics – Mass of oxygen in 24 g of the same copper oxide

JEE Advanced · Chemistry · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **60%** got it right · Campus: **49%** (31 of 63)

Within reach
You: Wrong

PARAGRAPH-I

An unknown copper oxide is analysed. A 12.0 g sample contains 9.6 g Cu, so the remaining 2.4 g is oxygen. Taking Cu = 64 and O = 16, the moles of Cu and O are equal

49. If the same oxide has the same composition, what mass of oxygen is present in a 24 g sample?

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q51 **Atomic Structure – Hydrogen-like ion – find Z from photon energies**

JEE Advanced · Chemistry · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **0%** got it right · Campus: **5%** (3 of 63)

Tough one

You: Wrong

PARAGRAPH-II

When electron jumps from lower orbit to higher orbit energy is gained and when electron jumps from higher orbit to lower orbit energy is released.

51. A hydrogen like species (atomic number Z) is present in a higher excited state of quantum number n. This excited atom can make a transition to the first excited state by successive emission of two photons of energies 10.20 eV and 17.0 eV respectively. Alternatively the atom from the same excited state can make a transition to the second excited state by successive emission of two photons of energy 4.25 eV and 5.95 eV respectively. Determine the value of Z.

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

10 How to read this report

Where the numbers come from. Marks, percentages, All-India / State / Campus ranks, subject marks, subject All-India ranks and the wrong / skipped questions are the official results in the institute's *All India Marks Analysis* (graded with the final answer key). Maximum marks leave out deleted questions — JEE Mains: Q20 — JEE Advanced: Q14, Q17, Q50. **Program rank** = rank in Pragya G8 (EXI) Achievers across all campuses, from the batch's question-wise sheet († JEE Advanced: that sheet was graded before the final key, so this rank may differ slightly). **Top 100** = the batch's top 100 across all campuses, from the same question-wise sheet; * where it has no data for a question, the campus figure decides the priority. In JEE Advanced that sheet predates the final key, so its figures for the revised questions Q25, Q48, Q51 use the old answers. **Campus** = all students of this program on campus. **Priority:** Easy miss = 70%+ of the top 100 got it right; Within reach = 40–69%; Tough one = below 40%. **Marks lost** on a question = its marks plus the negative mark if answered wrong, its marks if skipped. In JEE Advanced, 2 marks went on multiple-correct answers that were only partly correct. **Weekly-test average** = your average over the weekly Mains tests (WTM) or Advanced tests (WTA) before this Part Test, from the campus averages sheet. The weekly Advanced tests are marked out of 240; their averages are scaled to 180 to match this paper (× 180/240, so 80 per subject becomes 60). Percentages are unchanged.