CLASS 10 · PART TEST-1 · STUDENT PERFORMANCE REPORT

Vihaan Bussari

Pragya G10 (EXT) Achievers · Adm No **235417357**

Campus HYD/Nallakunta DD Colony
JEE Mains 08 Oct 2026
JEE Advanced 09 Oct 2026

JEE Mains

08 Oct 2026 · 74 questions

196 / 296 — **66.2%**

55	41	16/65	18
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **213.3 / 300** (71.1%) → this test **66.2%** ▼ **4.9**

100 marks lost: 60 on 12 wrong answers (incl. 12 negative) · 40 on 10 skipped

JEE Advanced

09 Oct 2026 · 50 questions

118 / 176 — **67.0%**

50	34	8/61	12
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **106.7 / 180** (59.3%) → this test **67.0%** ▲ **7.8**

58 marks lost: 58 on 14 wrong answers (incl. 8 negative) · 0 on 0 skipped

1 Subject scorecard

Your marks in each subject, against the 65 students of this program on campus and against your own weekly-test average.

SUBJECT	MARKS	SCORE VS CAMPUS	SUBJECT AIR	CAMPUS AVG / BEST	WEEKLY AVG	CHANGE	RIGHT / WRONG / SKIP
JEE Mains 196 / 296 · 66.2%							
Maths	63/100	63%	113	58.2 / 100	66.7/100 66.7%	▼ 3.7	16 / 1 / 8
Physics	62/100	62%	155	53.8 / 100	70.5/100 70.5%	▼ 8.5	17 / 6 / 2
Chemistry	71/96	74%	22	55.9 / 91	76.0/100 76.0%	▼ 2.0	19 / 5 / 0
JEE Advanced 118 / 176 · 67.0%							
Maths	44/60	73%	55	30.5 / 57	41.5/60 69.2%	▲ 4.1	14 / 3 / 0
Physics	25/56	45%	273	24.5 / 56	27.5/60 45.8%	▼ 1.2	8 / 8 / 0
Chemistry	49/60	82%	35	38.2 / 60	37.7/60 62.8%	▲ 18.9	14 / 3 / 0

Bar = your score % | campus average | best on campus Change = this test % − weekly-test average %

What this report says

1. **Chemistry** is your strongest JEE Mains subject: 71/96 (74%), All-India subject rank 22.
2. **Physics** needs the most work in JEE Mains: 62/100 (62%), subject rank 155; you lost 8 of its 25 questions.
3. **Chemistry** is your strongest JEE Advanced subject: 49/60 (82%), All-India subject rank 35.
4. **Physics** needs the most work in JEE Advanced: 25/56 (45%), subject rank 273; you lost 8 of its 16 questions.
5. Biggest fall against your weekly-test average: **JEE Mains Physics** – 62% here against 71% on average.
6. Biggest improvement: **JEE Advanced Chemistry** – 82% here against 63% on average.
7. **3 questions** you lost were answered correctly by 70%+ of the batch's top 100 – about **16 marks** within easy reach (Mains Q10, Q36 · Advanced Q24).
8. Negative marking cost you **12 in JEE Mains and 8 in JEE Advanced**.
9. You attempted almost everything (only 10 questions skipped across both papers), so the marks to gain are in **accuracy**: wrong answers – 12 in JEE Mains and 14 in JEE Advanced.

JEE Mains – question by question

196 / 296 · 66.2%
08 Oct 2026

2 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25
Physics	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50
Chemistry	51	52	53	54	55	~~56~~	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75

3 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 65 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40-69% · **Tough one** below 40%

Maths

9 of 25 questions lost · 37 marks · 1 easy miss · 4 within reach · 4 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q10	Easy miss	Differential Equations	Single correct · +4 / −1	Wrong	5	76%	85%
Q9	Within reach	Indefinite Integration	Single correct · +4 / −1	Skipped	4	66%	63%
Q8	Within reach	Definite Integration	Single correct · +4 / −1	Skipped	4	56%	38%
Q12	Within reach	Indefinite Integration	Single correct · +4 / −1	Skipped	4	51%	43%
Q5	Within reach	Definite Integration	Single correct · +4 / −1	Skipped	4	40%	32%
Q14	Tough one	Definite Integration	Single correct · +4 / −1	Skipped	4	39%	34%
Q18	Tough one	Definite Integration – $\int$ of min{|x − 3k|} – periodic zig-zag	Single correct · +4 / −1	Skipped	4	37%	40%
Q6	Tough one	Indefinite Integration – $\int \sqrt{(x^2+1)}\,[\ln(x^2+1) - 2\ln x] / x^4\,dx$	Single correct · +4 / −1	Skipped	4	33%	31%
Q11	Tough one	Definite Integration	Single correct · +4 / −1	Skipped	4	29%	26%

Physics

8 of 25 questions lost · 38 marks · 1 easy miss · 5 within reach · 2 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q36	Easy miss	Current Electricity	Single correct · +4 / −1	Wrong	5	73%	77%
Q41	Within reach	Kinematics	Single correct · +4 / −1	Wrong	5	65%	71%
Q37	Within reach	Current Electricity – Infinite ladder – voltage across the 2nd shunt resistor	Single correct · +4 / −1	Wrong	5	52%	46%
Q35	Within reach	Current Electricity	Single correct · +4 / −1	Wrong	5	50%	48%
Q34	Within reach	Current Electricity	Single correct · +4 / −1	Skipped	4	45%	37%
Q47	Within reach	Electrostatics	Numerical · +4 / −1	Skipped	4	42%	34%
Q46	Tough one	Electrostatics	Numerical · +4 / −1	Wrong	5	25%	25%
Q30	Tough one	Electrostatics – Field energy inside vs outside a charged sphere	Single correct · +4 / −1	Wrong	5	17%	11%




Chemistry

5 of 24 questions lost · 25 marks · 3 within reach · 2 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q53	Within reach	Alcohols, Phenols & Ethers	Single correct · +4 / −1	Wrong	5	68%	69%
Q71	Within reach	Alcohols, Phenols & Ethers	Numerical · +4 / −1	Wrong	5	55%	68%
Q55	Within reach	Alcohols, Phenols & Ethers	Single correct · +4 / −1	Wrong	5	54%	71%
Q52	Tough one	Alcohols, Phenols & Ethers – Ether vs butanol miscibility – assertion / reason	Single correct · +4 / −1	Wrong	5	24%	22%
Q74	Tough one	Alcohols, Phenols & Ethers – Site of benzylation of 2-naphthoxide	Numerical · +4 / −1	Wrong	5	12%	15%

4 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Definite Integration	9	16/36		29%	Q5, Q8, Q11, Q14, Q18
Indefinite Integration	7	16/28		61%	Q6, Q9, Q12
Differential Equations	4	11/16		84%	Q10
Area Under Curves	5	20/20		86%	– none
Physics					
Current Electricity	8	13/32		57%	Q34, Q35, Q36, Q37
Electrostatics	8	18/32		27%	Q30, Q46, Q47
Kinematics	9	31/36		75%	Q41
Chemistry					
Alcohols, Phenols & Ethers	10	15/40		48%	Q52, Q53, Q55, Q71, Q74
Aldehydes & Ketones	14	56/56		65%	– none

Bar = your % of the chapter's marks | campus average

5 JEE Mains · Maths – questions to revisit

9 questions · easy misses first
cropped from the question paper

Q10 Differential Equations

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5
Top 100: **76%** got it right · Campus: **85%** (55 of 65)

Easy miss
You: Wrong

10. The solution of $\frac{xdy}{x^2+y^2}=\left(\frac{y}{x^2+y^2}-1\right)dx$ is_____ (c is arbitrary constant)

A) $y=x\cot(c-x)$ B) $y=x\tan(c-x)$ C) $y=x\cos(c-x)$ D) $y=x\sin(c-x)$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q9 Indefinite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4
Top 100: **66%** got it right · Campus: **63%** (41 of 65)

Within reach
You: Skipped

9. $\int\frac{dx}{\sin x+\sec x}=$ (c is arbitrary constant)

A) $\frac{1}{2\sqrt{3}}\ln\left|\frac{\sin x-\cos x+\sqrt{3}}{\cos x-\sin x+\sqrt{3}}\right|+\tan^{-1}(\sin x+\cos x)+c$

B) $\frac{1}{2\sqrt{3}}\ln\left|\frac{\sin x-\cos x-\sqrt{3}}{\sin x-\cos x+\sqrt{3}}\right|+c$

C) $\frac{1}{2\sqrt{3}}\ln\left|(\sin x-\cos x+\sqrt{3})(\sin x-\cos x-\sqrt{3})\right|+c$

D) $\frac{1}{2\sqrt{3}}\ln\left|(\sin x-\cos x+\sqrt{3})(\sin x-\cos x-\sqrt{3})\right|+\tan^{-1}(\sin x+\cos x)+c$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q8

Definite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **56%** got it right · Campus: **38%** (25 of 65)

Within reach

You: Skipped

8. If $\int_0^{\frac{\pi}{2}} \frac{\tan x}{\tan x+\sec x}dx = m(\pi+n)$, then $m\times n$ is equal to

A) $-\frac{1}{2}$ B) 1 C) -1 D) $\frac{1}{2}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q12

Indefinite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **51%** got it right · Campus: **43%** (28 of 65)

Within reach

You: Skipped

12. $\int \sqrt{\sec x-1}\,dx =$ _____ (c is arbitrary constant)

A) $2\ln\left|\cos\frac{x}{2}+\sqrt{\cos^2\frac{x}{2}+\frac{1}{2}}\right|+c$ B) $\ln\left|\cos\frac{x}{2}-\sqrt{\cos^2\frac{x}{2}-\frac{1}{2}}\right|+c$

C) $-2\ln\left|\cos\frac{x}{2}+\sqrt{\cos^2\frac{x}{2}-\frac{1}{2}}\right|+c$ D) $\ln\left|\cos\frac{x}{2}-\sqrt{\cos^2\frac{x}{2}+\frac{1}{2}}\right|+c$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q5

Definite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **40%** got it right · Campus: **32%** (21 of 65)

Within reach

You: Skipped

5. $\int_{-1}^{1} \sin x \ln\left(1^x+2^x+3^x+6^x\right)dx =$

A) $(\ln 6)\left(\cos\left(\frac{\pi+4}{4}\right)\right)$ B) $(\ln 6)\left(\sin\left(\frac{\pi+4}{4}\right)\right)$

C) $\sqrt{2}(\ln 6)\left(\sin\left(\frac{4-\pi}{4}\right)\right)$ D) $\sqrt{2}(\ln 6)\left(\sin\left(\frac{\pi+4}{4}\right)\right)$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q14 — Definite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **39%** got it right · Campus: **34%** (22 of 65)

Tough one

You: Skipped

14. Let $I_1=\int_0^{\frac{\pi}{2}}\ln(\sin x)\ln(\sin 2x)\,dx$ and $I_2=\int_0^{\frac{\pi}{2}}(\ln(\sin 2x))^2\,dx$ then $\dfrac{\pi(\ln 2)^2}{2I_1-I_2}=$

A) $\frac{1}{2}$ B) 2 C) -2 D) $-\frac{1}{2}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q18 — Definite Integration – ∫ of min{|x − 3k|} – periodic zig-zag

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **37%** got it right · Campus: **40%** (26 of 65)

Tough one

You: Skipped

18. Let $f(x)=\min\{\ldots,|x+6|,|x+3|,|x|,|x-3|,|x-6|,\ldots\}$ then $\int_{-4}^{3}f(x)\,dx=$ _____

A) 4 B) $\frac{9}{2}$ C) 5 D) $\frac{11}{2}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q6 — Indefinite Integration – ∫ √(x²+1)[ln(x²+1) − 2 ln x] / x⁴ dx

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **33%** got it right · Campus: **31%** (20 of 65)

Tough one

You: Skipped

6. $\int\dfrac{\sqrt{x^2+1}\left[\ln(x^2+1)-2\ln x\right]}{x^4}\,dx$ is equal to (c is arbitrary constant)

A) $\dfrac{(x^2+1)^{3/2}}{x^3}\left[\dfrac{2}{3}-\ln\left(\dfrac{x^2+1}{x^2}\right)\right]+c$ B) $\dfrac{(x^2+1)^{3/2}}{3x^3}\left[\ln\left(\dfrac{x^2+1}{x^2}\right)-\dfrac{2}{3}\right]+c$

C) $\dfrac{2(x^2+1)^{3/2}}{3x^3}\left[\dfrac{1}{3}-\dfrac{1}{2}\ln\left(\dfrac{x^2+1}{x^2}\right)\right]+c$ D) $\dfrac{(x^2+1)^{3/2}}{3x^3}\left[\ln\left(\dfrac{x^2+1}{x^2}\right)+\dfrac{2}{3}\right]+c$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q11 **Definite Integration**

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **29%** got it right · Campus: **26%** (17 of 65)

Tough one

You: Skipped

11. Let $f(x)=\sin x+\sum_{r=0}^{2}\cos\left(x+\frac{2r\pi}{3}\right)$, $g(x)=\cos x+\sum_{r=0}^{2}\sin\left(x+\frac{2r\pi}{3}\right)$ then $\int_{0}^{\frac{\pi}{2}} f(g(x))f(x)\,dx=$

A) $1+\sin 1$ B) $1-\sin 1$ C) $1+\cos 1$ D) $1-\cos 1$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Mains · Physics – questions to revisit

8 questions · easy misses first
cropped from the question paper

Q36

Current Electricity
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **73%** got it right · Campus: **77%** (50 of 65)

Easy miss
You: Wrong

36. A copper wire carries a steady current I with drift velocity ‘v’. If the wire is replaced by one of the same material but double the diameter carrying the same current, the drift velocity becomes

A) $\frac{v}{2}$ B) $4v$ C) v D) $\frac{v}{4}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q41

Kinematics
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **65%** got it right · Campus: **71%** (46 of 65)

Within reach
You: Wrong

41. A swimmer can swim at 5 m/s in still water. A river 120 m wide flows at 3 m/s. If the swimmer heads so as to reach the point directly opposite on the other bank, the time taken is

A) 24 s B) 40 s C) 15 s D) 30 s

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q37 **Current Electricity – Infinite ladder – voltage across the 2nd shunt resistor**
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **52%** got it right · Campus: **46%** (30 of 65)

Within reach
You: Wrong

37. A 24 V battery of negligible internal resistance is connected to an infinite ladder network. In each section, both the top rail and the bottom rail contain a 1 Ω resistor, and a 4 Ω resistor is connected across the rails. The potential difference across the second 4 Ω resistor (counting from the battery) is

A) 12 V B) 8 V C) 6 V D) 3 V

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q35 **Current Electricity**
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **50%** got it right · Campus: **48%** (31 of 65)

Within reach
You: Wrong

35. A 12 V battery of negligible internal resistance is connected to the input terminals of an infinite ladder network. Each section of the ladder has a 1 Ω resistor in the top (series) arm and a 2 Ω resistor connected across the two rails. The current through the first 2 Ω shunt resistor is

A) 6 A B) 3 A C) 2 A D) 4 A

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q34 **Current Electricity**
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4
Top 100: **45%** got it right · Campus: **37%** (24 of 65)

Within reach
You: Skipped

34. In a Wheatstone bridge ABCD, a 12 V battery of negligible internal resistance is connected between A and C. The arms are AB = 2 Ω, BC = 3 Ω, AD = 3 Ω and DC = 3 Ω. An ammeter of resistance 0.3 Ω is connected between B and D. The current through the ammeter is

A) 0.2 A B) 0.6 A C) 0.4 A D) 1.2 A

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q47 **Electrostatics**
JEE Mains · Physics · Numerical · marking +4 / −1 · marks lost 4
Top 100: **42%** got it right · Campus: **34%** (22 of 65)

Within reach
You: Skipped

47. A non-conducting sphere of radius R has a volume charge density $\rho = \rho_0 \left(\frac{r}{R}\right)^2$, where 'r' is the distance from the centre. The electric field at its surface is $\rho_0 \frac{R}{n\varepsilon_0}$. The value of n is _____.

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q46 **Electrostatics**
JEE Mains · Physics · Numerical · marking +4 / −1 · marks lost 5
Top 100: **25%** got it right · Campus: **25%** (16 of 65)

Tough one
You: Wrong

46. In a region the electric field is $E = \left(5x\hat{i} + 3y\hat{j} + 4\hat{k}\right) N/C$, with x, y in meters. A cube of side 1m has one vertex at the origin and its edges along the positive x, y and z axis. The net electric flux through the cube (in Nm^2/C) is _____.

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q30 **Electrostatics – Field energy inside vs outside a charged sphere**
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **17%** got it right · Campus: **11%** (7 of 65)

Tough one
You: Wrong

30. A solid sphere of radius R carries a total charge Q distributed uniformly throughout its volume. The ratio of the electrostatic field energy stored inside the sphere to that stored outside the sphere is

A) 3 : 5 B) 1 : 5 C) 1 : 3 D) 1 : 2

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Mains · Chemistry — questions to revisit

5 questions · easy misses first
cropped from the question paper

Q53

Alcohols, Phenols & Ethers
JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5
Top 100: **68%** got it right · Campus: **69%** (45 of 65)

Within reach
You: Wrong

53. Phenol industrially prepared from

A) Benzene sulphonic acid
B) Cumene
C) Benzene diazonium chloride
D) Chlorobenzene

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q71

Alcohols, Phenols & Ethers
JEE Mains · Chemistry · Numerical · marking +4 / −1 · marks lost 5
Top 100: **55%** got it right · Campus: **68%** (44 of 65)

Within reach
You: Wrong

71. PCl_3 and an alcohol can react in x:y molar ratio to form corresponding alkyl halide and an oxyacid of basicity z. What is $\left(\frac{y}{x}+z\right)$?


Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q55 **Alcohols, Phenols & Ethers**
JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5
Top 100: **54%** got it right · Campus: **71%** (46 of 65)

Within reach
You: Wrong

55. Given below are two statements for the



$$C_2H_5O^- + H_2O \rightleftharpoons C_2H_5OH + OH^-; K_2$$

Statement I : The value of K_2 less than K_1

Statement II : Phenoxide is a weaker base than ethoxide

In the light of the above statement, choose the most appropriate answer from the options given below.

A) Both Statement I and Statement II are correct

B) Both Statement I and Statement II are incorrect

C) Statement I is correct but Statement II is incorrect

D) Statement II is correct but Statement I is incorrect

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q52 **Alcohols, Phenols & Ethers** – Ether vs butanol miscibility – assertion / reason
JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5
Top 100: **24%** got it right · Campus: **22%** (14 of 65)

Tough one
You: Wrong

52. **Assertion(A) :** Ethoxy ethane and butanol-1 have almost same miscibility with water.

Reason(R) : Ethoxy ethane and butanol-1 have form Hydrogen bond with water.

A) Both (A) and (R) are correct and (R) is the correct explanation of (A)

B) (A) is correct but (R) is not correct

C) Both (A) and (R) are not correct

D) Both (A) and (R) are correct and (R) is not the correct explanation of (A)


Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q74 **Alcohols, Phenols & Ethers – Site of benzylation of 2-naphthoxide**
JEE Mains · Chemistry · Numerical · marking +4 / −1 · marks lost 5
Top 100: **12%** got it right · Campus: **15%** (10 of 65)

Tough one
You: Wrong

74. Which site is predominately benzylated in the given reaction?



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced – question by question

118 / 176 · 67.0%
09 Oct 2026

6 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
Physics	18	19	20	21	22	23	24	~~25~~	26	27	28	29	30	31	32	33	34
Chemistry	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51

7 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 61 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40-69% · **Tough one** below 40%

Maths

3 of 17 questions lost · **16** marks · ● 3 within reach

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q6	Within reach	**Area Under Curves** – Line x = α halving the area between x^3 and x	Multiple correct · +4 / −2	Wrong	6	68%	69%
Q7	Within reach	**Differential Equations** – Curve from tangent-intercept ratio BP : AP = 3 : 1	Multiple correct · +4 / −2	Wrong	6	43%	26%
Q2	Within reach	**Definite Integration** – $f(a) = \int \ln(1+x^a)/(1+x)\,dx$ – find f(4) + f(1/4)	Single correct · +3 / −1	Wrong	4	42%	34%

Physics

8 of 16 questions lost · **31** marks · ● 1 easy miss · ● 4 within reach · ● 3 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q24	Easy miss	**Kinematics**	Multiple correct · +4 / −2	Wrong	6	71%	70%
Q19	Within reach	**Electrostatics**	Single correct · +3 / −1	Wrong	4	66%	56%
Q28	Within reach	**Electrostatics** – Flux through a cube with charges at its vertices	Integer · +4 / 0	Wrong	4	54%	44%
Q29	Within reach	**Electrostatics**	Integer · +4 / 0	Wrong	4	54%	34%
Q33	Within reach	**Electrostatics** – Flux through a ring – solid angle	Paragraph · +3 / 0	Wrong	3	42%	38%
Q27	Tough one	**Kinematics** – Projection angle that maximises area under the trajectory	Integer · +4 / 0	Wrong	4	27%	23%
Q31	Tough one	**Kinematics** – Hexagon projectile – maximum height	Paragraph · +3 / 0	Wrong	3	5%	5%
Q32	Tough one	**Kinematics** – Hexagon projectile – speed at the top	Paragraph · +3 / 0	Wrong	3	5%	5%









Chemistry

3 of 17 questions lost · **11** marks · ● 1 within reach · ● 2 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q44	Within reach	**Carboxylic Acids**	Integer · +4 / 0	Wrong	4	68%	80%
Q50	Tough one	**Carboxylic Acids** – Molar mass of an acid from NaOH titration	Paragraph · +3 / 0	Wrong	3	36%	52%
Q46	Tough one	**Alcohols, Phenols & Ethers** – Identifying polar aprotic solvents	Integer · +4 / 0	Wrong	4	24%	8%

8 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject. Chapter marks are approximate (≈): partly-correct answers cannot be traced to a question.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Area Under Curves	3	≈ 4/10		60%	Q6
Differential Equations	5	≈ 12/18		30%	Q7
Definite Integration	5	≈ 13/17		54%	Q2
Indefinite Integration	4	≈ 15/15		71%	– none
Physics					
Electrostatics	6	≈ 6/21		45%	Q19, Q28, Q29, Q33
Kinematics	8	≈ 12/28		41%	Q24, Q27, Q31, Q32
Current Electricity	2	≈ 7/7		68%	– none
Chemistry					
Carboxylic Acids	4	≈ 7/14		48%	Q44, Q50
Alcohols, Phenols & Ethers	7	≈ 22/26		62%	Q46
Aldehydes & Ketones	6	≈ 20/20		83%	– none

Bar = your % of the chapter's marks campus average

9 JEE Advanced · Maths – questions to revisit

3 questions · easy misses first
cropped from the question paper

Q6

Area Under Curves – Line x = α halving the area between x^3 and x

JEE Advanced · Maths · Multiple correct · marking +4 / −2 · marks lost 6

Top 100: **68%** got it right · Campus: **69%** (42 of 61)

Within reach

You: Wrong

6. If the line x=α divides the area of the region $R=\{(x,y)\in R^2 : x^3 \le y \le x, 0 \le x \le 1\}$ into two equal parts, then

A) $\frac{1}{2} \le \alpha \le 1$ B) $\alpha^4+4\alpha^2-1=0$ C) $0 \le \alpha \le \frac{1}{2}$ D) $2\alpha^4-4\alpha^2+1=0$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q7

Differential Equations – Curve from tangent-intercept ratio BP : AP = 3 : 1

JEE Advanced · Maths · Multiple correct · marking +4 / −2 · marks lost 6

Top 100: **43%** got it right · Campus: **26%** (16 of 61)

Within reach

You: Wrong

7. A curve $y=f(x)$ passes through the point (1,1) and tangent at P(x,y) cuts the x-axis and y-axis at A and B, respectively such that BP:AP=3:1, then

A) Equation of the curve is $xy^3=1$

B) Equation of normal at the point (1,1) is $x-3y+2=0$

C) Curve passes through the point $\left(\frac{1}{8},2\right)$

D) Equation of the curve is $x\frac{dy}{dx}+3y=0$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q2 **Definite Integration – f(a) = ∫ ln(1+x^a)/(1+x) dx – find f(4) + f(1/4)**

JEE Advanced · Maths · Single correct · marking +3 / −1 · marks lost 4

Top 100: **42%** got it right · Campus: **34%** (21 of 61)

Within reach

You: Wrong

2. Let $f(a)=\int_0^1 \frac{\ln\left(1+x^a\right)}{1+x}dx$ then $f(4)+f\left(\frac{1}{4}\right)$ is equal to

A) $\ln^2 1$ B) $\ln^2 2$ C) $\ln^2 3$ D) $\ln^2 4$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced · Physics – questions to revisit

8 questions · easy misses first
cropped from the question paper

Q24

Kinematics
JEE Advanced · Physics · Multiple correct · marking +4 / −2 · marks lost 6
Top 100: **71%** got it right · Campus: **70%** (43 of 61)

Easy miss
You: Wrong

24. Choose the correct statements from following

A) An accelerated body must always have non-zero velocity.

B) A body at rest may have non-zero acceleration.

C) A body moving with constant speed must have constant velocity.

D) A body moving with constant velocity must have constant speed.

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q19

Electrostatics
JEE Advanced · Physics · Single correct · marking +3 / −1 · marks lost 4
Top 100: **66%** got it right · Campus: **56%** (34 of 61)

Within reach
You: Wrong

19. The magnitude of electric field intensity at point B (2,0,0) due to a dipole moment $\vec{P} = \hat{i} + \sqrt{3}\hat{j}$ C-m kept at origin is (assume point B is at large distance from dipole)

$$k = \frac{1}{4\pi\varepsilon_0}$$

A) $\frac{\sqrt{13}k}{8}$ B) $\frac{\sqrt{13}k}{4}$ C) $\frac{\sqrt{7}k}{8}$ D) $\frac{\sqrt{7}k}{4}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q28

Electrostatics – Flux through a cube with charges at its vertices

JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **54%** got it right · Campus: **44%** (27 of 61)

Within reach

You: Wrong

28. A cube has charges each of value +Q placed at the vertices. The flux associated with the cube is $(X-1)\frac{Q}{\varepsilon_0}$, value of x is:

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q29

Electrostatics


JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **54%** got it right · Campus: **34%** (21 of 61)

Within reach

You: Wrong

29. An electron is projected from a distance d and with initial velocity u parallel to a uniformly charge flat conducting plate as shown. It strikes the plate after traveling a distance l along the direction of projection. The surface charge density of the conducting plate is $\sigma = \frac{nd\varepsilon_0 mu^2}{el^2}$, The value of n is:



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q33

Electrostatics – Flux through a ring – solid angle

JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **42%** got it right · Campus: **38%** (23 of 61)

Within reach

You: Wrong


PARAGRAPH-II

The electric flux linked with a surface is the number of electric field lines associated with that area. Gauss law helps determine the flux

$$\oint \vec{E}.\vec{ds} = \frac{Q_{inc}}{\varepsilon_0}$$

For an open surface the flux will change depending on the solid angle. Considering this answer the following questions.

33.



The flux associated with the ring due to the charge q is given by Xq/ε_0. The value of X is approx.

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q27

Kinematics – Projection angle that maximises area under the trajectory

JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **27%** got it right · Campus: **23%** (14 of 61)

Tough one

You: Wrong

27. A projectile is thrown with a speed 10 m/s, at an angle θ with the horizontal. For a certain value of angle of projection $\theta = \frac{(n+1)\pi}{12}$ the area bound by the trajectory with horizontal is maximum value of n is ($g=10\ m/s^2$). Assume no airdrag.

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q31 **Kinematics – Hexagon projectile – maximum height**


JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **5%** got it right · Campus: **5%** (3 of 61)

Tough one

You: Wrong

PARAGRAPH-I



A regular hexagon of side a=$10\sqrt{3}$ m is shown in figure. From any point a particle is projected such that it touches B,C,D and E. The velocity of particle is v m/s. O is the centre of the hexagon. x-axis is parallel to horizontal and y-axis is perpendicular to AE and CD (g=10 m/s^2) $\left(\sqrt{3}=1.73\right)$.

31. The maximum height from the horizontal surface X' attained by the projectile is ___ m.


Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q32 **Kinematics – Hexagon projectile – speed at the top**
JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **5%** got it right · Campus: **5%** (3 of 61)

Tough one
You: Wrong

PARAGRAPH-I



A regular hexagon of side a=$10\sqrt{3}$ m is shown in figure. From any point a particle is projected such that it touches B,C,D and E. The velocity of particle is v m/s. O is the centre of the hexagon. x-axis is parallel to horizontal and y-axis is perpendicular to AE and CD (g=10 m/s^2) $\left(\sqrt{3}=1.73\right)$.

32. The velocity of projectile at maximum height is _____ m/s

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced · Chemistry – questions to revisit


3 questions · easy misses first
cropped from the question paper

Q44 — Carboxylic Acids

JEE Advanced · Chemistry · Integer · marking +4 / 0 · marks lost 4
Top 100: **68%** got it right · Campus: **80%** (49 of 61)

Within reach
You: Wrong

44. How many of the following acids will show higher reactivity towards esterification reaction as compared to acetic acid?



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q50 — Carboxylic Acids – Molar mass of an acid from NaOH titration

JEE Advanced · Chemistry · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **36%** got it right · Campus: **52%** (32 of 61)

Tough one
You: Wrong

PARAGRAPH-II

Carboxylic acids are organic compounds containing the carbonyl(>C=O) and hydroxyl (-OH) functional groups combined as a carboxyl group(-COOH). Due to resonance stabilization of the carboxylate anion, carboxylic acids exhibit higher acidity compared to alcohols and phenols. They participate in several characteristic reactions, including nucleophilic acyl substitution (esterification, amide formation, acid halide synthesis), decarboxylation, and reduction to primary alcohols. Furthermore, the presence of the –COOH group strongly influences physical properties, enabling extensive intermolecular hydrogen bonding that results in higher boiling points and stable dimeric structures in non-polar solvents.

50. A 0.600 g sample of a pure monoprotic carboxylic acid (R-COOH) requires 25.00 mL of 0.200 M NaOH for complete neutralization. Calculate the molar mass of the carboxylic acid?

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q46 **Alcohols, Phenols & Ethers** – **Identifying polar aprotic solvents**


JEE Advanced · Chemistry · Integer · marking +4 / 0 · marks lost 4

Top 100: **24%** got it right · Campus: **8%** (5 of 61)

Tough one

You: Wrong

46. How many of the following are polar aprotic solvents?



[P]

Diethyl ether,


[Q]

Benzene,

[R]

Ethyl alcohol,

[S]



[T]

Petrol

[U]

Acetone

[Q]

DMSO (dimethyl sulphoxide

[W]

DMF (dimethyl formamide

[X]

formic acid

[Y]

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

10 How to read this report

Where the numbers come from. Marks, percentages, All-India / State / Campus ranks, subject marks, subject All-India ranks and the wrong / skipped questions are the official results in the institute's *All India Marks Analysis* (graded with the final answer key). Maximum marks leave out deleted questions — JEE Mains: Q56 — JEE Advanced: Q25. **Program rank** = rank in Pragya G10 (EXT) Achievers across all campuses, from the batch's question-wise sheet. **Top 100** = the batch's top 100 across all campuses, from the same question-wise sheet; * where it has no data for a question, the campus figure decides the priority. **Campus** = all students of this program on campus. **Priority:** Easy miss = 70%+ of the top 100 got it right; Within reach = 40–69%; Tough one = below 40%. **Marks lost** on a question = its marks plus the negative mark if answered wrong, its marks if skipped. **Weekly-test average** = your average over the weekly Mains tests (WTM) or Advanced tests (WTA) before this Part Test, from the campus averages sheet. The weekly Advanced tests are marked out of 240; their averages are scaled to 180 to match this paper (× 180/240, so 80 per subject becomes 60). Percentages are unchanged.