CLASS 10 · PART TEST-1 · STUDENT PERFORMANCE REPORT

Yapala Dhanush Reddy

Pragya G10 (EXT) Achievers · Adm No **235416325**

Campus HYD/Nallakunta DD Colony
JEE Mains 08 Oct 2026
JEE Advanced 09 Oct 2026

JEE Mains

08 Oct 2026 · 74 questions

186 / 296 — **62.8%**

70	52	20/65	24
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **204.1 / 300** (68.0%), 5 of 6 tests → this test **62.8%** ▼ **5.2**

110 marks lost: 50 on 10 wrong answers (incl. 10 negative) · 60 on 15 skipped

JEE Advanced

09 Oct 2026 · 50 questions

118 / 176 — **67.0%**

55	39	10/61	14
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **104.2 / 180** (57.9%) → this test **67.0%** ▲ **9.2**

58 marks lost: 28 on 8 wrong answers (incl. 3 negative) · 30 on 8 skipped

1 Subject scorecard

Your marks in each subject, against the 65 students of this program on campus and against your own weekly-test average.

SUBJECT	MARKS	SCORE VS CAMPUS	SUBJECT AIR	CAMPUS AVG / BEST	WEEKLY AVG	CHANGE	RIGHT / WRONG / SKIP
JEE Mains 186 / 296 · 62.8%							
Maths	42/100	42%	359	58.2 / 100	62.9/100 62.9%	▼ 20.9	11 / 2 / 12
Physics	77/100	77%	49	53.8 / 100	81.9/100 81.9%	▼ 4.9	20 / 3 / 2
Chemistry	67/96	70%	35	55.9 / 91	59.4/100 59.4%	▲ 10.4	18 / 5 / 1
JEE Advanced 118 / 176 · 67.0%							
Maths	23/60	38%	417	30.5 / 57	35.3/60 58.8%	▼ 20.4	7 / 3 / 7
Physics	46/56	82%	10	24.5 / 56	37.9/60 63.2%	▲ 18.9	13 / 3 / 0
Chemistry	49/60	82%	35	38.2 / 60	31.0/60 51.6%	▲ 30.0	14 / 2 / 1

Bar = your score % | campus average | best on campus Change = this test % − weekly-test average %

What this report says

1. **Physics** is your strongest JEE Mains subject: 77/100 (77%), All-India subject rank 49.
2. **Maths** needs the most work in JEE Mains: 42/100 (42%), subject rank 359; you lost 14 of its 25 questions.
3. **Physics** is your strongest JEE Advanced subject: 46/56 (82%), All-India subject rank 10.
4. **Maths** needs the most work in JEE Advanced: 23/60 (38%), subject rank 417; you lost 10 of its 17 questions.
5. Biggest fall against your weekly-test average: **JEE Mains Maths** – 42% here against 63% on average.
6. Biggest improvement: **JEE Advanced Chemistry** – 82% here against 52% on average.
7. **10 questions** you lost were answered correctly by 70%+ of the batch's top 100 – about **44 marks** within easy reach (Mains Q10, Q13, Q17, Q26, Q36, Q45, Q67 · Advanced Q4, Q5, Q36).
8. Negative marking cost you **10 in JEE Mains and 3 in JEE Advanced**.
9. You left **15 in JEE Mains and 8 in JEE Advanced** questions blank, and answered 10 in JEE Mains and 8 in JEE Advanced wrong – both are marks to recover.

JEE Mains – question by question

186 / 296 · 62.8%
08 Oct 2026

2 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25
Physics	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50
Chemistry	51	52	53	54	55	~~56~~	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75

3 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 65 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40-69% · **Tough one** below 40%

Maths

14 of 25 questions lost · **58** marks · 3 easy miss · 5 within reach · 6 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q13	Easy miss	Differential Equations	Single correct · +4 / −1	Wrong	5	86%	91%
Q10	Easy miss	Differential Equations	Single correct · +4 / −1	Skipped	4	76%	85%
Q17	Easy miss	Indefinite Integration	Single correct · +4 / −1	Skipped	4	74%	66%
Q19	Within reach	Differential Equations	Single correct · +4 / −1	Skipped	4	68%	77%
Q9	Within reach	Indefinite Integration	Single correct · +4 / −1	Skipped	4	66%	63%
Q23	Within reach	Area Under Curves	Numerical · +4 / −1	Wrong	5	66%	72%
Q8	Within reach	Definite Integration	Single correct · +4 / −1	Skipped	4	56%	38%
Q12	Within reach	Indefinite Integration	Single correct · +4 / −1	Skipped	4	51%	43%
Q18	Tough one	**Definite Integration** – ∫ of min{|x − 3k|} – periodic zig-zag	Single correct · +4 / −1	Skipped	4	37%	40%
Q6	Tough one	**Indefinite Integration** – ∫ $\sqrt{(x^2+1)}$ [ln(x^2+1) − 2 ln x] / x^4 dx	Single correct · +4 / −1	Skipped	4	33%	31%
Q11	Tough one	**Definite Integration**	Single correct · +4 / −1	Skipped	4	29%	26%
Q20	Tough one	**Definite Integration** – King's property with f(x) + f(y) = f(xy)	Single correct · +4 / −1	Skipped	4	29%	32%
Q22	Tough one	**Definite Integration** – Definite integral of a product × sum of reciprocals	Numerical · +4 / −1	Skipped	4	20%	20%
Q16	Tough one	**Definite Integration** – $\int_0^{100}$ ([$\tan^{-1}$x] + [$\cot^{-1}$x]) dx with GIF	Single correct · +4 / −1	Skipped	4	16%	11%

Physics

5 of 25 questions lost · **23** marks · 3 easy miss · 1 within reach · 1 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q45	Easy miss	Kinematics	Single correct · +4 / −1	Wrong	5	89%	88%
Q36	Easy miss	Current Electricity	Single correct · +4 / −1	Wrong	5	73%	77%
Q26	Easy miss	Electrostatics	Single correct · +4 / −1	Wrong	5	70%	69%
Q37	Within reach	**Current Electricity** – Infinite ladder – voltage across the 2nd shunt resistor	Single correct · +4 / −1	Skipped	4	52%	46%
Q30	Tough one	**Electrostatics** – Field energy inside vs outside a charged sphere	Single correct · +4 / −1	Skipped	4	17%	11%

Chemistry

6 of 24 questions lost · 29 marks · 1 easy miss · 3 within reach · 2 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q67	Easy miss	**Aldehydes & Ketones**	Single correct · +4 / −1	**Skipped**	4	78%	94%
Q55	Within reach	**Alcohols, Phenols & Ethers**	Single correct · +4 / −1	**Wrong**	5	54%	71%
Q69	Within reach	**Aldehydes & Ketones** – Order of hydration constants of carbonyls	Single correct · +4 / −1	**Wrong**	5	52%	49%
Q54	Within reach	**Alcohols, Phenols & Ethers**	Single correct · +4 / −1	**Wrong**	5	47%	49%
Q63	Tough one	**Aldehydes & Ketones** – Enol-ether hydrolysis; iodoform-positive products	Single correct · +4 / −1	**Wrong**	5	32%	45%
Q74	Tough one	**Alcohols, Phenols & Ethers** – Site of benzylation of 2-naphthoxide	Numerical · +4 / −1	**Wrong**	5	12%	15%

4 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Differential Equations	4	**3**/16		84%	Q10, Q13, Q19
Definite Integration	9	**12**/36		29%	Q8, Q11, Q16, Q18, Q20, Q22
Indefinite Integration	7	**12**/28		61%	Q6, Q9, Q12, Q17
Area Under Curves	5	**15**/20		86%	Q23
Physics					
Current Electricity	8	**23**/32		57%	Q36, Q37
Electrostatics	8	**23**/32		27%	Q26, Q30
Kinematics	9	**31**/36		75%	Q45
Chemistry					
Alcohols, Phenols & Ethers	10	**25**/40		48%	Q54, Q55, Q74
Aldehydes & Ketones	14	**42**/56		65%	Q63, Q67, Q69

Bar = your % of the chapter's marks | campus average

5 JEE Mains · Maths – questions to revisit

14 questions · easy misses first
cropped from the question paper

Q13 Differential Equations

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5
Top 100: **86%** got it right · Campus: **91%** (59 of 65)

Easy miss
You: Wrong

13. The solution of the differential equation $xy\frac{dy}{dx}=\frac{1+y^2}{1+x^2}\left(1+x+x^2\right)$ is $\sqrt{1+y^2}=c.x.e^{\tan^{-1}g(x)}$ then g(23) is

A) $\frac{1}{23}$ B) 24 C) 23 D) $\frac{1}{24}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q10 Differential Equations

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4
Top 100: **76%** got it right · Campus: **85%** (55 of 65)

Easy miss
You: Skipped

10. The solution of $\frac{xdy}{x^2+y^2}=\left(\frac{y}{x^2+y^2}-1\right)dx$ is_____ (c is arbitrary constant)

A) $y=x\cot(c-x)$ B) $y=x\tan(c-x)$ C) $y=x\cos(c-x)$ D) $y=x\sin(c-x)$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q17 Indefinite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4
Top 100: **74%** got it right · Campus: **66%** (43 of 65)

Easy miss
You: Skipped

17. $\int\left(1+x-x^{-1}\right)e^{x+x^{-1}}dx=$_____ (c is arbitrary constant)

A) $(1+x)e^{x+x^{-1}}+c$ B) $(x-1)e^{x+x^{-1}}+c$ C) $-xe^{x+x^{-1}}+c$ D) $xe^{x+x^{-1}}+c$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q19 Differential Equations

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **68%** got it right · Campus: **77%** (50 of 65)

Within reach

You: Skipped

19. The equation of the curve whose subnormal is equal to constant 'a' is

A) $y = ax + c$ B) $y^2 = 2ax + 2c$ C) $ay^2 = x^3 + ac$ D) $y^2 = ax + c$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q9 Indefinite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **66%** got it right · Campus: **63%** (41 of 65)

Within reach

You: Skipped

9. $\int \frac{dx}{\sin x + \sec x} =$ (c is arbitrary constant)

A) $\frac{1}{2\sqrt{3}} \ln \left| \frac{\sin x - \cos x + \sqrt{3}}{\cos x - \sin x + \sqrt{3}} \right| + \tan^{-1}(\sin x + \cos x) + c$

B) $\frac{1}{2\sqrt{3}} \ln \left| \frac{\sin x - \cos x - \sqrt{3}}{\sin x - \cos x + \sqrt{3}} \right| + c$

C) $\frac{1}{2\sqrt{3}} \ln \left| (\sin x - \cos x + \sqrt{3})(\sin x - \cos x - \sqrt{3}) \right| + c$

D) $\frac{1}{2\sqrt{3}} \ln \left| (\sin x - \cos x + \sqrt{3})(\sin x - \cos x - \sqrt{3}) \right| + \tan^{-1}(\sin x + \cos x) + c$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q23 Area Under Curves

JEE Mains · Maths · Numerical · marking +4 / −1 · marks lost 5

Top 100: **66%** got it right · Campus: **72%** (47 of 65)

Within reach

You: Wrong

23. The area bounded by the lines $y = ||x-1|-2|$ with x-axis is _____ (in sq.units)

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q8

Definite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **56%** got it right · Campus: **38%** (25 of 65)

Within reach

You: Skipped

8. If $\int_0^{\frac{\pi}{2}} \frac{\tan x}{\tan x+\sec x}dx = m(\pi+n)$, then $m\times n$ is equal to

A) $-\frac{1}{2}$ B) 1 C) -1 D) $\frac{1}{2}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q12

Indefinite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **51%** got it right · Campus: **43%** (28 of 65)

Within reach

You: Skipped

12. $\int \sqrt{\sec x-1}\,dx =$ _____ (c is arbitrary constant)

A) $2\ln\left|\cos\frac{x}{2}+\sqrt{\cos^2\frac{x}{2}+\frac{1}{2}}\right|+c$ B) $\ln\left|\cos\frac{x}{2}-\sqrt{\cos^2\frac{x}{2}-\frac{1}{2}}\right|+c$

C) $-2\ln\left|\cos\frac{x}{2}+\sqrt{\cos^2\frac{x}{2}-\frac{1}{2}}\right|+c$ D) $\ln\left|\cos\frac{x}{2}-\sqrt{\cos^2\frac{x}{2}+\frac{1}{2}}\right|+c$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q18

Definite Integration – ∫ of min{|x − 3k|} – periodic zig-zag

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **37%** got it right · Campus: **40%** (26 of 65)

Tough one

You: Skipped

18. Let $f(x)=\min\{\ldots,|x+6|,|x+3|,|x|,|x-3|,|x-6|,\ldots\}$ then $\int_{-4}^{3} f(x)\,dx =$ _____

A) 4 B) $\frac{9}{2}$ C) 5 D) $\frac{11}{2}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q6

Indefinite Integration – $\int \sqrt{(x^2+1)}[\ln(x^2+1) - 2\ln x] / x^4\, dx$

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **33%** got it right · Campus: **31%** (20 of 65)

Tough one

You: Skipped

6. $\int \frac{\sqrt{x^2+1}\left[\ln\left(x^2+1\right)-2\ln x\right]}{x^4}\,dx$ is equal to (c is arbitrary constant)

A) $\frac{\left(x^2+1\right)^{3/2}}{x^3}\left[\frac{2}{3}-\ln\left(\frac{x^2+1}{x^2}\right)\right]+c$

B) $\frac{\left(x^2+1\right)^{3/2}}{3x^3}\left[\ln\left(\frac{x^2+1}{x^2}\right)-\frac{2}{3}\right]+c$

C) $\frac{2\left(x^2+1\right)^{3/2}}{3x^3}\left[\frac{1}{3}-\frac{1}{2}\ln\left(\frac{x^2+1}{x^2}\right)\right]+c$

D) $\frac{\left(x^2+1\right)^{3/2}}{3x^3}\left[\ln\left(\frac{x^2+1}{x^2}\right)+\frac{2}{3}\right]+c$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q11

Definite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **29%** got it right · Campus: **26%** (17 of 65)

Tough one

You: Skipped

11. Let $f(x)=\sin x+\sum_{r=0}^{2}\cos\left(x+\frac{2r\pi}{3}\right)$, $g(x)=\cos x+\sum_{r=0}^{2}\sin\left(x+\frac{2r\pi}{3}\right)$ then $\int_0^{\frac{\pi}{2}} f\left(g(x)\right)f(x)\,dx=$

A) $1+\sin 1$ B) $1-\sin 1$ C) $1+\cos 1$ D) $1-\cos 1$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q20

Definite Integration – King's property with f(x) + f(y) = f(xy)

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **29%** got it right · Campus: **32%** (21 of 65)

Tough one

You: Skipped

20. Let $f:(0,\infty)\to R$ be a continuous function, satisfying $f(x)+f(y)=f(xy)$. Then the value of $\int_{\sqrt{f(e)}}^{\sqrt{f(e^2)}} \frac{x\tan x^2}{\tan x^2+\tan\left(f\left(e^3\right)-x^2\right)}\,dx$ is equal to

A) $\frac{1}{2}f\left(e^2\right)$ B) $\frac{1}{4}f\left(e^2\right)$ C) $\frac{1}{2}f(e)$ D) $\frac{f(e)}{4}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q22

Definite Integration – Definite integral of a product × sum of reciprocals

JEE Mains · Maths · Numerical · marking +4 / −1 · marks lost 4

Top 100: **20%** got it right · Campus: **20%** (13 of 65)

Tough one

You: Skipped

22. $\int_0^{102}(x-1)(x-2)....(x-100)\left(\frac{1}{x-1}+\frac{1}{x-2}+...+\frac{1}{x-100}\right)dx = p(p!)$ then the number of positive integral divisors of p is equal to_____ (where, $p \in N, p!$ means the product of first p natural numbers).

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q16

Definite Integration – $\int_0^{100}([\tan^{-1}x]+[\cot^{-1}x])\,dx$ with GIF

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **16%** got it right · Campus: **11%** (7 of 65)

Tough one

You: Skipped

16. If $\int_0^{100}\left(\left[\tan^{-1}x\right]+\left[\cot^{-1}x\right]\right)dx = ...$ [.] denotes the greatest integer function

A) $100-\cot 1$ B) $100+2\cot 1$ C) $100-2\cot 2$ D) $100+2\cot 2$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Mains · Physics – questions to revisit

5 questions · easy misses first
cropped from the question paper

Q45 Kinematics

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **89%** got it right · Campus: **88%** (57 of 65)

Easy miss
You: Wrong

45. The position vector of a particle is $r = 2t\hat{i} + t^2\hat{j}$ (SI units). Its speed at t = 2 s is

A) 4 *m/s* B) $2\sqrt{2}\ m/s$ C) 6 *m/s* D) $2\sqrt{5}\ m/s$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q36 Current Electricity

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **73%** got it right · Campus: **77%** (50 of 65)

Easy miss
You: Wrong

36. A copper wire carries a steady current I with drift velocity 'v'. If the wire is replaced by one of the same material but double the diameter carrying the same current, the drift velocity becomes

A) $\frac{v}{2}$ B) $4v$ C) v D) $\frac{v}{4}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q26 Electrostatics

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **70%** got it right · Campus: **69%** (45 of 65)

Easy miss
You: Wrong

26. A non-conducting sphere of radius R carries a uniform volume charge density ρ. A spherical cavity of radius $\frac{R}{2}$ is made in it such that the centre of the cavity is at a distance $\frac{R}{2}$ from the centre of the sphere. The magnitude of the electric field at any point inside the cavity is

A) $\rho\frac{R}{12\varepsilon_0}$ B) $\rho\frac{R}{3\varepsilon_0}$ C) $\rho\frac{R}{6\varepsilon_0}$ D) Zero

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q37 — Current Electricity – Infinite ladder – voltage across the 2nd shunt resistor

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4

Top 100: **52%** got it right · Campus: **46%** (30 of 65)

Within reach

You: Skipped

37. A 24 V battery of negligible internal resistance is connected to an infinite ladder network. In each section, both the top rail and the bottom rail contain a 1 Ω resistor, and a 4 Ω resistor is connected across the rails. The potential difference across the second 4 Ω resistor (counting from the battery) is

A) 12 V B) 8 V C) 6 V D) 3 V

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q30 — Electrostatics – Field energy inside vs outside a charged sphere

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4

Top 100: **17%** got it right · Campus: **11%** (7 of 65)

Tough one

You: Skipped

30. A solid sphere of radius R carries a total charge Q distributed uniformly throughout its volume. The ratio of the electrostatic field energy stored inside the sphere to that stored outside the sphere is

A) 3 : 5 B) 1 : 5 C) 1 : 3 D) 1 : 2

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q67 **Aldehydes & Ketones**


JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 4


Top 100: **78%** got it right · Campus: **94%** (61 of 65)


Easy miss

You: Skipped

67. Within each set in which case second compound is more reactive than first toward carbonyl addition reaction?

A) 

B) 

C) 

D) 

Why did I lose this question? (write the reason and the correct approach)


☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q55

Alcohols, Phenols & Ethers
JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5
Top 100: **54%** got it right · Campus: **71%** (46 of 65)

Within reach
You: Wrong

55. Given below are two statements for the



$$C_2H_5O^- + H_2O \rightleftharpoons C_2H_5OH + OH^-; K_2$$

Statement I : The value of K_2 less than K_1

Statement II : Phenoxide is a weaker base than ethoxide

In the light of the above statement, choose the most appropriate answer from the options given below.

A) Both Statement I and Statement II are correct

B) Both Statement I and Statement II are incorrect

C) Statement I is correct but Statement II is incorrect

D) Statement II is correct but Statement I is incorrect

Why did I lose this question? (write the reason and the correct approach)


☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q69

Aldehydes & Ketones – Order of hydration constants of carbonyls
JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5
Top 100: **52%** got it right · Campus: **49%** (32 of 65)

Within reach
You: Wrong

69. Arrange in their increasing order of equilibrium constants for hydration?



A) A < B < C < D < E

B) A < C < B < E < D

C) A < C < E < B < D

D) C < A < B < E < D


Why did I lose this question? (write the reason and the correct approach)


☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed


Q54 **Alcohols, Phenols & Ethers**
JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5
Top 100: **47%** got it right · Campus: **49%** (32 of 65)


Within reach
You: Wrong

54. Identify the correct acidic strength order of following pairs

A) 

B) 

C) 

D) 

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q63

Aldehydes & Ketones – Enol-ether hydrolysis; iodoform-positive products

JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 5

Top 100: **32%** got it right · Campus: **45%** (29 of 65)


Tough one


You: Wrong

63. $\underset{C_5H_{10}O}{(A)} \xrightarrow{H_3O^{\oplus}} B + C; (B) \text{ and } (C)$ both give +ve Iodoferm test.

Compound (A) is

A) $CH_3-CH=CH-O-CH_2-CH_3$

B) 

C) 

D) Both (B) and (C)

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q74

Alcohols, Phenols & Ethers – Site of benzylation of 2-naphthoxide


JEE Mains · Chemistry · Numerical · marking +4 / −1 · marks lost 5

Top 100: **12%** got it right · Campus: **15%** (10 of 65)

Tough one

You: Wrong

74. Which site is predominately benzylated in the given reaction?



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced – question by question

118 / 176 · 67.0%
09 Oct 2026

6 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
Physics	18	19	20	21	22	23	24	~~25~~	26	27	28	29	30	31	32	33	34
Chemistry	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51

7 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 61 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40–69% · **Tough one** below 40%

Maths

10 of 17 questions lost · **37** marks · 2 easy miss · 6 within reach · 2 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q5	Easy miss	**Indefinite Integration**	Multiple correct · +4 / −2	Skipped	4	93%	89%
Q4	Easy miss	**Differential Equations**	Single correct · +3 / −1	Wrong	4	92%	92%
Q6	Within reach	**Area Under Curves** – Line x = α halving the area between x^3 and x	Multiple correct · +4 / −2	Skipped	4	68%	69%
Q1	Within reach	**Indefinite Integration**	Single correct · +3 / −1	Skipped	3	66%	70%
Q8	Within reach	**Indefinite Integration**	Integer · +4 / 0	Skipped	4	64%	67%
Q13	Within reach	**Definite Integration**	Integer · +4 / 0	Skipped	4	64%	70%
Q14	Within reach	**Definite Integration**	Paragraph · +3 / 0	Skipped	3	60%	56%
Q17	Within reach	**Differential Equations**	Paragraph · +3 / 0	Wrong	3	42%	33%
Q12	Tough one	**Differential Equations** – Solve a first-order DE and match its constants	Integer · +4 / 0	Wrong	4	35%	26%
Q11	Tough one	**Differential Equations**	Integer · +4 / 0	Skipped	4	25%	21%

Physics

3 of 16 questions lost · **10** marks · 1 within reach · 2 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q33	Within reach	**Electrostatics** – Flux through a ring – solid angle	Paragraph · +3 / 0	Wrong	3	42%	38%
Q34	Tough one	**Electrostatics** – Flux through a cube from a wire along one edge	Paragraph · +3 / 0	Wrong	3	33%	23%
Q20	Tough one	**Kinematics** – Shell fired from a plateau – minimum landing distance	Single correct · +3 / −1	Wrong	4	29%	23%



Chemistry

3 of 17 questions lost · **11** marks · 1 easy miss · 1 within reach · 1 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q36	Easy miss	**Aldehydes & Ketones**	Single correct · +3 / −1	Wrong	4	70%	77%
Q49	Within reach	**Aldehydes & Ketones**	Paragraph · +3 / 0	Wrong	3	59%	70%
Q46	Tough one	**Alcohols, Phenols & Ethers** – Identifying polar aprotic solvents	Integer · +4 / 0	Skipped	4	24%	8%

8 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject. Chapter marks are approximate (≈): partly-correct answers cannot be traced to a question.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Differential Equations	5	≈ 3/18		30%	Q4, Q11, Q12, Q17
Indefinite Integration	4	≈ 4/15		71%	Q1, Q5, Q8
Definite Integration	5	≈ 10/17		54%	Q13, Q14
Area Under Curves	3	≈ 6/10		60%	Q6
Physics					
Electrostatics	6	≈ 15/21		45%	Q33, Q34
Kinematics	8	≈ 24/28		41%	Q20
Current Electricity	2	≈ 7/7		68%	– none
Chemistry					
Aldehydes & Ketones	6	≈ 13/20		83%	Q36, Q49
Alcohols, Phenols & Ethers	7	≈ 22/26		62%	Q46
Carboxylic Acids	4	≈ 14/14		48%	– none

Bar = your % of the chapter's marks | campus average

9 JEE Advanced · Maths – questions to revisit

10 questions · easy misses first
cropped from the question paper

Q5 Indefinite Integration

JEE Advanced · Maths · Multiple correct · marking +4 / −2 · marks lost 4
Top 100: **93%** got it right · Campus: **89%** (54 of 61)

Easy miss
You: Skipped

5. If $\int \frac{dt}{\left(1+\sqrt{t}\right)^8} = \frac{-1}{3\left(1+\sqrt{t}\right)^{p_1}} + \frac{2}{7\left(1+\sqrt{t}\right)^{p_2}} + C$ where C is constant of integration, then

A) $p_1=5$ B) $p_1=6$ C) $p_2=7$ D) $p_2=8$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q4 Differential Equations

JEE Advanced · Maths · Single correct · marking +3 / −1 · marks lost 4
Top 100: **92%** got it right · Campus: **92%** (56 of 61)

Easy miss
You: Wrong

4. Solution of the differential equation $\sin y\frac{dy}{dx}+\frac{1}{x}\cos y = x^4\cos^2 y$ is

(Where C is the constant of integration)

A) $x\sec y = x^6 + C$ B) $6x\sec y = x + C$ C) $6x\sec y = x^6 + C$ D) $x^6\sec y = x + C$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q6 Area Under Curves – Line x = α halving the area between x³ and x

JEE Advanced · Maths · Multiple correct · marking +4 / −2 · marks lost 4
Top 100: **68%** got it right · Campus: **69%** (42 of 61)

Within reach
You: Skipped

6. If the line x=α divides the area of the region $R = \{(x,y)\in R^2 : x^3 \le y \le x, 0 \le x \le 1\}$ into two equal parts, then

A) $\frac{1}{2} \le \alpha \le 1$ B) $\alpha^4 + 4\alpha^2 - 1 = 0$ C) $0 \le \alpha \le \frac{1}{2}$ D) $2\alpha^4 - 4\alpha^2 + 1 = 0$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q1

Indefinite Integration

JEE Advanced · Maths · Single correct · marking +3 / −1 · marks lost 3

Top 100: **66%** got it right · Campus: **70%** (43 of 61)

Within reach

You: Skipped

1. $\int \frac{2e^{2x}+e^{x}}{3e^{2x}-2e^{x}-1}dx$ is equal to

A) $\frac{1}{4}\ln\left|\frac{\left(e^{x}-1\right)^{1/3}}{\left(3e^{x}+1\right)^{3}}\right|+C$

B) $\frac{1}{4}\ln\left|\frac{\left(e^{x}-1\right)^{3}}{\left(3e^{x}+1\right)^{1/3}}\right|+C$

C) $\frac{1}{3}\ln\left|\frac{\left(e^{x}-1\right)^{4}}{\left(3e^{x}+1\right)^{1/4}}\right|+C$

D) $\frac{1}{3}\ln\left|\frac{\left(e^{x}-1\right)^{1/4}}{\left(3e^{x}+1\right)^{4}}\right|+C$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q8

Indefinite Integration

JEE Advanced · Maths · Integer · marking +4 / 0 · marks lost 4

Top 100: **64%** got it right · Campus: **67%** (41 of 61)

Within reach

You: Skipped

8. Let $I=\int\frac{\left(x^{6}-4\right)dx}{\left(x^{6}+2\right)^{1/4}x^{4}}=\frac{l\left(x^{6}+2\right)^{m}}{x^{n}}+C$ then $\frac{n}{lm}$ is equal to (where C is constant of integration).

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q13

Definite Integration

JEE Advanced · Maths · Integer · marking +4 / 0 · marks lost 4

Top 100: **64%** got it right · Campus: **70%** (43 of 61)

Within reach

You: Skipped

13. Let $t(x)=\sqrt{1+x\sqrt{1+(x+1)\sqrt{1+(x+2)(x+4)}}}$ then $\int_{0}^{100}t(x)dx$ is K. Find $\frac{K}{1700}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q14 Definite Integration

JEE Advanced · Maths · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **60%** got it right · Campus: **56%** (34 of 61)

Within reach

You: Skipped

PARAGRAPH-I

$$\int_0^a f(x)dx = \int_0^a f(a-x)dx\,; \quad \int_a^b f(x)dx = \int_a^b f(a+b-x)dx\,; \quad \int_0^{2a} f(x)dx = \int_0^a \left(f(x)+f(2a-x)\right)dx$$

14. If $\int_0^{\frac{\pi}{2}} \ln(\sin x)dx = \frac{-\pi}{q}\ln(p), (p,q \in N)$ then find the least possible value of p+q

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q17 Differential Equations

JEE Advanced · Maths · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **42%** got it right · Campus: **33%** (20 of 61)

Within reach

You: Wrong

PARAGRAPH-II

Let $f(x)$ be a differentiable function.

$$f(x) > 0 \forall x \in R \qquad f(x) = \left(2\int_0^x f(t)dt\right) + 1 \forall x \in R$$

17. Find the value of $\lim_{x\to 0} \frac{f(x)-2x-1}{1-\cos x}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q12 Differential Equations – Solve a first-order DE and match its constants

JEE Advanced · Maths · Integer · marking +4 / 0 · marks lost 4

Top 100: **35%** got it right · Campus: **26%** (16 of 61)

Tough one

You: Wrong

12. The differential equation $\left(xy^3(1+\cos x)-y\right)dx + xdy = 0$ represents the curve

$\frac{x^2}{ay^2} = \frac{x^3}{b} + x^2\sin x + cx\cos x - d\sin x + k$ .Then a+b+c+d is equal to (where 'k' is arbitrary constant).

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q11 **Differential Equations**

JEE Advanced · Maths · Integer · marking +4 / 0 · marks lost 4

Top 100: **25%** got it right · Campus: **21%** (13 of 61)

Tough one

You: Skipped

11. Let $e^{-x}\frac{d^2y}{dx^2}-\frac{dy}{dx}-y=0$ such that at the point $(0,1)$, $\frac{dy}{dx}=1$. Then the number of point of intersection of the given curve with $y=e^{e^x}$ is

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q33 **Electrostatics – Flux through a ring – solid angle**
JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **42%** got it right · Campus: **38%** (23 of 61)

Within reach
You: Wrong


PARAGRAPH-II

The electric flux linked with a surface is the number of electric field lines associated with that area. Gauss law helps determine the flux

$$\oint \vec{E}.\vec{ds} = \frac{Q_{inc}}{\varepsilon_0}$$

For an open surface the flux will change depending on the solid angle. Considering this answer the following questions.

33.



The flux associated with the ring due to the charge q is given by Xq/ε_0. The value of X is approx.

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q34 **Electrostatics – Flux through a cube from a wire along one edge**
JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **33%** got it right · Campus: **23%** (14 of 61)

Tough one
You: Wrong


PARAGRAPH-II

The electric flux linked with a surface is the number of electric field lines associated with that area. Gauss law helps determine the flux

$$\oint \vec{E}.\vec{ds} = \frac{Q_{inc}}{\varepsilon_0}$$

For an open surface the flux will change depending on the solid angle. Considering this answer the following questions.

34.



A cube of edge length a contains an infinite wire of linear charge density λ at one of edges the flux associated with cube is given by $P\lambda a/\varepsilon_0$.The value of P upto two decimal places is :


Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q20 **Kinematics – Shell fired from a plateau – minimum landing distance**
JEE Advanced · Physics · Single correct · marking +3 / −1 · marks lost 4
Top 100: **29%** got it right · Campus: **23%** (14 of 61)

Tough one
You: Wrong

20. A gun is mounted on a plateau 960m away from its edge as shown. Height of plateau is 960m. The gun can fire shells with a velocity 100 m/s at any angle of the following choices. What is the minimum distance OP(=x) from the edge of plateau where shell of



gun can reach? (given $g=10m/s^2$, $\sin 74^0 = \sin 106^0 \simeq 0.96$)

A) 480 m B) 720 m C) 360 m D) 960 m


Why did I lose this question? (write the reason and the correct approach)


☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed


Q36 **Aldehydes & Ketones**
JEE Advanced · Chemistry · Single correct · marking +3 / −1 · marks lost 4
Top 100: **70%** got it right · Campus: **77%** (47 of 61)


Easy miss
You: Wrong


36. The end products of the following sequence of reactions are



A)yellow precipitate of CHI_3, 

B)yellow precipitate of CHI_3, 

C)yellow precipitate of CHI_3, 

D)yellow precipitate of CHI_3, 

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q49 **Aldehydes & Ketones**
JEE Advanced · Chemistry · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **59%** got it right · Campus: **70%** (43 of 61)

Within reach
You: Wrong

PARAGRAPH-I

When a carbonyl compound reacts with hydroxylamine(NH_2OH), it undergoes nucleophilic addition followed by dehydration to form an oxime ($R_2C=N-OH$) and water. The rate of oxime formation is maximum at an optimum pH≈4.5. At lower pH, the hydroxylamine gets protonated into NH_3^+OH, losing its nucleophilicity, whereas at higher pH, protonation of the carbonyl oxygen is insufficient to activate the carbonyl group.


49. In a reaction vessel, 0.1 mol of benzaldehyde reacts completely with 0.1 mol of hydroxylamine to form benzaldoxime. Calculate the total volume of water (H_2O) vapor produced in liters at STP (1 mol of gas at STP = 22.4 L).

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q46 **Alcohols, Phenols & Ethers** – **Identifying polar aprotic solvents**
JEE Advanced · Chemistry · Integer · marking +4 / 0 · marks lost 4
Top 100: **24%** got it right · Campus: **8%** (5 of 61)

Tough one
You: Skipped

46. How many of the following are polar aprotic solvents?



[P]

Diethyl ether,


[Q]

Benzene,

[R]

Ethyl alcohol,

[S]



[T]

Petrol

[U]

Acetone

[Q]

DMSO (dimethyl sulphoxide

[W]

DMF (dimethyl formamide

[X]

formic acid

[Y]

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

10 How to read this report

Where the numbers come from. Marks, percentages, All-India / State / Campus ranks, subject marks, subject All-India ranks and the wrong / skipped questions are the official results in the institute's *All India Marks Analysis* (graded with the final answer key). Maximum marks leave out deleted questions — JEE Mains: Q56 — JEE Advanced: Q25. **Program rank** = rank in Pragya G10 (EXT) Achievers across all campuses, from the batch's question-wise sheet. **Top 100** = the batch's top 100 across all campuses, from the same question-wise sheet; * where it has no data for a question, the campus figure decides the priority. **Campus** = all students of this program on campus. **Priority:** Easy miss = 70%+ of the top 100 got it right; Within reach = 40–69%; Tough one = below 40%. **Marks lost** on a question = its marks plus the negative mark if answered wrong, its marks if skipped. **Weekly-test average** = your average over the weekly Mains tests (WTM) or Advanced tests (WTA) before this Part Test, from the campus averages sheet. The weekly Advanced tests are marked out of 240; their averages are scaled to 180 to match this paper (× 180/240, so 80 per subject becomes 60). Percentages are unchanged.