CLASS 10 · PART TEST-1 · STUDENT PERFORMANCE REPORT

Mano Nischal Reddy D

Pragya G10 (EXT) Achievers · Adm No **246418646**

Campus HYD/Nallakunta DD Colony
JEE Mains 08 Oct 2026
JEE Advanced 09 Oct 2026

JEE Mains

08 Oct 2026 · 74 questions

224 / 296 — **75.7%**

19	14	6/65	6
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **194.4 / 300** (64.8%) → this test **75.7%** ▲ **10.9**

72 marks lost: 40 on 8 wrong answers (incl. 8 negative) · 32 on 8 skipped

JEE Advanced

09 Oct 2026 · 50 questions

123 / 176 — **69.9%**

42	29	7/61	8
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **90.2 / 180** (50.1%), 5 of 6 tests → this test **69.9%** ▲ **19.8**

53 marks lost: 44 on 11 wrong answers (incl. 5 negative) · 6 on 2 skipped · 3 on partly-correct answers

1 Subject scorecard

Your marks in each subject, against the 65 students of this program on campus and against your own weekly-test average.

SUBJECT	MARKS	SCORE VS CAMPUS	SUBJECT AIR	CAMPUS AVG / BEST	WEEKLY AVG	CHANGE	RIGHT / WRONG / SKIP
JEE Mains 224 / 296 · 75.7%							
Maths	**57**/100	57%	**166**	58.2 / 100	**57.6**/100 57.6%	▼ **0.6**	15 / 3 / 7
Physics	**76**/100	76%	**55**	53.8 / 100	**61.4**/100 61.4%	▲ **14.6**	20 / 4 / 1
Chemistry	**91**/96	95% Best on campus	**1**	55.9 / 91	**75.3**/100 75.3%	▲ **19.5**	23 / 1 / 0
JEE Advanced 123 / 176 · 69.9%							
Maths	**34**/60	57%	**177**	30.5 / 57	**28.5**/60 47.5%	▲ **9.2**	11 / 5 / 1
Physics	**33**/56	59%	**115**	24.5 / 56	**27.4**/60 45.7%	▲ **13.3**	10 / 5 / 1
Chemistry	**56**/60	93%	**4**	38.2 / 60	**34.3**/60 57.2%	▲ **36.1**	16 / 1 / 0

Bar = your score % | campus average | best on campus Change = this test % − weekly-test average %

What this report says

1. **Chemistry** is your strongest JEE Mains subject: 91/96 (95%), All-India subject rank 1 — the best score on campus.
2. **Maths** needs the most work in JEE Mains: 57/100 (57%), subject rank 166; you lost 10 of its 25 questions.
3. **Chemistry** is your strongest JEE Advanced subject: 56/60 (93%), All-India subject rank 4.
4. **Maths** needs the most work in JEE Advanced: 34/60 (57%), subject rank 177; you lost 6 of its 17 questions.
5. Biggest improvement: **JEE Advanced Chemistry** — 93% here against 57% on average.
6. **2 questions** you lost were answered correctly by 70%+ of the batch's top 100 — about **10 marks** within easy reach (Mains Q24, Q25).
7. Negative marking cost you **8 in JEE Mains and 5 in JEE Advanced**.
8. You attempted almost everything (only 10 questions skipped across both papers), so the marks to gain are in **accuracy**: wrong answers — 8 in JEE Mains and 11 in JEE Advanced.

JEE Mains – question by question

224 / 296 · 75.7%
08 Oct 2026

2 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	2	3	4	**5**	**6**	7	8	9	10	**11**	**12**	13	14	15	**16**	17	**18**	19	**20**	21	22	**23**	**24**	**25**
Physics	26	27	28	29	**30**	31	32	33	**34**	35	36	**37**	38	39	40	**41**	42	43	44	45	**46**	47	48	49	50
Chemistry	51	52	53	54	55	~~56~~	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	**74**	75

3 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 65 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40–69% · **Tough one** below 40%

Maths

10 of 25 questions lost · **43** marks · ● 2 easy miss · ● 3 within reach · ● 5 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q24	Easy miss	**Indefinite Integration**	Numerical · +4 / −1	Wrong	5	86%	88%
Q25	Easy miss	**Differential Equations**	Numerical · +4 / −1	Wrong	5	78%	89%
Q23	Within reach	**Area Under Curves**	Numerical · +4 / −1	Wrong	5	66%	72%
Q12	Within reach	**Indefinite Integration**	Single correct · +4 / −1	Skipped	4	51%	43%
Q5	Within reach	**Definite Integration**	Single correct · +4 / −1	Skipped	4	40%	32%
Q18	Tough one	**Definite Integration** – ∫ of min{|x − 3k|} – periodic zig-zag	Single correct · +4 / −1	Skipped	4	37%	40%
Q6	Tough one	**Indefinite Integration** – ∫ $\sqrt{(x^2+1)}$ [ln(x^2+1) − 2 ln x] / x^4 dx	Single correct · +4 / −1	Skipped	4	33%	31%
Q11	Tough one	**Definite Integration**	Single correct · +4 / −1	Skipped	4	29%	26%
Q20	Tough one	**Definite Integration** – King's property with f(x) + f(y) = f(xy)	Single correct · +4 / −1	Skipped	4	29%	32%
Q16	Tough one	**Definite Integration** – $\int_0^{100}$ ([$\tan^{-1}$x] + [$\cot^{-1}$x]) dx with GIF	Single correct · +4 / −1	Skipped	4	16%	11%

Physics

5 of 25 questions lost · **24** marks · ● 3 within reach · ● 2 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q41	Within reach	**Kinematics**	Single correct · +4 / −1	Wrong	5	65%	71%
Q37	Within reach	**Current Electricity** – Infinite ladder – voltage across the 2nd shunt resistor	Single correct · +4 / −1	Skipped	4	52%	46%
Q34	Within reach	**Current Electricity**	Single correct · +4 / −1	Wrong	5	45%	37%
Q46	Tough one	**Electrostatics**	Numerical · +4 / −1	Wrong	5	25%	25%
Q30	Tough one	**Electrostatics** – Field energy inside vs outside a charged sphere	Single correct · +4 / −1	Wrong	5	17%	11%


Chemistry

1 of 24 questions lost · **5** marks · ● 1 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q74	Tough one	**Alcohols, Phenols & Ethers** – Site of benzylation of 2-naphthoxide	Numerical · +4 / −1	Wrong	5	12%	15%

4 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Definite Integration	9	**16**/36		29%	**Q5**, **Q11**, **Q16**, **Q18**, **Q20**
Indefinite Integration	7	**15**/28		61%	**Q6**, **Q12**, **Q24**
Differential Equations	4	**11**/16		84%	**Q25**
Area Under Curves	5	**15**/20		86%	**Q23**
Physics					
Electrostatics	8	**22**/32		27%	**Q30**, **Q46**
Current Electricity	8	**23**/32		57%	**Q34**, **Q37**
Kinematics	9	**31**/36		75%	**Q41**
Chemistry					
Alcohols, Phenols & Ethers	10	**35**/40		48%	**Q74**
Aldehydes & Ketones	14	**56**/56		65%	– none

Bar = your % of the chapter's marks | campus average

5 JEE Mains · Maths – questions to revisit

10 questions · easy misses first
cropped from the question paper

Q24 Indefinite Integration

JEE Mains · Maths · Numerical · marking +4 / −1 · marks lost 5
Top 100: **86%** got it right · Campus: **88%** (57 of 65)

Easy miss
You: Wrong

24. The value of $\int(\sin 101x).\sin^{99} x\ dx$ is $\frac{(\sin 100x).\sin^{100} x}{k+5}+C$ then $\frac{k}{19}$ is_____ (C is arbitrary constant)

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q25 Differential Equations

JEE Mains · Maths · Numerical · marking +4 / −1 · marks lost 5
Top 100: **78%** got it right · Campus: **89%** (58 of 65)

Easy miss
You: Wrong

25. If the solution $y(x)$ of the given differential equation $(e^y+1)\cos x dx+e^y \sin x dy=0$ passes through the point $\left(\frac{\pi}{2},0\right)$,then the value of $e^{y\left(\frac{\pi}{6}\right)}$ is equal to

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q23 Area Under Curves

JEE Mains · Maths · Numerical · marking +4 / −1 · marks lost 5
Top 100: **66%** got it right · Campus: **72%** (47 of 65)

Within reach
You: Wrong

23. The area bounded by the lines $y=||x-1|-2|$ with x-axis is _____ (in sq.units)

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q12 Indefinite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **51%** got it right · Campus: **43%** (28 of 65)

Within reach

You: Skipped

12. $\int \sqrt{\sec x - 1}\, dx =$ _____(c is arbitrary constant)

A) $2\ln\left|\cos\frac{x}{2}+\sqrt{\cos^2\frac{x}{2}+\frac{1}{2}}\right|+c$

B) $\ln\left|\cos\frac{x}{2}-\sqrt{\cos^2\frac{x}{2}-\frac{1}{2}}\right|+c$

C) $-2\ln\left|\cos\frac{x}{2}+\sqrt{\cos^2\frac{x}{2}-\frac{1}{2}}\right|+c$

D) $\ln\left|\cos\frac{x}{2}-\sqrt{\cos^2\frac{x}{2}+\frac{1}{2}}\right|+c$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q5 Definite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **40%** got it right · Campus: **32%** (21 of 65)

Within reach

You: Skipped

5. $\int_{-1}^{1} \sin x \ln\left(1^x+2^x+3^x+6^x\right) dx =$

A) $(\ln 6)\left(\cos\left(\frac{\pi+4}{4}\right)\right)$

B) $(\ln 6)\left(\sin\left(\frac{\pi+4}{4}\right)\right)$

C) $\sqrt{2}(\ln 6)\left(\sin\left(\frac{4-\pi}{4}\right)\right)$

D) $\sqrt{2}(\ln 6)\left(\sin\left(\frac{\pi+4}{4}\right)\right)$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q18 Definite Integration – ∫ of min{|x − 3k|} – periodic zig-zag

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **37%** got it right · Campus: **40%** (26 of 65)

Tough one

You: Skipped

18. Let $f(x)=\min\{\ldots,|x+6|,|x+3|,|x|,|x-3|,|x-6|,\ldots\}$ then $\int_{-4}^{3} f(x)\,dx =$ _____

A) 4

B) $\frac{9}{2}$

C) 5

D) $\frac{11}{2}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q6

Indefinite Integration – $\int \sqrt{(x^2+1)}[\ln(x^2+1) - 2\ln x] / x^4\, dx$

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **33%** got it right · Campus: **31%** (20 of 65)

Tough one

You: Skipped

6. $\int \frac{\sqrt{x^2+1}\left[\ln\left(x^2+1\right)-2\ln x\right]}{x^4}\,dx$ is equal to (c is arbitrary constant)

A) $\frac{\left(x^2+1\right)^{3/2}}{x^3}\left[\frac{2}{3}-\ln\left(\frac{x^2+1}{x^2}\right)\right]+c$

B) $\frac{\left(x^2+1\right)^{3/2}}{3x^3}\left[\ln\left(\frac{x^2+1}{x^2}\right)-\frac{2}{3}\right]+c$

C) $\frac{2\left(x^2+1\right)^{3/2}}{3x^3}\left[\frac{1}{3}-\frac{1}{2}\ln\left(\frac{x^2+1}{x^2}\right)\right]+c$

D) $\frac{\left(x^2+1\right)^{3/2}}{3x^3}\left[\ln\left(\frac{x^2+1}{x^2}\right)+\frac{2}{3}\right]+c$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q11

Definite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **29%** got it right · Campus: **26%** (17 of 65)

Tough one

You: Skipped

11. Let $f(x)=\sin x+\sum_{r=0}^{2}\cos\left(x+\frac{2r\pi}{3}\right)$, $g(x)=\cos x+\sum_{r=0}^{2}\sin\left(x+\frac{2r\pi}{3}\right)$ then $\int_0^{\frac{\pi}{2}} f\left(g(x)\right)f(x)\,dx =$

A) $1+\sin 1$ B) $1-\sin 1$ C) $1+\cos 1$ D) $1-\cos 1$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q20

Definite Integration – King's property with f(x) + f(y) = f(xy)

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **29%** got it right · Campus: **32%** (21 of 65)

Tough one

You: Skipped

20. Let $f:(0,\infty)\to R$ be a continuous function, satisfying $f(x)+f(y)=f(xy)$. Then the value of $\int_{\sqrt{f(e)}}^{\sqrt{f(e^2)}} \frac{x\tan x^2}{\tan x^2+\tan\left(f\left(e^3\right)-x^2\right)}\,dx$ is equal to

A) $\frac{1}{2}f\left(e^2\right)$ B) $\frac{1}{4}f\left(e^2\right)$ C) $\frac{1}{2}f(e)$ D) $\frac{f(e)}{4}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q16 — Definite Integration – $\int_0^{100}([\tan^{-1}x]+[\cot^{-1}x])\,dx$ with GIF

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **16%** got it right · Campus: **11%** (7 of 65)

Tough one

You: Skipped

16. If $\int_0^{100}\left(\left[\tan^{-1}x\right]+\left[\cot^{-1}x\right]\right)dx=\ldots$ [.] denotes the greatest integer function

A) $100-\cot 1$ B) $100+2\cot 1$ C) $100-2\cot 2$ D) $100+2\cot 2$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Mains · Physics – questions to revisit

5 questions · easy misses first
cropped from the question paper

Q41 Kinematics

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **65%** got it right · Campus: **71%** (46 of 65)

Within reach
You: Wrong

41. A swimmer can swim at 5 m/s in still water. A river 120 m wide flows at 3 m/s. If the swimmer heads so as to reach the point directly opposite on the other bank, the time taken is

A) 24 s B) 40 s C) 15 s D) 30 s

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q37 Current Electricity – Infinite ladder – voltage across the 2nd shunt resistor

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4
Top 100: **52%** got it right · Campus: **46%** (30 of 65)

Within reach
You: Skipped

37. A 24 V battery of negligible internal resistance is connected to an infinite ladder network. In each section, both the top rail and the bottom rail contain a 1 Ω resistor, and a 4 Ω resistor is connected across the rails. The potential difference across the second 4 Ω resistor (counting from the battery) is

A) 12 V B) 8 V C) 6 V D) 3 V

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q34 Current Electricity

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **45%** got it right · Campus: **37%** (24 of 65)

Within reach
You: Wrong

34. In a Wheatstone bridge ABCD, a 12 V battery of negligible internal resistance is connected between A and C. The arms are AB = 2 Ω, BC = 3 Ω, AD = 3 Ω and DC = 3 Ω. An ammeter of resistance 0.3 Ω is connected between B and D. The current through the ammeter is

A) 0.2 A B) 0.6 A C) 0.4 A D) 1.2 A

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q46 Electrostatics

JEE Mains · Physics · Numerical · marking +4 / −1 · marks lost 5

Top 100: **25%** got it right · Campus: **25%** (16 of 65)

Tough one

You: Wrong

46. In a region the electric field is $E = \left(5x\hat{i} + 3y\hat{j} + 4\hat{k}\right) N/C$, with x, y in meters. A cube of side 1m has one vertex at the origin and its edges along the positive x, y and z axis. The net electric flux through the cube (in Nm^2/C) is _____.

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q30 Electrostatics – Field energy inside vs outside a charged sphere

JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5

Top 100: **17%** got it right · Campus: **11%** (7 of 65)

Tough one

You: Wrong

30. A solid sphere of radius R carries a total charge Q distributed uniformly throughout its volume. The ratio of the electrostatic field energy stored inside the sphere to that stored outside the sphere is

A) 3 : 5 B) 1 : 5 C) 1 : 3 D) 1 : 2

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Mains · Chemistry — questions to revisit


1 question · easy misses first
cropped from the question paper

Q74

Alcohols, Phenols & Ethers — **Site of benzylation of 2-naphthoxide**
JEE Mains · Chemistry · Numerical · marking +4 / −1 · marks lost 5
Top 100: **12%** got it right · Campus: **15%** (10 of 65)

Tough one
You: Wrong

74. Which site is predominately benzylated in the given reaction?



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

6 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
Physics	18	19	20	21	22	23	24	~~25~~	26	27	28	29	30	31	32	33	34
Chemistry	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51

7 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 61 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40–69% · **Tough one** below 40%

Maths

6 of 17 questions lost · **26** marks · ● 6 within reach

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q6	Within reach	**Area Under Curves** – Line x = α halving the area between x^3 and x	Multiple correct · +4 / −2	Wrong	6	68%	69%
Q1	Within reach	**Indefinite Integration**	Single correct · +3 / −1	Wrong	4	66%	70%
Q13	Within reach	**Definite Integration**	Integer · +4 / 0	Wrong	4	64%	70%
Q7	Within reach	**Differential Equations** – Curve from tangent-intercept ratio BP : AP = 3 : 1	Multiple correct · +4 / −2	Wrong	6	43%	26%
Q2	Within reach	**Definite Integration** – $f(a) = \int \ln(1+x^a)/(1+x)\,dx$ – find f(4) + f(1/4)	Single correct · +3 / −1	Skipped	3	42%	34%
Q15	Within reach	**Definite Integration**	Paragraph · +3 / 0	Wrong	3	41%	39%

Physics

6 of 16 questions lost · **20** marks (+3 on partly-correct answers) · ● 1 within reach · ● 5 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q29	Within reach	**Electrostatics**	Integer · +4 / 0	Wrong	4	54%	34%
Q34	Tough one	**Electrostatics** – Flux through a cube from a wire along one edge	Paragraph · +3 / 0	Wrong	3	33%	23%
Q20	Tough one	**Kinematics** – Shell fired from a plateau – minimum landing distance	Single correct · +3 / −1	Skipped	3	29%	23%
Q27	Tough one	**Kinematics** – Projection angle that maximises area under the trajectory	Integer · +4 / 0	Wrong	4	27%	23%
Q31	Tough one	**Kinematics** – Hexagon projectile – maximum height	Paragraph · +3 / 0	Wrong	3	5%	5%
Q32	Tough one	**Kinematics** – Hexagon projectile – speed at the top	Paragraph · +3 / 0	Wrong	3	5%	5%




Chemistry

1 of 17 questions lost · **4** marks · ● 1 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q46	Tough one	**Alcohols, Phenols & Ethers** – Identifying polar aprotic solvents	Integer · +4 / 0	Wrong	4	24%	8%

8 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject. Chapter marks are approximate (≈): partly-correct answers cannot be traced to a question.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Area Under Curves	3	≈ **4**/10		60%	**Q6**
Definite Integration	5	≈ **7**/17		54%	**Q2**, **Q13**, **Q15**
Differential Equations	5	≈ **12**/18		30%	**Q7**
Indefinite Integration	4	≈ **11**/15		71%	**Q1**
Physics					
Kinematics	8	≈ **15**/28		41%	**Q20**, **Q27**, **Q31**, **Q32**
Electrostatics	6	≈ **14**/21		45%	**Q29**, **Q34**
Current Electricity	2	≈ **7**/7		68%	– none
Chemistry					
Alcohols, Phenols & Ethers	7	≈ **22**/26		62%	**Q46**
Aldehydes & Ketones	6	≈ **20**/20		83%	– none
Carboxylic Acids	4	≈ **14**/14		48%	– none

Bar = your % of the chapter's marks | campus average

9 JEE Advanced · Maths – questions to revisit

6 questions · easy misses first
cropped from the question paper

Q6 Area Under Curves – Line x = α halving the area between x³ and x

JEE Advanced · Maths · Multiple correct · marking +4 / −2 · marks lost 6
Top 100: **68%** got it right · Campus: **69%** (42 of 61)

Within reach
You: Wrong

6. If the line x=α divides the area of the region $R=\{(x,y)\in R^2 : x^3 \le y \le x, 0 \le x \le 1\}$ into two equal parts, then

A) $\frac{1}{2} \le \alpha \le 1$ B) $\alpha^4+4\alpha^2-1=0$ C) $0 \le \alpha \le \frac{1}{2}$ D) $2\alpha^4-4\alpha^2+1=0$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q1 Indefinite Integration

JEE Advanced · Maths · Single correct · marking +3 / −1 · marks lost 4
Top 100: **66%** got it right · Campus: **70%** (43 of 61)

Within reach
You: Wrong

1. $\int \frac{2e^{2x}+e^x}{3e^{2x}-2e^x-1}dx$ is equal to

A) $\frac{1}{4}\ln\left|\frac{(e^x-1)^{1/3}}{(3e^x+1)^3}\right|+C$ B) $\frac{1}{4}\ln\left|\frac{(e^x-1)^3}{(3e^x+1)^{1/3}}\right|+C$

C) $\frac{1}{3}\ln\left|\frac{(e^x-1)^4}{(3e^x+1)^{1/4}}\right|+C$ D) $\frac{1}{3}\ln\left|\frac{(e^x-1)^{1/4}}{(3e^x+1)^4}\right|+C$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q13 Definite Integration

JEE Advanced · Maths · Integer · marking +4 / 0 · marks lost 4
Top 100: **64%** got it right · Campus: **70%** (43 of 61)

Within reach
You: Wrong

13. Let $t(x)=\sqrt{1+x\sqrt{1+(x+1)\sqrt{1+(x+2)(x+4)}}}$ then $\int_0^{100} t(x)dx$ is K. Find $\frac{K}{1700}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q7

Differential Equations – Curve from tangent-intercept ratio BP : AP = 3 : 1

JEE Advanced · Maths · Multiple correct · marking +4 / −2 · marks lost 6

Top 100: **43%** got it right · Campus: **26%** (16 of 61)

Within reach

You: Wrong

7. A curve $y = f(x)$ passes through the point (1,1) and tangent at P(x,y) cuts the x-axis and y-axis at A and B, respectively such that BP:AP=3:1, then

A) Equation of the curve is $xy^3 = 1$

B) Equation of normal at the point (1,1) is $x - 3y + 2 = 0$

C) Curve passes through the point $\left(\frac{1}{8}, 2\right)$

D) Equation of the curve is $x\frac{dy}{dx} + 3y = 0$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q2

Definite Integration – f(a) = ∫ ln(1+x^a)/(1+x) dx – find f(4) + f(1/4)

JEE Advanced · Maths · Single correct · marking +3 / −1 · marks lost 3

Top 100: **42%** got it right · Campus: **34%** (21 of 61)

Within reach

You: Skipped

2. Let $f(a) = \int_0^1 \frac{\ln(1+x^a)}{1+x} dx$ then $f(4) + f\left(\frac{1}{4}\right)$ is equal to

A) $\ln^2 1$ B) $\ln^2 2$ C) $\ln^2 3$ D) $\ln^2 4$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q15 **Definite Integration**

JEE Advanced · Maths · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **41%** got it right · Campus: **39%** (24 of 61)

Within reach

You: Wrong

PARAGRAPH-I

$$\int_0^a f(x)dx = \int_0^a f(a-x)dx\,; \quad \int_a^b f(x)dx = \int_a^b f(a+b-x)dx\,; \quad \int_0^{2a} f(x)dx = \int_0^a \left(f(x)+f(2a-x)\right)dx$$

15. If $\left|\int_0^{\frac{\pi}{\pi}} \ln(1+\cos x)\,dx\right| = \frac{\pi}{q}\ln(p), (p,q \in N)$ then find the least possible value p+q

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q29

Electrostatics


JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **54%** got it right · Campus: **34%** (21 of 61)

Within reach

You: Wrong

29. An electron is projected from a distance d and with initial velocity u parallel to a uniformly charge flat conducting plate as shown. It strikes the plate after traveling a distance l along the direction of projection. The surface charge density of the conducting plate is $\sigma = \frac{nd\varepsilon_0 mu^2}{el^2}$, The value of n is:



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q34

Electrostatics – Flux through a cube from a wire along one edge

JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **33%** got it right · Campus: **23%** (14 of 61)

Tough one

You: Wrong


PARAGRAPH-II

The electric flux linked with a surface is the number of electric field lines associated with that area. Gauss law helps determine the flux

$$\oint \vec{E}.\vec{ds} = \frac{Q_{inc}}{\varepsilon_0}$$

For an open surface the flux will change depending on the solid angle. Considering this answer the following questions.

34.



A cube of edge length a contains an infinite wire of linear charge density λ at one of edges the flux associated with cube is given by $P\lambda a/\varepsilon_0$.The value of P upto two decimal places is :

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q20

Kinematics – Shell fired from a plateau – minimum landing distance


JEE Advanced · Physics · Single correct · marking +3 / −1 · marks lost 3

Top 100: **29%** got it right · Campus: **23%** (14 of 61)

Tough one

You: Skipped

20. A gun is mounted on a plateau 960m away from its edge as shown. Height of plateau is 960m. The gun can fire shells with a velocity 100 m/s at any angle of the following choices. What is the minimum distance OP(=x) from the edge of plateau where shell of



gun can reach? (given $g=10m/s^2$, $\sin 74^0 = \sin 106^0 \simeq 0.96$)

A) 480 m B) 720 m C) 360 m D) 960 m

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q27

Kinematics – Projection angle that maximises area under the trajectory

JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4

Top 100: **27%** got it right · Campus: **23%** (14 of 61)

Tough one

You: Wrong

27. A projectile is thrown with a speed 10 m/s, at an angle θ with the horizontal. For a certain value of angle of projection $\theta = \frac{(n+1)\pi}{12}$ the area bound by the trajectory with horizontal is maximum value of n is (g=10 m/s^2). Assume no airdrag.

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q31

Kinematics – Hexagon projectile – maximum height


JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **5%** got it right · Campus: **5%** (3 of 61)

Tough one

You: Wrong

PARAGRAPH-I



A regular hexagon of side a=$10\sqrt{3}$ m is shown in figure. From any point a particle is projected such that it touches B,C,D and E. The velocity of particle is v m/s. O is the centre of the hexagon. x-axis is parallel to horizontal and y-axis is perpendicular to AE and CD (g=10 m/s^2) $\left(\sqrt{3}=1.73\right)$.

31. The maximum height from the horizontal surface X' attained by the projectile is ___ m.


Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q32 **Kinematics – Hexagon projectile – speed at the top**
JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **5%** got it right · Campus: **5%** (3 of 61)

Tough one
You: Wrong

PARAGRAPH-I



A regular hexagon of side $a=10\sqrt{3}$ m is shown in figure. From any point a particle is projected such that it touches B,C,D and E. The velocity of particle is v m/s. O is the centre of the hexagon. x-axis is parallel to horizontal and y-axis is perpendicular to AE and CD ($g=10$ m/s^2) $\left(\sqrt{3}=1.73\right)$.

32. The velocity of projectile at maximum height is _____ m/s

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced · Chemistry — questions to revisit



1 question · easy misses first
cropped from the question paper

Q46

Alcohols, Phenols & Ethers – **Identifying polar aprotic solvents**
JEE Advanced · Chemistry · Integer · marking +4 / 0 · marks lost 4
Top 100: **24%** got it right · Campus: **8%** (5 of 61)

Tough one
You: Wrong

46. How many of the following are polar aprotic solvents?

 Diethyl ether, Benzene, Ethyl alcohol,  Petrol

[P] [Q] [R] [S] [T] [U]

Acetone | DMSO (dimethyl sulphoxide | DMF (dimethyl formamide | formic acid

[Q] [W] [X] [Y]

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

10 How to read this report

Where the numbers come from. Marks, percentages, All-India / State / Campus ranks, subject marks, subject All-India ranks and the wrong / skipped questions are the official results in the institute's *All India Marks Analysis* (graded with the final answer key). Maximum marks leave out deleted questions — JEE Mains: Q56 — JEE Advanced: Q25. **Program rank** = rank in Pragya G10 (EXT) Achievers across all campuses, from the batch's question-wise sheet. **Top 100** = the batch's top 100 across all campuses, from the same question-wise sheet; * where it has no data for a question, the campus figure decides the priority. **Campus** = all students of this program on campus. **Priority:** Easy miss = 70%+ of the top 100 got it right; Within reach = 40–69%; Tough one = below 40%. **Marks lost** on a question = its marks plus the negative mark if answered wrong, its marks if skipped. In JEE Advanced, 3 marks went on multiple-correct answers that were only partly correct. **Weekly-test average** = your average over the weekly Mains tests (WTM) or Advanced tests (WTA) before this Part Test, from the campus averages sheet. The weekly Advanced tests are marked out of 240; their averages are scaled to 180 to match this paper (× 180/240, so 80 per subject becomes 60). Percentages are unchanged.