CLASS 10 · PART TEST-1 · STUDENT PERFORMANCE REPORT

Poli Karthikeya Reddy

Pragya G10 (EXT) Achievers · Adm No **246554570**

Campus HYD/Nallakunta DD Colony
JEE Mains 08 Oct 2026
JEE Advanced 09 Oct 2026

JEE Mains

08 Oct 2026 · 74 questions

205 / 296 — **69.3%**

43	31	13/65	14
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **221.9 / 300** (74.0%) → this test **69.3%** ▼ **4.7**

91 marks lost: 35 on 7 wrong answers (incl. 7 negative) · 56 on 14 skipped

JEE Advanced

09 Oct 2026 · 50 questions

129 / 176 — **73.3%**

30	19	4/61	5
ALL-INDIA	STATE	CAMPUS	PROGRAM

Weekly-test average **113.4 / 180** (63.0%) → this test **73.3%** ▲ **10.3**

47 marks lost: 35 on 9 wrong answers (incl. 3 negative) · 12 on 4 skipped

1 Subject scorecard

Your marks in each subject, against the 65 students of this program on campus and against your own weekly-test average.

SUBJECT	MARKS	SCORE VS CAMPUS	SUBJECT AIR	CAMPUS AVG / BEST	WEEKLY AVG	CHANGE	RIGHT / WRONG / SKIP
JEE Mains 205 / 296 · 69.3%							
Maths	53/100	53%	210	58.2 / 100	72.1/100 72.1%	▼ 19.1	14 / 3 / 8
Physics	74/100	74%	74	53.8 / 100	76.5/100 76.5%	▼ 2.5	19 / 2 / 4
Chemistry	78/96	81%	8	55.9 / 91	73.3/100 73.3%	▲ 7.9	20 / 2 / 2
JEE Advanced 129 / 176 · 73.3%							
Maths	44/60	73%	55	30.5 / 57	34.6/60 57.7%	▲ 15.6	13 / 3 / 1
Physics	32/56	57%	129	24.5 / 56	39.5/60 65.8%	▼ 8.6	9 / 4 / 3
Chemistry	53/60	88%	12	38.2 / 60	39.3/60 65.5%	▲ 22.9	15 / 2 / 0

Bar = your score % | campus average | best on campus Change = this test % − weekly-test average %

What this report says

1. **Chemistry** is your strongest JEE Mains subject: 78/96 (81%), All-India subject rank 8.
2. **Maths** needs the most work in JEE Mains: 53/100 (53%), subject rank 210; you lost 11 of its 25 questions.
3. **Chemistry** is your strongest JEE Advanced subject: 53/60 (88%), All-India subject rank 12.
4. **Physics** needs the most work in JEE Advanced: 32/56 (57%), subject rank 129; you lost 7 of its 16 questions.
5. Biggest fall against your weekly-test average: **JEE Mains Maths** – 53% here against 72% on average.
6. Biggest improvement: **JEE Advanced Chemistry** – 88% here against 65% on average.
7. **2 questions** you lost were answered correctly by 70%+ of the batch's top 100 – about **9 marks** within easy reach (Mains Q25, Q67).
8. Negative marking cost you **7 in JEE Mains and 3 in JEE Advanced**.
9. You left **14 in JEE Mains and 4 in JEE Advanced** questions blank, and answered 7 in JEE Mains and 9 in JEE Advanced wrong – both are marks to recover.

JEE Mains – question by question

205 / 296 · 69.3%
08 Oct 2026

2 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	2	3	4	**5**	**6**	7	8	**9**	10	**11**	**12**	13	14	15	**16**	17	18	**19**	**20**	21	**22**	**23**	24	**25**
Physics	26	**27**	28	**29**	**30**	**31**	32	33	**34**	**35**	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50
Chemistry	51	52	53	54	55	~~56~~	57	58	59	60	61	62	63	64	65	66	**67**	68	**69**	70	71	72	**73**	**74**	75

3 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 65 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40–69% · **Tough one** below 40%

Maths

11 of 25 questions lost · **47** marks · ● 1 easy miss · ● 5 within reach · ● 5 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q25	Easy miss	**Differential Equations**	Numerical · +4 / −1	**Wrong**	5	**78%**	89%
Q19	Within reach	**Differential Equations**	Single correct · +4 / −1	**Skipped**	4	**68%**	77%
Q9	Within reach	**Indefinite Integration**	Single correct · +4 / −1	**Skipped**	4	**66%**	63%
Q23	Within reach	**Area Under Curves**	Numerical · +4 / −1	**Wrong**	5	**66%**	72%
Q12	Within reach	**Indefinite Integration**	Single correct · +4 / −1	**Skipped**	4	**51%**	43%
Q5	Within reach	**Definite Integration**	Single correct · +4 / −1	**Skipped**	4	**40%**	32%
Q6	Tough one	**Indefinite Integration** – $\int \sqrt{(x^2+1)}\,[\ln(x^2+1) - 2\ln x] / x^4\,dx$	Single correct · +4 / −1	**Wrong**	5	**33%**	31%
Q11	Tough one	**Definite Integration**	Single correct · +4 / −1	**Skipped**	4	**29%**	26%
Q20	Tough one	**Definite Integration** – King's property with f(x) + f(y) = f(xy)	Single correct · +4 / −1	**Skipped**	4	**29%**	32%
Q22	Tough one	**Definite Integration** – Definite integral of a product × sum of reciprocals	Numerical · +4 / −1	**Skipped**	4	**20%**	20%
Q16	Tough one	**Definite Integration** – $\int_0^{100} ([\tan^{-1}x] + [\cot^{-1}x])\,dx$ with GIF	Single correct · +4 / −1	**Skipped**	4	**16%**	11%

Physics

6 of 25 questions lost · **26** marks · ● 5 within reach · ● 1 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q29	Within reach	**Electrostatics**	Single correct · +4 / −1	**Skipped**	4	**61%**	48%
Q31	Within reach	**Electrostatics**	Single correct · +4 / −1	**Skipped**	4	**50%**	35%
Q35	Within reach	**Current Electricity**	Single correct · +4 / −1	**Wrong**	5	**50%**	48%
Q34	Within reach	**Current Electricity**	Single correct · +4 / −1	**Skipped**	4	**45%**	37%
Q27	Within reach	**Electrostatics** – Potential at the centre of a thick charged shell	Single correct · +4 / −1	**Skipped**	4	**42%**	32%
Q30	Tough one	**Electrostatics** – Field energy inside vs outside a charged sphere	Single correct · +4 / −1	**Wrong**	5	**17%**	11%

Chemistry

4 of 24 questions lost · **18** marks · ● 1 easy miss · ● 2 within reach · ● 1 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q67	Easy miss	**Aldehydes & Ketones**	Single correct · +4 / −1	**Skipped**	**4**	**78%**	94%
Q69	Within reach	**Aldehydes & Ketones** – Order of hydration constants of carbonyls	Single correct · +4 / −1	**Skipped**	**4**	**52%**	49%
Q73	Within reach	**Alcohols, Phenols & Ethers**	Numerical · +4 / −1	**Wrong**	**5**	**43%**	49%
Q74	Tough one	**Alcohols, Phenols & Ethers** – Site of benzylation of 2-naphthoxide	Numerical · +4 / −1	**Wrong**	**5**	**12%**	15%

4 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Differential Equations	4	**7**/16		84%	**Q19**, **Q25**
Definite Integration	9	**16**/36		29%	**Q5**, **Q11**, **Q16**, **Q20**, **Q22**
Indefinite Integration	7	**15**/28		61%	**Q6**, **Q9**, **Q12**
Area Under Curves	5	**15**/20		86%	**Q23**
Physics					
Electrostatics	8	**15**/32		27%	**Q27**, **Q29**, **Q30**, **Q31**
Current Electricity	8	**23**/32		57%	**Q34**, **Q35**
Kinematics	9	**36**/36		75%	– none
Chemistry					
Alcohols, Phenols & Ethers	10	**30**/40		48%	**Q73**, **Q74**
Aldehydes & Ketones	14	**48**/56		65%	**Q67**, **Q69**

Bar = your % of the chapter's marks | campus average

5 JEE Mains · Maths – questions to revisit

11 questions · easy misses first
cropped from the question paper

Q25

Differential Equations
JEE Mains · Maths · Numerical · marking +4 / −1 · marks lost 5
Top 100: **78%** got it right · Campus: **89%** (58 of 65)

Easy miss
You: Wrong

25. If the solution $y(x)$ of the given differential equation $(e^y+1)\cos x dx + e^y \sin x dy = 0$ passes through the point $\left(\frac{\pi}{2}, 0\right)$, then the value of $e^{y\left(\frac{\pi}{6}\right)}$ is equal to

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q19

Differential Equations
JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4
Top 100: **68%** got it right · Campus: **77%** (50 of 65)

Within reach
You: Skipped

19. The equation of the curve whose subnormal is equal to constant 'a' is

A) $y = ax + c$ B) $y^2 = 2ax + 2c$ C) $ay^2 = x^3 + ac$ D) $y^2 = ax + c$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q9

Indefinite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **66%** got it right · Campus: **63%** (41 of 65)

Within reach

You: Skipped

9. $\int \frac{dx}{\sin x + \sec x} =$ (c is arbitrary constant)

A) $\frac{1}{2\sqrt{3}} \ln \left| \frac{\sin x - \cos x + \sqrt{3}}{\cos x - \sin x + \sqrt{3}} \right| + \tan^{-1}(\sin x + \cos x) + c$

B) $\frac{1}{2\sqrt{3}} \ln \left| \frac{\sin x - \cos x - \sqrt{3}}{\sin x - \cos x + \sqrt{3}} \right| + c$

C) $\frac{1}{2\sqrt{3}} \ln \left| (\sin x - \cos x + \sqrt{3})(\sin x - \cos x - \sqrt{3}) \right| + c$

D) $\frac{1}{2\sqrt{3}} \ln \left| (\sin x - \cos x + \sqrt{3})(\sin x - \cos x - \sqrt{3}) \right| + \tan^{-1}(\sin x + \cos x) + c$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q23

Area Under Curves

JEE Mains · Maths · Numerical · marking +4 / −1 · marks lost 5

Top 100: **66%** got it right · Campus: **72%** (47 of 65)

Within reach

You: Wrong

23. The area bounded by the lines $y = ||x-1|-2|$ with x-axis is _____ (in sq.units)

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q12

Indefinite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **51%** got it right · Campus: **43%** (28 of 65)

Within reach

You: Skipped

12. $\int \sqrt{\sec x - 1}\, dx = _____$ (c is arbitrary constant)

A) $2\ln\left|\cos\frac{x}{2} + \sqrt{\cos^2\frac{x}{2} + \frac{1}{2}}\right| + c$

B) $\ln\left|\cos\frac{x}{2} - \sqrt{\cos^2\frac{x}{2} - \frac{1}{2}}\right| + c$

C) $-2\ln\left|\cos\frac{x}{2} + \sqrt{\cos^2\frac{x}{2} - \frac{1}{2}}\right| + c$

D) $\ln\left|\cos\frac{x}{2} - \sqrt{\cos^2\frac{x}{2} + \frac{1}{2}}\right| + c$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q5

Definite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **40%** got it right · Campus: **32%** (21 of 65)

Within reach

You: Skipped

5. $\int_{-1}^{1} \sin x \ln\left(1^x + 2^x + 3^x + 6^x\right) dx =$

A) $(\ln 6)\left(\cos\left(\frac{\pi+4}{4}\right)\right)$

B) $(\ln 6)\left(\sin\left(\frac{\pi+4}{4}\right)\right)$

C) $\sqrt{2}(\ln 6)\left(\sin\left(\frac{4-\pi}{4}\right)\right)$

D) $\sqrt{2}(\ln 6)\left(\sin\left(\frac{\pi+4}{4}\right)\right)$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q6

Indefinite Integration – $\int \sqrt{(x^2+1)}[\ln(x^2+1) - 2\ln x] / x^4\, dx$

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 5

Top 100: **33%** got it right · Campus: **31%** (20 of 65)

Tough one

You: Wrong

6. $\int \frac{\sqrt{x^2+1}\left[\ln\left(x^2+1\right)-2\ln x\right]}{x^4}\,dx$ is equal to (c is arbitrary constant)

A) $\frac{\left(x^2+1\right)^{3/2}}{x^3}\left[\frac{2}{3}-\ln\left(\frac{x^2+1}{x^2}\right)\right]+c$

B) $\frac{\left(x^2+1\right)^{3/2}}{3x^3}\left[\ln\left(\frac{x^2+1}{x^2}\right)-\frac{2}{3}\right]+c$

C) $\frac{2\left(x^2+1\right)^{3/2}}{3x^3}\left[\frac{1}{3}-\frac{1}{2}\ln\left(\frac{x^2+1}{x^2}\right)\right]+c$

D) $\frac{\left(x^2+1\right)^{3/2}}{3x^3}\left[\ln\left(\frac{x^2+1}{x^2}\right)+\frac{2}{3}\right]+c$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q11

Definite Integration

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **29%** got it right · Campus: **26%** (17 of 65)

Tough one

You: Skipped

11. Let $f(x)=\sin x+\sum_{r=0}^{2}\cos\left(x+\frac{2r\pi}{3}\right)$, $g(x)=\cos x+\sum_{r=0}^{2}\sin\left(x+\frac{2r\pi}{3}\right)$ then $\int_0^{\frac{\pi}{2}} f(g(x))f(x)\,dx =$

A) $1+\sin 1$ B) $1-\sin 1$ C) $1+\cos 1$ D) $1-\cos 1$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q20

Definite Integration – King's property with f(x) + f(y) = f(xy)

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **29%** got it right · Campus: **32%** (21 of 65)

Tough one

You: Skipped

20. Let $f:(0,\infty)\to R$ be a continuous function, satisfying $f(x)+f(y)=f(xy)$. Then the value of

$$\int_{\sqrt{f(e)}}^{\sqrt{f(e^2)}} \frac{x\tan x^2}{\tan x^2+\tan\left(f\left(e^3\right)-x^2\right)}\,dx \text{ is equal to}$$

A) $\frac{1}{2}f\left(e^2\right)$ B) $\frac{1}{4}f\left(e^2\right)$ C) $\frac{1}{2}f(e)$ D) $\frac{f(e)}{4}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q22

Definite Integration – Definite integral of a product × sum of reciprocals

JEE Mains · Maths · Numerical · marking +4 / −1 · marks lost 4

Top 100: **20%** got it right · Campus: **20%** (13 of 65)

Tough one

You: Skipped

22. $\int_0^{102}(x-1)(x-2)....(x-100)\left(\frac{1}{x-1}+\frac{1}{x-2}+...+\frac{1}{x-100}\right)dx = p(p!)$ then the number of positive integral divisors of p is equal to_____ (where, $p \in N, p!$ means the product of first p natural numbers).

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q16

Definite Integration – $\int_0^{100}([\tan^{-1}x]+[\cot^{-1}x])\,dx$ with GIF

JEE Mains · Maths · Single correct · marking +4 / −1 · marks lost 4

Top 100: **16%** got it right · Campus: **11%** (7 of 65)

Tough one

You: Skipped

16. If $\int_0^{100}\left(\left[\tan^{-1}x\right]+\left[\cot^{-1}x\right]\right)dx = ...$ [.] denotes the greatest integer function

A) $100-\cot 1$ B) $100+2\cot 1$ C) $100-2\cot 2$ D) $100+2\cot 2$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Mains · Physics – questions to revisit

6 questions · easy misses first
cropped from the question paper

Q29

Electrostatics
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4
Top 100: **61%** got it right · Campus: **48%** (31 of 65)

Within reach
You: Skipped

29. A point charge q is placed on the axis of a circular disc of radius R at a distance $\frac{R}{\sqrt{3}}$ from its centre. The electric flux through the disc is

A) $\frac{q\left(2-\sqrt{3}\right)}{4\varepsilon_0}$ B) $\frac{q}{6\varepsilon_0}$ C) $\frac{q}{2\varepsilon_0}$ D) $\frac{q}{4\varepsilon_0}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q31

Electrostatics
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4
Top 100: **50%** got it right · Campus: **35%** (23 of 65)

Within reach
You: Skipped

31. Three thin concentric conducting shells of radii R, 2R and 3R carry charges +Q, +2Q and +3Q respectively. The innermost and the outermost shells are now joined by a thin wire. The charge that flows through the wire is

A) $\frac{Q}{2}$ B) Q C) 2Q D) $\frac{3Q}{2}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q35 **Current Electricity**
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **50%** got it right · Campus: **48%** (31 of 65)

Within reach
You: Wrong

35. A 12 V battery of negligible internal resistance is connected to the input terminals of an infinite ladder network. Each section of the ladder has a 1 Ω resistor in the top (series) arm and a 2 Ω resistor connected across the two rails. The current through the first 2 Ω shunt resistor is

A) 6 A B) 3 A C) 2 A D) 4 A

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q34 **Current Electricity**
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4
Top 100: **45%** got it right · Campus: **37%** (24 of 65)

Within reach
You: Skipped

34. In a Wheatstone bridge ABCD, a 12 V battery of negligible internal resistance is connected between A and C. The arms are AB = 2 Ω, BC = 3 Ω, AD = 3 Ω and DC = 3 Ω. An ammeter of resistance 0.3 Ω is connected between B and D. The current through the ammeter is

A) 0.2 A B) 0.6 A C) 0.4 A D) 1.2 A

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q27 **Electrostatics – Potential at the centre of a thick charged shell**
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 4
Top 100: **42%** got it right · Campus: **32%** (21 of 65)

Within reach
You: Skipped

27. A thick spherical shell of inner radius ‘a’ and outer radius ‘b’ carries a uniform volume charge density ρ. The electric potential at the centre of the shell (taking V = 0 at infinity) is

A) $\rho\frac{\left(b^2-a^2\right)}{12\varepsilon_0}$ B) $\rho\frac{\left(b^3-a^3\right)}{3\varepsilon_0 b}$ C) $\rho\frac{\left(b^2-a^2\right)}{2\varepsilon_0}$ D) $\rho\frac{\left(b^3-a^3\right)}{3\varepsilon_0 a}$

Why did I lose this question? (write the reason and the correct approach)
☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q30 **Electrostatics – Field energy inside vs outside a charged sphere**
JEE Mains · Physics · Single correct · marking +4 / −1 · marks lost 5
Top 100: **17%** got it right · Campus: **11%** (7 of 65)

Tough one
You: Wrong

30. A solid sphere of radius R carries a total charge Q distributed uniformly throughout its volume. The ratio of the electrostatic field energy stored inside the sphere to that stored outside the sphere is

A) 3 : 5 B) 1 : 5 C) 1 : 3 D) 1 : 2

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q67

Aldehydes & Ketones


JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 4


Top 100: **78%** got it right · Campus: **94%** (61 of 65)


Easy miss


You: Skipped

67. Within each set in which case second compound is more reactive than first toward carbonyl addition reaction?

A) 

B) 

C) 

D) 

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q69

Aldehydes & Ketones – Order of hydration constants of carbonyls


JEE Mains · Chemistry · Single correct · marking +4 / −1 · marks lost 4

Top 100: **52%** got it right · Campus: **49%** (32 of 65)

Within reach

You: Skipped

69. Arrange in their increasing order of equilibrium constants for hydration?



A) A < B < C < D < E

B) A < C < B < E < D

C) A < C < E < B < D

D) C < A < B < E < D

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q73

Alcohols, Phenols & Ethers

JEE Mains · Chemistry · Numerical · marking +4 / −1 · marks lost 5

Top 100: **43%** got it right · Campus: **49%** (32 of 65)

Within reach

You: Wrong

73. How many chiral alcohols, including stereoisomers can give 2-chloro-2-methylbutane as the major product on reaction with $NaCl / H_2SO_4$ by hydride ion shift during its path?

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q74

Alcohols, Phenols & Ethers – Site of benzylation of 2-naphthoxide


JEE Mains · Chemistry · Numerical · marking +4 / −1 · marks lost 5

Top 100: **12%** got it right · Campus: **15%** (10 of 65)

Tough one

You: Wrong

74. Which site is predominately benzylated in the given reaction?



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced – question by question

129 / 176 · 73.3%
09 Oct 2026

6 Every question at a glance

Right · Wrong · Skipped · Deleted (no marks) · Easy miss · Within reach · Tough one

Maths	1	**2**	3	4	5	6	**7**	8	9	10	**11**	12	13	14	15	16	**17**
Physics	18	19	**20**	21	22	23	24	~~25~~	26	27	**28**	**29**	30	**31**	**32**	**33**	**34**
Chemistry	35	36	37	38	39	40	41	42	43	44	45	46	**47**	48	49	50	**51**

7 Questions you lost – what to fix first

Every question you got wrong or skipped, one table per subject, easiest marks first. **Top 100** = the batch's top 100 across all campuses; **Campus** = all 61 students of this program on campus.

Easy miss 70%+ of the top 100 got it right – easiest marks to win back · **Within reach** 40–69% · **Tough one** below 40%

Maths

4 of 17 questions lost · **16** marks · 3 within reach · 1 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q7	Within reach	**Differential Equations** – Curve from tangent-intercept ratio BP : AP = 3 : 1	Multiple correct · +4 / −2	Wrong	6	43%	26%
Q2	Within reach	**Definite Integration** – $f(a) = \int \ln(1+x^a)/(1+x)\,dx$ – find f(4) + f(1/4)	Single correct · +3 / −1	Skipped	3	42%	34%
Q17	Within reach	**Differential Equations**	Paragraph · +3 / 0	Wrong	3	42%	33%
Q11	Tough one	**Differential Equations**	Integer · +4 / 0	Wrong	4	25%	21%

Physics

7 of 16 questions lost · **24** marks · 3 within reach · 4 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q28	Within reach	**Electrostatics** – Flux through a cube with charges at its vertices	Integer · +4 / 0	Wrong	4	54%	44%
Q29	Within reach	**Electrostatics**	Integer · +4 / 0	Wrong	4	54%	34%
Q33	Within reach	**Electrostatics** – Flux through a ring – solid angle	Paragraph · +3 / 0	Wrong	3	42%	38%
Q34	Tough one	**Electrostatics** – Flux through a cube from a wire along one edge	Paragraph · +3 / 0	Skipped	3	33%	23%
Q20	Tough one	**Kinematics** – Shell fired from a plateau – minimum landing distance	Single correct · +3 / −1	Wrong	4	29%	23%
Q31	Tough one	**Kinematics** – Hexagon projectile – maximum height	Paragraph · +3 / 0	Skipped	3	5%	5%
Q32	Tough one	**Kinematics** – Hexagon projectile – speed at the top	Paragraph · +3 / 0	Skipped	3	5%	5%


Chemistry

2 of 17 questions lost · **7** marks · 2 tough one

Q	PRIORITY	CHAPTER – WHAT IT TESTS	TYPE	YOU	MARKS LOST	TOP 100 RIGHT	CAMPUS RIGHT
Q47	Tough one	**Carboxylic Acids**	Integer · +4 / 0	Wrong	4	32%	33%
Q51	Tough one	**Carboxylic Acids** – Moles of $LiAlH_4$ to reduce RCOOH to RCH_2OH	Paragraph · +3 / 0	Wrong	3	17%	23%

8 Chapter scorecard

Your marks on each chapter's questions against the campus average for the same questions. Weakest chapters first within each subject. Chapter marks are approximate (≈): partly-correct answers cannot be traced to a question.

CHAPTER	QS	YOUR MARKS	YOU VS CAMPUS AVERAGE	CAMPUS AVG	QUESTIONS YOU LOST
Maths					
Differential Equations	5	≈ **5**/18		30%	**Q7**, **Q11**, **Q17**
Definite Integration	5	≈ **14**/17		54%	**Q2**
Area Under Curves	3	≈ **10**/10		60%	– none
Indefinite Integration	4	≈ **15**/15		71%	– none
Physics					
Electrostatics	6	≈ **7**/21		45%	**Q28**, **Q29**, **Q33**, **Q34**
Kinematics	8	≈ **18**/28		41%	**Q20**, **Q31**, **Q32**
Current Electricity	2	≈ **7**/7		68%	– none
Chemistry					
Carboxylic Acids	4	≈ **7**/14		48%	**Q47**, **Q51**
Alcohols, Phenols & Ethers	7	≈ **26**/26		62%	– none
Aldehydes & Ketones	6	≈ **20**/20		83%	– none

Bar = your % of the chapter's marks | campus average

9 JEE Advanced · Maths — questions to revisit

4 questions · easy misses first
cropped from the question paper

Q7

Differential Equations – **Curve from tangent-intercept ratio BP : AP = 3 : 1**
JEE Advanced · Maths · Multiple correct · marking +4 / −2 · marks lost 6
Top 100: **43%** got it right · Campus: **26%** (16 of 61)

Within reach
You: Wrong

7. A curve $y = f(x)$ passes through the point (1,1) and tangent at P(x,y) cuts the x-axis and y-axis at A and B, respectively such that BP:AP=3:1, then

A) Equation of the curve is $xy^3 = 1$

B) Equation of normal at the point (1,1) is $x - 3y + 2 = 0$

C) Curve passes through the point $\left(\frac{1}{8}, 2\right)$

D) Equation of the curve is $x\frac{dy}{dx} + 3y = 0$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q2

Definite Integration – **f(a) = ∫ ln(1+x^a)/(1+x) dx – find f(4) + f(1/4)**
JEE Advanced · Maths · Single correct · marking +3 / −1 · marks lost 3
Top 100: **42%** got it right · Campus: **34%** (21 of 61)

Within reach
You: Skipped

2. Let $f(a) = \int_0^1 \frac{\ln(1+x^a)}{1+x} dx$ then $f(4) + f\left(\frac{1}{4}\right)$ is equal to

A) $\ln^2 1$ B) $\ln^2 2$ C) $\ln^2 3$ D) $\ln^2 4$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q17 **Differential Equations**
JEE Advanced · Maths · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **42%** got it right · Campus: **33%** (20 of 61)

Within reach
You: Wrong

PARAGRAPH-II

Let $f(x)$ be a differentiable function.

$$f(x) > 0 \forall x \in R \quad f(x) = \left(2\int_0^x f(t)dt \right) + 1 \forall x \in R$$

17. Find the value of $\lim_{x \to 0} \frac{f(x) - 2x - 1}{1 - \cos x}$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q11 **Differential Equations**
JEE Advanced · Maths · Integer · marking +4 / 0 · marks lost 4
Top 100: **25%** got it right · Campus: **21%** (13 of 61)

Tough one
You: Wrong

11. Let $e^{-x}\frac{d^2y}{dx^2} - \frac{dy}{dx} - y = 0$ such that at the point $(0,1)$, $\frac{dy}{dx} = 1$. Then the number of point of intersection of the given curve with $y = e^{e^x}$ is

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced · Physics – questions to revisit

7 questions · easy misses first
cropped from the question paper

Q28

Electrostatics – Flux through a cube with charges at its vertices
JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4
Top 100: **54%** got it right · Campus: **44%** (27 of 61)

Within reach
You: Wrong

28. A cube has charges each of value +Q placed at the vertices. The flux associated with the cube is $(X-1)\frac{Q}{\varepsilon_0}$, value of x is:

Why did I lose this question? (write the reason and the correct approach)


☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q29

Electrostatics
JEE Advanced · Physics · Integer · marking +4 / 0 · marks lost 4
Top 100: **54%** got it right · Campus: **34%** (21 of 61)

Within reach
You: Wrong

29. An electron is projected from a distance d and with initial velocity u parallel to a uniformly charge flat conducting plate as shown. It strikes the plate after traveling a distance l along the direction of projection. The surface charge density of the conducting plate is $\sigma = \frac{nd\varepsilon_0 mu^2}{el^2}$, The value of n is:



Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q33 **Electrostatics – Flux through a ring – solid angle**
JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **42%** got it right · Campus: **38%** (23 of 61)

Within reach
You: Wrong


PARAGRAPH-II

The electric flux linked with a surface is the number of electric field lines associated with that area. Gauss law helps determine the flux

$$\oint \vec{E}.\vec{ds} = \frac{Q_{inc}}{\varepsilon_0}$$

For an open surface the flux will change depending on the solid angle. Considering this answer the following questions.

33.



The flux associated with the ring due to the charge q is given by Xq/ε_0. The value of X is approx.

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q34

Electrostatics – Flux through a cube from a wire along one edge

JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3

Top 100: **33%** got it right · Campus: **23%** (14 of 61)

Tough one

You: Skipped


PARAGRAPH-II

The electric flux linked with a surface is the number of electric field lines associated with that area. Gauss law helps determine the flux

$$\oint \vec{E}.\vec{ds} = \frac{Q_{inc}}{\varepsilon_0}$$

For an open surface the flux will change depending on the solid angle. Considering this answer the following questions.

34.



A cube of edge length a contains an infinite wire of linear charge density λ at one of edges the flux associated with cube is given by $P\lambda a/\varepsilon_0$.The value of P upto two decimal places is :

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q20

Kinematics – Shell fired from a plateau – minimum landing distance


JEE Advanced · Physics · Single correct · marking +3 / −1 · marks lost 4

Top 100: **29%** got it right · Campus: **23%** (14 of 61)

Tough one

You: Wrong

20. A gun is mounted on a plateau 960m away from its edge as shown. Height of plateau is 960m. The gun can fire shells with a velocity 100 m/s at any angle of the following choices. What is the minimum distance OP(=x) from the edge of plateau where shell of



gun can reach? (given $g=10m/s^2$, $\sin 74^0 = \sin 106^0 \simeq 0.96$)

A) 480 m B) 720 m C) 360 m D) 960 m


Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q31 **Kinematics – Hexagon projectile – maximum height**
JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **5%** got it right · Campus: **5%** (3 of 61)

Tough one
You: Skipped

PARAGRAPH-I



A regular hexagon of side a=$10\sqrt{3}$ m is shown in figure. From any point a particle is projected such that it touches B,C,D and E. The velocity of particle is v m/s. O is the centre of the hexagon. x-axis is parallel to horizontal and y-axis is perpendicular to AE and CD (g=10 m/s^2) $\left(\sqrt{3}=1.73\right)$.

31. The maximum height from the horizontal surface X' attained by the projectile is ___ m.


Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q32 **Kinematics – Hexagon projectile – speed at the top**
JEE Advanced · Physics · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **5%** got it right · Campus: **5%** (3 of 61)

Tough one
You: Skipped

PARAGRAPH-I



A regular hexagon of side a=$10\sqrt{3}$ m is shown in figure. From any point a particle is projected such that it touches B,C,D and E. The velocity of particle is v m/s. O is the centre of the hexagon. x-axis is parallel to horizontal and y-axis is perpendicular to AE and CD (g=10 m/s^2) $\left(\sqrt{3}=1.73\right)$.

32. The velocity of projectile at maximum height is _____ m/s

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

JEE Advanced · Chemistry – questions to revisit


2 questions · easy misses first
cropped from the question paper


Q47 Carboxylic Acids


JEE Advanced · Chemistry · Integer · marking +4 / 0 · marks lost 4
Top 100: **32%** got it right · Campus: **33%** (20 of 61)


Tough one
You: Wrong

47. At room temperature how many of the following compounds can decompose $NaHCO_3$ to liberate CO_2?

a) 

b) 

c) 

d) 

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

Q51 Carboxylic Acids – Moles of $LiAlH_4$ to reduce RCOOH to RCH_2OH

JEE Advanced · Chemistry · Paragraph · marking +3 / 0 · marks lost 3
Top 100: **17%** got it right · Campus: **23%** (14 of 61)

Tough one
You: Wrong

PARAGRAPH-II

Carboxylic acids are organic compounds containing the carbonyl(>C=O) and hydroxyl (-OH) functional groups combined as a carboxyl group(-COOH). Due to resonance stabilization of the carboxylate anion, carboxylic acids exhibit higher acidity compared to alcohols and phenols. They participate in several characteristic reactions, including nucleophilic acyl substitution (esterification, amide formation, acid halide synthesis), decarboxylation, and reduction to primary alcohols. Furthermore, the presence of the –COOH group strongly influences physical properties, enabling extensive intermolecular hydrogen bonding that results in higher boiling points and stable dimeric structures in non-polar solvents.

51. The number of moles of $LiAlH_4$ required theoretically to reduce eight moles of carboxylic acid to primary alcohol

$$RCOOH \xrightarrow[2.H_3O^+]{1.LiAlH_4} RCH_2OH$$

Why did I lose this question? (write the reason and the correct approach)

☐ Did not know the concept ☐ Calculation mistake ☐ Misread the question ☐ Ran out of time ☐ Guessed

10 How to read this report

Where the numbers come from. Marks, percentages, All-India / State / Campus ranks, subject marks, subject All-India ranks and the wrong / skipped questions are the official results in the institute's *All India Marks Analysis* (graded with the final answer key). Maximum marks leave out deleted questions — JEE Mains: Q56 — JEE Advanced: Q25. **Program rank** = rank in Pragya G10 (EXT) Achievers across all campuses, from the batch's question-wise sheet. **Top 100** = the batch's top 100 across all campuses, from the same question-wise sheet; * where it has no data for a question, the campus figure decides the priority. **Campus** = all students of this program on campus. **Priority:** Easy miss = 70%+ of the top 100 got it right; Within reach = 40–69%; Tough one = below 40%. **Marks lost** on a question = its marks plus the negative mark if answered wrong, its marks if skipped. **Weekly-test average** = your average over the weekly Mains tests (WTM) or Advanced tests (WTA) before this Part Test, from the campus averages sheet. The weekly Advanced tests are marked out of 240; their averages are scaled to 180 to match this paper (× 180/240, so 80 per subject becomes 60). Percentages are unchanged.