

Eigenvalue Problem

Mingmin Chi

SCS Fudan University, Shanghai, China

1 Why $\mathbf{Ax} = \lambda\mathbf{x}$

2 How to Solve $\mathbf{Ax} = \lambda\mathbf{x}$

1 Why $\mathbf{Ax} = \lambda \mathbf{x}$

2 How to Solve $\mathbf{Ax} = \lambda \mathbf{x}$

Eigenvalue

- Eigenvalues have their best importance in dynamic problems
- The solution is changing with time-growing or decaying or oscillating
- Suppose we need the hundredth power \mathbf{A}^{100}

$$\mathbf{A} = \begin{bmatrix} .8 & .3 \\ .2 & .7 \end{bmatrix}$$

Eigenvalue

- Eigenvalues have their best importance in dynamic problems
- The solution is changing with time-growing or decaying or oscillating
- Suppose we need the hundredth power \mathbf{A}^{100}

$$\begin{bmatrix} .8 & .3 \\ .2 & .7 \end{bmatrix} \quad \begin{bmatrix} .7 & .45 \\ .3 & .55 \end{bmatrix}$$

$\mathbf{A} \qquad \mathbf{A}^2$

Eigenvalue

- Eigenvalues have their best importance in dynamic problems
- The solution is changing with time-growing or decaying or oscillating
- Suppose we need the hundredth power \mathbf{A}^{100}

$$\begin{array}{ccc} \begin{bmatrix} .8 & .3 \\ .2 & .7 \end{bmatrix} & \begin{bmatrix} .7 & .45 \\ .3 & .55 \end{bmatrix} & \begin{bmatrix} .65 & .525 \\ .35 & .475 \end{bmatrix} \\ \mathbf{A} & \mathbf{A}^2 & \mathbf{A}^3 \end{array}$$

Eigenvalue

- Eigenvalues have their best importance in dynamic problems
- The solution is changing with time-growing or decaying or oscillating
- Suppose we need the hundredth power \mathbf{A}^{100}

$$\begin{array}{cccc}
 \begin{bmatrix} .8 & .3 \\ .2 & .7 \end{bmatrix} & \begin{bmatrix} .7 & .45 \\ .3 & .55 \end{bmatrix} & \begin{bmatrix} .65 & .525 \\ .35 & .475 \end{bmatrix} & \dots \begin{bmatrix} .6 & .6 \\ .4 & .4 \end{bmatrix} \\
 \mathbf{A} & \mathbf{A}^2 & \mathbf{A}^3 & \dots \mathbf{A}^{100}
 \end{array}$$

Eigenvalue

- Eigenvalues have their best importance in dynamic problems
- The solution is changing with time-growing or decaying or oscillating
- Suppose we need the hundredth power \mathbf{A}^{100}

$$\begin{bmatrix} .8 & .3 \\ .2 & .7 \end{bmatrix} \quad \begin{bmatrix} .7 & .45 \\ .3 & .55 \end{bmatrix} \quad \begin{bmatrix} .65 & .525 \\ .35 & .475 \end{bmatrix} \quad \dots \quad \begin{bmatrix} .6 & .6 \\ .4 & .4 \end{bmatrix}$$

$\mathbf{A} \qquad \mathbf{A}^2 \qquad \mathbf{A}^3 \qquad \dots \qquad \mathbf{A}^{100}$

- \mathbf{A}^{100} was found using the eigenvalues of \mathbf{A} , not by multiplying 100 matrices

Eigenvector

- Multiplied by \mathbf{A} , almost all vectors change direction
- Certain exceptional vectors \mathbf{x} are in the same direction as \mathbf{Ax} .

Eigenvector

- Multiplied by \mathbf{A} , almost all vectors change direction
- Certain exceptional vectors \mathbf{x} are in the same direction as \mathbf{Ax} . Those are the “eigenvectors”
- An eigenvector of a linear transformation is a vector that is either left unaffected or simply multiplied by a scale factor after the transformation (the former corresponds to a scale factor of 1)

$$\mathbf{Ax} = \lambda \mathbf{x}$$

Eigenvector

- Multiplied by \mathbf{A} , almost all vectors change direction
- Certain exceptional vectors \mathbf{x} are in the same direction as \mathbf{Ax} . Those are the “eigenvectors”
- An eigenvector of a linear transformation is a vector that is either left unaffected or simply multiplied by a scale factor after the transformation (the former corresponds to a scale factor of 1)

$$\mathbf{Ax} = \lambda \mathbf{x}$$

where the number λ is the “eigenvalue”

- We can also say that \mathbf{Ax} is parallel to \mathbf{x}

Steady State

$$\mathbf{A} = \begin{bmatrix} .8 & .3 \\ .2 & .7 \end{bmatrix} \quad \mathbf{x}_1 = \begin{bmatrix} .6 \\ .4 \end{bmatrix} \quad \mathbf{x}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

Steady State

$$\mathbf{A} = \begin{bmatrix} .8 & .3 \\ .2 & .7 \end{bmatrix} \quad \mathbf{x}_1 = \begin{bmatrix} .6 \\ .4 \end{bmatrix} \quad \mathbf{x}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \quad \lambda_1 = 1 \quad \lambda_2 = .5$$

$$\text{Col}_1[\mathbf{A}] = \begin{bmatrix} .8 \\ .2 \end{bmatrix} = \mathbf{x}_1 +$$

Steady State

$$\mathbf{A} = \begin{bmatrix} .8 & .3 \\ .2 & .7 \end{bmatrix} \quad \mathbf{x}_1 = \begin{bmatrix} .6 \\ .4 \end{bmatrix} \quad \mathbf{x}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \quad \lambda_1 = 1 \quad \lambda_2 = .5$$

$$\text{Col}_1[\mathbf{A}] = \begin{bmatrix} .8 \\ .2 \end{bmatrix} = \mathbf{x}_1 + .2\mathbf{x}_2$$

Steady State

$$\mathbf{A} = \begin{bmatrix} .8 & .3 \\ .2 & .7 \end{bmatrix} \quad \mathbf{x}_1 = \begin{bmatrix} .6 \\ .4 \end{bmatrix} \quad \mathbf{x}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \quad \lambda_1 = 1 \quad \lambda_2 = .5$$

$$\text{Col}_1[\mathbf{A}] = \begin{bmatrix} .8 \\ .2 \end{bmatrix} = \mathbf{x}_1 + .2\mathbf{x}_2$$

$$\text{Col}_1[\mathbf{A}^{99}] = \mathbf{x}_1 + .2$$

Steady State

$$\mathbf{A} = \begin{bmatrix} .8 & .3 \\ .2 & .7 \end{bmatrix} \quad \mathbf{x}_1 = \begin{bmatrix} .6 \\ .4 \end{bmatrix} \quad \mathbf{x}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \quad \lambda_1 = 1 \quad \lambda_2 = .5$$

$$\text{Col}_1[\mathbf{A}] = \begin{bmatrix} .8 \\ .2 \end{bmatrix} = \mathbf{x}_1 + .2\mathbf{x}_2$$


$$\text{Col}_1[\mathbf{A}^{99}] = \mathbf{x}_1 + .2(.5)^{99}\mathbf{x}_2 = \begin{bmatrix} .6 \\ .4 \end{bmatrix} + \begin{bmatrix} \text{very} \\ \text{small} \\ \text{vector} \end{bmatrix}$$

- the eigenvector \mathbf{x}_1 is a “steady state” that does not change
- the eigenvector \mathbf{x}_2 is a “decaying mode” that visually disappears
- the higher the power of \mathbf{A} , the closer its columns approach the steady state
- this particular \mathbf{A} is a [Markov matrix](#)


Eigenvector - An Example

After a shear transformation

$$\mathbf{A} = \begin{bmatrix} 1 & 0 \\ -\frac{1}{2} & 1 \end{bmatrix}$$




Eigenvector - An Example



The shear transformation of the Mona Lisa

- the central vertical axis (red vector) was not modified, but the diagonal vector (blue) has changed direction

Eigenvector - An Example



The shear transformation of the Mona Lisa

- the central vertical axis (red vector) was not modified, but the diagonal vector (blue) has changed direction
- the red vector is an eigenvector of the transformation with eigenvalue 1
- all vectors with the same vertical direction - i.e. parallel to this vector - are also eigenvectors, with the same eigenvalue

1 Why $\mathbf{Ax} = \lambda \mathbf{x}$

2 How to Solve $\mathbf{Ax} = \lambda \mathbf{x}$

Eigenvalue Equation

So far, we could try to solve $\mathbf{Ax} = \lambda\mathbf{x}$ by trial and error, but ... Note that

$$\mathbf{Ax} = \lambda\mathbf{x} \Rightarrow (\mathbf{A} - \lambda\mathbf{I})\mathbf{x} = \mathbf{0}$$

- $\mathbf{A} - \lambda\mathbf{I}$ is not invertible
- the determinant of $\mathbf{A} - \lambda\mathbf{I}$ must be zero, i.e.,

$$\det(\mathbf{A} - \lambda\mathbf{I}) = 0$$

involves only λ , not \mathbf{x}

- then, find the roots of this polynomial

Properties of Eigenvalues

- the product of the n eigenvalues equals the determinant \mathbf{A}
- the sum of the n eigenvalues equals the sum of the n diagonal entries - the **trace** of \mathbf{A} :

$$\lambda_1 + \dots + \lambda_n = \text{trace} = a_{11} + \dots + a_{nn}$$

Diagonalization

- suppose the n by n matrix \mathbf{A} has n linearly independent eigenvectors and put them into the columns of an eigenvector matrix \mathcal{S} . We have

$$\mathcal{S}^{-1}\mathbf{A}\mathcal{S} = \Lambda = \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}$$

the matrix \mathbf{A} is “diagonalized”

- without n independent eigenvectors, we cannot diagonalize
- the matrices \mathbf{A} and Λ have the same eigenvalues $\lambda_1, \dots, \lambda_n$, but the eigenvectors are different

Markov Matrix: a Revisit

$$\mathbf{A} = \begin{bmatrix} .8 & .3 \\ .2 & .7 \end{bmatrix}$$

- its entries are positive and every column adds to 1
- the largest eigenvalue is $\lambda_1 = 1$ and the corresponding eigenvector is the steady state - which all columns of \mathbf{A}^k will approach
- \mathbf{A}^k has the same eigenvectors as \mathbf{A} but λ_i^k of the \mathbf{A} 's

Markov Matrix: a Revisit

$$\mathbf{A} = \begin{bmatrix} .8 & .3 \\ .2 & .7 \end{bmatrix}$$

- its entries are positive and every column adds to 1
- the largest eigenvalue is $\lambda_1 = 1$ and the corresponding eigenvector is the steady state - which all columns of \mathbf{A}^k will approach
- \mathbf{A}^k has the same eigenvectors as \mathbf{A} but λ_i^k of the \mathbf{A} 's

$$\mathbf{A}^2 = \mathbf{S} \mathbf{\Lambda} \mathbf{S}^{-1} \mathbf{S} \mathbf{\Lambda} \mathbf{S}^{-1} = \mathbf{S} \mathbf{\Lambda}^2 \mathbf{S}^{-1}$$

$$\mathbf{A}^k = \mathbf{S} \mathbf{\Lambda}^k \mathbf{S}^{-1}$$