

16 methods and comparators over 58 evaluations; blue clears that column's floor, orange falls below.

	cell-context												donor				unseen perturbation						
	T1 · Kang								T5c · OP3				T2	T3 · CRISPR				T4	T5u				
ctrl-pred	-.02	.00	.02	.00	.01	-.02	.01	.00	-.03	.01	-.01	-.01	.00										
cell-mean floor	.67	.62	.62	.71	.53	.46	.53	.58	.07	.02	.00	.01	.26										
donor-shift	.83	.79	.72	.82	.60	.82	.88	.78	.28	.19	.26	.32	.02										
linear-PCA floor	.83	.78	.72	.82	.60	.82	.88	.78	.30	.18	.26	.33	.04										
scGen	.77	.78	.77	.73	.54	.84	.82	.75	.13	.16	.24	.19	.15										
CPA/chemCPA	.36	.34	.53	.50	.27	.41	.26	.41	.11	-.02	.10	.04	.21										
biolord																							
GEARS																							
AttnPert																							
scGPT	.67	.64	.62	.69	.54	.40	.54	.58	.38	.28	.35	.38	.25										
scFound.	.75	.73	.68	.70	.52	.69	.69	.65	.28	.28	.38	.38	.12										
STATE	.59	.63	.68	.70	.41	.76	.79	.58	.08	.00	.03	.05	.14										
PertAdapt																							
CellOT	.81	.82	.76	.80	.56	.91	.90	.74	.15	.15	.26	.23	.37										
scPRAM	.80	.73	.69	.81	.54	.90	.86	.68	.19	.12	.22	.20	.16										
PerturbNet	.79	.79	.78	.82	.55	.94	.88	.75	.12	.10	.22	.15	.25										
PRnet									.13	.04	.08	.13											
CellFlow	.83	.82	.79	.83	.56	.93	.88	.77	.10	.07	.09	.15	.24										
FP-ridge									.42	.31	.39	.44											
lin-shift-KO																							

method (grouped by role)

cell fill = margin over that column's binding floor (Pearson-Δ)

-.43 .00 +.13

blue = above the floor; white = at the floor; orange = below it

clears both floor members and survives multiplicity correction

point-estimate clearance, without statistical support

adapted interface

diagnostic comparator

not evaluated

row accent:

baseline

conditioned

diagnostic comparators

93 of 106 donors
over the cell-mean floor
mean gap +0.107 [+0.086, +0.128]
paired Wilcoxon $p = 1.6 \times 10^{-13}$
mean +0.107

(Pearson- Δ)

cell-mean floor

donor (Soskic CD4 leave-one-donor-out), sorted by gap