
MARROW ED8

Paediatrics

Comprehensive Question Bank

Contents

Chapter	Title	Page
1	Basics of Neonatology and Routine Newborn Care	4
2	Disorders of Newborn	20
3	Diseases in Neonates requiring Special Care	39
4	Apgar score and Neonatal Resuscitation	56
5	Developmental Milestones	68
6	Facets of Growth and Development	83
7	Nutrition and Breastfeeding	101
8	Protein Energy Malnutrition	117
9	Deficiency of Fat Soluble Vitamins	131
10	Deficiency of Water-soluble Vitamins & Trace El...	151
11	Fluid and Electrolyte Disorders	167
12	Mendelian and Non-Mendelian Disorders	181
13	Chromosomal Disorders	213
14	Metabolic Disorders of Amino Acids	238
15	Metabolic Disorders of Urea Cycle, Complex Mole...	260
16	Polio and AIDS	280
17	Pediatric Bacterial and Parasitic Infections	296
18	Measles, Mumps, Rubella and Other Viral Infections	325
19	Surgical GI Disorders	349
20	Medical GI disorders	371

Chapter	Title	Page
21	Disorders of the Liver	384
22	Neonatal Respiratory Disorders	402
23	Childhood Respiratory Disorders	422
24	Fetal Circulation	448
25	Acyanotic Congenital Heart Diseases	457
26	Cyanotic Congenital Heart Diseases	479
27	Pediatric Nephrology	502
28	Pediatric Urology	525
29	Disorders of the Nervous System	538
30	Disorders of Thyroid	579
31	Congenital Adrenal Hyperplasia and Related Diso...	595
32	Disorders of the Pituitary Gland	617
33	Disorders of Puberty	627
34	Paediatric Hemato-oncology	640
35	Solid Neoplasms of Childhood	661
36	Musculoskeletal Disorders	691
37	Paediatric Rheumatology	705
38	Pediatric Hematology - Introduction, Bleeding, ...	719
39	Pediatric Anemias	734

Basics of Neonatology and Routine Newborn Care

Question 1:

Which of the following timelines would be included in the perinatal period as per WHO?

- a) 37th week of gestation to 28 days after birth
- b) 20th week of gestation to 7 days after birth
- c) 28th week of gestation to 28 days after birth
- d) 22nd week of gestation to 7 days after birth

Question 2:

You would classify a baby as post-term if it is born after _____

- a) 40 weeks of completed gestation
- b) 40 weeks and 2 days
- c) 41 weeks of completed gestation
- d) 42 weeks of completed gestation

Question 3:

You would classify a baby as a very low birth weight baby if the weight of the baby is found to be less than _____ kg.

- a) 1
- b) 1.5
- c) 2
- d) 2.5

Question 4:

Reena, a 34-year-old lady with gestation diabetes mellitus delivers an LGA baby at 37 weeks. What would the birth weight of this child be?

- a) >75th Percentile
- b) >97th Percentile

- c) >90th Percentile
- d) >95th Percentile

Question 5:

As the intern assisting in the labor room, you are tasked with administering Vitamin K to the term newborns with no complications. Which of the following dose and route would you use for administration?

- a) 0.5 - 1 mg IM
- b) 5 - 10 mg IV
- c) 1 - 5 mg IV
- d) 0.1 - 0.5 mg IM

Question 6:

A term neonate born by normal vaginal delivery is diagnosed with hypothermia. Which of the following is not a physiological response for heat regulation in this baby?

- a) Shivering
- b) Breakdown of brown fat with noradrenaline secretion
- c) Increased muscular activity
- d) Cutaneous vasoconstriction

Question 7:

You are conducting the routine newborn assessment of a 2.3 kg baby born 10 minutes back. You would classify the neonate as having cold stress if you notice the body temperature to be:

- a) 35.4-36.0 °C
- b) 33.4-34.0 °C
- c) 34.4-35.4 °C
- d) 36.0-36.4 °C

Question 8:

A 24-year-old primigravida delivers a preterm baby at 33 weeks of gestation with a birth weight of 2 kg. She is advised to hold her child as shown below. Which of the following statements about this method is false?



- a) Can be continued at home
- b) Safety and thermal protection is equivalent to incubators
- c) Reduces severe morbidity in the newborns
- d) Should be performed only by the mother

Question 9:

You are the intern posted in the labor room and are tasked with calculating the gestational maturity of the infants delivered. As per the expanded New Ballard scoring, what will be the lowest and highest scores you can obtain?

- a) 0 and 40
- b) -10 and 50
- c) -20 and 44
- d) 10 and 32

Question 10:

Which of the following is a component of the neuromuscular criteria for maturity according to the expanded New Ballard score?

- a) Heel-knee test
- b) Finger-nose test
- c) Scarf sign
- d) Cubital angle

Question 11:

A 1.8 Kg neonate is born to a 24-year-old primigravida with no antenatal checkups. While calculating the physical maturity of the neonate by the new Ballard scoring, which of the following criteria would you not take into account?

- a) Skin
- b) Lanugo hair
- c) Palmar surface
- d) Breast

Question 12:

While examining a healthy term neonate, you notice that the baby passes greenish-black colored stools. This color is due to -

- a) Bilirubin
- b) Biliverdin
- c) Meconium
- d) Gut flora of neonate

Question 13:

When does the parachute reflex disappear?

- a) At 3 months
- b) At 6 months
- c) At birth and reappears at 9 months
- d) Never

Question 14:

A 6-day old infant is being examined. While checking the reflexes, a brisk Moro's reflex is noted. Which of the following conditions is a likely cause for this?

- a) Kernicterus
- b) Stage III hypoxic ischemic encephalopathy (HIE)
- c) Hypoglycemia
- d) Erb's palsy

Question 15:

You are demonstrating findings in a pediatric OSCE examination on a 4-day old neonate. On turning the baby's head to the left, you obtain the finding as shown below. At what age does this reflex normally disappear?



- a) 6 months
- b) 9 months
- c) 4 months
- d) 5 months

Question 16:

A first-time mother rushes to the OPD with her 4-day old infant and worriedly points to the rash that appears as shown below. You reassure her of its harmless, transient nature. On microscopic examination, these papular lesions on an erythematous base will reveal



- a) Eosinophils
- b) Neutrophils
- c) Giant Cells
- d) Lymphocytes

Question 17:

A 5-month-old infant is brought to the OPD for a well-baby check-up. A lesion that was present since birth is noticed as given below. What is the most likely diagnosis?



- a) Dermal melanocytosis

- b) Salmon patch
- c) Neonatal pustular melanosis
- d) Harlequin colour change

Answer Key

Question No.	Correct Option
1	d
2	d
3	b
4	c
5	a
6	a
7	d
8	d
9	b
10	c
11	c
12	b
13	d
14	c
15	a
16	a
17	a

Detailed Explanations

Solution to Question 1:

The WHO states that the perinatal period commences at 22 completed weeks and ends at seven completed days after birth - 154 days. It corresponds to when the fetus weighs 500g or more.


This definition is not to be confused with the period used for the calculation of the perinatal mortality rate (PMR). To calculate the PMR, a fetal weight above 1000 g is considered, which is achieved at 28 weeks of gestation. Therefore, for all statistical purposes, international comparisons, and calculation of perinatal mortality rate, the period from the 28th week of life is considered.

Perinatal period: from the 22nd week.

Perinatal mortality rate calculation: from the 28th week.

The neonatal period is defined as the first 28 days after birth. Infancy is defined as the first year of life.

The image below shows various time periods. Only the time period for perinatal mortality and the perinatal period are different as shown in the image. For others, corresponding mortality rates are calculated using the same time periods shown.



Solution to Question 2:

Post-term infants are those born after 42 completed weeks of gestation, as calculated from the mother's last menstrual period, regardless of weight at birth.

Solution to Question 3:


A very low birth weight baby is one whose birth weight is <1.5 Kg. They are often premature (<37 weeks of gestation).

A low birth weight baby is one whose birth weight is <2.5 Kg.

An extremely low birth weight baby is one whose birth weight is <1 kg.

Solution to Question 4:

Infants with birth weight greater than the 90th percentile for their gestational age are called large for gestational age (LGA) babies.



Maternal diabetes and obesity are predisposing factors for LGA babies. They have a higher incidence of:

- Birth injuries:
- Brachial plexus injury
- Phrenic nerve injury
- Cephalohematomas
- Hypoglycemia
- Polycythemia
- Congenital heart disease

Solution to Question 5:

All newborn infants are to be administered vitamin K at a dose of 0.5–1 mg intramuscularly, to prevent hemorrhagic disease of the newborn (HDN).

However, for newborns with vitamin K deficiency, the dose of vitamin K administered is 1–5 mg intravenously.

Solution to Question 6:

Neonates do not generate heat by shivering.

Heat production in neonates happens by increasing the metabolic rate and oxygen consumption by releasing norepinephrine, which results in non-shivering thermogenesis through beta-oxidation of brown fat. This sympathetic surge also causes cutaneous vasoconstriction.

Increased physical activity - crying, increased body movements, and muscular activity also contribute to thermogenesis.

Preterm and small-for-gestational-age (SGA) infants have scanty brown fat stores.

Solution to Question 7:

A temperature of 36.0 to 36.4°C categorizes a neonate as having cold stress.

An infant's normal body temperature is 36.5 to 37.5 °C. It is measured in the apex of the baby's axilla by holding the thermometer for at least 3 minutes.

Hypothermia is an axillary temperature of ≤ 36.5 °C.

- Cold stress: 36.0 to 36.4 °C
- Moderate hypothermia: 32 to 35.9 °C
- Severe hypothermia: ≤ 32 °C

Hyperthermia is an axillary temperature of ≥ 37.5 °C.

Solution to Question 8:

Kangaroo Mother Care (KMC) can be carried out by the mother or any other caregiver.

KMC was introduced for the prevention of neonatal hypothermia in low birth weight/ premature newborns. It has 4 components—

- Skin-to-skin positioning of the newborn on the mother's chest.
- Adequate nutrition through breastfeeding.
- Early discharge and ambulatory care.
- Support for mother and family for child care.

Evidence shows that-

- It is as good as incubators, in terms of safety and thermal protection (measured by mortality)
- Reduces severe morbidity by way of facilitating breastfeeding
- Humanization of neonatal care
- Promoted bonding between mother and child

Eligibility criteria for KMC:

It is indicated for all hemodynamically stable low birth weight babies. Short sessions of KMC can be initiated in babies on medical treatment like IV fluids or oxygen therapy. Initiation of KMC can vary with the birth weight of the baby.

- Birth weight ≤ 1200 g - It may take days to weeks for initiation of KMC
- Birth weight =1200-1800 g - Only a few days may be taken before KMC initiation

- Birth weight ≥ 1800 g - KMC can be initiated immediately after birth

All mothers without any serious illness can provide KMC.

Positioning for KMC: The baby is placed between the mother's breast in an upright position. The hips are abducted and flexed in the frog position and the arms are also flexed. The abdomen of the baby should be at the level of the mother's epigastrium. In this position, the mother's breathing would stimulate the baby and prevent the development of apnea.

The duration of KMC should be gradually increased from short sessions of more than one hour to 24 hours a day.

KMC is not necessary once the baby attains 2500 g of weight or 37 weeks of gestation.

Note:

Incubatory care, feeding via nasal catheter, and prevention of infection are part of intensive care for low birthweight babies.

Solution to Question 9:

The lowest score on the expanded New Ballard Score is -10, which corresponds to 20 weeks gestational age, and the highest score is 50, which corresponds to 44 weeks gestational age.

An increase in the score by 5 points corresponds to an increase of 2 weeks gestational age.

Maturity Rating on the expanded New Ballard Score:


Score	Weeks
-10	20
-5	22
0	24
5	26
10	28
15	30
20	32
25	34
30	36
35	38
40	40
45	42
50	44

Solution to Question 10:

Scarf sign is a component of the neuromuscular criteria for maturity according to the expanded new Ballard score.

Scarf sign: If the infant's arm is pulled medially across the chest (as shown in the image below),

- In a hypotonic infant, the elbow will cross the midline.
- In a term infant with a normal tone, the elbow will not reach the midline.



Neuromuscular criteria for maturity (according to the expanded New Ballard score) include:

- Posture
- Square window (wrist)
- Arm recoil
- Popliteal angle
- Scarf sign
- Heel to ear

Physical criteria for maturity (according to the expanded New Ballard score) include:

- Skin
- Lanugo
- Eye/ear
- Plantar crease
- Breast
- Genitals

The expanded new Ballard score is used to assess the physical and neuromuscular maturity of the newborn by scoring six criteria and thereby calculating the gestational age. It is accurate between 20 to 44 weeks.

The lowest score on the expanded New Ballard Score is -10, which corresponds to 20 weeks gestational age, and the highest score is 50, which corresponds to 44 weeks gestational age. An increase in the score by 5 points corresponds to an increase of 2 weeks gestational age

Given in the image below is the scoring system for the physical and neuromuscular criteria:

Neuromuscular Maturity							
Score	-1	0	1	2	3	4	5
Posture							
Square window (wrist)							
Arm recoil							
Popliteal angle							
Scarf sign							
Heel to ear							

Physical Maturity							
Skin	Sticky, friable, transparent	Gelatinous, red, translucent	Smooth, pink, visible veins	Superficial peeling and/or rash; few veins	Cracking, pale areas; rare veins	Parchment, deep cracking; no vessels	Leathery, cracked, wrinkled
Lanugo	None	Scarse	Abundant	Thinning	Bald areas	Mostly bald	
Plantar surface	Heel size 40-50 mm, < 40 mm: -2	> 50 mm, no crease	Faint red marks	Anterior transverse crease only	Creases, anterior 2/3	Creases over entire sole	
Breast	Imperceptible	Barely perceptible	Flat areola, no bud	Stippled areola, 1-2 mm bud	Raised areola, 3-4 mm bud	Full areola, 5-10 mm bud	
Eye/Ear	Lids fused, loosely -1 tightly: -2	Lids open, pinna flat, stays folded	Slightly curved pinna, soft	Well curved pinna, soft but ready recoil	Formed and firm, instant recoil	Thick cartilage, ear 95°	
Genitals (male)	Scrotum flat, smooth	Scrotum empty, faint rugae	Testes in upper canal, little rugae	Testes descending, few rugae	Testes down, good rugae	Testes pendulous, deep rugae	
Genitals (female)	Clitoris prominent, labia flat	Clitoris prominent, small labia minora	Clitoris prominent, enlarging minora	Majora and minora equally prominent	Majora large, minora small	Majora cover clitoris and minora	

Maturity Rating	
Score	Weeks
-10	20
-5	22
0	24
5	26
10	28
15	30
20	32
25	34
30	36
35	38
40	40
45	42
50	44

Solution to Question 11:

Palmar surface is not a component of the physical criteria for maturity according to the expanded New Ballard score.

Physical criteria for maturity (according to the expanded New Ballard score):

- Skin
- Lanugo
- Eye/ear
- Plantar crease (Not palmar surface)
- Breast
- Genitals

Solution to Question 12:

The greenish-black color of the first stool in the newborn is due to biliverdin.

Meconium is a dark, sticky material that is normally present in the intestine at birth and passed in the feces after birth. It is detected in the ileum at about 70–85 days of fetal life and begins to fill the colon and lower ileum late in the 4th month of gestation. The greenish color of the meconium is the consequence of the bile pigments excreted by the biliary tract. The main component

responsible for the greenish color of the meconium is biliverdin (a pigment formed from the oxidation of bilirubin).

Up to the 22nd week of gestation, the intestinal content is not greenish and for this reason, fetal defecation would not stain the amniotic fluid. The production of biliverdin starts over the next few weeks or at a time closer to labor, probably due to a biochemical switch related to the bilirubin oxidation system. Bilirubin is responsible for the yellow color of the stools.

Neonates do not harbor gut flora at birth.

Solution to Question 13:

The parachute reflex has its onset at 7–8 months of age. It is fully developed by 10–11 months of age and remains throughout life.

It can be evoked by holding the infant's trunk and then suddenly lowering the infant as if he or she were falling. The arms will spontaneously extend to break the infant's fall.

This reflex is a prerequisite to walking.

Solution to Question 14:

A brisk Moro reflex can be seen in neonates with hypoglycemia.

In hypoglycemic babies, there is an activation of the autonomic system, which results in jitteriness. Newborns often exhibit this as an exaggerated Moro reflex.

The Moro reflex is a primitive reflex that is very sensitive to the infant's level of alertness. It is elicited by supporting the infant in the semierect position and allowing the infant's head to fall backward onto the examiner's hand.

An animated gif of normal response to Moro's reflex where there is symmetrical extension and abduction of the upper extremities, followed by flexion of the upper extremities is shown below



The Moro reflex disappears by 6 months of age in normal infants.

Absent Moro reflex: caused by significant generalized disturbance of the CNS

- Stage III HIE
- Kernicterus

Asymmetric Moro reflex: feature of the root, plexus, or nerve disease

- Fractured clavicle
- Brachial plexus injury (Erb's palsy)
- Hemiparesis

Brisk Moro reflex:

- Neonatal hypoglycemia
- Can also be a normal variant in healthy infants
- Stage I HIE

Exaggerated, stereotyped, non-habituating Moro reflex is a feature of severe bilateral intrauterine cerebral disturbance, perhaps because of the release of the brain stem from inhibitory cortical influences. It is seen in conditions such as the following:

- Hydranencephaly
- Severe micrencephaly

Moro reflex in different stages of hypoxic-ischemic encephalopathy (HIE):

- Stage I HIE (mild): Brisk
- Stage II HIE (moderate): Weak/incomplete
- Stage III HIE (severe): Absent

Option A: In kernicterus, there is a deposition of unconjugated bilirubin in the basal ganglia and brain stem. Hence, the Moro reflex is absent

Option C: In stage III HIE, the muscles are flaccid and the infant is comatose. Thus, the Moro reflex is absent.

Option D: Erb's palsy results from an injury to the brachial plexus. So, the reflex is absent on the side of injury as the nerve supplying the muscles involved (efferent of the reflex) is affected.

Solution to Question 15:

Asymmetric tonic neck reflex (ATNR) disappears by 6 months postnatally.

The tonic neck reflex is produced by manually rotating the infant's head to 1 side and observing for the characteristic fencing posture (extension of the arm on the side to which the face is rotated and flexion of the contralateral arm).

An obligatory tonic neck response, in which the infant becomes “stuck” in the fencing posture, is always abnormal and implies a CNS disorder.



Solution to Question 16:

The given clinical complex describes a case of erythema neonatorum. Wright stained smear of the contents of the pustule will show an abundance of eosinophils.

They are papular lesions on an erythematous base, distributed over the trunk and face predominantly. They are seen 1-3 days after birth and may persist for up to a week, after which they resolve spontaneously.

Neutrophils are seen in pustular melanosis.

Solution to Question 17:

The above image shows Mongolian spots or dermal melanocytosis.


Mongolian spots are benign blue or slate-grey macular lesions, with clearly defined margins, that appear over the presacral region (most common), posterior thighs, legs, back, and forearms.

These spots occur due to the presence of melanin-containing melanocytes in the dermis, which is arrested during the migration from neural crest cells to the epidermis. They are non-tender and do not rapidly change in color or size.

Disorders of Newborn

Question 1:

A male infant born at 34 weeks of gestation by emergency LSCS was found to have the following finding on histopathological examination of his umbilical cord. Anomalies of which of the following systems are most likely to be found in this infant?



- a) Renal system
- b) Central nervous system
- c) Gastro-intestinal system
- d) Endocrine system

Question 2:

On a particularly busy day in the labor room, you are requested by the PG to take anthropometric measurements and calculate the ponderal index of a baby while she attends to another baby. You find the ponderal index to be 1.45. Which of the following organs is unlikely to be affected in this neonate?

- a) Subcutaneous fat
- b) Muscle
- c) Liver
- d) Brain

Question 3:

The ponderal index of a term baby delivered by elective cesarean section is 1.75. Which of the following is unlikely to be associated?

- a) Gestational diabetes
- b) Diabetes with vasculopathy
- c) Hypertension
- d) Sickle cell anemia

Question 4:

An unbooked primigravida, with no antenatal records, delivers a 2.5kg male baby by normal delivery. While ascertaining the maturity of the child, which of the following would be against a diagnosis of prematurity?

- a) No creases on sole
- b) Insignificant breast nodule
- c) Thick ear cartilage
- d) Empty scrotum

Question 5:

A lady with wrong dates and a corrected gestational age of 43 weeks presents with labor pains. The CTG suggests placental insufficiency, and she is taken up for emergency LSCS. Which of the following physical signs is unlikely to be seen in this neonate?

- a) Long nails
- b) Pale skin
- c) Alert faces
- d) Scanty hair

Question 6:

A 34-year-old lady with an obstetric score of G3P2L2 delivers a baby boy at 43 weeks + 2 days period of gestation. Which of the following are unlikely to be associated in the given scenario?

- a) Anemia
- b) Hypocalcemia
- c) Meconium aspiration
- d) Persistent pulmonary hypertension


Question 7:

Immediately following vaginal delivery, a baby is noted to have a diffuse ecchymotic swelling on the scalp. What is the most likely diagnosis?

- a) Cephalohematoma
- b) Caput succedaneum
- c) Subgaleal hemorrhage
- d) Epidural hematoma

Question 8:

Which of the following is not a feature of marking B?



- a) Swelling of the soft tissues of the scalp
- b) Does not cross suture lines
- c) Resorbed within 2 weeks to 3 months
- d) Appears only several hours after birth

Question 9:

A baby in the postnatal ward is noted to have paralysis of the left hand along with ipsilateral ptosis and miosis. What is the most likely cause?

- a) Erb-Duchenne paralysis
- b) Phrenic nerve palsy
- c) Horner syndrome
- d) Klumpke's paralysis associated with Horner syndrome

Question 10:

Following a delivery complicated by shoulder dystocia, the baby is noted to have a deformity in the right upper limb. On inspection, the orthopedics JR notes an adducted, internally rotated arm and a pronated forearm. Which nerve roots are affected here?

- a) Only C5
- b) C5 and C6
- c) C6 and C7
- d) C7 and C8

Question 11:

A term infant develops jaundice at day 2 of life. It first appeared on the sclera and progressed to the trunk by day 4. It completely disappeared by day 7 following phototherapy. Which of the following statements is false regarding the cause of this pattern of jaundice?

- a) Due to a rise in unconjugated bilirubin
- b) Due to limitation in the conjugation of bilirubin by the immature neonatal liver
- c) Due to Rh incompatibility between maternal and fetal red blood cells
- d) Due to increased breakdown of fetal red blood cells

Question 12:

A male infant with a birth weight of 3.2 Kg is admitted to the phototherapy unit with a total bilirubin of 11mg/dl at 36 hours of life. Which of the following statements does not describe pathological jaundice?

- a) Jaundice appears within first 24 hours after birth

- b) Jaundice present at or beyond age 3 weeks
- c) Jaundice associated with rise in bilirubin level at a rate of ≥ 5 mg/dl/24 hour
- d) Jaundice seen in the mid-abdomen and soles

Question 13:

The peripheral smear of a newborn baby girl sent on day 2 of life shows mild hemolysis. What is the most common cause of this finding in a newborn?

- a) Rh incompatibility
- b) ABO incompatibility
- c) Kell antigen incompatibility
- d) Hemolytic diseases due to infections

Question 14:

A 3.3 Kg term neonate is admitted on day 3 of life with yellowish discoloration of the skin and is going to undergo phototherapy. What is the most effective wavelength of light used for treatment?

- a) 370 to 390 nm
- b) 390 to 420 nm
- c) 460 to 490 nm
- d) 410 to 450 nm

Question 15:

A mother noticed yellowish discoloration of her term newborn within the first 6 hours of life. On examination, jaundice extended up to the lower abdomen and thighs. Which of the following drugs can be administered along with phototherapy?

- a) Sn Mesoporphyrin
- b) Tolazoline
- c) Tromethamine
- d) Sucrose

Question 16:

While on rounds in the postnatal ward, the senior PG shows you a baby with dark, grayish-brown skin discoloration. It is a term, AGA baby on day 7 of life. She tells you that it is a case of bronze baby syndrome. This syndrome is caused by:

- a) Phototherapy
- b) Hemosiderosis
- c) Vancomycin
- d) Chloramphenicol

Question 17:

A 3.2 Kg term neonate born to a primigravida with gestational diabetes presented to the OPD with complaints of lethargy, hypotonia, and difficulty in feeding. Periodic monitoring of blood glucose is being done. What blood glucose levels indicate a need for the treatment of neonatal hypoglycemia?

- a) < 45 mg/dL
- b) < 40 mg/dL
- c) < 50 mg/dL
- d) < 55 mg/dL

Question 18:

After the delivery of an infant of a diabetic mother, the blood glucose of the Infant was 60 mg/dl. Which other investigation would you do?

- a) Serum potassium
- b) Serum sodium
- c) Serum chloride
- d) Serum calcium

Question 19:

A 31-year-old lady with diabetes mellitus delivers a term neonate. Which of the following problems may be associated with the baby born to this lady?

- a) 1, 2, 3
- b) 2, 3, 5

c) 1, 4, 5

d) 2, 3, 4

Answer Key


Question No.	Correct Option
1	a
2	d
3	a
4	c
5	d
6	a
7	b
8	a
9	d
10	b
11	c
12	c
13	b
14	c
15	a
16	a
17	b
18	d
19	d

Detailed Explanations

Solution to Question 1:

The image shows the presence of a single umbilical artery which is associated with an increased risk for an occult renal anomaly.

The histopathological image of the cross-section of the umbilical cord shows 2 vessels as marked, a thin-walled umbilical vein and a thicker-walled umbilical artery.



The normal umbilical cord contains 2 arteries and 1 vein. A single umbilical artery is the most common cord abnormality. Anomalies are seen in 30% of cases with a single umbilical artery. The incidence is higher in twin deliveries.

Trisomy 18 is a common association.

Solution to Question 2:

A ponderal index of < 2 suggests asymmetric intrauterine growth restriction. Asymmetric IUGR is associated with a brain-sparing effect, unlike symmetric IUGR, where there is no sparing of head growth.

Ponderal index = $\frac{\text{Birth weight (gm)}}{\text{Length (cm)}^3} \times 100$.

It is a weight-height-related parameter used as an indicator of fetal growth status.

Symmetric IUGR	Asymmetric IUGR
Insult on fetal growth occurs early	Insult occurs in late gestation
Head circumference, length, and weight are equally affected	Relative sparing of brain growth
A ponderal index greater than or equal to 2	A ponderal index less than 2
Caused by genetic, chromosomal disorders, or TORCH infections	Caused by placental insufficiency, pregnancy-induced hypertension, or maternal medical diseases

Solution to Question 3:

A ponderal index of 1.75 suggests intrauterine growth restriction. Gestational diabetes usually presents with fetal macrosomia.

Maternal factors associated with intrauterine growth restriction (IUGR):

- Hypertension/renal disease
- Diabetes with placental vasculopathy
- Hypoxemia
- Malnutrition
- Chronic illness
- Sickle cell anemia
- Drugs (narcotics, alcohol, cigarettes, cocaine)
- Maternal infection and parasite infestations

Solution to Question 4:

Thick cartilage with a stiff ear in a newborn is a sign of maturity (not prematurity).

No creases on the sole, insignificant breast nodule, and empty scrotum are all physical signs of prematurity in a newborn.

Note:

Abundant lanugo hair is not discriminatory evidence of prematurity. This is because neonates of diabetic mothers born as a term or even post-term have abundant lanugo hair, especially over the pinna and upper back.

Score	Ear
0	Pinna flat and stays folded; no recoil
1	Slightly curved pinna; soft; slow recoil
2	Well-curved pinna; soft but ready recoil
3	Formed & firm, instant recoil
4	Thick cartilage, ear stiff

Expanded New Ballard Scoring -
Physical Maturity

Skin	Sticky, friable, transparent	Gelatinous, red, translucent	Smooth pink; visible veins	Superficial peeling and/or rash; few veins	Cracking pale areas; rare veins	Parchment, deep cracking; no vessels	Leathery, cracked, wrinkled
Lanugo	None	Sparse	Abundant	Thinning	Bald areas	Mostly bald	
Plantar surface	Heel-toe 40-50 mm: -1 <40 mm: -2	>50 mm, no crease	Faint red marks	Anterior transverse crease only	Creases anterior 2/3	Creases over entire sole	
Breast	Imperceptible	Barely perceptible	Flat areola, no bud	Stippled areola, 1-2 mm bud	Raised areola, 3-4 mm bud	Full areola, 5-10 mm bud	
Eye/Ear	Lids fused loosely: -1 tightly: -2	Lids open, Pinna flat, stays folded	Slightly curved pinna, soft, slow recoil	Well curved pinna, soft but ready recoil	Formed and firm, instant recoil	Thick cartilage, ear stiff	
Genitals (male)	Scrotum flat, smooth	Scrotum empty, faint rugae	Testes in upper canal, rare rugae	Testes descending, few rugae	Testes down, good rugae	Testes pendulous, deep rugae	
Genitals (female)	Clitoris prominent, labia flat	Clitoris prominent, small labia minora	Clitoris prominent, enlarging minora	Majora and minora equally prominent	Majora large, minora small	Majora cover clitoris and minora	

©MARROW

Solution to Question 5:

The physical sign of an infant born post-term (43 weeks) in association with presumed placental insufficiency is abundant hair (not scanty hair).

Infants born post-term in association with presumed placental insufficiency may have various physical signs, together termed as postmaturity syndrome (or dysmaturity syndrome).

These include:

- Desquamation, pale skin, and long nails.
- Loose skin around the thighs and buttocks.
- Meconium-stained nails, skin, vernix, and umbilical cord.
- Presence of placental membranes.

Solution to Question 6:

Anemia is not a complication of post-maturity. Polycythemia may be seen in neonates born after 42 weeks of gestation.

Complications of post-maturity include:

- Perinatal depression
- Meconium aspiration
- Persistent pulmonary hypertension
- Hypoglycemia
- Hypocalcemia

- Polycythemia

Solution to Question 7:


The most likely diagnosis is caput succedaneum.

Caput succedaneum is a diffuse, sometimes ecchymotic, edematous collection of serosanguinous fluid in the subcutaneous layer of the scalp, involving the presenting part in a vertex delivery. It may extend across the midline and crosses suture lines.

Option A: Cephalohematoma is a subperiosteal hemorrhage that is always limited to the surface of 1 cranial bone and does not cross suture lines. The swelling appears only several hours after birth. There is no discoloration of the overlying scalp.

Option C: Subgaleal hemorrhages occur with the rupture of emissary veins, which connect the dural sinuses with the superficial veins of the scalp. Blood accumulates between the epicranial aponeurosis of the scalp and the periosteum and resolves over 2-3 weeks. Often associated with vacuum-assisted delivery.

Option D: Epidural hemorrhages, usually traumatic in origin, are associated with cephalo-pelvic disproportion, prolonged labor, breech, precipitous labors, or occur as a result of mechanical assistance in labor. Majority resolve with no intervention.




Solution to Question 8:

The area marked B represents a cephalohematoma. It is not a swelling of the soft tissues of the scalp.

Cephalohematoma is a subperiosteal hemorrhage that is always limited to the surface of 1 cranial bone and does not cross suture lines. The swelling appears only several hours after birth. There is no discoloration of the overlying scalp.

Most cephalohematomas are resorbed within 2 weeks to 3 months, depending on their size. Cephalohematomas require no treatment, although phototherapy may be necessary to treat the associated hyperbilirubinemia (resorption of blood takes place by the breakdown of RBCs).



Other markings-

A- Caput succedaneum. It is a diffuse ecchymotic or edematous collection in the soft tissues of the scalp with vertex presentation. It can extend across skull sutures and disappears with the first week of life.

C- Sub-galeal hemorrhage. This is a collection of blood under the aponeurosis of the scalp. It is a large potential space, thus bleeding can be extensive. It occurs as a result of ventouse deliveries due to tears of the emissary veins which connect to the dural venous sinuses. It can also occur in hereditary coagulopathies and in association with skull fractures.

D- Epidural hemorrhage. Usually traumatic in origin, these are associated with cephalo-pelvic disproportion, prolonged labor, breech, precipitous labors, or as a result of mechanical assistance in labor. Majority resolve with no intervention.

Solution to Question 9:

Left-hand paralysis with ipsilateral ptosis and miosis in a newborn suggests a diagnosis of Klumpke's paralysis associated with Horner syndrome.

There is an injury to the 7th-8th cervical nerves and the 1st thoracic nerve. This produces a paralyzed hand where the intrinsic muscles of the hand (T1) and the ulnar flexors of the wrist and fingers (C8) are mainly affected, and the baby lies with the forearm supinated and the elbow flexed. There may be a unilateral Horner's syndrome, characterized by ipsilateral ptosis and miosis if the sympathetic fibers of the 1st thoracic root are also injured.


Option A: Erb-Duchenne paralysis affects the 5th and 6th cervical nerves. There is loss of abduction of the shoulder, external rotation, and supination. Moro's reflex is absent on the affected side.

Option B: Phrenic nerve palsy (3, 4, and 5 cervical nerves) results in diaphragmatic paralysis. It is characterized by cyanosis and irregular and labored breathing.

Option C: Horner's syndrome is characterized by miosis, ptosis, enophthalmos, anhydrosis, and loss of ciliospinal reflex.



Klumpke's paralysis-Damage to C8, T1



Solution to Question 10:

The clinical vignette suggests a diagnosis of Erb–Duchenne paralysis. The injury is limited to the 5th and 6th cervical nerve roots.

The characteristic position consists of adduction and internal rotation of the arm with pronation of the forearm. There is loss of abduction of the shoulder, external rotation, and supination.

There is an asymmetric Moro's Reflex

Note:

In shoulder dystocia, the most common nerve injury is to the C5-6 nerve root. The most common bone to fracture is the clavicle.




Solution to Question 11:

The above pattern of jaundice is suggestive of physiological jaundice/icterus neonatorum. It is due to increased bilirubin production from the breakdown of fetal red blood cells combined with transient limitation in the conjugation of bilirubin by the immature neonatal liver.

It first appears on the 2nd or 3rd day, usually peaks on the 4th day at 5–6 mg/dL, and decreases to <2 mg/dL between the 5th and 7th days after birth.

Rh incompatibility between maternal and fetal red blood cells leads to pathological jaundice.

Kramer's Zones for Neonatal Jaundice

	Kramer's zones	Approximate TSB level	
		Mild jaundice (Lemon yellow color)	Deep jaundice (Orange-yellow color)
1	1. Face and neck	5–7 mg/dL	7–9 mg/dL
2	2. Chest and upper abdomen	7–9 mg/dL	9–11 mg/dL
3	3. Lower abdomen and thighs	9–11 mg/dL	11–13 mg/dL
4	4. Legs and arms/forearms	11–13 mg/dL	14–16 mg/dL
5	5. Palms and soles	13–15 mg/dL	17 mg/dL or more

©Marrow

Note:

Depth of jaundice (degree of yellowness) should be carefully noted (light staining as lemon-yellow; deep staining as orange-yellow), as it is an important indicator of the level of jaundice. Earlier, the involvement of soles and palms indicated severe jaundice but deep yellow staining (even in absence of yellow soles or palms) may be associated with severe jaundice.

Solution to Question 12:

Jaundice associated with a rise in bilirubin level at a rate of ≤ 5 mg/dL/24 hours suggests physiological jaundice.

Features suggestive of pathological jaundice:

- Jaundice appears within the first 24 hours after birth
- Jaundice present at or beyond age 3 weeks or sick infant
- Elevated direct (or conjugated) bilirubin level
- Total serum bilirubin levels above the expected normal range; jaundice seen in the mid-abdominal region (≈ 15 mg/dL) and soles (≈ 20 mg/dL)
- Total serum bilirubin concentration not responding to phototherapy
- Total serum bilirubin rising rapidly (i.e., crossing percentiles) in an infant receiving phototherapy

Solution to Question 13:

ABO incompatibility is the most common cause of hemolytic disease of newborns and affects the firstborn. It is usually mild and subclinical.

This incompatibility between the mother and fetus generally results in milder disease than Rh incompatibility does.

It arises when a mother with blood group type O becomes pregnant with a fetus of different blood type (A/B). The mother's serum contains naturally occurring anti-A and anti-B, which can cross the placenta and cause hemolysis of fetal RBCs.


Although antibodies against A and B factors occur without previous immunization (natural antibodies), they are usually IgM, which do not cross the placenta. However, IgG antibodies to A antigen may be present, and these do cross the placenta, resulting in A-O iso-immune hemolytic disease in the first-born infant (A-O is more common than B-O).

Solution to Question 14:

The most effective range of wavelength of light for phototherapy in neonatal indirect hyperbilirubinemia is 460–490 nm (blue-green light).

Clinical jaundice and indirect hyperbilirubinemia are reduced by exposure to the high intensity of light in the visible spectrum by 3 mechanisms:

- Irreversible structural isomerization: Most important mechanism; acts by forming lumirubin from native bilirubin. It is then excreted by the kidneys in the unconjugated state.
- Reversible photo-isomerization reaction: converts toxic unconjugated bilirubin into an unconjugated configurational isomer. It is then excreted in bile without conjugation.
- Photo-oxidation: minor method



Solution to Question 15:

Sn-mesoporphyrin (SnMP), a synthetic metalloporphyrin is a potential alternative medical therapy for pathological jaundice in neonates.

It acts by competitive inhibition of the conversion of heme protein to biliverdin by heme oxygenase. A single intramuscular dose on the 1st day of life reduces the need for subsequent phototherapy. Complication from metalloporphyrins includes transient erythema if the infant is receiving phototherapy.

Tolazoline is an alpha-adrenergic antagonist that produces systemic and pulmonary vasodilatation, which is used to correct pulmonary artery vasospasm when it causes severe right-left shunting soon after birth.

Tromethamine is an organic buffer used in the management of metabolic acidosis.

Solution to Question 16:

The bronze baby syndrome is a dark, grayish-brown skin discoloration in infants undergoing phototherapy; it mainly occurs in infants with direct hyperbilirubinemia as an adverse effect of

phototherapy.

Despite the bronze baby syndrome, phototherapy can continue if needed.

Other complications of Phototherapy:

- Loose stools
- Erythematous macular rash
- The purpuric rash associated with transient porphyrinemia
- Overheating
- Dehydration (increased insensible water loss, diarrhea)
- Hypothermia from exposure

Gray baby syndrome is an adverse effect of chloramphenicol.

Redman syndrome is an adverse effect of vancomycin.

Solution to Question 17:

The cut-off for the treatment of neonatal hypoglycemia to start IV glucose is a blood glucose level of <40 mg/dL.

Neonatal hypoglycemia can be suspected when the blood glucose is <45 mg/dL. Clinically in neonates, hypoglycemia can be asymptomatic or symptomatic. The neonate presents with tremors, cyanosis, convulsions, apneic spells, tachypnea, weak and high-pitched cry, lethargy, difficulty in feeding, episodes of sweating, sudden pallor, hypothermia, and rarely cardiac arrest.

Management in symptomatic neonates:

- Blood glucose <40 mg/dL: IV glucose infusion
- Blood glucose <25 mg/dL: Bolus IV dextrose 1-2 mL/kg followed by infusion at the rate of 5-8mL/kg/min.

Management in asymptomatic neonates:

- Breastfeeding within 30 minutes
- Start IV glucose therapy if blood glucose <35 mg/dL
- Recheck blood glucose before the next feeding
- Continue skin-to-skin care
- Evaluation for other underlying illnesses

The desired target glucose value is >50 mg/dL and blood glucose should be rechecked until three normal values.

Solution to Question 18:

In an infant of a diabetic mother with normal blood glucose levels, the next investigation to consider is serum calcium due to the high risk of developing hypocalcemia. Hypocalcemia in these infants is attributed to the delayed response of the parathyroid hormone system.

Hypocalcemia is defined as a total serum calcium concentration of less than 8 mg/dL for a term baby (ionized calcium below 4mg/dL) and less than 7 mg/dL for a preterm baby. Asymptomatic neonates can be managed conservatively with early nutrition. Symptomatic neonates should receive iv or oral calcium replacement.

Maternal hyperglycemia causes fetal hyperglycemia, and the fetal pancreatic response leads to fetal hyperinsulinemia, which causes neonatal hypoglycemia. Neonatal hypoglycemia is defined as blood glucose $\leq$ 40 mg/dL (plasma glucose $\leq$ 45 mg/dL). The infant tends to be jittery, tremulous, hyperexcitable, hypotonic, lethargic, and poor sucking during the 1st three days after birth. These symptoms can also be related to hypocalcemia and hypomagnesemia which occur in the 1st 24 to 72 hours of life.

Respiratory distress syndrome, macrosomia, hyperbilirubinemia, polycythemia, renal vein thrombosis, visceromegaly, heart failure, and persistent pulmonary hypertension are other features seen in infants of diabetic mothers.

Congenital anomalies are seen in infants of mothers with pregestational diabetes. Neural tube defects (Encephalocele, meningomyelocele, anencephaly), transposition of great vessels (TGA), ventricular septal defect (VSD), atrial septal defect (ASD), hypoplastic left heart, aortic stenosis and coarctation of the aorta are commonly seen.

Caudal regression syndrome is rare, but a specific anomaly seen in infants of diabetic mothers. Small left colon syndrome is a rare anomaly that develops in the second and third trimesters.

Solution to Question 19:

Infants born to diabetic mothers may present with macrosomia (large for gestational age), hypocalcemia, and/ or neural tube defects.

Other common features include:

- Hypoglycemia
- Respiratory distress syndrome
- Cardiomegaly
- Heart failure
- Lumbosacral agenesis
- Cardiac malformations
- Ventricular or atrial septal defect
- Transposition of the great vessels (most specific)
- Truncus arteriosus
- Double-outlet right ventricle
- Tricuspid atresia

- Coarctation of the aorta

Additional anomalies include neural tube defects, hydronephrosis, renal agenesis and dysplasia, duodenal or anorectal atresia, situs inversus, double ureter, and holoprosencephaly.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Diseases in Neonates requiring Special Care

Question 1:

A preterm neonate delivered one hour back in a PHC is rushed to the district hospital. There is an evisceration of bowel loops from a defect in the abdominal wall to the right of the umbilicus. What is the likely diagnosis?

- a) Omphalocele
- b) Umbilical hernia
- c) Gastroschisis
- d) Ectopia vesicae

Question 2:

While performing the head-to-toe examination in a term 2.6kg neonate, a large, wide anterior fontanelle is noted. Which of the following congenital anomalies is least likely?

- a) Achondroplasia
- b) Osteogenesis Imperfecta
- c) Trisomy 21
- d) Craniosynostosis

Question 3:

An anxious new mother brings her 18-day-old baby, to the casualty with complaints of poor feeding and yellowish discoloration of the skin. In the casualty, the baby had an episode of seizure. His total bilirubin is 32 mg/dl and he is diagnosed to have kernicterus. Which of the following will not be seen as a late feature of the disease?

- a) Defective upward gaze
- b) Sensorineural hearing loss
- c) Choreoathetosis
- d) Hypotonia


Question 4:

What is the most common cause of seizures in a term neonate?

- a) Hypoglycemia
- b) Sepsis
- c) Hypoxic ischemic encephalopathy
- d) Intraventricular hemorrhage

Question 5:

A 2-day-old neonate was admitted to the NICU after he was observed to have movements as shown in the clip. His APGAR scores were 8 and 9 at 1 and 5 minutes, respectively. His blood sugar was within the normal range in the first 24 hours. What is the likely diagnosis?



©MARROW

- a) Focal tonic seizure
- b) Focal clonic seizure
- c) Spasms
- d) Subtle seizure

Question 6:

A term neonate born to a lady with chorioamnionitis is diagnosed with neonatal sepsis. Which of the following statements are false?

- a) A, B, C and E
- b) B, C and E

- c) A, B and D
- d) C and E only

Question 7:

Blood cultures drawn from a late preterm baby on day 8 of life show the growth of streptococcus species within 18 hours. An umbilical line tip culture also shows the same bacteria. Which among the following is the most common manifestation of the described type of neonatal sepsis?

- a) Meningitis
- b) Pneumonia
- c) Disseminated intravascular coagulation
- d) Shock

Question 8:

A focus of Klebsiella is detected in one of the ventilators during a routine quality and maintenance check in the NICU. Sepsis protocol is enacted and a sepsis screen is ordered for all the neonates at risk. Which of the following investigations is not sent as part of a neonatal sepsis screen?

- a) Immature to total neutrophil ratio
- b) C-reactive protein
- c) Absolute eosinophil count
- d) Micro-erythrocyte sedimentation rate


Question 9:

Which of the following findings in a healthy term child would warrant further evaluation?

- a) Sub-conjunctival hemorrhage
- b) Buphthalmos
- c) Erythema toxicum
- d) Mongolian spots

Question 10:

A neonate undergoing routine ophthalmological examination was found to have the finding shown below. Which among the following can be ruled out as the cause of this finding?



- a) Congenital cataract
- b) Persistent hyperplastic primary vitreous
- c) Retinopathy of prematurity
- d) Congenital glaucoma

Question 11:

A 26-year-old primigravida delivers a 1.5 kg baby girl at 27 weeks of gestation. When is an ophthalmological examination recommended for the initial screening of retinopathy of prematurity (ROP) in this baby?

- a) At the time of birth
- b) At age equivalent to 32 weeks of gestation
- c) 4 weeks postnatal
- d) 2 weeks postnatal

Question 12:

An infant born 48 hours back by vaginal delivery with birth asphyxia develops seizures. What stage of hypoxic-ischemic encephalopathy is the child likely to be in?

- a) Stage 1

- b) Stage 2
- c) Stage 3
- d) Stage 4

Question 13:

A 27-year-old primigravida delivers a 3kg term baby after a cesarean section done in view of prolonged labor. The child cries 5 mins after birth and has a seizure 8 hours later. What is the imaging of choice for evaluation?

- a) Ultrasonography
- b) Computed tomography
- c) Amplitude-integrated electroencephalography
- d) Diffusion-weighted magnetic resonance imaging (MRI)

Question 14:

A 10-day old female neonate with hypoxic-ischemic encephalopathy is hyper-alert with mydriasis. Examination shows hyperactive tendon reflexes and clonus. On investigations, a normal EEG is obtained. What stage of HIE is she in?

- a) Stage 1
- b) Stage 2
- c) Stage 3
- d) Options A and B

Question 15:

A 1.4 Kg neonate is born at 29 weeks of gestation. The sonologist performing a neurosonogram is most likely to find an intracranial hemorrhage in the:

- a) Germinal matrix
- b) Corpus callosum
- c) Basal ganglia
- d) Choroid plexus

Question 16:

A 2-week old infant with a birth weight of 1.8 Kg is being treated in the NICU for necrotizing enterocolitis. Which of the following statements is false regarding this disease?

- a) Proximal part of the ileum is involved most frequently
- b) Pneumatosis intestinalis may be seen
- c) It is associated with intestinal ischemia
- d) It is more common in preterm infants


Question 17:

A 12-day old preterm neonate was diagnosed to have necrotizing enterocolitis. She had been switched over from breastmilk to formula feed 5 days after birth as the mother died of COVID-associated complications. Which of the following features will not be seen in the baby?

- a) Vomiting
- b) Abdominal mass
- c) Erythema of the abdominal wall
- d) Metabolic alkalosis

Question 18:

A 1.9 Kg neonate delivered at 32 weeks has abdominal distention and hematochezia on the 11th day of life. The abdomen x-ray is as shown below. What is the stage of disease according to Modified Bell's Staging Criteria?



- a) Stage IB
- b) Stage IIA
- c) Stage IIIA
- d) Stage IIIB

Answer Key

Question No.	Correct Option
1	c
2	d
3	d
4	c
5	b
6	b
7	a
8	c
9	b
10	d
11	d
12	b
13	d
14	a
15	a
16	a
17	d
18	b

Detailed Explanations



Solution to Question 1:

The most probable diagnosis for the evisceration of bowel loops, coming out of a defect in the abdominal wall to the right of the umbilicus is gastroschisis. It occurs when there is a failure in the closure of the body wall in the abdominal region.

The defects seen in the ventral body wall include:

- **Gastroschisis**- The intestinal loops herniate through the defect in the abdominal wall. The defect usually lies to one side of the umbilicus (right side commonly). The loops of the bowel are not covered by amnion as they herniate through the abdominal wall directly into the amniotic cavity. The affected bowel loops can be damaged by exposure to amniotic fluid.
- **Omphalocele**- The bowel loops fail to return from the physiologic umbilical hernia. Since the umbilicus is covered by a reflection of the amnion, the omphalocele is also covered by amnion.
- **Congenital umbilical hernia**- This is due to a failure of the umbilical ring to close after physiologic herniation of the intestines.

Ectopia Vesicae/ Bladder exstrophy is a congenital abnormality that occurs when the lower abdominal wall does not form properly and the bladder is open and exposed.



Solution to Question 2:

In craniosynostosis, infants have small fontanelles. Craniosynostosis is the premature fusion of cranial sutures

An infant has 2 fontanelles at birth, the diamond-shaped anterior fontanelle and the triangular posterior fontanelle.

- Closure of the anterior fontanelle occurs at 18 months
- Closure of the posterior fontanelle occurs by the 6th - 8th week after birth; its persistence suggests underlying hydrocephalus or congenital hypothyroidism.
- The presence of a third fontanelle may suggest trisomy 21 or prematurity.

The persistence of excessively large anterior fontanelles and small fontanelles has been associated with several disorders:

Large fontanelles	Small fontanelles
Achondroplasia	Microcephaly
Cleidocranial dysostosis	Craniosynostosis
Congenital rubella syndrome	Congenital hyperthyroidism
Hydrocephaly	Wormian bones
Osteogenesis imperfecta	
Prematurity	
Russell–Silver syndrome	
Trisomies 13, 18, and 21	
Vitamin D deficiency, rickets	

Solution to Question 3:

Hypotonia is seen in the early phase of kernicterus.

Kernicterus or bilirubin encephalopathy is a neurological syndrome resulting from the deposition of unconjugated (indirect) bilirubin in the basal ganglia and brainstem nuclei. It can result in permanent brain damage.

Early Phase (Acute)	Late Phase
Poor suck	Movement disorders- choreo athetosis, ballismus, tremor
Stupor	Sensorineural hearing loss
Hypotonia	Defective upward gaze
Seizures	

Solution to Question 4:

Hypoxic-ischemic encephalopathy (HIE) is the most common cause of neonatal seizures in term babies. Seizures due to HIE typically appear within the first 12-24 hours of birth. Subtle seizures are the most common type of seizures in neonates with HIE.

Neonatal seizures:

- Most common type - Subtle seizures
- Least common type - Myoclonic seizures
- Most common metabolic cause - Hypoglycemia
- Investigation of choice for the diagnosis of seizures - Electroencephalogram (EEG)
- Neuroimaging modality of choice for etiological evaluation - MRI
- Drug of choice - Phenobarbitone

Solution to Question 5:

The video clip of a 2-day-old neonate exhibiting repetitive, rhythmic movement in unilateral upper and lower limbs suggests a neonatal seizure of focal clonic type.

A focal clonic seizure is characterized by multiple, repetitive, rhythmic contractions of muscle groups of the limbs, face, or trunk, maybe unifocal or multifocal. It may occur synchronously or asynchronously in muscle groups on one side of the body. It may occur simultaneously but asynchronously on both sides. It cannot be suppressed by restraint.

Other options:

Option A: Focal tonic seizures are characterized by sustained posturing of a single limb, sustained asymmetrical posturing of the trunk, and sustained eye deviation. They cannot be provoked by stimulation or suppressed by restraint.

Option C: Spasms are sudden generalized jerks lasting 1-2 sec and are usually associated with a single, very brief, generalized discharge. They may be flexor, extensor, or mixed extensor/flexor and can occur in clusters. They too cannot be provoked by stimulation or suppressed by restraint.

Option D: Subtle seizures include transient eye deviations, nystagmus, blinking, mouthing, abnormal extremity movements (rowing, swimming, cycling, pedalling), fluctuations in heart rate, hypertension episodes, and apnea. They are more common in preterm than in term babies.

Myoclonic seizures are random, single, rapid muscle contractions of the limbs, face, or trunk. They are not repetitive or may recur at a slow rate. They can be generalized, focal, or fragmentary. They can be provoked by stimulation.

Clinically apparent seizures in a neonate should be treated if they last ≥ 3 minutes or are brief serial seizures. In all neonates with seizures, hypoglycemia, CNS infection, and hypocalcemia should be ruled out before starting anti-epileptic drug treatment.

Causes of neonatal seizure according to the age of presentation:

1 to 4 days	4 to 14 days	2 to 8 weeks
Hypoxic ischemic encephalopathy Maternal use of narcotics or barbiturates Drug toxicity: lidocaine, penicillin Intraventricular hemorrhage Hypocalcemia, sepsis, hypoglycemia, hypomagnesemia, hypo or hypernatremia Galactosemia, urea cycle disorders	Meningitis (bacterial) Encephalitis (enteroviral, herpes simplex) Hypocalcemia, hypoglycemia Galactosemia, fructosemia Hyperinsulinism, hyperammonemia syndrome	Herpes simplex or enteroviral encephalitis, bacterial meningitis Subdural hematoma, child abuse Aminoaciduria, urea cycle defects Malformations of cortical development Tuberous sclerosis Sturge-Weber syndrome

Solution to Question 6:

Late-onset sepsis usually presents after 72 hours of age (not 48 hours).

Neutropenia appears to be a better predictor of neonatal sepsis (not neutrophilia).

Neonatal sepsis is a clinical syndrome characterized by signs and symptoms of infection, with or without accompanying bacteremia in the first month of life.

Escherichia coli, Staphylococcus aureus, and Klebsiella spp. are the predominant organisms involved. Organisms such as Acinetobacter, Pseudomonas, and coagulase-negative staphylococci are also important pathogens in hospital-acquired infections.

	Early-onset sepsis	Late-onset sepsis
Presentation	Within the first 72 hours of life	After 72 hours of life
Source of infection	Maternal genital tract	Nosocomial or community-acquired
Manifestation	Respiratory distress and pneumonia	Septicemia, pneumonia, or meningitis

Solution to Question 7:

Bacterial growth in the umbilical tip and blood cultures on day 8 suggests a hospital-acquired infection after 72 hours. This is late-onset neonatal sepsis.

The most common manifestation of late-onset neonatal sepsis is meningitis.

	Early-onset sepsis	Late-onset sepsis
Presentation	Presents within first 72 hours of life	Presents after 72 hours of life
Source of infection	Maternal genital tract	Nosocomial or community-acquired

	Early-onset sepsis	Late-onset sepsis
Manifestation	Respiratory distress and pneumonia	Septicemia, pneumonia, meningitis

NOTE:

Traditionally, neonatal sepsis had been classified as EOS (<7 days) and LOS (>7days), but recently, experts have come to believe that Early-onset neonatal sepsis manifests within the first 72 hours of life and Late-onset neonatal sepsis after 72 hours of life.

Solution to Question 8:

Absolute eosinophil count is not a component of neonatal sepsis screen.

The various components of the septic screen include:

- Total leukocyte count (TLC)
- Absolute neutrophil count (ANC)
- Immature: Total neutrophil ratio (IT),
- Micro-erythrocyte sedimentation rate
- C-reactive protein.

Blood culture: Gold standard for diagnosis.

All neonates suspected to have sepsis should have a septic screen to corroborate the diagnosis.

Note:

Most common fungus causing neonatal sepsis: Candida species.

Solution to Question 9:

Buphthalmos is not a normal finding in a newborn; it is suggestive of congenital glaucoma and should, therefore, be investigated.

The sclera and cornea are more elastic in early childhood and an increase in intraocular pressure (IOP), therefore, leads to an expansion of the globe and the development of buphthalmos ("ox eye").

Recommended screening for eye diseases at birth:

- Buphthalmos
- Cataract
- Ophthalmia Neonatorum
- Developmental anomalies - Microphthalmos, Anophthalmos
- Nystagmus

- Squint
- Retinopathy of prematurity



Solution to Question 10:

The image shows leukokoria or white pupillary reflex. It is not seen in congenital glaucoma.

Leukokoria is seen in:

- Congenital cataract
- Chorioretinitis
- Retinoblastoma
- Retinopathy of prematurity
- Persistent hyperplastic primary vitreous.

Solution to Question 11:

The first Retinopathy of Prematurity (ROP) screening in this baby should be conducted 2 weeks postnatal.

Retinopathy of Prematurity (ROP) is a vaso-proliferative disorder of the developing retinal blood vessels of the preterm infant, which may lead to poor visual acuity or blindness.

Criteria for screening:

- Birth weight ≤ 2000 g OR
- Gestational age ≤ 34 weeks OR

- Gestational age between 34-36 weeks with the following risk factors - prolonged oxygen therapy (respiratory distress syndrome, apnea, cardiorespiratory support), neonatal sepsis, blood/exchange transfusion.

According to IAP guidelines, ROP screening should be conducted according to the table given below:

Period of gestation (POG) / birth weight	Screening
<28 weeks OR <1200g	2-3 weeks of postnatal age
>28 weeks OR >1200g	4 weeks of postnatal age

According to Nelson's 22nd edition, ROP screening should be conducted according to the table given below:

Annual follow-up is carried out unless indicated otherwise in cases where cryotherapy/laser treatment or cycloplegic refraction has been done.

POG	Screening
<28 weeks	31 weeks of POG
28-32 weeks	4 weeks of postnatal age

Solution to Question 12:

Seizures are common in stage 2 of hypoxic-ischemic encephalopathy (HIE).

HIE is an important cause of permanent damage to CNS tissues, resulting in the following:

- Neonatal Death
- Cerebral Palsy
- Developmental Delay

Stages	Seizures
Stage 1	Nil
Stage 2	Common
Stage 3	Decerebration

Solution to Question 13:

The clinical vignette of prolonged labor, delayed cry after birth (birth asphyxia), and seizure suggest hypoxic-ischemic injury. The preferred imaging modality in term neonates with HIE is diffusion-weighted magnetic resonance imaging (MRI). This is because of its increased sensitivity and specificity early in the process and its ability to outline the topography of the lesion.

Computed Tomography (CT) is helpful in identifying focal hemorrhagic lesions, diffuse cortical injury, and damage to the basal ganglia, but limited ability during the first few days of life.

Ultrasonography (USG) - Neurosonogram is the initial preferred modality for preterm infants.

Amplitude-integrated electroencephalography (aEEG) may help to determine which infants are at increased risk for long-term brain injury.

Solution to Question 14:

Hyperactive tendon reflexes may be seen in stages 1 and 2; however, the infant is hyperalert with mydriasis and normal EEG findings in stage 1.

Sarnat staging of hypoxic-ischemic encephalopathy (HIE):

Signs	Stage 1	Stage 2	Stage 3
Level of consciousness	Hyperalert	Lethargic	Stuporous, comatose
Muscle tone	Normal	Hypotonic	Flaccid
Posture	Normal	Flexion	Decerebrate
Tendon reflex/clonus	Hyperactive	Hyperactive	Absent
Myoclonus	Present	Present	Absent
Moro reflex	Strong	Weak	Absent
Pupils	Mydriasis	Miosis	Unequal, poor light reflex
Seizures	None	Common	Decerebration
EEG findings	Normal	Low voltage changing to seizure activity	Burst suppression to isoelectric
Duration	<24 hour if progresses; otherwise, may remain normal	24 hour to 14 days	Days to weeks
Outcome	Good	Variable	Death, severe deficits

Solution to Question 15:

Intraventricular hemorrhage (IVH) in premature infants occurs in the gelatinous subependymal or germinal matrix.

In preterm infants, the germinal matrix (situated ventrolateral to lateral ventricles) is extensive and highly vascular which makes them prone to bleeding. Germinal matrix normally involutes during the third trimester.

Major neuropathologic lesions associated with very-low-birth-weight (VLBW) infants are intraventricular hemorrhage (IVH) and periventricular leukomalacia (PVL).

Periventricular leukomalacia (PVL) is characterized by focal necrotic lesions in the periventricular white matter.

Solution to Question 16:

The distal ileum and the proximal colon are involved most frequently in necrotizing enterocolitis.

Necrotizing enterocolitis is a multifactorial disease primarily associated with intestinal immaturity characterized by mucosal or transmural necrosis of the intestine. It is more common in preterm infants, and prematurity is the biggest risk factor.

The triad of intestinal ischemia (injury), enteral nutrition (metabolic substrate), and bacterial translocation have classically been linked to necrotizing enterocolitis.

Pathologic findings include necrotic segment(s) of intestine, gas accumulation in the submucosa of the bowel wall (pneumatosis intestinalis), and progression of the necrosis to perforation, peritonitis, sepsis, and death.

Breast milk protects against necrotizing enterocolitis.

Solution to Question 17:

Necrotizing enterocolitis can cause metabolic or respiratory acidosis (not alkalosis).

Other Signs and Symptoms

Gastrointestinal	Systemic
Abdominal distention	Lethargy
Abdominal tenderness	Apnea/respiratory distress
Vomiting	Temperature instability
Occult/gross blood in stools,	Acidosis- metabolic and/or re spiratory
Change in stool pattern/diarr hea	Hyponatremia
Abdominal mass	Glucose instability
	Poor perfusion/shock
	DIC

Solution to Question 18:

The above X-ray image shows gas accumulation in the submucosa of the bowel wall- pneumatosis intestinalis, a pathognomonic finding of necrotizing enterocolitis (NEC), seen in stage IIA, according to the Modified Bell Staging Criteria.

Modified Bell's Staging Criteria:

Stage I - Suspected NEC

Stage II - Proven NEC

Stage III - Advanced NEC

Stage		Intestinal signs	Systemic signs	Radiological signs	Treatment
IA	Suspected NEC	Mild abdominal distention, emesis, guaiac-positive stool	Temperature instability, apnea, bradycardia, lethargy	Normal or intestinal dilatation, mild ileus	NPO, Antibiotic for 3 days
IB		IA + Bright red blood from the rectum	Same as IA	Same as IA	Same as IA
IIA	Mildly ill	Absent bowel sounds	Same as IA	Pneumatosis intestinalis, ileus	NPO, Antibiotics for 7-10 days
IIB	Moderately ill	IIA + definite abdominal tenderness	IIA + Metabolic acidosis and thrombocytopenia	IIA + portal venous gas, with or without ascites	NPO, Antibiotics for 14 days
IIIA	Severely ill, bowel intact	IIB + peritonitis with marked tenderness, and abdominal distension	IIB + hypotension, bradycardia, acidosis, DIC, neutropenia	IIB + definite ascitis	NPO, Antibiotics for 14 days, fluid resuscitation +/- inotropes, ventilator, paracentesis
IIIB	Severely ill, bowel perforated	Same as III A	Same as IIIA	IIB + pneumoperitoneum	Same as III A, surgery

Apgar score and Neonatal Resuscitation

Question 1:

While assessing the Apgar score of the newborns, you would assess and score all of the following except:

- a) Color
- b) Heart rate
- c) Respiratory rate
- d) Muscle tone


Question 2:

A baby assessed 5 minutes after birth, is found to be cyanosed with irregular gasping respiration. Heart rate is 60 beats/min, and shows minimal response to stimulation with some flexion of extremities. The Apgar score for this newborn is _____

- a) 2
- b) 5
- c) 3
- d) 4

Question 3:

Where would you attach the oxygen inlet in the instrument given below?



©Marrow

- a) 4
- b) 3
- c) 2
- d) 1

Question 4:

A 35-week neonate with grade 2 meconium-stained amniotic fluid is delivered by emergency LSCS. There is no spontaneous cry and the baby is unresponsive to tactile stimulation also. As the intern assisting in resuscitation, what is the order of suctioning that you will follow?

- a) Mouth-nose
- b) Nose-mouth
- c) Mouth-nose-oesophagus
- d) Trachea-nose-mouth

Question 5:

What is the ideal ratio of compression to ventilation that you would use in the resuscitation of a newborn?

- a) 30:2
- b) 5:1
- c) 3:1

d) 2:1

Question 6:

A neonate with suspected birth asphyxia is intubated in the labor room after bag and mask ventilation did not yield successful results. All of the following parameters would indicate successful ventilation except _____

- a) Zero reading of end-tidal CO₂ measurement
- b) Heart rate more than 100 beats/min
- c) Symmetric breath sounds
- d) Good chest rise

Question 7:

A panicked primigravida, at 38 weeks of gestation, rushes to the ER at 3 am as she had not appreciated any fetal movements since 10 pm. She is taken up for an emergency c-section in view of fetal distress. Following delivery, the infant was cyanotic and apneic. What is the preferred initial gas for neonatal resuscitation?

- a) 30% oxygen
- b) Hyperbaric oxygen
- c) 70% oxygen
- d) Room air

Question 8:

During neonatal resuscitation, when would you consider administering IV adrenaline?

- a) Heart rate $\leq$ 60 beats/minute despite effective compressions and ventilations.
- b) Heart rate $\leq$ 100 beats/minute despite effective compressions and ventilations.
- c) Heart rate does not improve after 30 seconds with bag and mask ventilation.
- d) Severe respiratory depression, and failure to respond to positive-pressure ventilation with bag and mask ventilation.

Question 9:

A 34-year-old lady is taken up for an emergency C-section due to meconium-stained amniotic fluid. The neonate does not cry after tactile stimulation and the heart rate is 60 beats/min. In order to determine the course of resuscitation, you assess if the baby is vigorous. For this, you would consider all of the following parameters except:

- a) Color
- b) Tone
- c) Respiratory effort
- d) Heart rate

Question 10:

A term primiparous lady with moderate mitral stenosis received 2 doses of meperidine, 2 hours before delivery. She delivers a female baby vaginally, but the baby is apneic at birth. What is the most important immediate management of the newborn in this case?

- a) Obtain a cord blood pH
- b) Perform bag-mask ventilation
- c) Administer naloxone in the endotracheal tube
- d) Begin chest compressions

Question 11:

Antenatal imaging in an infant delivered late preterm, showed severe congenital diaphragmatic hernia with tracheal occlusion. An airway is secured before the placenta is removed from the circulation. What is this procedure called?

- a) Bilevel airway positive pressure (BiPAP)
- b) Ex utero intrapartum treatment procedure (EXIT)
- c) Extracorporeal membrane oxygenation (ECMO)
- d) Continuous positive airway pressure (CPAP)

Question 12:

A newborn baby has respiratory distress in the delivery room. Attempts to pass a nasogastric tube through both the nares are unsuccessful. Genital and ear abnormalities are also noted on the preliminary examination. What is the likely diagnosis?

- a) Potter sequence

- b) VATER syndrome
- c) CHARGE syndrome
- d) Pierre-Robbin syndrome

Answer Key

Question No.	Correct Option
1	c
2	d
3	b
4	a
5	c
6	a
7	d
8	a
9	a
10	b
11	b
12	c

Detailed Explanations

Solution to Question 1:

Respiratory effort, not rate is a component of Apgar score.

The Apgar score is an objective method of evaluating the newborn's condition using the 5 parameters explained below. It is performed at 1 minute and 5 minutes after birth.

- Color (A, appearance)
- Heart rate (P, pulse)
- Response to a catheter in nostril (G, grimace)
- Muscle tone (A, activity)
- Respiratory effort (R)

Apgar score is not used to guide resuscitation, and resuscitation must be initiated before the 1-minute score. The change of score at sequential time points following birth reflects how well the

baby is responding to resuscitative efforts.

Solution to Question 2:

The Apgar score in this baby is 4. It is calculated as:

Appearance: blue - score 0

Pulse: <100 beats/min - score 1

Grimace: with minimal response to stimulation - score 1

Activity: some flexion of the extremities - score 1

Respiratory effort: irregular - score 1

The total score is 4.

Apgar score is not used to guide resuscitation, and resuscitation must be initiated before the 1-minute score. The change of score at sequential time points following birth reflects how well the baby is responding to resuscitative efforts.

Causes for low Apgar score are:

- Fetal distress
- Prematurity
- Drugs like analgesics, narcotics, sedatives, magnesium sulfate
- Acute cerebral trauma
- Precipitous delivery
- Airway obstruction (choanal atresia)
- Congenital pneumonia or sepsis
- Haemorrhage- hypovolemia

False-negative Apgar score is seen in maternal acidosis and high fetal catecholamine levels.

A total score of 10 indicates that the infant is in the best possible condition. An infant with a score of 0-3 requires immediate resuscitation. If the score at 5 minutes remains less than 7, the infant should be assessed again every 5 minutes for the next 20 minutes.



Solution to Question 3:

The above image shows a self-inflating bag and mask. It is used in neonatal resuscitation; it inflates automatically, without a compressed gas source.

- Marking 1 is the pressure release valve.
- Marking 2 is the 360-degree swivel.
- Marking 3 is the oxygen inlet.
- Marking 4 is the air inlet.

The parts of the self-inflating bag and mask are labeled in the image below:



©MARROW

Absolute contraindication for using the bag and mask ventilation is congenital diaphragmatic hernia

Solution to Question 4:

During neonatal resuscitation, the mouth is suctioned first and then the nares, in order to decrease the risk of aspiration. A 12 or 14 French suction catheter is used, with the suction pressure kept between 80-100 mmHg.


Routine suctioning of the trachea during neonatal resuscitation is not recommended. The suctioning of the esophagus and stomach is avoided if not indicated specifically. It may produce a vagal response - apnea and/or bradycardia due to stimulation of the nerve in the posterior pharynx.

Solution to Question 5:

The ratio of compressions to ventilation in the resuscitation of the newborn is 3:1 (90 compressions: 30 breaths in one minute).

A 3:1 compression-to-ventilation ratio is used for neonatal resuscitation where compromise of gas exchange is nearly always the primary cause of the cardiovascular collapse.

Chest compressions are indicated if the heart rate is below 60/min even after 30 seconds of positive pressure ventilation. Once the HR is 60/min or more, chest compressions should be discontinued.




Solution to Question 6:

End-tidal CO₂ should be present (EtCO₂ should not be zero). This is an important sign of successful ventilation.

Successful ventilation is signified by the following:

- Adequate chest rise
- Symmetric breath sounds
- Improved pink color
- Heart rate $\geq$ 100 beats/min
- Spontaneous respirations
- Presence of end-tidal CO₂ (EtCO₂ should not be zero)
- Improved tone




Solution to Question 7:

Room air is the preferred initial gas for neonatal resuscitation in term infants.

Resuscitation with room air in term infants is as effective as using 100% oxygen. By using room air, we may reduce the risk of hyperoxia which is associated with decreased cerebral blood flow and generation of oxygen free radicals.

If the neonate does not achieve normal oxygen saturation levels within 90 seconds, increased concentrations of oxygen are administered until normal saturation levels are achieved.


If pulmonary hypertension is suspected (meconium aspiration, diaphragmatic hernia), one may consider 100% oxygen as the initial gas for resuscitation. Resuscitation of most preterm infants can be initiated with a slightly higher fraction of inspired oxygen (21–30%).



Solution to Question 8:

If the heart rate remains at ≤ 60 beats/minute despite effective compressions and ventilation, consider administering epinephrine in neonatal resuscitation.

Note: Umbilical venous catheter is the preferred method of obtaining emergency vascular access in the delivery room.



Solution to Question 9:

Color is not a parameter considered to determine if the child is vigorous.

Vigorous babies have (HRT):

- Heart rate of ≥ 100 ,
- Respiratory effort: good
- Tone: good

If the infant is vigorous with good respiratory effort and a heart rate ≥ 100 beats/min; the mouth and nose may be suctioned with a bulb or suction catheter if needed. A newborn is defined as non-vigorous if any of the above three signs are absent.

Resuscitation of non-vigorous newborns with meconium-stained amniotic fluid includes positive-pressure ventilation and in severe cases, endotracheal intubation and chest compressions.

Note: Even for non-vigorous infants, endotracheal intubation to aspirate meconium is not recommended anymore.

Solution to Question 10:

The most important immediate management is to maintain adequate ventilation via bag and mask, because the baby is apneic.

Meperidine is a narcotic and the administration of narcotics such as opioids or barbiturates, to the mother shortly before delivery (within 4 hours prior) result in narcosis and respiratory depression in the newborn.

In these babies, naloxone hydrochloride is given while adequate ventilation (bag and mask) is maintained. Breathing in the depressed infant should be maintained until a response to naloxone is noted.

Naloxone hydrochloride, 0.1 mg/kg, can be given intravenously or intramuscularly.

Solution to Question 11:

Ex-utero intrapartum treatment procedure (EXIT) is a procedure by which the airway is secured before the infant is separated from the placenta.

It allows time to secure the airway in an infant known to have airway obstruction due to various causes, including laryngeal atresia or stenosis, teratomas, hygromas, and oral tumors before the infant is separated from the placenta. Uteroplacental gas exchange is maintained throughout the procedure.

Option A: BiPAP machines provide positive airway pressure during exhalation and additional positive pressure during inspiration. It allows for adjusting of inspiratory and expiratory pressures. The additional pressures augment tidal volume and improve alveolar ventilation. Children with neuromuscular weakness and intrathoracic airway obstruction benefit from this.

Option C: Extracorporeal membrane oxygenation (ECMO) is a form of cardiopulmonary bypass that augments systemic perfusion and provides gas exchange. This may be helpful to prevent and treat hypoxia and respiratory acidosis and decrease the work of breathing in ARDS, central apnea, refractory shock, severe sepsis, pulmonary hypertension, etc.

Option D: CPAP is provided via a high-flow nasal cannula, with a gas flow of 4-16 L/min. It is used in diseases with mildly decreased lung compliance, and low functional residual capacity.

Solution to Question 12:

Inability to pass the nasogastric tube through both the nares along with the presence of respiratory distress, suggests choanal atresia. The presence of ear and genital anomalies along with choanal atresia suggests the diagnosis of CHARGE syndrome.

It consists of

- Coloboma of the eye
- Heart anomaly
- Choanal Atresia
- Retardation
- Genital anomalies
- Ear anomalies

VATER syndrome: Vertebral defects, imperforate anus, tracheoesophageal fistula, as well as radial and renal dysplasia.

Potter's sequence: Renal agenesis, oligohydramnios, and pulmonary hypoplasia

Pierre–Robin syndrome: Micrognathia, retrognathia, and glossoptosis.

Developmental Milestones

Question 1:

Which of the following gross motor milestone would you expect to see in a 3-year-old child?

- a) Skips without falling
- b) Hop on 1 foot
- c) Walk upstairs with alternate foot
- d) Walk downstairs with alternate foot

Question 2:

You notice a child pulling to the standing position and then shuffling around by holding onto furniture. At what age is this milestone typically attained?

- a) 8 months
- b) 10 months
- c) 13 months
- d) 15 months


Question 3:

In the pediatric wards, you notice a baby, totally occupied with his little game and a mother trying hard to spoonfeed him. He is busy stacking toy cubes on top of each other and starts gleefully clapping once he makes a tower of 6. What is the likely age that this milestone corresponds to?

- a) 15 months
- b) 24 months
- c) 18 months
- d) 36 months

Question 4:

By what age is the milestone shown in the image typically attained?



- a) 4 months
- b) 6 months
- c) 9 months
- d) 12 months

Question 5:

A final-year medical student is examining a baby in the ward. As the infant is pulled to sit, he notices a head lag. By what age should this infant ideally start to hold his head steady without any bobbing motion?

- a) 2 months
- b) 4 months
- c) 5 months
- d) 6 months

Question 6:

Your sister tells you that your nephew has now started using his crayons. He easily copies circles, but still has trouble with squares. Other than a minor ASD noted incidentally at birth, he is a perfectly healthy child. Which of the following would you expect him to be able to do at this stage?

- a) Tell a story
- b) Skip

- c) Ask meaning of words
- d) Know his name and gender

Question 7:

Match the following milestones with corresponding age.

- a) A-1, B-4, C-5, D-3, E-2
- b) A-2, B-1, C-4, D-3, E-5
- c) A-2, B-1, C-3, D-5, E-4
- d) A-1, B-3, C-4, D-2, E-5

Question 8:

You are taking a detailed history from the parents of a child with a suspected autism spectrum disorder. Parents are worried that their child does not smile or maintain eye contact and is not interested in toys. At what ages should a typical child exhibit "social smile" and "stranger anxiety"?

- a) 2 months and 6 months respectively
- b) 1 month and 6 months respectively
- c) 6 months and 12 months respectively
- d) 6 months and 9 months respectively

Question 9:

A pediatrician is evaluating a toddler for mental retardation. She hands a bright red ball to the baby and lets him play with it. She then takes the ball and hides it under a blanket in front of him. The child confusedly stares at the blanket and cries instead of searching for it. What is being tested here?

- a) Separation anxiety
- b) Transitional object space
- c) Symbolic transformation
- d) Object permanence

Question 10:

Parents of 9-month-old infant, with a history of preterm delivery and prolonged NICU stay, are concerned that child is lagging behind in his milestones. You would suspect a developmental delay in this child if his developmental quotient (DQ) is below _____ %?

- a) 80
- b) 77
- c) 70
- d) 65

Question 11:

Which of the following milestones is expected to be seen in a child at 5 years of age?

- a) Dressing and undressing himself
- b) Hopping and telling stories
- c) Riding a tricycle and making a tower of 9 blocks
- d) Establishment of handedness and parallel play

Question 12:

While you are evaluating a baby, you show him a bright pink teddy bear, that he reaches out to with both hands. What is the earliest age by which this milestone is typically achieved?

- a) 4 months
- b) 5 months
- c) 6 months
- d) 7 months

Question 13:

Which of the following is not a definitive test for developmental surveillance?

- a) Bayley-II
- b) Stanford-Binet
- c) Denver-II
- d) Wechsler

Question 14:

A child was able to count 4 pennies but was unable to name 4 colors. The probable age of the child is _____.

- a) 3 years
- b) 4 years
- c) 5 years
- d) 6 years

Question 15:

A term, 4-month-old infant, is developing normally. By what age do you expect her to walk without support and use polysyllabic jargon?

- a) 9 months
- b) 2 years
- c) 6 months
- d) 15 months

Question 16:

One of the kids has scribbled all over a newly painted wall inside an Anganwadi. You notice there are some triangles and squares as well, in this "drawing". This child must be older than _____

- a) 3 years
- b) 5 years
- c) 6 years
- d) 4 years

Question 17:

The absence of which of the following milestones in a 3-year-old child is called as developmental delay?

- a) Hopping on one leg
- b) Drawing a square

- c) Feeding by spoon
- d) Passing a ball to someone

Question 18:

After assessing a young child treated for unilateral retinoblastoma, you note significant delays in the cognitive and fine motor domains. The baby can stand without support and creep well. Normally, the highest fine motor skill a baby of this age is expected to have is _____

- a) Scribble with crayons
- b) Unidextrous reach and transfer objects
- c) Immature pincer grasp
- d) Mature pincer grasp

Question 19:

You are a pediatrics resident handling the well-child clinic for the week and you are teaching undergraduates how to do a simple developmental assessment. A preterm baby born at 32 weeks, is brought in for a follow-up visit at 6 months. What is the age you would ask the students to assume when undertaking developmental assessment in this child?

- a) 24 weeks
- b) 20 weeks
- c) 16 weeks
- d) 19 weeks

Question 20:

All of the following statements are true regarding principles of development except?

- a) It is dependent on the functional maturation of the nervous system
- b) Specific responses replace mass activity as the child matures
- c) It progresses in cephalocaudal direction
- d) In the extremities, development occurs distal to proximal

Question 21:

A young infant is brought to you with a suspected developmental delay. The child is intently looking at his own hands. You would consider referring this child to a specialist if this milestone persists after _____.

- a) 12 weeks
- b) 16 weeks
- c) 20 weeks
- d) 24 weeks

Answer Key

Question No.	Correct Option
1	c
2	b
3	b
4	d
5	b
6	d
7	b
8	a
9	d
10	c
11	a
12	a
13	c
14	b
15	d
16	b
17	c
18	d
19	c
20	d
21	c

Detailed Explanations

Solution to Question 1:

A three-year-old should normally be able to walk upstairs with alternate feet.

Solution to Question 2:

Cruising is an activity in which a child pulls to the standing position and then shuffles around holding onto a piece of furniture. Cruising is attained by 10-11 months.

Solution to Question 3:

A child can usually make a tower of 6 blocks by 24 months.

A child makes a tower of:

- 2 blocks by 15 months
- 3 blocks by 18 months
- 6 blocks by 2 years
- 9 blocks by 3 years


Solution to Question 4:

The milestone shown in the image is a mature pincer grasp, which is seen at the age of 12 months. It is identified as a mature pincer grasp as the object is secured between the opposed thumb and the tip of the index finger.

Developmental progression of the grasp:

- At 6 months, the infant reaches and holds a larger object in a crude manner using the ulnar aspect of his hand (Ulnar palmar grasp). At 8-9 months, the object is grasped by the radial side of the palm (radial palmar grasp).
- At 9 months, the immature pincer grasp develops. The object is secured between the adducted thumb and the extended index finger.
- At 12 months, the mature pincer grasp develops. The object is secured between the opposed thumb and the tip of the index finger.

Immature pincer grasp (9 months)



©MARROW

Mature pincer grasp (12 months)



©MARROW

Solution to Question 5:

An infant can hold his head steady when pulled to sit, without head bobbing, by the age of 4 months.

Regarding head steadiness:

- Up till 2 months of age, there is a significant head lag when the infant is pulled to sit.

Head Lag in Infants Upto 2 Months of Age



©MARROW

- At 3 months, head lag is partially compensated but still present; the infant attains early head control with bobbing motion and sits with back rounded.
- At 4 months, there is no head lag when pulled to sit and the infant sits with full truncal support.

Solution to Question 6:

Normally, a child is able to copy a circle at 3 years and a square at 4 years. This child is a 3-year-old, and he should be able to know his name and gender.

- A child is able to tell a story at 4 years.
- Asks meaning of words at 5 years.
- A child can skip at 5 years.

Solution to Question 7:

Handedness is established during the 3rd year (24 to 36 months).

Taste perception is present since birth.

Monosyllabic babble is seen at 6 months.

Response to faces and voices with a smile is seen at 1.5 months.

The child creeps upstairs by 15 months.

Solution to Question 8:

Social smile is typically seen at 2 months and stranger anxiety at 6 months.

Social smile: At 1 month of age, a child smiles involuntarily. At 2 months of age social smile develops; that is, the child smiles voluntarily and reciprocally, in response to someone else's smile.

Stranger anxiety: Before 6 months, the child is sociable and friendly even with strangers. At 6 months, there is an onset of stranger anxiety; infants look back and forth between an approaching stranger and parent and may cling or cry anxiously.

Solution to Question 9:

Understanding that objects continue to exist even when not seen is called object permanence. This is due to an ability to form a mental representation of the object. It is attained at the age of 8–9 months.

Option A: Separation anxiety begins at about 9 months of age. Children wake up crying from their naps as they realize that their parents are not in the same room as them. This is different from stranger anxiety displayed by the child at the age of 6-7 months, wherein they react to approaching strangers by crying or clinging to their parent.

Option B: A transitional object is an object, such as a blanket or a stuffed toy, that is used by many children as a symbol of the absent parent. It is seen at around 18 months.

Option C: Symbolic transformations in play:

- At 12 months - make-believe (symbolic) play centres on the child's own body, e.g., the child pretends to drink from an empty cup.
- At 18 month - symbolic transformations in play are not limited to the child's body, e.g., a doll can be fed from a plate.

Solution to Question 10:

A developmental delay should be suspected if the development quotient (DQ) is below 70%.

It is calculated as:

$$DQ = \frac{\text{Developmental age} \times 100}{\text{Chronological age}}$$

From the highest milestones achieved by the child, the developmental age in a particular domain is calculated. The chronological age is the actual age in term babies. However, in preterm infants, a corrected age rather than postnatal age is used till two years of age. This is used to account for the small gap in neurological development that premature children have yet to catch up. For example, a child born at 32 weeks gestation (gestational age) seen at 12 weeks of age (postnatal age), should be considered as a 4-week-old (corrected age) child.

If there is a delay ($DQ < 70\%$) in more than 2 domains, then global developmental delay (GDD) is suspected.

Solution to Question 11:

Developmental milestones at the age of 5 years include:

- Motor - skips
- Adaptive - Draws a triangle
- Language - Asks meaning of words
- Social - Dresses and undresses

Undressing is easier and hence learned before dressing.

- 2 years - Helps to undress
- 3 years - Helps in dressing (if helped with buttons)
- 5 years - Dress and undress fully (without any help)

Solution to Question 12:

The earliest age of appearance of bidextrous reach is 4 months.

Fine motor reflexes begin to appear only after the palmar grasp reflex disappears at 3-4 months. The development of unidextrous reach occurs by 6 months of age.

Solution to Question 13:

Denver-II is a developmental screening test (not a definitive test) that assesses child development in 4 domains:

- Gross motor
- Fine motor adaptive
- Language and
- Personal social behaviour

Tests for the developmental quotient

Definitive Tests	Screening tests
Bayley Scales of Infant and Toddler development (Bayley-I) Stanford-Binet Intelligence Scale Wechsler Intelligence Scale Developmental Activities Screening Inventory (DASI-II) Vineland Adaptive Behavior Scale-II	Denver-II Developmental screening test Phatak's Baroda Screening test Trivandrum Development Screening Chart Cognitive Adaptive Test/Clinical Linguistic and Auditory Milestone Scale (CAT/CLAMS) Gothenburg Harris Draw-a-man Test Gesell figures and block skills

Solution to Question 14:

A 4-year-old child can count 4 pennies but cannot name 4 colours. A child can name 4 colours at 5 years of age.

Note:

An easy way to remember this:

- No. of objects counted = age
- No. of colours identified = age - 1

Solution to Question 15:

A 15-month-old child can walk without support and use polysyllabic jargon.

Walking: A child begins to cruise at 10 months, walk with support at 12 months, and walk without support at 15 months.

Jargon is seen around 15 months of age. It is a pre-linguistic vocalization in which the child babbles using strings of word-like sounds and produces words that do not make much sense to others.

Solution to Question 16:

The milestones at 5 years of age are:

- Motor - skips
- Adaptive - draws triangle
- Language - asks meaning of words, names 4 colors
- Social - dresses and undresses

Solution to Question 17:

Feeding by spoon is a social milestone that typically appears by 18 months. If this is absent, then the child should be termed as having a developmental delay. Hopping on one leg, drawing a square, and passing a ball to someone appear by 48 months of age.

Red flag signs in developmental screening and surveillance are indicators that suggest a serious disorder in a child's development, requiring referral to a pediatrician. These are given below:

Positive indicators: Presence of any of the following

- Loss of developmental skills at any age
- Concerns about vision or confirmed visual impairment

- Hearing loss at any age
- Persistently low muscle tone
- Lack of speech by 18 months without other means of communication
- Asymmetry of movements or features suggestive of cerebral palsy
- Persistent toe walking
- Complex disabilities
- Unusual head circumference compared to norms
- Clinician uncertainty about the developmental disorder.

Negative Indicators:

- Inability to sit unsupported by 12 months
- Inability to walk by 18 months in boys or 2 years in girls.
- Inability to walk normally
- Inability to run by 2.5 years
- Inability to hold objects in hand by 5 months or reach for them by 6 months.
- Inability to point at objects to share interest by 2 years

Solution to Question 18:

A baby that stands without support is most likely to be 12 months. The highest fine motor skill she is expected to have is mature pincer grasp (12 months).

Bidextrous reach is seen at 4 months and unidextrous reach at 6 months. Immature pincer grasp is seen at 9 months. Child begins to scribble after 15 months.

Fine motor development is critically linked with hand-eye coordination. Children who have disease-related loss of vision in early infancy develop difficulties in visuomotor integration and lags behind in fine motor milestones.

Solution to Question 19:

The corrected age (in weeks) to be assumed while undertaking developmental assessment is 16 weeks.

Corrected age (in weeks) = (Gestational age + postnatal age) - 40 weeks

In this case, since the baby is born 8 weeks preterm, the 8 weeks of intrauterine growth and development she missed, is accounted for post-natally. Thus, for assessing the correct developmental age, 8 weeks has to be subtracted from the chronological age.

In babies born preterm, corrected age is used for their developmental assessment until 2 years of age.

Solution to Question 20:

In the extremities, development occurs proximal to distal.

Rules of development:

- It is a continuous process
- It depends on the functional maturation of the nervous system
- The sequence of attainment of milestones is the same in all children
- It progresses in a cephalocaudal direction and proximal to distal.
- Certain primitive reflexes have to be lost before relevant milestones are attained
- The initial disorganized mass activity is gradually replaced by specific actions.

Growth	Development
Denotes a net increase in size /mass of tissues	Denotes maturation of functions
Cephalocaudal Head $\Rightarrow$ Neck $\Rightarrow$ Trunk & upper limbs $\Rightarrow$ Lower limbs	
Distal $\Rightarrow$ Proximal Hands increase in size before upper arms	Proximal $\Rightarrow$ Distal Hand use is learnt before the use of fingers

Solution to Question 21:

Between 12 and 20 weeks, the child observes his own hands very intently. This milestone is called hand regard and it should disappear by 20 weeks. Its persistence after 20 weeks is considered abnormal.

In a blind child, however, it may persist beyond 20 weeks normally and is called apparent hand regard.

Facets of Growth and Development

Question 1:

Which of the following fetal hormones does not play any role in fetal growth?

- a) Insulin
- b) Glucocorticoid
- c) Thyroxine
- d) Growth hormone

Question 2:

A child is brought to the pediatrician for vaccination. The child is able to speak in 2-3 word sentences and is able to make a tower of 6 cubes. What is the expected weight of the child, if his development is normal and birth weight was 3 Kgs?

- a) 12 kgs
- b) 4 kgs
- c) 8 kgs
- d) 10 kgs

Question 3:

Which of the following statements is false?

- a) Head circumference should be measured within 24 hours after birth
- b) Head circumference equals chest circumference at 1 year of age
- c) A boys height at 2 and half years is half of his adult height
- d) A girls height at 2 years is half of her adult height

Question 4:

Which of the following statements regarding the eruption of teeth is false?

- a) 1 and 3
- b) 2 and 4

- c) 2, 4 and 5
- d) 5 only

Question 5:

A 3-year-old boy presented to the clinic for a missed vaccination. His routine anthropometric measurements were noted. He was found to be 90 cm tall with a lower segment of 45 cm. Which among the following is the most likely diagnosis in this child?

- a) Congenital hypothyroidism
- b) Achondroplasia
- c) Rickets
- d) Spondyloepiphyseal dysplasia

Question 6:

A team participating in an Under-14 national athletics sports tournament suspects foul play by a rival team member. He claims to fit the criteria for the age requirements for the tournament, but the opposing team does not agree. Which of the following X-ray would be most useful to verify the claims of the player?

- a) X-ray of the hip
- b) X-ray of the hands and wrist
- c) X-ray of the shoulder
- d) X-ray of the elbow and hip

Question 7:

Which of the following statements is true?

- a) Head reaches 90% of adult size at 1 year
- b) Lymphoid growth follows a sigmoid-shaped pattern
- c) Perinatal period is between 28 weeks of gestation to 7 days after birth
- d) Growth of child is cephalocaudal and distal to proximal


Question 8:

A young boy is brought to the clinician by his parents with complaints of impulsive behavior and decreased levels of attention at school and everyday activities. The clinician is suspecting a case of ADHD. Which of the following is not included in the criteria for diagnosis of ADHD?

- a) Symptoms begin before 7 years of age
- b) Symptoms should be present for at least 6 months
- c) Should be reported by independent observers
- d) Behaviour must be developmentally inappropriate in two or more settings

Question 9:

In the image given below, the weight and height of a child are plotted against his age in years. Identify the correct statement.



- a) WHO growth charts used now are developed by National Center for Health Statistics (NCHS).
- b) Short stature is defined as a height below 5th percentile.
- c) Puberty in this child will be normal.
- d) Bone age of this child is less than chronological age.

Question 10:

Identify the correct statement from the following.

- a) Children with pica are at increased risk of megaloblastic anemia
- b) Breath-holding spells are associated with hypertonemia

- c) Thumbsucking in a child is a way of discharging tension
- d) Bedwetting after a period of 3 months of dryness at night is called secondary enuresis

Question 11:

As part of the annual school health check-up camp, the visual acuity of a 6-year-old girl is being tested by the optometrist. What is the expected visual acuity in this age?

- a) 20/30
- b) 20/40
- c) 20/20
- d) 20/50

Question 12:

A 3-year-old boy was brought to the clinic by his mother who complains that for the past few months he has developed body-rocking movements and repeats words he hears on the television. He doesn't respond immediately on calling his name and also makes limited eye contact during interactions. Based on clinical examination you suspect autism in the child. Which of the following screening tools can be used to detect the boy's condition?

- a) Conner rating scale
- b) MCHAT
- c) CRAFFT
- d) EPDS

Question 13:

A 17-year-old girl presented with primary amenorrhoea. On physical examination, her breast and areola are enlarged without separation of their contours. What is the sexual maturity rating in this girl?

- a) SMR stage 1
- b) SMR stage 2
- c) SMR stage 3
- d) SMR stage 4

Question 14:

A 14-year-old boy is concerned about the change in colour of his scrotum since the last few weeks. On examination, the boy has a dark scrotum with normal-sized testis, with coarse and curly pubic hair. What is the sexual maturity rating of this boy?

- a) Stage 2
- b) Stage 4
- c) Stage 3
- d) Stage 5

Question 15:

Identify the correct statement.

- a) Macrocephaly is defined as an occipitofrontal circumference greater than 3 SD above the mean for age and sex
- b) Monosomy 1p36 is associated with macrocephaly
- c) Microcephaly is defined as an occipitofrontal circumference more than 2 SD below the mean for age and sex
- d) Infants with benign familial megalencephaly have normal head circumference at birth

Question 16:

A mother brings her baby of 14 months to the clinic after being concerned about the shape of the baby's head. You are suspecting a case of craniosynostosis. The premature fusion of which of the following suture is seen in this child?



- a) Metopic suture
- b) Both coronal sutures
- c) Lambdoid suture
- d) Sagittal suture

Question 17:

A 7-year-old boy was found to have a height for age 2 standard deviations below the median. Furthermore, his bone age was found to be equal to chronological age. Which of the following is most likely?

- a) Familial short stature
- b) Constitutional delay
- c) Undernutrition
- d) Growth hormone deficiency

Answer Key

Question No.	Correct Option
1	d
2	a
3	a

4	b
5	d
6	d
7	d
8	a
9	d
10	c
11	c
12	b
13	c
14	b
15	c
16	c
17	a

Detailed Explanations

Solution to Question 1:

Growth hormone does not play any role in fetal growth, although it is present in high levels during the fetal period. However, it does have a role in the postnatal period.

Insulin and thyroxine hormones are required for normal growth and development during late gestation. Glucocorticoid has a role in the prepartum maturation of organs such as liver, lungs, and gastrointestinal tract.

Solution to Question 2:

The child is able to make a tower of 6 cubes and speaks in 2-3 word sentences which indicates the child is of 2 years of age. Birth weight quadruples by the age of 2 years. Hence the child's weight should be $3 \text{ kgs} \times 4 = 12 \text{ Kgs}$.

Infants may lose up to 10% of their birth weight in the 1st week of life and regain it by their 2nd week.

Solution to Question 3:

Head circumference should not be measured within 24 hours because the head is subjected to molding during delivery, which makes the measurements unreliable.

The maximum circumference of the head from the occipital protuberance to the supraorbital ridges on the forehead is recorded using the cross-tape method.

- Head circumference increases by the following:
- 2 cm/month in first 3 months
- 1 cm/month in next 3 months
- 0.5 cm/month in next 6 months

Head circumference is more than chest circumference by 2.5 cm at birth.

- At birth: Head circumference > Chest circumference
- At 9-12 months: Head circumference = Chest circumference
- After 1 year: Head circumference < Chest circumference

Age	Head circumference
At birth	33 to 35cm
3 m	40 cm
6 m	43 cm
1 year	46 cm
2 years	48 cm
3 years	49 cm
4 years	50 cm
12 years	52 cm (adult size)

Solution to Question 4:

Primary teeth in the upper jaw erupt earlier than their counterpart in the lower jaw, except for lower central incisors and second molar. No primary teeth till the age of 13 months is called delayed eruption.

As seen in the image below, the first primary tooth to erupt is the lower (mandibular) central incisor, at 5-7 months. All primary teeth erupt by 1.5–2.5 years of age.



The first secondary tooth to erupt is the first molar (maxillary or mandibular), at 6-7 years, followed by the lower central incisor.

Causes of delayed eruption:

- Idiopathic: Most common
- Hypothyroidism
- Hypoparathyroidism
- Familial
- Mechanical blockage of individual teeth


Solution to Question 5:

The upper segment to lower segment (U/L) ratio in this child is 1. The U/L ratio is low for his age which makes spondyloepiphyseal dysplasia most likely among the given options.

The lower segment is defined as the length from the symphysis pubis to the floor. The upper segment is measured by subtracting the lower segment from the height.

The U/L ratio is

- 1.7 at birth
- 1.3 at 3 years
- 1 after 7 years
- 0.9 in adults



High U/L ratio:

- Rickets
- Achondroplasia
- Untreated congenital hypothyroidism

Low U/L ratio:

- Spondyloepiphyseal dysplasia
- Vertebral anomalies

Arm span:

- Length between the tips of the middle finger of the outstretched hands is the arm span.
- It is shorter than the length by 2.5 cm at birth; it equals height at 11 y and thereafter is slightly greater than height (usually ± 1 cm).

Solution to Question 6:

The X-ray of choice to determine the skeletal age of a 14-year-old boy is X-ray of the elbow and hip.

The bone age of a child indicates his/her level of biological and structural maturity. Methods used for bone age estimation are Gruelisch–Pyle atlas (more common) and Tanner–Whitehouse method.

Note: There is overlap in the age ranges for X-ray of choice.

Age	X-ray of choice
3 to 9 months	X-ray shoulder

Age	X-ray of choice
1 to 13 years	Single film of X-ray hands and wrist
12 to 14 years	X-ray elbow and hips


Solution to Question 7:

Growth of child is cephalocaudal and distal to proximal.

During fetal life, the growth of the head occurs before that of the neck, and arms grow before legs. The distal parts of the body such as hands increase in size before the upper arms.

Growth rate of different tissues:

Growth	Development
Denotes a net increase in size /mass of tissues	Denotes maturation of functions
Cephalocaudal Head ⇒ Neck ⇒ Trunk & upper limbs ⇒ Lower limbs	
Distal ⇒ Proximal Hands increase in size before upper arms	Proximal ⇒ Distal Hand use is learnt before the use of fingers



- Brain growth: At age of 1 month, the head size is about 36% of the expected head size in adults. It doubles in the first year of life to 72% and becomes 83% of adult size at 2 years.
- Lymphoid growth: It is prominent in mid-childhood. It peaks at 4–8 years and regresses subsequently.

- Gonadal growth: It stays dormant during childhood and shows a rise with the onset of puberty.
- Somatic growth: General body growth shows an S-shaped curve. It is rapid during the fetal period and the first 3–4 years. It is steady in the following period, followed by a growth spurt during puberty. Later, it slows down considerably.

Note: WHO defines the perinatal period from 22 weeks of gestation (or ≥ 500 g) up to 7 days after birth.

Solution to Question 8:

Attention-deficit hyperactive disorder (ADHD) must begin before 12 years of age, according to DSM-5 criteria. It was 7 years as per the previous DSM-4 criteria.

Treatment: Presynaptic dopaminergic agonists, commonly called psychostimulant medications, including methylphenidate, dexamethylphenidate, amphetamine, and various amphetamine and dextroamphetamine preparations are used.

DSM-5 Diagnostic Criteria for attention-deficit/Hyperactivity Disorder (ADHD)-

1. A persistent pattern of inattention and/or hyperactivity/impulsivity present for at least 6 months, that interferes with functioning or development.
2. Several inattentive or hyperactive/impulsive symptoms were present before 12 years of age.
3. Several inattentive or hyperactive/impulsive symptoms are present in 2 or more settings (e.g., at school [or work] or at home) and is documented independently.
4. There is clear evidence of clinically significant impairment in social, academic, or occupational functioning.
5. Symptoms do not occur exclusively during the course of schizophrenia, or another psychotic disorder, and are not better accounted for by another mental disorder (e.g., mood disorder, anxiety disorder, dissociative disorder, personality disorder, substance intoxication or withdrawal).

Solution to Question 9:

Bone age of this child is less than chronological age, as seen in cases of constitutional delay.

As seen in the graph, the child was short initially, but the final height is within normal range. Hence, this child has constitutional delay.

Other options:

- The WHO growth charts are based on Multicentre Growth Reference Study (MGRS). National Center for Health Statistics (NCHS) developed growth charts that were used in the past.
- Short stature is defined as a height below the third percentile or more than 2 standard deviations below the median height for age and gender (<-2 SD) according to the population standard.
- Puberty will be delayed in this child.

Solution to Question 10:

Thumbsucking in a child is a way of discharging tension. It is a self-limiting habit disorder, usually seen between 6 months to 2 years.

Option A: Pica is persistent ingestion of non-nutritive substances for at least 1 month. It hampers the development of the child and requires clinical attention. Children with pica are at an increased risk of lead poisoning, iron-deficiency anemia, and parasitic infections.

Option B: Breath-holding spells: It is preceded by a provocative event such as anger. Crying stops at full expiration. Child may have apnoea, cyanosis, pallor, hypotonia, or even loss of consciousness, and generalized tonic-clonic seizure (GTCS). Subtypes include cyanotic, pallid, and mixed episodes. Seizures and cardiac causes should be ruled out. Parents should be reassured that breath-holding spells are self-limiting.

Option D: Enuresis:

- When the child has never been dry, it is called primary enuresis.
- When the bedwetting starts after a minimum period of six months of dryness at night, it is termed as secondary enuresis.
- Nocturnal enuresis is caused due to maturational delay. In a normal individual, antidiuretic hormone (ADH) levels usually peak between 4 am and 8 am. This circadian rhythm is impaired in nocturnal enuresis, leading to bedwetting.
- Treatment - Behavioural therapy with positive reinforcement, bladder alarms, and desmopressin reserved for cases resistant to behavioural therapy.

Solution to Question 11:

According to Parson's diseases of the eye, visual acuity of 20/20 is attained by the age of 6 years. Nelson's Textbook of Paediatrics mentions that by the age of 5 or 6 years most children attain 20/20 vision.

Generally accepted visual acuity according to age:

- Visual acuity of 20/40 at 3 years of age
- Visual acuity of 20/30 at 4 years of age
- Visual acuity of 20/20 at 5 or 6 years of age

Solution to Question 12:

The symptoms are suggestive of Autism - which is a pervasive developmental disorder characterized by:

- Impairment in quality of social interaction
- Impairment in quality of communication
- Restricted, repetitive, and stereotyped patterns of behaviors and interests

Common symptoms of Autism Spectrum Disorder -

- Repetition of set words and phrases (echolalia)
- Absent or delayed response to name being called, despite normal hearing
- Reduced or absent responsive social smiling
- Reduced or absent social use of eye contact
- Reduced social interactions
- Lacking the ability to initiate or sustain an interaction with peers or adults
- Stereotypic movements and repetitive behaviors may include hand flapping, finger movements, body rocking, and lunging
- Overly sensitive to noise, smells, or texture and may scream when they hear a siren or vacuum

Modified Checklist for Autism in Toddlers (MCHAT) is a screening tool for the detection of autism.

Test	Disorder
Modified Checklist for Autism in Toddlers (MCHAT)	Autism
Conner Rating Scale	Attention-deficit/hyperactivity disorder (ADHD)
CRAFFT(Car, Relax, Alone, Forget, Friends, Trouble)	Substance abuse in adolescents
Edinburgh Postnatal Depression Scale (EPDS)	Postpartum depression

Solution to Question 13:

Breast and areola enlargement without the separation of the contour is seen in sexual maturity rating (SMR) stage 3.

SMR Stage	PUBIC HAIR	BREASTS
1	Preadolescent	Preadolescent
2	sparse, lightly pigmented, straight, medial border of labia	Breast and papilla elevated as a small mound; diameter of areola increased
3	Darker, beginning to curl, increased amount	Breast and areola enlarged; no contour separation
4	Coarse, curly, abundant, but less than that in adults	Areola and papilla form a secondary mound
5	Adult feminine triangle, spread to the medial surface of thighs	Mature, nipple projects, areola a part of a general breast contour

SMR Stage	PUBIC HAIR	PENIS	TESTES
1	None	Preadolescent	Preadolescent
2	Scanty, long, slightly pigmented	Minimal change/enlargement	Enlarged scrotum, pink, texture altered
3	Darker, starting to curl, small amount	Lengthens	Larger
4	Resembles adult-type but less quantity, coarse, curly	Larger; glans and breadth increase in size	Larger, scrotum dark
5	Adult distribution, spread to the medial surface of thighs	Adult size	Adult size

Solution to Question 14:

The stage at which the pubic hair is coarse, curly and scrotum become dark is SMR stage 4.

The first visible sign of puberty in males is testicular enlargement at 9.5 years, which is a hallmark of SMR stage 2. This is followed by penile growth during SMR 3.

SMR Stage	PUBIC HAIR	PENIS	TESTES
1	None	Preadolescent	Preadolescent

SMR S TAGE	PUBIC HAIR	PENIS	TESTES
2	Scanty, long, slightly pigmented	Minimal change/enlargement	Enlarged scrotum, pink, texture altered
3	Darker, starting to curl, small amount	Lengthens	Larger
4	Resembles adult type, but less quantity; coarse, curly	Larger; glans and breadth increase in size	Larger, scrotum dark
5	Adult distribution; spread to the medial surface of thighs	Adult size	Adult size

Solution to Question 15:

Microcephaly is defined as an occipitofrontal circumference more than 2 SD below the mean for age and sex.

Macrocephaly is defined as an occipitofrontal circumference more than 2 SD above the mean for age and sex.

Monosomy 1p36 is associated with microcephaly.

Infants with benign familial megalencephaly have increased head circumference at birth.

Important causes of macrocephaly:

- Megalencephaly (enlargement of the brain parenchyma)
- Benign familial
- Neurocutaneous syndromes such as neurofibromatosis, tuberous sclerosis
- Lysosomal storage diseases: Tay–Sachs, mucopolysaccharidosis, gangliosidosis
- Increased cerebrospinal fluid
- Hydrocephalus
- Benign enlargement of subarachnoid space
- Hydranencephaly (absence of cerebral hemispheres; the cranial cavity is filled with cerebrospinal fluid [CSF])
- Enlarged vascular compartment
- Arteriovenous malformation
- Subdural, epidural, subarachnoid, or intraventricular hemorrhage
- Increase in the bony compartment
- Bone disease: Achondroplasia, osteogenesis imperfecta
- Bone marrow expansion: Thalassemia major
- Miscellaneous causes
- Intracranial mass lesions: Cyst, abscess, or tumor

- Raised intracranial pressure: Idiopathic pseudotumor cerebri

Solution to Question 16:

The picture is suggestive of a child with left posterior plagiocephaly (as the left side of the head is flattened) which involves the premature fusion of the left lambdoid suture.


In the case of a left anterior plagiocephaly, the premature fusion of the left coronal suture is seen.

Other deformities of the skull:

Craniosynostosis consists of premature fusion of one or more cranial sutures, resulting in an abnormal head shape.

- Primary craniosynostosis: It results from a primary defect of ossification.
- Secondary craniosynostosis: It results from the failure of brain growth.

Deformity	Suture involved
Anterior Plagiocephaly	Respective side coronal suture
Posterior Plagiocephaly	Respective side lambdoid suture
Deformation Plagiocephaly	Positional skull flattening (without suture fusion)
Brachycephaly	Bilateral Coronal suture
Trigonocephaly	Metopic suture
Scaphocephaly	Sagittal suture



Solution to Question 17:

Familial short stature is most likely as bone age is equal to chronological age.

Bone age is more than chronological age in precocious puberty.

Bone age is less than chronological age in all other causes of short stature such as constitutional delay, undernutrition, genetic diseases, and endocrine causes.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Nutrition and Breastfeeding

Question 1:

The most calorically dense macronutrient is _____

- a) Protein
- b) Simple carbohydrates
- c) Fat
- d) Complex carbohydrates

Question 2:

The mother of a school-going child is worried that he eats a lot of junk food. She is anxious to learn about the healthy diet to be followed in the child. You would ideally advise the mother to restrict the dietary intake of saturated fat to less than _____ percentage of total energy intake.

- a) 10%
- b) 5%
- c) 15%
- d) 30%

Question 3:

Your school friend, the mother of a four-year-old child gives you a casual call. She had recently noticed an ad on margarine being a better alternative over butter/ghee and wishes to seek your opinion about adding it to her son's diet. What percentage of total daily energy requirement in children can come from margarine?

- a) < 2
- b) < 10
- c) < 5
- d) 0

Question 4:

A newborn baby is ready to be discharged. As a part of your discharge advice, for how many months would you recommend exclusive breastfeeding?

- a) 4 months
- b) 6 months
- c) 9 months
- d) 1 year

Question 5:

You are taking a neonatal case for presentation during your postings. The mother tells you that the child is exclusively breastfed. Which of the following is least likely to corroborate with this history of being exclusively breastfed?

- a) Hemorrhagic disease of newborn
- b) Breast milk jaundice
- c) Golden color stools
- d) Evening colic

Question 6:

Which of the following mothers can initiate breastfeeding?

- a) 3 only
- b) 2 and 5 only
- c) 1 and 4 only
- d) 2 only

Question 7:

As a part of your neonatal rotation, you are asked to assess the adequacy of breastfeeding and breastfeeding techniques in a primi mother. Which of the following would be an indicator of inadequate breastfeeding in the neonate?

- a) Baby sleeping for 2-3 hours after feed
- b) Weight loss $>10\%$ of birth weight
- c) Urine 5-7 times/day
- d) Stool 4 times/day

Question 8:

For how long can freshly expressed breastmilk be stored at room temperature?

- a) 4 hours
- b) 6 hours
- c) 12 hours
- d) 24 hours

Question 9:

Shrija, a working mother planning to resume her work wishes to know more about expressed breast milk. Within how many hours would you advise her to use the refrigerated expressed breast milk?

- a) 48 hours
- b) 24 hours
- c) 12 hours
- d) 6 hours

Question 10:

An anxious new mother feeding her child with expressed breastmilk (EBM) feels that the consistency and colour of her milk have changed recently. She adds that when she expresses breastmilk, the colour and consistency of the first portion and the ending portion is completely different and that she is worried about the same. Which statement is not true about different breast milk?

- a) Preterm milk has more proteins
- b) Mature milk is thicker with all the essential nutrients for growth
- c) Foremilk is watery and is rich in proteins and sugar
- d) Hindmilk is richer in fat and provides more energy

Question 11:

Following the delivery of a healthy baby boy, his COVID positive mother slipped into ARDS and passed away. The relatives have arranged a breast milk donor and wish to feed the baby

using the device shown in the image. In newborns of which gestational age, would you recommend this as a method of primary feeding?



- a) 28-31 weeks
- b) 32-34 weeks
- c) >34 weeks
- d) Any gestational age

Question 12:

Bifidus factor present in breast milk protects against _____

- a) Enteroamoeba histolytica
- b) Giardia lamblia
- c) Salmonella
- d) Vibrio cholerae

Question 13:

A 2-month-old child is brought to your OPD with complaints of diarrhoea. The mother wants to know if she should continue breastfeeding. Which of these components in breast milk, beneficial in intestinal cell maturation and repair, prompt you to advise her to continue breastfeeding?

- a) Lactoferrin

- b) Epidermal growth factor
- c) Para aminobenzoic acid
- d) Glutathione peroxidase

Question 14:

A young working mother wants to add formula feeds to the diet of her 5-month-old child as she would be unable to breastfeed her baby frequently and satiate her. Which of the following fatty acid should the mother look for in the formula to ensure optimal brain development in the child?

- a) B only
- b) B and C
- c) A,B and C
- d) A and D

Question 15:

Breast milk is superior to cow's milk because _____

- a) Saturated fat content is high
- b) Sodium concentration is high
- c) Casein is 60% of the protein content
- d) Whey is 70% of the protein content

Question 16:

What is found more in human milk as compared to cow's milk?

- a) Lactose
- b) Protein
- c) Vitamin K
- d) Fat

Question 17:

An exclusively breastfed neonate in his 4th week of life was found to be jaundiced. His stools are yellow and urine appears pale straw, he is otherwise well. A provisional diagnosis of breastmilk jaundice is made. Which of the following is false about this condition?

- a) Labs will reveal unconjugated hyperbilirubinemia
- b) Some babies need treatment with phototherapy
- c) Breast feeding is contraindicated
- d) It is common in the second week of life

Question 18:

A 26-year-old nursing mother presents with fever and pain in her left breast. She has had difficulty with breastfeeding previously, and now examination reveals her left breast is warm, red, swollen, and painful to palpation. She was started on dicloxacillin. What advice should be given to the mother regarding breastfeeding?

- a) Complete cessation of breastfeed
- b) Continue feeding from both the breasts
- c) Continue feeding from only right breast
- d) Continue feeding from only left breast

Question 19:

Which of the following statements are true with respect to the WHO Baby-Friendly Hospital Initiative for small, sick, and preterm newborns (2020)?

- a) Paladai feeding is indicated in very preterm infants with no suck-swallow-breath coordination
- b) Finger-feeding is optimally indicated for short-term use and may improve breastfeeding rates at discharge
- c) Feeding tubes are indicated only in neurologically compromised infants for long-term use
- d) Bottle feeding can lead to overfeeding but has a reduced risk of aspiration in infants with immature suck-swallow-breathe reflex

Answer Key

Question No.	Correct Option
--------------	----------------

1	c
2	a
3	d
4	b
5	d
6	b
7	b
8	b
9	b
10	b
11	b
12	c
13	b
14	b
15	d
16	a
17	c
18	b
19	b

Detailed Explanations

Solution to Question 1:

Fat is the most calorically dense macronutrient providing approximately 9 kcal/g.

The nutrients that provide energy intake in the child's diet are referred to as macronutrients. They include the following:

- Fats (~9 kcal/g)
- Carbohydrates (4 kcal/g)
- Proteins (~4 kcal/g)

The main dietary sources of fat are as follows:

- For infants: human milk/formula
- For older children: fat from animal products, vegetable oils, and margarine.

Solution to Question 2:

According to current dietary guidelines for children, energy from saturated fats should be <10% of total energy intake.

Current dietary guidelines for children and adolescents:

- Total fat should account for <30% of total daily energy
- Saturated fat less than 10% of total daily energy
- Dietary cholesterol <300 mg/day
- No trans-fat

Solution to Question 3:

Margarine and other hydrogenated oils are trans fat. Trans fat is not recommended in children according to current dietary guidelines.

Current dietary guidelines for children and adolescents:

- Total fat should account for <30% of total daily energy
- Saturated fat less than 10% of total daily energy
- Dietary cholesterol <300 mg/day
- No trans-fat

Solution to Question 4:

Exclusive breastfeeding is recommended for 6 months.

Complementary foods rich in iron and other micronutrients should be introduced at about 6 months of age. Breastfeeding should continue for at least the first year.

Solution to Question 5:

Exclusively breastfed infants are least likely to have evening colic. The incidence of diarrhoea, allergy to bovine milk, and atopic eczema are least reported in exclusively breastfed infants.

Option A: Vitamin K content of milk is low and may contribute to hemorrhagic disease of the newborn.

Option B: Breast milk contains inhibitors of glucuronyl transferase and enhances absorption of bilirubin from the gut. This leads to the development of breast milk jaundice.

Option C: During the first few days, the breastfed infant has a black-green coloured stool. It changes to golden yellow by the end of the first week.

Solution to Question 6:

The correct answer is 2 and 5 only. Breastfeeding is not contraindicated if the mother has hepatitis B infection or if she is a chronic smoker.

Infants born to an HbsAg positive mother routinely receive hepatitis B immunoglobulin and hepatitis B vaccine. So, a delay in the initiation of breastfeeding is not required.

Alcohol consumption and smoking are NOT contraindications of breastfeeding. However, smoking and alcohol consumption should be discouraged.

Note: In a mother with untreated, active tuberculosis:

- Temporarily withhold breastfeeding but can feed with expressed breastmilk
- Resume breastfeeding once she has been adequately treated for 2 weeks and is no longer contagious.

Contraindications of breastfeeding:	
Do not Breastfeed Do not give expressed breastmilk	Infant factors: Classic galactosemia Maternal factors: Human T-cell lymphocyte virus (HTLV) infection Ebola virus infection PCP/cocaine abuse
Do not Breastfeed (Temporarily) Do not give expressed breastmilk (Temporarily)	Maternal factors: Brucellosis undergoing diagnostic imaging with radiopharmaceuticals Active herpes simplex virus (HSV) infection with lesions on the breasts
Do not Breastfeed (Temporarily) Can give expressed breastmilk	Maternal factors: Untreated, active Tuberculosis Active Varicella infection

Solution to Question 7:

Weight loss of >7-10% of birth weight is an indicator of inadequate breastfeeding in a neonate.

Indicators of inadequate breastfeeding in the newborn:

- Weight loss >7-10 % of birth weight
- Lethargy (sleeping longer than 2–3 hours)
- Delayed passage of stools (Less than 4 times/day)
- Decreased urine output (urinating less than 5–7 times/day)
- Hypernatremic dehydration
- Inconsolable crying and increased hunger

Insufficient milk intake may be caused by insufficient milk production, failure of established breastfeeding, and health conditions in the infant that prevent proper breast stimulation.

Breastfed neonates usually feed 8–12 times a day, with a minimum of 8 times per day. However, the neonate should be breastfed only on demand.

Solution to Question 8:

Expressed breast milk can be stored at room temperature for 6-8 hours.

Note: This is a controversial question.

A difference of opinion among authors is noted regarding the refrigeration of expressed breast milk. Nelson's (21st ed, page 324) mentions that breast milk can be refrigerated and used within 48 hours and frozen for around 6 months. However, AIIMS in collaboration with WHO recommends that breast milk can be refrigerated up to 24 hours and frozen at -20 degree C for about 3 months.

Expressed breast milk	Stored for / Used within(maxi mum time)
Room Temperature	6 hours
In refrigerator	24 hours
Frozen breastmilk at -20° C	3 months

Solution to Question 9:

Breast milk stored in the refrigerator should be used within 24 hours.

Refrigerated or frozen milk should be thawed rapidly by holding under running tepid water and used completely within 24 h after thawing. Milk should never be microwaved.

Note: This is a controversial question.

A difference of opinion among authors is noted regarding the refrigeration of expressed breast milk. Nelson's (21st ed, page 324) mentions that breast milk can be refrigerated and used within 48 hours and frozen for around 6 months. However, AIIMS in collaboration with WHO recommends that breast milk can be refrigerated up to 24 hours and frozen at -20 degree C for about 3 months.

Expressed breast milk	Stored for / Used within(maxi mum time)
Room Temperature	6 hours
In refrigerator	24 hours

Expressed breast milk	Stored for / Used within(maximum time)
Frozen breastmilk at -20° C	3 months

Solution to Question 10:

Mature milk is thinner (not thicker) with all essential nutrients for growth.

Various transitional forms of breastmilk:

Colostrum

- In the initial 3 to 4 days
- Small quantity, thick, and yellow in colour
- A large number of antibodies, immune component cells; and vitamins A, D, E, and K.

Transitional milk

- Secreted after 3 to 4 days until 2 weeks
- Immunoglobulin and protein content decreases, while the fat and sugar content increases.

Mature milk

- Follows transitional milk.
- Thinner and more watery.
- Contains all the nutrients essential for the optimal growth of the baby.

Foremilk

- Secreted at the start of the feed
- Watery and rich in proteins, sugar, vitamins, minerals, and water that quenches the baby's thirst.

Hindmilk

- Comes towards the end of the feed
- Provides more energy and a sense of satiety and it is richer in fat.

Preterm milk

- Milk of mother who delivers before 37 weeks.
- Contains more proteins, sodium, iron, immunoglobins, and calories as per the requirement of the preterm infant.

The given image shows watery foremilk and hindmilk which is rich in fat.



Solution to Question 11:

The given image shows paladai. It can be used as a primary feeding method during routine feeding in neonates born at gestational age 32-34 weeks only. Sick mothers or mothers with contraindications to breastfeeding can feed their child using paladai. The milk can either be expressed breastmilk or they can resort to using undiluted cow milk or formula feeds.

Choice of initial feeding methods in low-birth-weight (LBW) infants:

Gestational Age	Feeding Skill Matured	Method of Feeding
<28 weeks	No proper sucking effort. No propulsive motility of the gut	IV Fluids
28-32 weeks	Sucking Bursts developNo coordination between suck/swallow and breathing	Oro-gastric/ Nasogastric tube feeding with occasional spoon/ paladai feeding
32-34 weeks	Slightly mature sucking pattern.Coordination between breathing and swallowing begins	Feeding by spoon/paladai/cup
>34 weeks	Mature sucking pattern. More coordination between breathing and swallowing	Breastfeeding

Solution to Question 12:

Bifidus factor present in breast milk protects against Salmonella.

Components of breast milk	Protection against
Lysozyme	Escherichia coli, Salmonella, Entamoeba histolytica
Lactoferrin	E. coli, Shigella, Salmonella, E. histolytica, Giardia lamblia, Plasmodium falciparum, Rotavirus
Bifidus factor	E. coli, Shigella, Salmonella
Low levels of para-aminobenzoic acid	Plasmodium

Solution to Question 13:

The epidermal growth factor present in breastmilk helps in intestinal cell maturation and repair. This is why mothers are advised to continue breastfeeding in an infant affected with diarrhoea. It is also beneficial in luminal surveillance.

Lactoferrin (Option A) helps in immunomodulation, iron chelation, antimicrobial action. It is also an antiadhesive and trophic for intestinal growth.

Para-aminobenzoic acid (Option C) is present in low levels and offers protection against plasmodium.

Glutathione peroxidase (Option D) prevents lipid oxidation.

Solution to Question 14:

Docosahexenoic acid (DHA) and arachidonic acid (ARA), both play an important role in the growth and CNS development in babies.

DHA and ARA are essential poly unsaturated fatty acids (PUFA) present in breast milk.

Both are essential for cognition functions, visual acuity, and immune responses. They have optimal effects on brain development when included together, in efficacious levels, in an infant's diet.

Almost all infant formulae are now fortified with pre-formed DHA and ARA.

It is advisable that all infants upto 6 months of age are to be exclusively breastfed and not formula fed. However, in unavoidable situations such as lactation failure or maternal infections where breastfeeding is contraindicated, the use of a formula feed that contains both DHA and ARA is recommended.

Other options:

- Eicosapentaenoic acid (EPA) and Linolenic acid are other essential fatty acids that are present in breast milk.

- These are required for the overall protection of the cardiovascular system as well as to reduce systemic inflammation.

Solution to Question 15:

Whey constitutes 70-80% of breast milk's protein content. The Whey: Casein ratio in human milk fluctuates between 70:30 and 80:20. This ratio is 20:80 in cow's milk. The high fraction of soluble whey protein in breast milk makes it easily digestible.

The predominant whey protein is α -lactalbumin in human milk and β -globulin in cow milk. Human milk contains amino acids like taurine and cysteine (lacking in cow's milk), which are necessary for neurotransmission and neuromodulation.

Breast milk is rich in polyunsaturated fat (not saturated fat), which is necessary for myelination. Human breast milk is generally rich in oleic acid and linoleic acid.

Sodium concentration is not high in breast milk. It has a low osmolality, and hence, the solute load on the kidneys is low.

Differences between human milk and cow's milk are shown below:

Nutrient	Human milk (per 100 mL)	Cow's milk (per 100 mL)
Protein	1.1 g	3.5 g
Calories	67 kcal	62 kcal
Casein	30–40% (rich in whey protein)	80%
Lactose	7.4 g	4.5 g
Fat	3.5 g	4g
Na ⁺ (mEq/L)	6.5	25
K ⁺ (mEq/L)	14	35
Osmolality (mOsm/L)	290	350
calcium/phosphate	>2	<2
Vitamin K	1.5 μ g	6 μ g
Iron	0.29–0.45 mg	0.01-0.38 mg
Vitamin D	0.5-10 IU	0.5-4.5 IU

Solution to Question 16:

The percentage of lactose in human milk is greater than that in cow's milk.

Differences between human milk and cow's milk are shown below:

Human breast milk is generally rich in oleic acid and linoleic acid.

Nutrient	Human milk (per 100 mL)	Cow's milk (per 100 mL)
Protein	1.1 g	3.5 g
Calories	67 kcal	62 kcal
Casein	30–40% (rich in whey protein)	80%
Lactose	7.4 g	4.5 g
Fat	3.5 g	4g
Na ⁺ (mEq/L)	6.5	25
K ⁺ (mEq/L)	14	35
Osmolality (mOsm/L)	290	350
calcium/phosphate	>2	<2
Vitamin K	1.5 µg	6 µg
Iron	0.29–0.45 mg	0.01-0.38 mg
Vitamin D	0.5-10 IU	0.5-4.5 IU

Solution to Question 17:

Breastfeeding is not contraindicated in breast milk jaundice. Mothers are advised to continue breastfeeding at frequent intervals.

Breastmilk jaundice is seen in about 2–4% of exclusively breastfed babies. There is significant unconjugated hyperbilirubinemia after the 7th day and may last for 2–3 weeks. This may be caused by the breast milk that might contain β -glucuronidase, which deconjugates intestinal bilirubin and promotes its absorption and other inhibitors of glucuronyl transferase. Persistently high bilirubin levels may require changing from breast milk to infant formula for 24–48 h and/or treatment with phototherapy, without cessation of breastfeeding.

Breastfeeding jaundice (BFJ)

It is an early-onset, unconjugated hyperbilirubinemia of breastfed infants. Occurs in the 1st week after birth (usually appears after 24 - 72 hours of birth). It is due to decreased milk intake with dehydration and/or reduced caloric intake. The treatment would include continuing breastfeeding and supplementation with formula feeds if needed.

Solution to Question 18:

Lactational mastitis is not a contraindication to breastfeeding and the mother should continue feeding on both breasts.

It is most commonly caused due to *Staphylococcus aureus* present in the infant's nasopharynx. Dicloxacillin is safe to be given in nursing mothers.

Solution to Question 19:

Finger-feeding is optimally indicated for short-term use and may improve breastfeeding rates at discharge. It is indicated in neurologically compromised and preterm infants for a short period of time.

Paladai feeding is indicated in preterm, late preterm, and term infants for supplementation. It is economic, cheap, and is easy to use and clean. It has a risk of aspiration and is a slower method of feeding.

Feeding tubes- orogastric or nasogastric tubes are an efficient way of measuring intake. It is optimally used in very preterm infants with no suck-swallow-breathe coordination for long-term use. It requires frequent replacement.

Bottle and teat feeding, though not recommended is an efficient way to feed. Slow-flow teats mimic breastfeeding and are preferred to rapid-flow teats. They allow overfeeding and there is a risk of aspiration.

Protein Energy Malnutrition

Question 1:

A 9-month-old boy was brought to the well-baby clinic for routine immunization. He has a weight for height less than 2 standard deviations below the median. What is the most likely diagnosis?

- a) Acute malnutrition
- b) Chronic malnutrition
- c) Acute on chronic malnutrition
- d) Stunting

Question 2:

A child is brought to the OPD for regular follow-up. Which of the following anthropometric record would prompt you to suspect a diagnosis of moderate malnutrition?

- a) Weight for age ≤ -2 SD
- b) Weight for height ≤ -2 SD to ≥ -3 SD
- c) Weight for age ≤ -3 SD
- d) Weight for height $+2$ SD to $+3$ SD

Question 3:

Which of the following is an age-independent index?

- a) Kanawati and McLaren index
- b) Jelliffe ratio
- c) Rao and Singh index
- d) All of the above

Question 4:

A child in your unit is diagnosed with severe acute malnutrition (SAM). According to the WHO criteria, SAM is defined best by:

- a) Weight for age less than - 2SD
- b) Weight for height less than + 2SD
- c) Weight for age less than + 3SD
- d) Weight for height less than - 3SD


Question 5:

You are an intern evaluating a 2-year-old girl child for malnutrition. The measurement with the WHO standard mid-upper-arm-circumference (MUAC) tape shows yellow. This reading would correspond to?

- a) 10.5 to 11.5 cm
- b) 11.5 to 12.5 cm
- c) 12.5 to 13.5 cm
- d) 13.5 to 14.5 cm

Question 6:

A young child is brought to you for a general check-up. The physician is examining as shown below in the image. What would be the cut-off to label the child as malnourished?



- a) 11.5 mm
- b) 10 mm
- c) 6 mm

d) 4 mm

Question 7:

A laborer's second child is brought to the OPD with a swollen belly and dull face. He has been fed rice water (rice milk) in his diet mostly. On investigations, the child is found to have low serum protein and low albumin. What is the probable diagnosis?

- a) Kwashiorkor
- b) Marasmus
- c) Indian childhood cirrhosis
- d) Kawasaki disease

Question 8:

You are sent for active field surveillance of malnourished children to a rural district in central India. Which of the following specific finding would prompt you to make a diagnosis of marasmus?

- a) Fatty change in liver
- b) Voracious appetite
- c) Hypoalbuminemia
- d) Edema

Question 9:

Sheila, a widowed maid brings her youngest child to your OPD with complaints of cough and cold. She has not been able to feed her kids after the lockdown affected her work. The image of the child is shown below. Which of the following is least likely to be seen in this case?



- a) Baggy pant appearance
- b) Reduced mid arm circumference
- c) Apathy
- d) Pitting not seen

Question 10:

Shubham, a young infant was found to have a mid-upper arm circumference of 11cms. On examination, the child appeared stunted with loss of muscle mass and subcutaneous tissue including the buccal pad of fat. What is the grade of marasmus in this child?

- a) 1
- b) 2
- c) 3
- d) 4

Question 11:

A child from an urban slum area is admitted to your unit and is receiving treatment for Kwashiorkor. Which of the following is not true regarding the treatment?

- a) Start iron therapy on admission
- b) Correct dehydration slowly over 12 hours
- c) Suspect primary treatment failure if appetite is not regained by day 4

- d) Suspect secondary treatment failure if oedema reappears after 4 weeks

Question 12:

A 20-month-old child diagnosed with severe acute malnutrition (SAM) is receiving treatment in your clinic. Given below are five statements regarding the treatment. Pick out the incorrect statements.

- a) 1, 2, 3, 4, 5
- b) 1, 3, 5
- c) 2 and 4 only
- d) 3 only

Question 13:

A 3 year old child is brought to the hospital with complaints of multiple episodes of loose stools and progressive weight loss. She was found to have a mid-upper arm circumference of 10.5 cm. Which of the following is true regarding the management of this child?

- a) Initiate IV fluids immediately
- b) Administer 10 mL of 50% glucose IV, if unconscious
- c) Continue aggressive hydration, even after dehydration resolves
- d) Use ReSoMal for rehydration 5 mL/kg every 30 min for the first 2 hours

Question 14:

A young girl who was grossly emaciated and undernourished was started on partial parenteral nutrition together with enteral nutrition via a nasogastric tube. In the first week, which among the following would be the hallmark for the development of refeeding syndrome?

- a) Hyperphosphatemia
- b) Hypophosphatemia
- c) Hypokalemia
- d) Hyperkalemia

Answer Key

Question No.	Correct Option
1	a
2	b
3	d
4	d
5	b
6	c
7	a
8	b
9	c
10	d
11	a
12	d
13	d
14	b

Detailed Explanations

Solution to Question 1:

A low weight-for-height, or wasting, indicates acute malnutrition.


Nutritional status is often assessed in terms of anthropometry

- Low weight-for-height, or wasting, indicates acute malnutrition.
- High weight-for-height indicates overweight.
- Low height-for-age, or stunting, indicates chronic malnutrition. This is the best indicator of long-term nutritional status in the pediatric population.
- Low weight for the age indicates acute or chronic malnutrition. It does not differentiate between wasting and stunting. Hence, it has limited clinical significance.

Note:

Mid-upper arm circumference (MUAC) is an age-independent parameter used for screening wasting in children.

Anthropometric indicators of different types of malnutrition in Children




©Marrow

Solution to Question 2:

An anthropometric score of Weight for height ≤ -2 to $\geq +3$ SD indicates moderate malnutrition.

The indicators of undernutrition include:

Anthropometric indicators of different types of malnutrition in Children



©Marrow

Solution to Question 3:

All the indices listed, namely, Kanawati and McLaren index, Jelliffe ratio, and Rao and Singh index, are age-independent indices.

Age-independent indices:

- Kanawati and McLaren index: Mid-arm circumference/Head circumference
- Jelliffe Ratio: Head circumference/Chest circumference
- Rao and Singh Index: $\text{Weight (kg)} \times 100/\text{height}^2 \text{ (cm)}$
- Dugdale Index: $\text{Weight (kg)}/\text{height}^{1.6} \text{ (cm)}$

Solution to Question 4:

Severe acute malnutrition (SAM) is best defined as weight for height less than median minus -3SD

Severe acute malnutrition (SAM) in children of ages 6 to 59 months is defined by the WHO as any of the following:

- Weight-for-length (or height) below -3 SD of the median WHO child growth standards.
- Presence of bipedal edema
- Mid-upper arm circumference ≤ 11.5 cm

Once the child is diagnosed with severe acute malnutrition, the doctor must assess for severe edema (+++); low appetite (failed appetite test); medical complications (severe anemia, pneumonia, diarrhea, dehydration, cerebral palsy, tuberculosis, HIV, heart disease, etc.) and one or more danger signs as per IMNCI

If any of the complications are present, the child is diagnosed with complicated SAM and is managed inpatient in the facility.

If the above-mentioned signs are absent, the child is diagnosed with uncomplicated SAM and supervised management at home is given.


Solution to Question 5:

The yellow colour on the new WHO standard mid-upper arm circumference of (MUAC) tape correlates to a MUAC of 11.5 to 12.5 cm.

MUAC tape cut-offs:

Colours	Measurements	Indications
Red	< 11.5 cm	Severe Acute Malnutrition (SAM)
Yellow	11.5 to 12.5 cm	Under-nourished
Green	> 12.5 cm	Normal

Mid-upper arm circumference (MUAC) tape



©MARROW

Mid-upper arm circumference is used in children between 6 to 60 months. It is measured between the acromion and olecranon process and is an age-independent parameter.

Alternative methods for evaluating malnutrition:

Bangle Test - It has an internal diameter of 4 cms. Normally, it should not cross the elbow.

QUAC stick test - It is given by the formula - Mid-upper arm circumference/ Height of child

Skinfold thickness: It is an age-independent parameter. Routinely, we measure the triceps skinfold thickness. It is measured with the help of Harpenden calliper.

- Normal: 10 mm
- Severe malnutrition: < 6 mm

Solution to Question 6:

The image shows the measurement of skinfold thickness using Harpenden calipers. The cut-off to label the child as malnourished would be the triceps skinfold thickness value of less than 6mm. This is an age-independent parameter.

Alternative methods for evaluating malnutrition:

Bangle Test - It has an internal diameter of 4 cms. Normally, it should not cross the elbow.

QUAC stick test - It is given by the formula - mid-upper arm circumference/ height of child.

Solution to Question 7:

The given clinical scenario of a child with a swollen belly, dull face with hypoproteinemia along with poor diet history suggests the diagnosis of kwashiorkor.

Kwashiorkor is a form of severe acute malnutrition (SAM) characterized by edema and apathy. In edematous malnutrition, the initial appearance of edema is commonly observed in the feet and lower legs, progressing into generalized edema affecting the hands, arms, and face. The skin over the swollen areas may show dark, crackled, peeling patches (flaky paint dermatosis) with pale skin underneath, making it susceptible to infections.

There may be sparse and easily pluckable hair. Hepatomegaly with fatty infiltration is a common finding. Associated symptoms of misery, apathy, and loss of appetite are seen in children with edema.

Other options:

Option B: Marasmus is a severe form of wasting characterized by significant loss of fat and muscle tissue as the body utilizes these resources for energy. It is often triggered by acute starvation or illness in individuals with borderline nutritional status.

Option C: Indian childhood cirrhosis (ICC) is a chronic liver disease that affects infants and young children. Common symptoms include jaundice, pruritus, lethargy, and hepatosplenomegaly. The disease progresses rapidly, leading to the development of cirrhosis.

Option D: Kawasaki disease (KD) is a systemic inflammatory disorder with arteritis of medium to large-sized vessels. It presents with fever, rash, conjunctivitis, and cervical lymph node enlargement.

Solution to Question 8:

Children with marasmus have a voracious appetite, whereas children with kwashiorkor often refuse to eat.

Marasmus is a calorie deficit state and kwashiorkor is a protein deficit state. The signs of low protein manifesting as hypoalbuminemia and edema are characteristically seen in kwashiorkor.

Solution to Question 9:

The image shows a child with marasmus which is not associated with apathy. Apathy is a psychomotor change seen in kwashiorkor.

Solution to Question 10:

Wasting of buccal pad of fat is classified as grade 4 marasmus.

Grading of marasmus (depending on wasting)

Grading of kwashiorkor (depending on edema)


Grade	Feature
1	Loose skin fold in groin and axilla
2	Wasting of the thighs and buttocks
3	Wasting of the chest and back
4	Wasting of the buccal pad of fat

Grade	Feature
1	Pedal edema
2	Facial edema
3	Chest and body edema
4	Ascites

Solution to Question 11:

In a child with kwashiorkor, iron therapy (3 mg/kg/day) should be started only once the child starts gaining weight. Iron should not be given in the stabilization phase because excess iron can lead to free radical formation.

The image below explains the treatment protocol to be followed in the case of PEM.



©MARROW

The following should strike a suspicion of treatment failure in PEM:

Inpatient cases:

Outpatient cases:

Primary Failure	Secondary failure
Failure to regain appetite by day 4.Failure to start losing oedema by day 4.Presence of oedema on day 10.Failure to gain at least 5 g/kg/day by day 10.	Failure to gain 5 g/kg/day for 3 consecutive days during rehabilitation.Reappearance of oedema after 4 weeks.

Primary failure	Secondary failure
Failure to gain any weight by 2 weeksFailure to lose oedema in 4 weeks	No weight gain in two visits

Solution to Question 12:

Vitamin A is not given daily in the treatment of SAM. It is given on day 1, day 2 and day 14.

The general treatment involves 10 steps in 2 phases:

Initial stabilization phase: It focuses on restoring homeostasis and treating medical conditions and requires 2-7 days of inpatient treatment.

Mnemonic: SHIELDED

- Sugar deficiency(hypoglycemia)
- Hypothermia
- Infection
- Electrolyte imbalance
- Dehydration
- Deficiency of nutrients

Rehabilitation phase: It focuses on rebuilding wasted tissue and generally takes several weeks.

Steps:

- Treat hypoglycemia (blood glucose level < 54 mg/dl): if symptomatic, give 10% Dextrose IV at 5ml/kg. Feeding should be initiated with starter F-75 (formula with 75 kcal/100 ml of feed)
- Treat hypothermia (axillary $< 35^{\circ}\text{C}$ or 95°F): Skin-to-skin contact with care called "kangaroo technique" or dress in warmer clothes, cover head, wrap in a warmed blanket, and provide indirect heat.
- Treat dehydration: Hydrate with ReSoMal (45 mmol Na/L). It is a low-sodium rehydration solution for malnutrition.

- Correct electrolyte imbalance: There is a deficit of potassium and magnesium and excess sodium. Give extra potassium (3 to 4 mEq/kg/day) and magnesium (4 mEq/ kg/day) for at least 2 weeks.
- Treat an infection: Starting on the first day, give broad-spectrum antibiotics to all children.
- Correct micronutrient deficiencies:
- Do not give iron in the stabilization phase, because excess iron leads to free radical injury. Give iron 3 mg/kg/day once the child starts gaining weight after the stabilization phase.
- Give vitamin A on day 1. Repeat on day 2 and day 14.
- Folic acid 1 mg (5 mg on day 1)
- Zinc (2 mg/kg/day)
- Copper (0.3 mg/kg/day)
- Start cautious feeding: Give 8 to 12 small feeds of F75 feeds. It is low-lactose milk that contains 75 kcal and 0.9 g protein per 100 mL. After 3 days, transition to F100 feeds is made which is called a catch-up diet. This provides 100 kcal and 2.9 g protein per 100 mL. After 1 week, calories are further increased to 150–200 kcal/kg/day and protein to 4–6 g/kg/day.
- Rehabilitation: The indications of entry to this phase are reduced/minimal oedema and the return of appetite.
- Beginning of feeding.
- After 3 days, the transition to energy-dense feeding
- Stimulation of emotional and sensorial development
- Transfer to home-based diet

Management of protein energy malnutrition

Activity	Stabilization		Rehabilitation
	Days 1-2	Days 3-7	Weeks 2-6
Treat or prevent: Hypoglycaemia Hypothermia Dehydration	----->	----->	
Correct electrolyte imbalance	----->		
Treat infection	----->		
Correct micronutrient deficiencies	-----> Without iron ----->		-----> With iron ----->
Begin feeding	----->		
Increase feeding to recover lost weight ("catch-up growth")			----->
Stimulate emotional and sensorial development			----->
Prepare for discharge			----->

©MARROW

Solution to Question 13:

Progressive weight loss and mid-upper arm circumference of <11.5 cm denote severe acute protein-energy malnutrition (PEM). When accompanied by loose stools, management includes the use of low sodium ORS for rehydration of 5mL/kg every 30 min for the first 2 hours.

Management of fluid/electrolyte status in SAM:

- Do not give IV fluids unless the child is in shock.
- Give ReSoMal (low-sodium rehydration solution for malnutrition) 5 mL/kg every 30 min for the first 2 hr orally or NG tube.
- Then give 5-10 mL/kg in alternate hours for up to 10 hour. The amount depends on stool loss and eagerness to drink. Feed every other alternate hour.
- Give ReSoMal after each watery stool as a prevention strategy.
- Do not continue aggressive hydration even after dehydration resolves.

Treatment of hypoglycemia in malnutrition:

- If conscious: 10% glucose (50 mL) orally or 1 teaspoon sugar under the tongue, whichever is the quickest.
- If unconscious: Immediately give sterile 10% glucose (5 mL/kg) by IV.

Note:

50% glucose is never used in children as it is extremely viscous and concentrated. This could lead to injection site pain, severe thrombophlebitis, and irritation. Also, this can cause osmotic overload in a three-year-old.

Solution to Question 14:

The hallmark of refeeding syndrome is the development of severe hypophosphatemia.

Refeeding syndrome is a complication of the acute nutritional rehabilitation of children who are undernourished from any cause, and it may follow overly aggressive enteral or parenteral alimentation.

When excessive carbohydrates are administered, the resultant increase in serum insulin levels may produce the following:

- Hypophosphatemia (hallmark)
- Hypokalemia
- Hypomagnesemia

A controlled transition over 3 days from F-75 to F-100 is recommended by the WHO for treating malnutrition, to prevent refeeding syndrome.

Deficiency of Fat Soluble Vitamins

Question 1:

Which of the following is an 'Anti-infective' vitamin?

- a) Vitamin E
- b) Vitamin A
- c) Vitamin D
- d) Vitamin C


Question 2:

What would be the ideal dose of vitamin - A that you should prescribe to a 9-month-old infant along with the measles vaccine?

- a) 10,000 IU
- b) 100,000 IU
- c) 50,000 IU
- d) 200,000 IU

Question 3:


A 2-year-old unvaccinated girl is brought to the OPD with high-grade fever and maculopapular rashes for 2 days. On examination, she has blue-white spots on her buccal mucosa as shown in the image attached below. Which of the following vitamin deficiency is associated with this condition?



- a) Vitamin A
- b) Vitamin E
- c) Vitamin C
- d) Vitamin K

Question 4:

A young girl of class 4 complains of a burning sensation in the eye. Clinical examination of eyes reveals the finding shown below. What could be the mechanism for the development of such lesions?




- a) Deposition of calcium in the conjunctiva

- b) Scarring of the conjunctiva
- c) Excessive growth of conjunctival tissue
- d) Keratinization of the conjunctival epithelium

Question 5:

A 6-year-old boy with complaints of dry, itchy eyes and night blindness had the following ophthalmic finding as given in the image. What is the stage of vitamin A deficiency as per WHO classification?



- a) Stage X1B
- b) Stage X3A
- c) Stage X2
- d) Stage XF

Question 6:

A 6-year-old boy is brought to the OPD with complaints of headache, desquamation of the skin, and bone pain. Pathological examination revealed lysosomal rupture within the cells, prompting a suspicion of hypervitaminosis A. Which of the following statement is false about the principle vitamin that is excess in the given condition?

- a) 1, 3, 4, 5
- b) 1 and 4 only
- c) 2 and 5 only

d) 4 and 5 only

Question 7:

Your best friend is recently worried that she has a lot of hair fall. Which of the following vitamin capsules if added to her diet can help prevent free radical injury and thus be beneficial for her??

- a)** A
- b)** E
- c)** K
- d)** D

Question 8:

An anxious mother brings her 6-month-old child with concerns about not gaining weight and height. On examination, stunted growth, craniotabes, frontal bossing, and craniosynostosis are noted. What is the most likely diagnosis?

- a)** Scurvy
- b)** Rickets
- c)** Rhizomelic chondrodysplasia punctata
- d)** Achondroplasia

Question 9:

The X-ray image of a child with muscle weakness and failure to thrive is attached below. What is the likely diagnosis?



- a) Scurvy
- b) Rickets
- c) Osteomalacia
- d) Osteopetrosis

Question 10:

An 8-year-old child from a rural slum area is brought to your OPD with deformity as shown in the image. Which of the following conditions is characteristically associated with this deformity?



- a) Scurvy

- b) Rickets
- c) Achondroplasia
- d) Rheumatoid arthritis

Question 11:

Your school friend brings his 5-year-old son to your OPD with small swellings on the chest that he had felt while giving the child a bath. On examination, you note swellings at the costochondral junctions. Which of the following conditions would be your differentials to this clinical presentation?

- a) 1 and 2 only
- b) 2 and 3 only
- c) 2 only
- d) 1, 2, 3

Question 12:

A 22-month-old girl presents with failure to thrive and delayed achievement of motor milestones. On examination, she seems irritable and is noted to have bowed legs, thick wrists, and dental caries. Which of the following is not a radiographic sign of this condition?

- a) Increase in the width of the growth plate
- b) Decreased bone density
- c) Splaying
- d) Subperiosteal hemorrhage

Question 13:

A 2-year-old child from an urban slum is diagnosed with nutritional rickets. Which of the following biochemical markers can be seen in this child?

- a) Increased alkaline phosphatase
- b) Hyperphosphatemia
- c) Reduced parathyroid hormone
- d) Alkalosis

Question 14:

A child in your unit is admitted with alopecia, oligodontia, and features of rickets with elevated serum levels of alkaline phosphatase activity and 1,25-dihydroxy vitamin D. What could be the pathology?

- a) Nutritional deficiency of vitamin D
- b) Defect in 1 α hydroxylase
- c) Autoantibodies against vitamin D
- d) End-organ resistance to vitamin D

Question 15:

Stoss therapy is used in the treatment of which of the following micronutrient deficiencies?

- a) Vitamin A
- b) Vitamin E
- c) Vitamin D
- d) Vitamin K

Question 16:

A child with rickets not responding to treatment is being evaluated. Investigations show serum calcium level of 9 mg/dL, low phosphate level of 2 mg/dL, elevated alkaline phosphatase, and normal parathyroid hormone levels with normal arterial blood gas levels. What is the diagnosis?

- a) Renal tubular acidosis
- b) Hypophosphatemic rickets
- c) Chronic kidney disease
- d) Vitamin D-dependant rickets, type 2

Question 17:

A concerned mother of an infant, known case of cystic fibrosis, presents to your OPD with complaints of abdominal pain, loose stools not settling with antibiotics, and other medications. She recently stopped breastfeeding and switched to formula feeds. On examination, you find petechial rashes. Which of the following below mentioned causes

would not prompt you to suspect a deficiency of Vitamin K in this child?

- a) Biliary obstruction
- b) Artificial milk feeds
- c) Oral antibiotic therapy
- d) Cystic fibrosis

Question 18:

What is the recommended dietary allowance (RDA) of vitamin D in a 4-year-old child?

- a) 400 IU
- b) 600 IU
- c) 200 IU
- d) 1000 IU

Question 19:

Rickets in an infant presents as all except _____

- a) Craniotables
- b) Widened fontanelle
- c) Rachitic rosary
- d) Bow legs

Question 20:

Which of the following is false regarding phosphatonins?

- a) It is a phosphate-regulating factor
- b) It increases the renal reabsorption of phosphate
- c) It is produced by osteocytes
- d) FGF 23 is an important phosphatonin

Answer Key

Question No.	Correct Option
1	b
2	b
3	a
4	d
5	b
6	d
7	b
8	b
9	b
10	b
11	d
12	d
13	a
14	d
15	c
16	b
17	b
18	b
19	d
20	b

Detailed Explanations

Solution to Question 1:

Vitamin A is required for the maintenance of epithelial functions, creating an effective barrier against pathogens; hence, it is called as an anti-infective vitamin. So, vitamin A deficiency can lead to changes in the epithelia of all mucosal surfaces, leading to increased chances of infections.

Vitamin E (Tocopherol) helps is an anti-oxidant. Deficiency manifests as skin changes and hair loss.

Vitamin D maintains calcium and phosphorus homeostasis. Deficiency can lead to rickets in children and osteomalacia in adults.

Vitamin C helps in the formation of connective tissues and collagen in the skin, bone, and blood vessels. Deficiency leads to their fragility and causes scurvy.

Solution to Question 2:

For a child aged 9 months, the dose of vitamin A given is 100,000 IU.

Solution to Question 3:

The girl is suffering from measles. It is associated with vitamin A deficiency.

Vitamin A deficiency is associated with an increased risk of infection and increased mortality with a variety of infectious diseases particularly measles. Hence, vitamin A therapy is indicated in all patients with measles.

The main functions of vitamin A are as follows:

- Maintenance of vision, especially night vision
- Maintenance of epithelial tissues
- Differentiation of various tissues, particularly during reproduction and gestation, by regulating gene expression.

Features of vitamin A deficiency:

- Eyes: Defective dark adaptation is a characteristic early clinical feature, resulting in night blindness (earliest manifestation). This can further progress to xerophthalmia, Bitot's spots, conjunctivitis, or keratomalacia, leading to blindness.
- Poor overall growth, anaemia
- Diarrhoea, susceptibility to infections
- Increased intracranial pressure
- Keratinization of mucous membranes and skin (hyperkeratosis)

Solution to Question 4:

The given clinical image is showing Bitot spots, which occur due to keratinization of the conjunctival epithelium. Bitot spots are triangular, frequently bilateral, and present on the bulbar conjunctiva.

Normally, bulbar conjunctiva is a stratified, columnar, non-keratinized epithelium. In vitamin A deficiency, there is squamous metaplasia of the conjunctival epithelium and tangles of keratin admixed with gas-forming bacteria producing a dry, scaly patch with a foamy appearance called Bitot spots.

Option A: Deposition of calcium in the conjunctiva is termed conjunctival concretions, which usually occurs over the palpebral conjunctiva, in chronic inflammatory conditions of the eye.

Option B: Scarring of the conjunctiva is seen in chlamydial infection of the eye.

Option C: Excessive growth of conjunctival tissue is seen in the pterygium of the eye.

Solution to Question 5:

The image depicts a corneal ulcer occupying $\frac{1}{3}$ of the cornea under a slit lamp with a fluorescein stain. This corresponds to stage X3A of vitamin A deficiency.

Solution to Question 6:

Both hypervitaminosis A and deficiency of vitamin A cause raised intracranial pressure (ICP) (not just hypervitaminosis A). 11-cis retinal(not trans) is required for vision.

Hypervitaminosis A causes headache, vomiting, desquamation of the skin, and hyperostosis of long bones. Rupture of lysosomes in the cells is the pathological feature of hypervitaminosis A.

All-trans retinoic acid is the most bioactive form of vitamin A, and it is required for cell differentiation and the regulation of transcription. On the other hand, 11-cis retinal is required for vision. Protein-energy malnutrition can cause vitamin A deficiency because of the impaired synthesis of retinol-binding protein (RBP).

Solution to Question 7:

Vitamin E/ Tocopherol is a potent antioxidant and is helpful in preventing hair fall. It promotes hair growth and prevents hair roots from getting damaged by free radical injury.

Alpha-tocopherol is the active component of vitamin E and it is mainly located within cell membranes, preventing lipid peroxidation and the formation of free radicals.

Deficiency of this can also lead to neurological defects, hemolysis, and posterior column defect.

Treatment includes vitamin E supplementation (25–50 units/day for a week).

Note:

The other vitamins having antioxidant properties are vitamins A, C, and E (vitamin E is the most important).

The other vitamin beneficial in the treatment of hair fall is biotin.

Solution to Question 8:

Craniotabes (soft skull), frontal bossing, stunted growth, and craniosynostosis are clinical features of rickets.

General features include failure to thrive, listlessness, protruding abdomen, proximal muscle weakness, and fractures.

Head to toe-examination would include the following features:

Head examination would include craniotables, frontal bossing, delayed fontanel closure, delayed dentition, and craniosynostosis.

Chest examination would reveal rachitic rosary, Harrison's sulcus, and repeated respiratory infections and atelectasis.

Spinal deformities could include kyphosis, scoliosis, lordosis.

Limb examination and extremities may show windswept deformity, anterior bowing of tibia, coxa vara, and enlargement of wrists and ankles. Bowings of legs (genu varum) or knock knees (genu valgum) may be noted. These changes can be seen only in children of the weight-bearing age group/ toddlers.


The image shows frontal bossing:



The image below shows the rachitic rosary:



The image below shows craniotables:



Solution to Question 9:

The clinical vignette with the above X-ray image shows fraying and cupping of the metaphysis, suggestive of rickets.

Radiological features in rickets:

- Fraying (Widening and irregularity of the growth plate)
- Cupping (Concavity of the metaphysis)
- Splaying at metaphysis (widening of metaphyseal ends)
- Generalized rarefaction

Solution to Question 10:

Windswept deformity (combination of valgus deformity of one leg with varus deformity of the other leg) is seen in rickets. It is also seen in rheumatoid arthritis but characteristically seen in rickets.

Other deformities of extremities seen in rickets:

- Enlargement of wrists and ankles
- Valgus or varus deformities of limbs
- Anterior bowing of the tibia and femur
- Coxa vara

Solution to Question 11:

Swelling of the costochondral junctions is seen in all the 3 conditions mentioned above.

Rachitic rosary	Scorbutic rosary
Beaded (rounded) in appearance	Angulated in appearance
Non-tender	Tenderness on palpation (sub periosteal haemorrhage)
Due to metaphyseal widening	Due to epiphyseal separation

Note:

Tietze syndrome is the benign inflammation of the costochondral junction.

Image: Given below is a photo of the chest cage with pectus excavatum.




Image: The above image shows a rachitic rosary type of appearance.

Solution to Question 12:

The child is suffering from rickets. Subperiosteal hemorrhage is not a radiographic sign of rickets. On the contrary, it is seen in scurvy.

Radiographic features of rickets:

- Widening of the growth plate: Decreased calcification leads to reduced bone density
- Fraying: The edge of the metaphysis loses its sharp border.
- Cupping: The edge of the metaphysis changes from a convex or flat surface to a more concave surface.
- Splaying: Weight-bearing and stress on the uncalcified bone gives rise to the widening of the metaphyseal end.
- Enlargement of the wrists and ankles due to the widening of the distal end of the metaphysis.
- Coarse trabeculation of the diaphysis
- Generalized rarefaction



Solution to Question 13:

In nutritional rickets serum, alkaline phosphatase levels are increased.

Biochemical findings in nutritional rickets (nutritional vitamin D deficiency):

- Normal or low calcium (since due to low calcium and vitamin D deficiency, PTH increases which will try to normalize the low calcium)

- Parathyroid hormone levels are elevated (in order to increase the serum calcium)
- Hypophosphatemia (because of increased parathyroid hormone [PTH])
- Metabolic acidosis (secondary to PTH induced renal bicarbonate wasting)

Solution to Question 14:

Alopecia and oligodontia (reduced number of teeth) with features of rickets in a child is classically seen in vitamin D-dependant rickets type 2 where the pathology is end-organ resistance (receptor hyposensitivity) to vitamin D.

Rickets causes

- Nutritional vitamin D deficiency
- Secondary vitamin D deficiency
- Vitamin D-dependant rickets, type 1 (defect in 1-alpha hydroxylase)
- Vitamin D-dependant rickets, type 2 (end organ resistance)
- Chronic kidney disease
- Hypophosphatemic rickets
- X-linked dominant hypophosphatemic rickets (PHEX gene mutation)
- Autosomal dominant (fibroblast growth factor [FGF] resistance to endopeptidases)
- Oncogenous rickets (increased FGF 23)

Note:

Vitamin D antibodies are found in most patients of systemic sclerosis.

Solution to Question 15:

Stoss therapy is used in the treatment of vitamin D deficiency.

There are 2 strategies for the administration of vitamin D:

- Stoss therapy: Here, 300,000-600,000 IU of vitamin D are administered orally or intramuscularly as 2 or 4 doses over 1 day.

•

The alternative strategy is daily vitamin D with a minimum dose of 2,000 IU/day for a minimum of 3 months.

Both strategies should be followed by a daily vitamin D intake of 400 IU/day if the child is <1-year-old or 600 IU/day if the child is ≥1-year-old. Children must also receive adequate dietary calcium and phosphorus.


Solution to Question 16:

This clinical scenario of a child with a history of treatment-resistant rickets, normal levels of calcium (9 to 11 mg/dL), parathyroid hormone, arterial blood gas (ABG), with decreased phosphate (normal, 2.5 to 4.3 mg/dL), and elevated levels of alkaline phosphatase (normal, 33 to 96 U/L) is suggestive of hypophosphatemic rickets.

Hypophosphatemic rickets:

X-linked dominant hypophosphatemic rickets is the most common genetic cause of rickets while the most common, overall is nutritional vitamin D deficiency. PHEX gene (phosphate-regulating endopeptidase on x chromosome) products are responsible for the inactivation of phosphatonin. Mutation of this PHEX leads to increased phosphatonin (FGF 23) levels, and hence, the excretion of phosphorous is increased. This is also associated with impaired renal tubular reabsorption of phosphate leading to its excess excretion.

The chart attached below summarizes the evaluation of hypophosphatemic rickets.



Treatment-resistant rickets :

Patients respond well to a combination of oral phosphorus and 1,25-D (calcitriol). Frequent dosing helps prevent prolonged decrements in serum phosphorus levels because there is a rapid decline after each dose. In addition, frequent dosing decreases diarrhea, which is a complication of high-dose oral phosphorus.

Solution to Question 17:

Vitamin K deficiency occurs mostly in breastfed infants not in artificial milk-fed babies, as the formula milk is fortified.

The use of broad-spectrum antibiotics that eliminate the intestinal vitamin K2- producing bacteria can cause vitamin K deficiency. Biliary atresia causes malabsorption of vitamin K, due to defective fat absorption.

There are 3 forms of vitamin K deficiency bleeding (VKDB) of the newborn:

Early VKBD of the newborn is prevented by the administration of a single intramuscular (IM) injection of vitamin K (1 mg) at birth.

	Secondary to maternal intake of medications	Early VKDB(classic hemorrhagic disease of the newborn)	Late VKDB
Onset	At birth	1 to 14 days	2 to 12 weeks
Cause	Drugs that cross the placenta and interferes with Vit K function eg. phenobarbital, phenytoin, warfarin	Low stores of Vitamin K at birth, poor transfer of Vitamin K across the placenta, Inadequate intake during the first few days of life.	In breastfed babies and in those with malabsorption of vitamin K (cystic fibrosis, biliary atresia)

Solution to Question 18:

In a child of age 4 years, the recommended dietary allowance (RDA) of vitamin D is 600 IU.

Note:

- Vitamin A: 1 IU = 1 IU retinol = 0.3 mcg of Retinol Equivalents (RE)
- Vitamin C: 1 IU = 50 mcg of ascorbic acid
- Vitamin D: 1 IU = 0.025 mcg cholecalciferol

Solution to Question 19:

Bowing of legs is not a feature of rickets in infants, because it appears only after weight-bearing. Thus, bow legs are a feature of rickets in toddlers.

Image: Rachitic rosary (widening of the costochondral junctions)



Image: Wrist X-ray image



Solution to Question 20:

Phosphatonin is a group of phosphate-regulating factors produced by osteocytes, and FGF 23 is the most important phosphatonin. Phosphatonin increases renal excretion (not reabsorption) of phosphate.

Mechanism of phosphatonin action:

Vitamin D receptor activation by 1,25 dihydroxy D (active vit D)



FGF 23 release by osteocytes



Reduces expression of sodium-phosphate transporter in proximal convoluted tubules of the kidney



Increased urinary phosphate excretion

Meanwhile, phosphatonin decreases the activity of 1 alpha hydroxylase and, therefore, the production of 1,25-D as well. So, the absorption of phosphate from the intestine is also reduced.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Deficiency of Water-soluble Vitamins & Trace Elements

Question 1:

Li Jing, a young tourist boy from the Keshan district of China visits India for a novel scientific exchange program. He presents to the ER with severe breathlessness. His previous medical records give the diagnosis of congestive cardiomyopathy. Which micronutrient deficiency is most likely seen in this case?

- a) Zinc
- b) Copper
- c) Magnesium
- d) Selenium

Question 2:

Anika, a 6-year-old girl good at sprint events, has been avoiding her track practices over the recent few weeks. Her mother informs that she skips her classes owing to her knee pain. Her X-ray reveals subperiosteal haemorrhages. What is the likely diagnosis?

- a) Vitamin C deficiency
- b) Vitamin B6 deficiency
- c) Vitamin B1 deficiency
- d) Vitamin B3 deficiency

Question 3:

A four-year-old boy presents with petechiae, perifollicular haemorrhages, and swollen, bleeding gums. On examination, you identify the child to be malnourished and suspect scurvy. Which of the following is/are part of biochemical functions of the vitamin deficient in this condition?

- a) 2, 3, 4
- b) 1 and 4 only
- c) 1, 3, 5
- d) 1, 2, 3, 4, 5

Question 4:

A child in your unit is receiving treatment for chronic diarrhea. His diet history is remarkable for a high phytate-rich diet. His labs reveal low serum zinc. Which of the following findings is least likely to be seen in this child?

- a) Dwarfism
- b) Incomplete development of gonads
- c) Dermatitis of extremities
- d) Predominant impairment of humoral immunity


Question 5:

The X-ray of a 7-year-old child diagnosed with scurvy showed the white line of Frankel. This would be best described as?

- a) Lucent metaphyseal band
- b) Calcified cartilage at metaphysis
- c) Opaque radiologic shadow surrounding epiphyseal centres of ossification
- d) Lateral prolongation of metaphyseal calcification

Question 6:

The eldest child of a ragpicker from the urban slums of Bandra presents with severe lower limb pain. Examination reveals swelling of costochondral junctions. The X-ray image obtained is given below. What is the most probable diagnosis?



- a) Scurvy
- b) Achondroplasia
- c) Osteomyelitis
- d) Rickets

Question 7:

A farmer from Hoshiarpur, Punjab rushes his 10-year-old boy to the paediatric ER. The father complained of intractable nausea and diarrhoea in the child. On examination, the child has a garlicky odour and is severely dehydrated. You notice neurologic manifestations along with nail and hair changes in the child. Which of the following mineral toxicities would you suspect?

- a) Selenium toxicity
- b) Chromium toxicity
- c) Molybdenum toxicity
- d) Iodine toxicity

Question 8:

A 2-month-old infant is brought to your OPD. The mother complains that the child refuses to feed and is not gaining adequate weight. She has recently noticed changes in her hair. On examination, the hair seems to resemble steel to wood. Labs reveal low serum copper levels. What is the diagnosis?

- a) De Vivo disease
- b) Menkes disease
- c) Keshan disease
- d) Kashin beek disease


Question 9:

The anxious mother of a 5-month-old baby informs you that the baby incessantly cries on touching or lifting. The baby is comfortable when it is made to lie down. Diet history is significant for the child being fed exclusively on cow's milk. What could be the probable cause?

- a) Scurvy
- b) JRA
- c) Rickets
- d) Caffey's disease

Question 10:

A child is born with the following abnormality. What advice do you give to the mother, with regards to the next pregnancy?



- a) 4 mcg of folic acid periconceptionally
- b) 400 mcg of folic acid periconceptionally
- c) 400 mcg of folic acid after confirmation of pregnancy

d) 4000 mcg of folic acid periconceptionally

Question 11:

A 10-year-old girl presented with the triad of diarrhoea, dermatitis, and delayed mental development. The child had neutral aminoaciduria and a positive Obermeyer test. Which of the following statement is false regarding the vitamin deficient in this child?

- a) Carcinoid syndrome can cause secondary deficiency of this vitamin
- b) Tyrosine is its precursor
- c) It decreases the levels of lipoprotein (a)
- d) Most fatal complication with high dose of this vitamin is fulminant hepatitis

Question 12:

A 7-month-old child was brought by her mother with complaints of diarrhoea and mucocutaneous lesions as shown in the image. The symptoms have been present since weaning was started. What is the diagnosis?



- a) Wiskott Aldrich syndrome
- b) Acrodermatitis enteropathica
- c) Pediatric epidermolysis bullosa
- d) Atopic dermatitis

Question 13:

A young school-going girl, a known case of seizure disorder controlled with valproic acid, complains of constant bullying by her classmates for her patchy hair loss. Local scalp examination reveals scaly dermatitis and hair loss with intermittent constrictions. Which of the following deficiencies is associated with this condition?

- a) Vitamin B3
- b) Vitamin H
- c) Vitamin B6
- d) Vitamin B1

Question 14:

A 2-day-old neonate is noted to have multifocal-clonic seizures. These seizures are unresponsive to pharmacologic therapy. Which vitamin dependency is responsible for such seizures?

- a) Pyridoxine
- b) Thiamine
- c) Niacin
- d) Folic acid


Question 15:

Cardiac edema and neuropathy are seen in which of the following vitamin deficiencies?

- a) Biotin
- b) Thiamine
- c) Pyridoxine
- d) Riboflavin

Question 16:

During the routine ophthalmic examination of a 6-year- old child as a part of a school health survey, you notice the following finding as shown in the image. This is characteristically associated with deficiency of _____



- a) Thiamine
- b) Riboflavin
- c) Biotin
- d) Pyridoxine

Answer Key

Question No.	Correct Option
1	d
2	a
3	c
4	d
5	b
6	a
7	a
8	b
9	a
10	d
11	b
12	b
13	b
14	a

15	b
16	b

Detailed Explanations

Solution to Question 1:

Keshan disease is due to the deficiency of selenium. It is a congestive cardiomyopathy that is caused due to dietary deficiency of selenium.

Keshan is a county of China where the symptoms were noted first. The disease is also endemic to that province. The patients will mostly present with acute breathlessness, paroxysms of exacerbated attacks.

Selenium is an enzyme cofactor for glutathione peroxidase, which prevents oxidative damage.

Note:

Kashin Beck disease is a chronic, endemic osteochondropathy due to selenium deficiency.

Solution to Question 2:

The clinical vignette is suggestive of pseudoparalysis which is a feature of scurvy - vitamin C deficiency

The subperiosteal haemorrhages in the lower limb bones can cause acute swelling and pain; the condition might mimic acute osteomyelitis or arthritis. The child refuses to walk because of the pain; hence, it is called pseudoparalysis.

Solution to Question 3:

Scurvy is a deficiency of vitamin C. The correct answer is 1, 3, 5.


Vitamin C is a water-soluble vitamin, which cannot be synthesized in the human body due to deficient L-gulonolactone oxidase enzyme. Breast milk contains adequate amounts of ascorbic acid, whereas the pasteurisation of animal milk at high temperature destroys vitamin C.

Functions of Vitamin C / Ascorbic Acid :

- Synthesis of collagen (Hydroxylation of proline, which is vital for collagen; in the formation of the bone matrix).
- Neurotransmitter metabolism (conversion of dopamine to norepinephrine and tryptophan to serotonin)
- Cholesterol metabolism (conversion of cholesterol to steroid hormones and bile acids)
- Biosynthesis of carnitine

- Antioxidation (electron donor)
- For the cellular and immunologic functions of the hematopoietic system

The image below shows peri-follicular haemorrhages.



Solution to Question 4:

The given clinical vignette is suggestive of chronic zinc deficiency. Zinc deficiency affects both cell-mediated and humoral immunity. The cell-mediated immunity is impaired predominantly, as zinc is essential for T-cell division, maturation, and differentiation.

In a child with a phytate-rich diet, a chronic deficiency of zinc and iron usually co-exist. Isolated chronic zinc deficiency can be seen in the setting of a child suffering from chronic diarrhoea, repeated pneumonia or chest infections, and congenital malabsorption disorders.

Chronic zinc deficiency manifests as:

- Dwarfism
- Hypogonadism
- Dermatitis of extremities and around orifices
- T-cell immunodeficiency
- Diarrhoea

The features of acute zinc deficiency include:

- Decreased growth
- Poor wound healing
- Diarrhea

Solution to Question 5:

Calcified cartilage at metaphysis is known as the white line of Frankel and it is seen in scurvy.

Solution to Question 6:

The clinical vignette involving a neglected child (eldest usually) from a poor socio-economic background with bone pains due to subperiosteal haemorrhages and scorbutic rosary (swelling of costochondral junctions) is suggestive of scurvy.

Radiological features of scurvy:

Early changes

- The white line of Frankel on the ends of the shafts of the tibia and fibula
- Wimberger ring sign: Sclerotic rings around the epiphyses of the femur and tibia.


Late changes

- Trummerfeld zone: Zones of destruction in the femur and tibia
- Pelkan spur at the cortical end

The image sums up the X-ray features noted in scurvy.



The image attached below shows a child with the scorbutic rosary.



Solution to Question 7:

The clinical vignette is suggestive of acute selenium toxicity.

Selenium toxicity is characterised by nausea, diarrhoea, neurologic manifestations, nail and hair changes, and garlicky odour. It is commonly seen in areas where drinking water and the collated soil is extremely rich in selenium. In India, selenium toxicity is endemic to agricultural sites located in the states of Punjab and Haryana. The presentations are usually acute in nature rather than chronic.

Chromium toxicity (Option - B) is commonly seen only in adults with occupational exposure. Chromium is a pulmonary irritant and the symptoms are predominantly respiratory complaints similar to bronchitis. They also cause chronic allergic dermatitis and is carcinogenic too.

Molybdenum toxicity (Option C) is very rarely seen in humans. It manifests usually as acute diarrhoea.

Iodine toxicity (Option D) can range from mild GI illness to chronic thyroid dysfunction varying from hyperthyroidism to papillary thyroid carcinoma.

Solution to Question 8:

The clinical vignette is suggestive of Menkes disease or Tricholipodystrophy or Steely hair disease caused by copper deficiency.

It is an X-linked recessive disorder caused by a mutation in ATP7A which encodes for a copper-transporting ATPase α polypeptide, leading to the disruption of copper transport in the intestine.

The clinical features include:

- Failure to thrive (First presenting complaint)


- Neurological degeneration (myeloneuropathy)
- Infections
- Hair abnormalities (kinky hair with pili troiti seen on a Trichogram).
- These infants have a characteristic cherubic appearance with a depressed nasal bridge, pudgy cheeks, 'cupids bow' upper lip, and doughy skin.

A therapeutic trial with copper-histidinase may be helpful.

Option A: De Vivo disease is a metabolic disorder due to GLUT1 deficiency.

Options C and D: Keshan disease (cardiomyopathy) and Kashin-beck disease (endemic osteochondropathy) are caused by selenium deficiency.

The image attached below shows the cherubic appearance of the child with kinky hair. The subsequent picture attached shows pili troiti seen under the trichogram.



Solution to Question 9:

The above scenario is suggestive of scurvy. A child who presents with a cry on touching or lifting is usually due to subperiosteal haemorrhages.

Scurvy/Barlow's Disease is caused due to the deficiency of vitamin c, especially in children fed predominantly on heat-treated cow's milk. High temperature destroys vitamin C.

JRA - Juvenile Rheumatoid arthritis (Option B) rarely presents in infancy.

Rickets (Option C) presents with growth retardation or other features such as craniotables, craniosynostosis in infancy. It is generally associated with individuals who have a milk deficient diet.

Caffey's disease (Option D) is infantile cortical hyperostosis. It usually presents at 6 months of age with painful soft tissue swelling and bone thickening. The mandible is the most common bone involved.

Solution to Question 10:

The given image shows spina bifida, the most common neural tube defect (NTD). 4000 mcg of folic acid periconceptionally is given to prevent NTDs in subsequent pregnancies.

The normal dose of folic acid in pregnancy is 400 mcg (0.4 mg). However, the history of a previous pregnancy with an NTD raises the risk of another NTD in subsequent pregnancies. An increased dose of folic acid is thus prophylactically given during the periconceptional period to prevent NTD in the next child.

Solution to Question 11:

The clinical vignette is suggestive of Pellagra secondary to Hartnup's disease. The vitamin deficient in this condition is niacin. The precursor of niacin is tryptophan and not tyrosine.

Niacin/vitamin B3 is a constituent of NAD and NADP. Its deficiency occurs in a maize-based diet (maize is deficient in tryptophan), carcinoid syndrome (all the tryptophan is converted to serotonin), and hartnup disease (defective absorption from intestine and kidney).

Niacin is the only lipid-lowering agent that lowers lipoprotein(a) levels significantly. It also results in a decrease in LDL, VLDL, and triglycerides along with an increase in HDL cholesterol.

Niacin deficiency causes pellagra (3Ds)-

- Diarrhoea
- Dermatitis (Cassal's necklace)
- Dementia
- Finally death.

Niacin toxicity manifests as:

- Prostaglandin mediated cutaneous flushing

- Glucose intolerance
- Fulminant hepatitis (most fatal complication)

Hartnup's disease is due to impaired renal and intestinal absorption of tryptophan and other neutral amino acids. It is characterized by a varied clinical presentation including ataxia (lack of muscle control), delayed mental development, diarrhoea and dermatitis (pellagra-like symptoms as tryptophan is a precursor of niacin). Diagnosis of niacin deficiency secondary to Hartnup's disease is based on neutral aminoaciduria and increased excretion of indole compounds detected by the Obermeyer test.

Image: Casal's necklace



Solution to Question 12:

A child with eczematous rash typically in the perioral and perineal region is characteristic of acrodermatitis enteropathica which is due to secondary zinc deficiency.

It is an autosomal recessive disorder seen in infants due to a defect in the intestinal zinc-specific transporter gene. Oral zinc therapy is the preferred management. It classically presents while weaning from breast milk to formula, which has lower zinc bioavailability than breast milk.

The symptoms are perioral and perineal rashes, periungual dermatitis- Beau's lines on nails, chronic diarrhea, and alopecia. Children commonly present with symmetrical, eczematous plaques that become vesicular, bullous, pustular, or erosive with characteristic crusting at the edges. The perioral eruption usually spares the upper lip, giving it a U-shaped or horseshoe-shaped appearance.

Investigations would reveal low serum zinc. Oral zinc therapy is the treatment of choice (3 mg/kg/24 hour)

Option A: Wiskott Aldrich syndrome is the classical triad of thrombocytopenia, eczema and recurrent pyogenic infections. No history of bleeding rules out this option.

Option C: Epidermolysis bullosa presents as bullous lesions in pressure handling areas.

Option D: In atopic dermatitis usually the lesions will be itchy and distributed in extensor aspects (flexural in adults).



Solution to Question 13:

Biotin (Vitamin H or Vitamin B7) deficiency can cause alopecia, dermatitis (periorificial), and thinning of hair. It is commonly seen in individuals who are on long term valproate therapy.

Scarring alopecia with islands of dense hair referred to as intermittent constrictions is classically seen in biotin deficiency.

Solution to Question 14:

Pyridoxine is one of the causes of refractory seizures in the neonatal period and infancy.

Pyridoxine and pyridoxal dependency disorders can cause severe seizures. These seizures, which are often multifocal clonic, usually start during the first few hours of life. It involves a mutation in the ALDH7A1 gene causing a deficiency of an enzyme involved in the degradation of I-alpha-aminoacidipic semialdehyde.

Solution to Question 15:

Cardiac edema and neuropathy are associated with the deficiency of thiamine (vitamin B1).

Early symptoms of thiamine deficiency include peripheral neuropathy, irritability, loss of deep tendon reflexes, and proprioception. As the disease progresses, increased intracranial pressure, meningismus, and coma develop.

Wet beriberi (cardiac) is associated with tachycardia, cardiomegaly, cardiac failure, and edema. It is commonly seen in the refugee camps where people consume polished rice based monotonous diets. Diagnosis of thiamine deficiency can be established by erythrocyte transketolase activity assessment.

Dry beriberi is associated with neurological deficits such as optic atrophy, ataxia, and irritability. Complications include lactic acidosis and Wernicke encephalopathy.

Solution to Question 16:

The given image shows corneal vascularization, which is characteristic of Riboflavin deficiency. Other features include glossitis, cheilosis, and photophobia collectively called ariboflavinosis.

The image below shows Cheilosis and glossitis



Fluid and Electrolyte Disorders

Question 1:

A 6-month-old infant was brought to the OPD by his mother with a history of passing watery stools more than 4 times in the past day. On examination, the baby is alert and on pinching his skin, it goes back quickly. The infant weighs 5.7 Kg. What is the initial fluid of choice for treating diarrhoea in this infant?

- a) Tender coconut water
- b) Oral rehydration solution (ORS)
- c) Salted yoghurt
- d) ReSoMal

Question 2:

The amount of total body water (TBW) in a term neonate weighing 3 Kg is approximately

- a) 1.2 L
- b) 1.8 L
- c) 2.25 L
- d) 2.75 L

Question 3:

What is the clinical procedure being performed?



- a) Pus drainage in acute osteomyelitis
- b) Femoral IV access for fluid transfusion
- c) Intraosseous fluid transfusion in shock
- d) Fixed suction catheter

Question 4:

A 9-year-old boy who has been operated on for craniopharyngioma presents with polyuria and polydipsia. On investigation, the serum osmolality is 305 mOsm/kg and urine osmolality is 250 mOsm/kg. The atrial natriuretic peptide (ANP) level is normal. What could be the cause?

- a) Post-operative diuresis
- b) Syndrome of inappropriate antidiuretic hormone secretion
- c) Post-surgical central diabetes insipidus
- d) Cerebral salt-wasting syndrome

Question 5:

All the following statements are true regarding zinc supplementation in children with diarrhoea except:

- a) It leads to reduced duration, severity, and recurrence of diarrhea
- b) It reduces the inappropriate use of antimicrobials
- c) Excessive zinc can cause copper deficiency syndrome

- d) It is given at a dose of 20 mg/day for 4 weeks

Question 6:

A 5-year-old child brought to the hospital with a history of loose stools. The mother says he is irritable and drinks water hastily when given. On examination, eyes are sunken and during the skin pinch test, the skin retracted within two seconds but not immediately. What is the treatment for the child?

- a) Administer the first dose of IV antibiotic and immediately refer to a higher center
- b) Give oral fluids and ask the mother to continue the same and visit again the next day
- c) Give zinc supplementation and oral rehydration solution only and ask mother to come back if some danger signs develop
- d) Consider severe dehydration, start intravenous (IV) fluids, IV antibiotics, and refer to a higher center

Question 7:

Which of the following is not true about reduced(low)-osmolarity oral rehydration solution (ORS)?

- a) It contains 20 mmol of potassium per litre
- b) Glucose is an essential component
- c) Total osmolarity is 254 mmol/litre
- d) Citrate is provided to prevent or correct base deficit acidosis

Question 8:

The amount of sodium present in ReSoMal is _____

- a) 60 mmol/L
- b) 75 mmol/L
- c) 45 mmol/L
- d) 85 mmol/L

Question 9:

A 2-year-old child weighing 7 Kg, presents to the emergency with a history of vomiting and diarrhea for the last 3 days. On examination, the skin pinch over the anterior abdominal wall goes quickly to its original position. What is the interpretation of the skin pinch test in this child?

- a) No dehydration
- b) Some dehydration
- c) Severe dehydration
- d) Skin pinch cannot be evaluated in this child

Question 10:

Which of the following is the most dangerous type of dehydration?

- a) Hyponatremic
- b) Hypernatremic
- c) Isonatremic
- d) Non-diarrheal causes

Question 11:

What is the maximum concentration of dextrose that can be given through a peripheral vascular line in a neonate?

- a) 5%
- b) 10%
- c) 12.5%
- d) 25%

Question 12:

A 7-year-old boy weighing 30 Kgs has been posted for surgical resection of a tumor. What is the daily maintenance fluid required for this child while he's NPO?

- a) 1000 mL
- b) 1700 mL
- c) 2500 mL
- d) 3000 mL

Question 13:

Which of the following is the most common type of shock in children?

- a) Cardiogenic shock
- b) Hypovolemic shock
- c) Obstructive shock
- d) Septic shock

Question 14:

A 3-year-old baby was brought by his parents to the ER. The parents complained that the child had been lethargic for a few hours and unresponsive for the past 30 mins. They give a history of 4-5 episodes of vomiting in the past one day. On examination, the baby had tachycardia, tachypnea, and very slow capillary refill time. The pediatrician asks you to initiate IV fluid therapy with a 24G IV cannula. Which colour IV cannula would you use?

- a) Pink
- b) Blue
- c) Yellow
- d) Purple


Question 15:

All of the following are causes of normal anion gap metabolic acidosis, except _____

- a) Renal tubular acidosis
- b) Diarrhea
- c) Acidosis due to intestinal bacterial overgrowth
- d) Urinary tract diversions

Question 16:

Rani, a 6-year-old girl, who is known to have thalassemia, presented to the emergency with weakness, chest pain, and palpitations. There is a history of recent blood transfusion. The ECG of the child shows tall and peaked T-waves and flat P-waves as shown below. What could be the cause?



- a) Acute pericarditis
- b) Iron-overload cardiomyopathy
- c) Pulmonary hypertension
- d) Hyperkalemia

Answer Key

Question No.	Correct Option
1	b
2	c
3	c
4	c
5	d
6	c
7	c
8	c
9	d
10	b
11	c
12	b
13	b
14	c

15	c
16	d

Detailed Explanations

Solution to Question 1:

In an infant with diarrhoea, the initial fluid of choice is oral rehydration solution (ORS).

The glucose-dependent sodium and water absorption is the principle behind replacing glucose and sodium in a 1:1 molar ratio in the WHO oral rehydration solution (ORS). The low osmolarity of the ORS in the lumen as compared to the blood facilitate absorption of fluid and hence the electrolytes due to the solvent drag.

Studies have shown that low osmolarity ORS causes reduction of stool output, decrease in vomiting and decrease in the use of unscheduled intravenous fluids without increasing the risk of hyponatremia. Other home available fluids that can be used are salted rice water/salted yoghurt drinks, vegetable/chicken soup with salt.

ReSoMal is given to severely malnourished children. In this case, since the infant's weight is appropriate for its age ORS is given.

Solution to Question 2:

The amount of total body water (TBW) in a term neonate is approximately 75% of the birth weight.

So, in this case, 75% of 3kg is 2.25 litres.

Note - TBW is 90% in early fetal life and becomes 60% by the end of the first year and stays the same till puberty.

Solution to Question 3:

The above image shows intraosseous cannulation for fluid transfusion in the anterior proximal tibia.

The intraosseous route is an alternative to the intravenous (IV) route in the transfusion of fluids in shock.

- The marrow of long bones has a rich network of vessels that drain into a central venous canal, emissary veins, and ultimately the central circulation.
- So, the bone marrow functions as a non-collapsible venous access route.

Intraosseous cannulation is recommended for patients in whom IV access proves difficult or unattainable. (e.g., in cardiopulmonary arrest). In such a case, if venous access is not available

within 1 min, an intraosseous needle should be placed in the anterior proximal tibia.

Insertion sites for intraosseous infusion:

- In the proximal tibia, 1-2 cm anteromedial from the tibial tuberosity
- Distal tibia at the junction of the medial malleolus and the shaft of the tibia
- Distal one-third of the femur

Solution to Question 4:

The given clinical scenario with ↑serum osmolality and ↓urinary osmolality, after a cranial surgery, suggests a diagnosis of post-surgical central diabetes insipidus (DI). Irreversible central DI follows most cases of complete resections of craniopharyngioma.

Serum osmolality of ≥ 295 mOsm/L (normal: 280-295 mOsm/L) and the urine osmolality of < 300 mOsm/L (normal: 500-850 mOsm/L) are suggestive of DI. A water deprivation test is done to further classify the types of DI.

Option A: In postoperative patients, because of IV fluids there is an acute expansion of ECF volume. This suppresses high ADH activity and causes diuresis with normal serum osmolality.

Option B: In SIADH, serum osmolality is decreased.

Option D: In cerebral salt-wasting syndrome there is hypersecretion ANP and their levels are usually elevated.

Solution to Question 5:

Zinc supplementation is usually given for a duration of 10 to 14 days.

- Age < 6 months: 10 mg/day
- Age ≥ 6 months: 20 mg/day

Zinc administration for diarrhoea management can significantly reduce mortality and hospital admission.

In addition, it improves diarrhoea recovery rates and reduces the inappropriate use of antimicrobials.

Excessive zinc therapy can lead to copper deficiency syndrome, hence requires the regular monitoring of plasma zinc and copper concentrations.

Solution to Question 6:

The child in this scenario has some degree of dehydration, since the child is restless and irritable, drinks water readily, has sunken eyes, and the skin goes back slowly on the skin pinch test (< 2 seconds) with sunken eyes.

Treatment includes oral rehydration therapy and zinc supplementation according to WHO Integrated Management of Neonatal and Childhood Diseases (IMNCI) Protocol, Plan B.

WHO IMNCI PLANS:

SIGNS	CLASSIFY AS	IDENTIFY TREATMENT
Two of the following signs: - Lethargic or unconscious - Sunken eyes - Not able to drink or drinking poorly - Skin pinch goes back very slowly	SEVERE DEHYDRATION	(Urgent pre-referral treatments are in bold print) - If child has no other severe classification - Give fluid for severe dehydration (Plan C). - OR - If child also has another severe classification: - Refer URGENTLY to hospital with mother giving frequent sips of ORS on the way. Advise the mother to continue breastfeeding. - If child is 2 years or older, and there is cholera in your area, give antibiotic for cholera.
Two of the following signs: - Restless, irritable - Sunken eyes - Drinks eagerly, thirsty - Skin pinch goes back slowly.	SOME DEHYDRATION	- Give fluid, Zinc supplements and food for some dehydration (Plan B) If Child also has a severe classification: - Refer URGENTLY to hospital with mother giving frequent sips of ORS on the way. Advise the mother to continue breastfeeding. - Advise mother when to return immediately. - Follow-up in 5 days if not improving. - If confirmed/symptomatic HIV, follow-up in 2 days if not improving.
Not enough signs to classify as some or severe dehydration	NO DEHYDRATION	- Give fluid, Zinc supplements and food to treat diarrhoea at home (Plan A) - Advise mother when to return immediately. - Follow-up in 5 days if not improving. - If confirmed/symptomatic HIV, follow-up in 2 days if not improving.

Does the child have diarrhoea? IF YES, ASK: LOOK AND FEEL: - For how long? - Look at the child's general condition. Is the child: - Lethargic or unconscious? Restless and irritable? - Look for sunken eyes. - Offer the child fluid. Is the child: - Not able to drink or drinking poorly? Drinking eagerly, thirsty? - Pinch the skin of the abdomen. Does it go back: - Very slowly (longer than 2 seconds)? Slowly?		for DEHYDRATION Classify DIARRHOEA		Two of the following signs: - Lethargic or unconscious - Sunken eyes - Not able to drink or drinking poorly - Skin pinch goes back very slowly	SEVERE DEHYDRATION	- If child has no other severe classification: - Give fluid for severe dehydration (Plan C). - OR - If child also has another severe classification: - Refer URGENTLY to hospital with mother giving frequent sips of ORS on the way. Advise the mother to continue breastfeeding. - If child is 2 years or older and there is cholera in your area, give antibiotic for cholera.
				Two of the following signs: - Restless, irritable - Sunken eyes - Drinks eagerly, thirsty - Skin pinch goes back slowly	SOME DEHYDRATION	- Give fluid, zinc supplements and food for some dehydration (Plan B) If child also has a severe classification: - Refer URGENTLY to hospital with mother giving frequent sips of ORS on the way. Advise the mother to continue breastfeeding. - Advise mother when to return immediately.
				Not enough signs to classify as some or severe dehydration.	NO DEHYDRATION	- Give fluid, zinc supplements and food to treat diarrhoea at home (Plan A) - Advise mother when to return immediately.
		and if diarrhoea 14 days or more		Dehydration present.	SEVERE PERSISTENT DIARRHOEA	- Treat dehydration before referral unless the child has another severe classification. - Refer to hospital.
				No dehydration.	PERSISTENT DIARRHOEA	- Advise the mother on feeding a child who has PERSISTENT DIARRHOEA. - Give multivitamin and minerals (including zinc) for 14 days. - Follow-up in 5 days.
		and if blood in stool		Blood in the stool.	BLOOD IN STOOL	- Treat for 5 days with an oral antimicrobial recommended for Shigella in your area. Treat dehydration and give zinc - Follow-up in 2 days.

Solution to Question 7:

The total osmolarity of low-osmolarity oral rehydration solution (ORS) is 245 mmol/L, and it is the most commonly used ORS.

- Potassium is provided to replace the large potassium losses, thus preventing serious hypokalemia.
- Citrate corrects base-deficit acidosis.
- Glucose promotes the absorption of sodium and water in the small intestine.

Solution to Question 8:

Rehydration solution for malnourished (ReSoMal) contains 45 mmol/L of sodium.

Severely malnourished children have excess intracellular sodium (leaky cell membranes and reduced activity of the sodium–potassium pump). So, sodium intake should be reduced and a low-sodium oral rehydration solution (ReSoMal) is advised. Because severely malnourished children are deficient in potassium, ReSoMal contains extra potassium, as well as extra glucose or sucrose and selected minerals (Mg, Cu, Zn).

Solution to Question 9:

In the given scenario, the child is malnourished, so the skin pinch cannot be evaluated.

The child is of age 2 years, and the expected weight is 12 Kg. But, the weight of the child here is 7 Kg, which suggests that the child is malnourished. The child may be having edematous malnutrition (kwashiorkor), where skin may go back to the original position even in the presence of dehydration. Hence, the skin pinch is not reliable for the assessment of dehydration in this child.

Reliable indicators of dehydration in the malnourished child include

- Eagerness to drink
- Cool and moist extremities
- Weak radial pulse
- Reduced urine flow

Solution to Question 10:

Hypernatremic dehydration is the most dangerous type of dehydration.

Hypernatremia causes movement of water from brain cells into hypertonic extracellular fluid, causing brain cell shrinkage and shearing of blood vessels within brain. This leads to central nervous system hemorrhages and thrombosis.

The fluid deficit is corrected slowly as rapid treatment of hypernatremic dehydration causes movement of water from the extracellular space into brain cells, leading to cerebral edema, seizures, brain herniation, and even death.

Solution to Question 11:

Through a peripheral vascular line, a maximum of 12.5% dextrose can be infused.

The dextrose concentration greater than 12.5% has an acidic pH and can cause irritation to the peripheral veins, which can lead to thrombophlebitis.

Note:

Central vein lines can be used for 25% D if needed.

Solution to Question 12:

The daily maintenance fluid required for a child weighing 30 Kg is 1700 ml.

Bodyweight method for calculating the daily maintenance fluid volume (also known as Holliday-Segar method) is used.

Calculation:

In the question, the weight of the child is given as 30 Kgs.

So, the formula for >20 Kg is used.

$1500 \text{ mL} + (20 \text{ mL} \times \text{weight more than } 20 \text{ Kgs})$

Since the child weight is 30 Kgs, he/she is 10 Kgs more.

Therefore, fluid required is

$= 1500 \text{ mL} + (20 \text{ mL} \times 10)$

$= 1500 \text{ mL} + 200 \text{ mL}$

$= 1700 \text{ mL}$.

Bodyweight	Fluid required per day
0–10 Kg	100 mL/Kg
11–20 Kg	$1000 \text{ mL} + 50 \text{ mL/Kg for each Kg } >10 \text{ Kg}$
$>20 \text{ Kg}$	$1500 \text{ mL} + 20 \text{ mL/Kg for each Kg } >20 \text{ kg}$

Solution to Question 13:

Hypovolemic shock is the most common type of shock in children.

Cardiogenic shock: cardiac pump failure due to cardiomyopathies, myocarditis or congenital heart diseases.

Hypovolemic shock: loss of body fluid due to diarrhea, vomiting or hemorrhage.

Obstructive shock: mechanical barrier which obstructs the cardiac outflow like pericardial tamponade, tension pneumothorax, pulmonary embolism.

Septic shock: It is the combination of hypovolemic (intravascular fluid loss because of capillary leak) and cardiogenic shock (because of myocardial depression by circulating inflammatory mediators) in association with infection.

Solution to Question 14:

The color of the 24 G cannula is yellow. 24 G cannula is usually used for paediatric age group.

Solution to Question 15:

Acidosis due to intestinal bacterial overgrowth causes high anion gap metabolic acidosis because of excessive bacterial metabolism of glucose into lactic acid.

Anion gap

Normally, in the serum, the total number of cations is equal to the total number of anions.

But, few of the anions (lactate, phosphate, and sulphate) cannot be measured and this is called an anion gap.

Anion gap = $[Na^+] - ([Cl^-] + [HCO_3^-])$

Metabolic acidosis

Metabolic acidosis is of two types:


(1) Normal anion gap metabolic acidosis (NAGMA)

- Diarrhea (Loss of bicarbonate is compensated by chloride reabsorption)
- Renal tubular acidosis
- Urinary tract diversions (reabsorption of ammonium and hydrogen by the bowel)

(2) High anion-gap metabolic acidosis (HAGMA)

- Lactic acidosis (hypoxia, malignancy)
- Ketoacidosis (diabetic, starvation)
- Poisoning (ethylene glycol, methanol, salicylate)

Anion gap



©Marrow

Solution to Question 16:

The given electrocardiograph (ECG) characteristics show tall and peaked T-waves, flat P waves and since the girl has a recent history of massive blood transfusion, the most likely diagnosis is hyperkalemia.

Hyperkalemia may be due to blood transfusion (stored blood has increased potassium concentration, due to the failure of the $\text{Na}^+ - \text{K}^+$ ATPase pump).

Causes of Hyperkalemia:

- Spurious laboratory value
- Sampling error
- Thrombocytosis
- Leucocytosis
- Increased intake
- Oral or intravenous
- Blood transfusions
- Transcellular shifts
- Acidosis
- Tumor lysis syndrome
- Hemolysis
- Decreased excretion
- Renal failure
- Hyporeninemic hypoaldosteronism


- Drugs
- ACE inhibitors
- ARB (angiotensin receptor blockers)
- Potassium-sparing diuretics

ECG Changes of Hyperkalemia

- Tall and peaked T waves
- Increased PR interval
- Flattening of the P wave
- ST-segment depression
- Widening of the QRS complex

Note:

Tall T waves are also seen in myocardial ischemia.



Option A:

- There will be history of fever and ECG shows diffuse ST elevation.

Option B:

- Patient presents with features of cardiac failure and ECG shows ventricular arrhythmias.


Option C:

- Pulmonary hypertension is a known complication of hemolytic diseases (free Hb, angiotrophic factors causing vasoconstriction) and ECG shows right ventricular hypertrophy (RVH).

Mendelian and Non-Mendelian Disorders

Question 1:


A pregnant female arrives at your hospital for her antenatal check-up. As an intern, you were required to draw her family pedigree chart while taking history. She has previously delivered full-term healthy dizygotic twins 5 years back. Which of the following symbol will you use to denote dizygotic twins in a pedigree chart?



- a) Option A
- b) Option B
- c) Option C
- d) Option D

Question 2:

Which pattern of inheritance does the pedigree chart denote?



©MARROW

- a) Autosomal-dominant disorder
- b) X- linked dominant disorder
- c) Autosomal-recessive disorder
- d) X- linked recessive disorder

Question 3:

While volunteering at the blood donation camp, Ananya came across a case of a thalassemic infant who has an AB-positive blood group. Her father is B-positive and her mother is A-positive. In order to understand more about donor eligibility to explain the concerned parents, she wishes to understand the inheritance pattern of ABO blood grouping. Can you help her out by choosing the correct option?

- a) X-linked inheritance
- b) Recessive inheritance
- c) Mitochondrial inheritance
- d) Codominance

Question 4:

An MBBS student, after attending a class on genetic disorders, noticed that a particular disorder appears in three consecutive generations of his family without any sex predilection. While drawing his pedigree chart, it also struck him that phenotypically normal family members were having healthy offspring. Based on this information, he concluded the pattern of inheritance of this disorder. What is the correct pattern?

- a) Autosomal recessive
- b) Autosomal dominant
- c) Mitochondrial inheritance
- d) Uniparental disomy

Question 5:

A young boy presented with a history of a genetic disorder. On further enquiry, his father and grandmother had the same disease as well. Prior to his birth, the doctor had counseled the father that there is a 50% chance of passing the deleterious gene to his offspring irrespective of sex. Which of the following diseases is an example of this inheritance pattern?

- a) Albinism
- b) Sickle cell anemia
- c) Thalassemia
- d) Tuberous sclerosis

Question 6:

Aryan is the first child born to parents suffering from achondroplasia. Although he is not suffering from this condition, the parents wish to know the chances of developing achondroplasia in the second child. What is the correct response?

- a) 0%
- b) 75%
- c) 25%
- d) 100%

Question 7:


A 3-year-old boy presented with short stature and delayed motor milestones. An x-ray showed a trident hand as given below. Which of the following is the most likely diagnosis?



- a) Achondroplasia
- b) Mucopolysaccharidosis
- c) Osteopetrosis
- d) Cleidocranial dysostosis

Question 8:

A 14-year-old child presented with multiple hyperpigmented lesions of around size 15-20 mm in the trunk and extremities. Similar lesions of size 2-3 mm were found in his axillary and inguinal regions. A photograph of his eye showing a characteristic lesion is shown below. What is the likely diagnosis?



- a) Tuberous sclerosis
- b) Sturge-Weber syndrome
- c) Neurofibromatosis
- d) Wilson disease

Question 9:

Which one of the following is not included in the criteria for neurofibromatosis 1?

- a) Two or more iris hamartomas
- b) Acoustic neuromas
- c) Dysplasia of the sphenoidal and tibial bone
- d) Cafe-au-lait spots

Question 10:

Consider the following statements about café- au- lait macules associated with neurofibromas:

- a) Statements A, C and E true, B and D false
- b) Statements B and C true, A, D and E false
- c) Statements B, D and E true, A and C false
- d) Statements A and B true, C, D and E false

Question 11:

A 9-month-old child presents with a history of a condition seen in the image below. On examination, a characteristic raised skin lesion of orange peel consistency known as shagreen patch is present in his lumbosacral region. Which one of the following is not seen in this autosomal dominant neurocutaneous syndrome?



- a) Astrocytoma
- b) Rhabdomyoma
- c) Optic glioma
- d) Subependymal nodule

Question 12:

The first child of a normal healthy couple was diagnosed to have cystic fibrosis. The parents are anxious and wish to seek advice for future pregnancies. What is the percentage of chances for the second child to be affected?

- a) 50 %
- b) 0%
- c) 25%
- d) 100 %

Question 13:

A 29-year-old woman is homozygous for an autosomal-recessive gene and her husband is heterozygous for the same. They are planning to start a family and wish to know the outcome of their pregnancy. Which out of the following options correctly represent the percentages of the outcome?

- a) 75% children affected
- b) No child affected, but all are carriers

- c) 50% children affected, rest are carriers
- d) 25% children affected, rest are carriers

Question 14:

What is the mode of inheritance of Menke's disease?

- a) X- linked recessive
- b) X- linked dominant
- c) Autosomal recessive
- d) Autosomal dominant

Question 15:

A 41-year-old father of two has an X-linked genetic disorder. His daughter is a carrier for the same and his son is unaffected. His wife is healthy with no inherited disease. Which of the following is an example of this disorder?

- a) Wiskott-Aldrich syndrome
- b) Rett Syndrome
- c) Thalassemia
- d) Alkaptonuria

Question 16:

Which of the following is an X-linked disorder?

- a) Color blindness
- b) Thalassemia
- c) Sickle cell anemia
- d) Cystic fibrosis

Question 17:

Which of the following is not true regarding ataxia-telangiectasia?

- a) Mutations in 11q gene is implicated

- b) Follows autosomal dominant mode of inheritance
- c) Humoral and cellular immunodeficiency
- d) Linked to adenocarcinomas

Question 18:

A 12-year-old boy presented with progressive muscular weakness showing the following finding. It was found that the child is unable to get up from a squatting position. He was diagnosed with a hereditary muscular disorder. What is the mode of inheritance in this disorder?



- a) Autosomal dominant
- b) X- linked recessive
- c) Mitochondrial inheritance
- d) X- linked dominant

Question 19:


Your second cousin, Prakriti, has been recently diagnosed with Wiskott-Aldrich syndrome. Your relatives are worried and keep calling you to confirm the new information that they read on the internet. In an attempt to settle their doubts about the credibility of online resources, you inform them that all of the following are true except:

- a) X- linked recessive mode of inheritance
- b) Atopic dermatitis and eczema during infancy

- c) Thrombocytopenia
- d) IgM is increased

Question 20:


A 5-year-old girl has been diagnosed with a genetic disorder that follows a Mendelian pattern of inheritance illustrated by the pedigree chart below. Which of the following can be a likely diagnosis?



- a) Wiskott-Aldrich syndrome
- b) Lesh-Nyhan syndrome
- c) Hypophosphatemic rickets
- d) Fabry disease

Question 21:

A 3-year-old girl presented with poor growth and social interaction as well as midline stereotypical movements of the hands as seen in the image below. All milestones were normal for the first 2 years. After that, the milestones delayed and her development slowed down. On examination, her gait was slightly clumsy, her speech had regressed to mere babbling and her anthropometry was below the 50th percentile. Following a battery of investigations, the diagnosis of a rare inherited disease was made. What is the affected gene for this condition?



- a) P53
- b) Rb
- c) MeCP2
- d) FGFR3

Question 22:

While reviewing a junior's research manuscript on trinucleotide repeat sequences, you come across the following list of examples and spot an error. Which of the following diseases will you mark for omission?

- a) Huntington's disease
- b) Friedreich's ataxia
- c) Myotonic dystrophy
- d) Hereditary motor-sensory neuropathy

Question 23:

A 3-year-old child presents with previously diagnosed fragile X syndrome. As an intern posted in the department, you are asked to follow up on his cytogenetic analysis report. What is the trinucleotide repeat mutation seen in the report?

- a) CGG
- b) GAA
- c) CAG

d) CTG


Question 24:

An 8-year-old boy was brought for psychiatric evaluation for intellectual disability. On examination, his IQ was 45 and a previous report in his records show a trinucleotide repeat mutation of CGG. Which of the following is not seen in it?

- a) Large head
- b) Large nose
- c) Large ear
- d) Large testis

Question 25:

As a participant in the Indian Academy of Paediatrics quiz, you are asked to mark the type of inheritance in MELAS. The pedigree chart shown in the image below is provided to explain which family members may be affected (coloured in red). Which of the following will you choose?



- a) X- linked Recessive
- b) Autosomal Recessive
- c) Mitochondrial
- d) X- linked Dominant

Question 26:

In order to understand the topic 'Non-Mendelian disorders', you have prepared flashcards. But, while browsing through them before your viva, the order of cards have been jumbled up. Which of the following option will you choose to correctly match the phenomena with their definitions?

- a) 1-C, 2-A, 3-D, 4-B
- b) 1-A, 2-B, 3-C, 4-D
- c) 1-B, 2-D, 3-A, 4-C
- d) 1-D, 2-C, 3-B, 4-A

Question 27:

A 6-month-old presented with a genetic disorder attributed to multifactorial inheritance. This type of inheritance is most likely to play a significant role in which of the following disorder?

- a) Achondroplasia
- b) Lysosomal storage disease
- c) Cleft lip
- d) Huntington disease

Answer Key

Question No.	Correct Option
1	b
2	c
3	d
4	b
5	d
6	b
7	a
8	c
9	b
10	d
11	c

12	c
13	c
14	a
15	a
16	a
17	b
18	b
19	d
20	c
21	c
22	d
23	a
24	b
25	c
26	a
27	c

Detailed Explanations

Solution to Question 1:

Pedigree charts are graphical representations of a person's family tree and are useful for identifying the pattern of inheritance of genetic disorders. Thus, they are vital in the history-taking of patients, especially in pregnancy. Option B is the symbol denoting dizygotic twins in a pedigree chart.

Option A denotes monozygotic twins in the pedigree chart.

Option C denotes offsprings.

Option D denotes consanguineous marriage.

Solution to Question 2:

The above pedigree chart denotes an autosomal recessive pattern. In the above pedigree chart, both males and females are carriers (equally affected) which indicates an autosomal pattern. Parents are unaffected (both are carriers/heterozygotes), and no affected members in other generations. 25% of the children are normal, 50% are carriers, and only 25% are affected. This indicates a recessive pattern.

Pedigree charts in various modes of inheritance:

Solution to Question 3:

The inheritance pattern of the ABO blood grouping system is codominance.

Codominance:

Gene expression and mendelian traits are usually described as dominant or recessive; in some cases, both of the alleles of a gene pair contribute to the phenotype—a condition called codominance. Examples include histocompatibility and ABO blood group antigens.

Classification of genetic disorders

Mendelian disorders (Classic inheritance)

- Autosomal dominant
- Autosomal recessive
- X-linked dominant
- X-linked recessive

Non-mendelian disorders (Non-classic inheritance)

- Trinucleotide repeat mutations
- Fragile X syndrome
- Friedreich ataxia
- Huntington's disease
- Mutations in mitochondrial genes: MELAS, MERRF
- Genomic imprinting
- Angelman syndrome
- Prader–Willi syndrome
- Silver–Russell syndrome
- Gonadal mosaicism

Chromosomal disorders:

- Microdeletions: Cri du chat syndrome
- Down's syndrome, Turner's syndrome, Klinefelter's syndrome


Solution to Question 4:

The pattern of inheritance in this genetic disorder is autosomal dominant.

Points in favour:

- Autosomal dominant disorders have a vertical mode of transmission (individuals of two generations are affected—parent and children affected).

- It appears in three consecutive generations of a family without any sex predilection.
- An affected individual has a 50% chance of passing on the deleterious gene in each pregnancy.
- Males and females are equally affected.
- Unaffected individuals do not pass on the disorder to their children.



Solution to Question 5:

Autosomal dominance is the inheritance pattern seen in this question due to three-generation of affected family members with a 50% chance of passing the deleterious gene to the next generation irrespective of sex. Tuberous sclerosis is an example of an autosomal dominant (AD) disorder.

Note:

- All Ehlers–Danlos syndrome types are autosomal dominant, except types V, VI, and VII.
- All types of osteogenesis imperfecta are autosomal dominant, except types II, III, and VII.

Solution to Question 6:


The chances of having an affected baby when both parents have achondroplasia (autosomal-dominant disorder) is 75%.

- Autosomal dominant (AD) disorders are manifested in the heterozygous state.
- If both of the parents are affected, the chances of having an affected baby are 75% (50% - 'Aa', and 25% - 'AA').

Note:

Here, we consider both parents as 'Aa' (heterozygous) because 'AA' (homozygous) is usually fatal in the first few months of life. The odds of having an affected baby does not change with

subsequent pregnancies.



Solution to Question 7:

The x-ray shows short stubby fingers, with a separation between middle and ring fingers. This is called a trident hand and is seen in achondroplasia.

Achondroplasia

It is an autosomal-dominant chondrodysplasia and the most common condition characterized by disproportionate short stature. It is caused by mutations in the gene coding for fibroblast growth factor receptor type 3 (FGFR3).

Clinical features (see image below):

Short stature, extreme joint laxity, bowing of legs, macrocephaly, frontal bossing, midfacial hypoplasia, small chest, redundant skin folds, recurrent infections, cor pulmonale, and restrictive pulmonary disease



Radiological features

- Champagne glass pelvic inlet
- Trident hand
- Shortening of tubular bones with metaphyseal flaring.
- Chevron deformity

Solution to Question 8:

The features of café-au-lait macules, axillary and inguinal freckling along with Lisch nodules points toward a diagnosis of neurofibromatosis-1 (NF-1).

The multiple hyperpigmented lesions of more than 15 in the trunk and extremities are known as café-au-lait macules ($>5\text{mm}$ in prepubertal, $>15\text{mm}$ in postpubertal) and are present in 100% of NF1 cases. The lesions of around 2-3mm in the axillary and inguinal regions are known as axillary/inguinal freckling and are pathognomonic of NF1. The given image shows discrete nodules over the iris, which are referred to as Lisch nodules, and are commonly seen in patients with NF-1. Lisch nodules are iris hamartomas.

Neurofibromatosis: It is an autosomal-dominant neurocutaneous syndrome with the following types-

Axillary freckling is shown in the image below-

NF1 (von Recklinghausen disease)	NF2 (Central neurofibromatosis)
Mutation on chromosome17	Mutation on chromosome22

NF1 (von Recklinghausen disease)	NF2 (Central neurofibromatosis)
2 or more of the following should be present: 6 or more café-au-lait spots 2 or more neurofibromas or one plexiform neurofibroma - Freckling in the axillary or inguinal region 2 or more Lisch nodules - Dysplasia of the sphenoidal and tibial bone - A first degree relative with NF1 - Optic gliomas	Predisposes to multiple tumours of the nervous system. Most common are bilateral vestibular schwannomas; intracranial and spinal meningiomas, gliomas, neurofibromas, and posterior subcapsular lenticular opacities. Vestibular schwannomas are benign tumours that arise from the intracanalicular segment of the inferior division of the vestibulocochlear nerve. Extracanalicular extension into the cerebellopontine angle can lead to an "ice cream cone" appearance in MRI.



Solution to Question 9:

In neurofibromatosis 1 (NF1), two or more of the following clinical manifestations are present:

- Six or more café-au-lait macules
- Axillary or inguinal freckling
- Two or more iris Lisch nodules (hamartomas of iris)
- Two or more neurofibromas or 1 plexiform neurofibroma
- A distinctive osseous lesion such as sphenoidal and tibial bone dysplasia
- Optic gliomas

- A 1st-degree relative with NF-1

Solution to Question 10:

Statement C: The spots are scattered over the body surface, with a predilection for the trunk and extremities; sparing the face.

Statement D: For diagnosis, the macules should be ≥ 5 mm in prepubertal and ≥ 15 mm in postpubertal individuals.

Statement E: 6 or more spots should be present for the clinical diagnosis of neurofibromatosis

Café-au-lait macules:

- Hallmark of neurofibromatosis, present in around 100% of the patients.
- 6 or more café-au-lait spots should be present for the clinical diagnosis of neurofibromatosis along with other criteria.
- Present at birth, but increase in size, number, and pigmentation, especially during the first few years of life.

Note:

Other diseases associated with multiple café-au-lait spots: Ataxia telangiectasia, Bloom syndrome, Fanconi's anaemia, McCune–Albright syndrome, tuberous sclerosis, Maffucci syndrome, Hunter syndrome.

Other neurocutaneous syndromes:

- Sturge–Weber syndrome: port-wine stain over the trigeminal distribution, contralateral hemiparesis, focal seizures contralateral to the side of facial capillary malformations, calcification of the cerebral cortex, mental retardation, and congenital glaucoma
- von Hippel–Lindau disease: retinal angiomas; cerebellar and spinal hemangioblastomas
- Ataxia telangiectasia
- Tuberous sclerosis

Café-au-lait spots are shown in the image given below:



Solution to Question 11:

Optic glioma is not seen in tuberous sclerosis. Tuberous sclerosis/Bourneville–Pringle Disease is an autosomal-dominant neurocutaneous syndrome, which may present during infancy with infantile spasms (as seen in the image), and has the characteristic skin lesion known as shagreen patch.

It is characterized by Vogt's triad (EPILOA), which includes

- Epilepsy
- Low intelligence (IQ)
- Adenoma sebaceum

The most common manifestations are skin lesions such as ashleaf (hypomelanotic) macules, adenoma sebaceum, shagreen patch, confetti lesions, and periungual fibromas. Koenen's tumours (periungual fibromas) are small fleshy tumors that grow under the toe nails/finger nails.

Cortical tubers and subependymal nodules are the characteristic brain lesions. Other lesions include bilateral renal angiomyolipomas, facial angiofibroma, cardiac rhabdomyoma, and retinal hamartomas.

Vigabatrin is the first-line therapy for infantile spasms in tuberous sclerosis.

Note: Drug of choice for infantile spasms: adrenocorticotrophic hormone (ACTH)

Image below: Adenoma sebaceum



Image below: Shagreen patch



Image below: Periungual fibroma



Solution to Question 12:

Cystic fibrosis follows autosomal recessive inheritance. Hence, the disease has a 25% chance of affecting the offspring.

If parents are unaffected (both are carriers/heterozygotes):


- 25% chance of normal genotype
- 50% chance of carriers
- 25% chance of homozygous

In autosomal recessive disorders, the disease manifests itself when the offspring has a homozygous state, which means each child has a 25% chance of being affected by the disease.

Note:

There are 2 alleles for this, say for example "A" = normal allele and "a" = abnormal allele. So, if each parent is phenotypically normal and their child is affected, then they are carriers, i.e., "Aa". So, each parent has a 0.5 (50%) probability of passing the affected allele to the next generation. Also, having each child is an independent event, and the chance that the child receives an abnormal allele from the father (0.5), as well as a mother (0.5), is equal to $0.5 \times 0.5 = 0.25$ i.e. 25%.

Autosomal recessive




Solution to Question 13:

If the husband is heterozygous and the wife is homozygous for an autosomal recessive gene, 50% of the children will be affected; the rest will be carriers.

This is pseudo-dominant inheritance, which refers to the observation of apparent dominant (parent to child) transmission of a known autosomal-recessive disorder, or it can also mean the inheritance of a recessive trait mimics a dominant pattern.

This occurs when a homozygous affected individual has a partner who is a heterozygous carrier, and their children have a 1 in 2 chance to get affected (50%).



©MARROW

Solution to Question 14:


The mode of inheritance of Menke's disease is X-linked recessive.

X-linked recessive disorders have the following features:

- Males are affected more commonly and more severely than females.
- Female carriers have a 25% risk for having an affected son, a 25% risk for a carrier daughter, and a 50% chance of having a child that does not inherit the mutated X-linked gene.
- The affected father will have all daughters (100%) to be carriers and unaffected sons because they pass their X chromosome to all of their daughters and their Y chromosome to all of their sons.

X- linked recessive pattern of inheritance is shown in the image given below:

Kinky hair disease (Menkes disease)	Wilson's disease
X- linked recessive	Autosomal recessive
Mutation of copper-transporting, P-type adenosine triphosphatase 7A leading to copper deficiency	Mutation of adenosine triphosphatase 7B on chromosome 13q14 leading to excessive copper accumulation.
Low serum copper and ceruloplasmin levels	Low ceruloplasmin levels; low serum-free copper levels and increased urinary copper excretion



Solution to Question 15:

The mode of inheritance is X- linked recessive and Wiskott-Aldrich syndrome is an example of such a genetic disorder.


Solution to Question 16:

Color blindness is primarily inherited as an X-linked recessive trait. The genes for color vision are located on the X chromosome.

Males have a single X chromosome inherited from mothers. If the X chromosome inherited carries a mutation, males exhibit the disease. However, females become carriers of the color blindness gene and have a 50% chance of passing it on to their children.

Female carriers usually are protected because of the random inactivation of one X chromosome. But there is a possibility that the mutant X chromosome remains active if the normal alleles are inactivated in some of the cells so the heterozygous female can express the disorder partially.

The image shows the pattern of X-linked inheritance:



In color blindness, the mechanism to appreciate one or more primary colors is either defective (anomalous) or absent (anopia). It may be congenital or acquired. Congenital color blindness can be dyschromatopsia (defective color vision of any one of the RGB colors) or achromatopsia (complete color blindness of all three RGB colors). RGB colors are red, green, and blue.

Tests for color blindness include the Ishihara test, Hardy–Rand–Rittler test, and Farnsworth–Munsell 100-hue test.

Solution to Question 17:

Ataxia telangiectasia (AT) or Louis–Bar syndrome is an autosomal recessive disorder.

The defect in AT is in the ATM gene localized to chromosome 11q22-23. The ATM gene is involved in the repair of double-strand DNA breaks. This defect in DNA repair leads to combined immunodeficiency (impaired immunoglobulin isotype switching and association with thymic hypoplasia) and predisposition to an increased risk of lymphoreticular tumors such as lymphoma or leukemia.

AT is characterized by progressive ataxia (beginning during infancy), telangiectasia (initially on bulbar conjunctiva), sinopulmonary infections, hypogonadism, sensitivity to ionizing radiation, insulin resistance, and raised alpha-fetoprotein levels.

Solution to Question 18:

This clinical vignette includes an image of a child with pseudohypertrophy of calf muscles along with a history of inability to get up from a squatting position. This is characteristically seen in Duchenne muscular dystrophy, which is an X-linked recessive disorder.

Salient features of Duchenne muscular dystrophy:

- Most common hereditary neuromuscular disease.
- X-linked recessive mutations of dystrophin gene on chromosome Xp21.

Clinical features:


- Progressive weakness, developmental delay, and intellectual impairment
- In toddlers: delayed walking, falling, toe walking, and trouble running or walking upstairs
- Common sites of muscular hypertrophy: calf & tongue & forearm
- Pseudohypertrophy of calf muscles due to fibrofatty replacement of muscle
- Gower's sign: Patients use their arms and hands to walk up their own body from a squatting position due to the reduced strength in the proximal muscles(hip & thigh) of the lower limbs.
- Trendelenburg gait/hip waddle
- Inability to walk develops by the age of 10-14 years.
- Complications: Contractures, scoliosis, cardiomyopathy, malignant hyperthermia after anaesthesia.
- Death occurs by 3rd decade of life due to respiratory failure.

Note:

Becker's Muscular Dystrophy (BMD) vs. Duchenne Muscular Dystrophy (DMD)

Mnemonic-

- BMD: Badly makes dystrophin (defective)
- DMD: Does not make dystrophin (absent)



Solution to Question 19:

IgM is decreased in Wiskott–Aldrich syndrome, not increased. It is a rare genetic immunodeficiency that keeps a child's immune system from functioning properly. It also makes it difficult for a child's bone marrow to produce platelets, making a child prone to bleeding. This syndrome is associated with decreased IgM, increased IgA, and increased IgE levels.

Wiskott-Aldrich syndrome:

Mnemonic WAXTIE

- WA: Wiskott–Aldrich
- X: X-linked recessive
- T: Thrombocytopenia
- I: Infections
- E: Eczema

Solution to Question 20:

The given pedigree chart illustrates an inheritance pattern that is suggestive of an X-linked dominant disorder. Hence, hypophosphatemic rickets, which is an X-linked dominant disorder, is a likely diagnosis here.

X-linked dominant disorders:

- Females are affected twice as often as males.
- An affected female has a 50% probability of transmitting the disorders to her son or daughter.
- An affected male transmits the disorder to all of his daughters and none of his sons.

- Some X-linked dominant conditions are lethal in a high percentage of males, e.g., incontinentia pigmenti.

Other options:

- Lesh–Nyhan syndrome: X-linked recessive disorder affects young boys due to the deficiency of HGPRT enzyme in the purine pathway. It is marked by compulsive self-mutilation of the head and hands, together with learning difficulties and involuntary muscular movements.
- Fabry disease: X-linked recessive lysosomal storage disease due to the deficiency of alpha-galactosidase A. Affected males have angiokeratomas, hypohidrosis, acroparesthesias, and corneal-lenticular opacities.

Note:

Mnemonic to remember X-linked dominant disorders: Ribbons For All Children

- Ribbons - Rett's syndrome/ Hypophosphatemic Rickets
- For - Fragile X tremor/ataxia syndrome (FXTAS)
- All - Alport syndrome
- Children - Charcot–Marie–Tooth disease

Solution to Question 21:

The given features of poor growth & social interaction, gradual loss of speech, clumsy gait along with the characteristic hand movements point towards a likely diagnosis of Rett's syndrome. The gene involved in Rett's syndrome is the MeCP2 gene located on the X chromosome (Xq28 predominantly of paternal origin).

Rett's syndrome:

- Almost always affects girls (X-linked dominant)
- Development may proceed normally until 1 year of age slowing down to the extent of regression.
- After 1 year of age, rapid regression in language and hand skills, motor milestones, and acquired microcephaly become apparent.
- Clinical features can include repetitive hand wringing movements, ataxia, breathing dysfunction, bruxism, scoliosis, and profound mental retardation.
- Autistic behaviour is a typical finding in all patients.

Note:

- Rb: Retinoblastoma: 13 chromosome
- P53: Li–Fraumeni syndrome, Sarcomas: 17
- NF1: Neurofibromatosis 1-17
- NF2: Neurofibromatosis 2-22
- WT1: Wilms's tumour: 11
- FGFR3: Achondroplasia

Solution to Question 22:

Hereditary motor-sensory neuropathy, commonly known as Charcot-Marie-Tooth neuropathy, is a type of X-linked dominant disorder. The rest of the options are associated with trinucleotide repeat sequences.

Triple-repeat mutations (Allelic expansion disorders).

- Due to an increase in the number of nucleotide repeats above a certain threshold.
- Length of the nucleotide repeat often correlates with the severity of the disease.
- Anticipation: When repeat length increases from one generation to the next, disease manifestations may worsen or be observed at an earlier age.

Disease	Repeats	Clinical features
Fragile X- syndrome	CGG	Mental retardation, large ears, and jaws, macroorchidism in males
Myotonic Dystrophy(chromosome 19)	CTG	Muscle loss, cardiac arrhythmia, cataracts, frontal balding
Huntington's disease	CAG	Affective disorder
Spinocerebellar ataxia type 1	CAG	Progressive ataxia, dysarthria, dysmetria.
Friedreich's ataxia	GAA	Progressive ataxia, dysarthria, hypertrophic cardiomyopathy.

Solution to Question 23:

The trinucleotide repeat mutation associated with fragile X syndrome is CGG in the familial mental retardation -1 (FMR1) gene.

Other common trinucleotide repeat mutations and their associated conditions:

- GAA- Friedreich ataxia
- CAG- Huntington disease
- CTG- Myotonic dystrophy

Solution to Question 24:

In fragile-X syndrome, there would be a small nose, not large.

- Fragile X syndrome or Martin–Bell syndrome: Trinucleotide repeat mutation of CGG.

Remember mnemonic Hello Mr. FRAGILE X Syndrome:

- Hello: Hyperextensible joint or high-arched palate
- Mr: Mitral valve prolapse
- F: FMR1 P deficiency (fragile X mental retardation 1 protein) - upregulates the metabotropic glutamate receptor (mGluR) 5 pathway and alters the expression of matrix metalloproteinase (MMP) 9.
- R: Repeating the sequence of 3 nucleotides CGG, recurrent otitis media.
- A: Autism/Attention deficit–hyperactivity disorder/Anxiety
- G: Giant brain (macrocephaly) and giant testis (macroorchidism)
- I: Intellectual disability, i.e., mental retardation
- L: Long face with a large mandible
- E: Everted ear
- X: X-linked disorder (controversial as to whether it is X-linked dominant or recessive)

The most common genetic cause of mental retardation is Down's syndrome > Fragile X Syndrome.

Sherman's paradox: Risk of expressing mental retardation in fragile X syndrome increases in later generations (anticipation).

Fragile X-associated tremor or ataxia: Intention tremors and cerebellar ataxia, which may progress to parkinsonism.

Note:

Inhibitors of the mGluR (overexpressed in fragile X) are undergoing clinical trials. Minocycline (lowers MMP9) has resulted in short-term improvements in anxiety, mood, and the Clinical Global Impression Scale.

Image: A child with affected fragile X- syndrome; note the everted ears, long face, and large mandible.




Solution to Question 25:

The type of inheritance in MELAS is mitochondrial, which manifests as mitochondrial myopathy, encephalopathy, lactic acidosis, and stroke-like episodes.

Mitochondrial inheritance:

- Inherited exclusively from the affected mother (ovum) as an individual's mitochondrial genome is entirely derived from the mother because sperms contain few mitochondria, which are typically shed upon fertilization.
- Both male and female children born to an affected mother will inherit the disease.
- 80% of mitochondrial DNA codes for mitochondrial RNA, and therefore most mitochondrial DNA mutations lead to functional problems, which may be manifested as muscle disorders.

The following pedigree shows the mitochondrial inheritance pattern-



Solution to Question 26:

The phenomena are correctly matched with their definitions as follows-

- Genomic imprinting: Phenotypic expression of a gene depending on the parent of origin.
- Mosaicism: Single organism possessing more than one genetic line as the result of genetic mutation.
- Anticipation: Symptoms become more severe and/or appear at an earlier age as the disorder is passed from one generation to the next.
- Non-penetrance: A genetic trait that is present in the genome but does not manifest itself in the individual.

Genomic imprinting: epigenetic process, that selectively inactivates either the maternal or paternal allele. These defects include:

- Beckwith- Wiedemann syndrome
- Silver Russell syndrome
- Angelman syndrome
- Prader Willi syndrome

Note:

Uniparental disomy: This refers to the situation in which two copies of a chromosome come from the same parent, instead of one from each parent. Eg., Angelman syndrome and Prader Willi syndrome.

	Prader–Willi syndrome	Angelman syndrome
Deletions (70% cases)	Deletion of Paternal genes after maternal gene silencing	Deletion of Maternal genes after paternal gene silencing
Uniparental disomy (UPD) (30%)	UPD of maternal chromosome 15	UPD of paternal chromosome 15

Solution to Question 27:

Multifactorial inheritance plays a significant role in the appearance of cleft lip.

Multifactorial inheritance- It refers to traits that are caused by a combination of inherited, environmental, and stochastic factors. These include the following:

Children (congenital malformation)	Adults
Pyloric stenosis Neural tube defects Congenital heart defects Congenital dislocation of the hip Cleft lip and cleft palate	Diabetes Mellitus Hypertension Coronary heart disease Gout Schizophrenia

Chromosomal Disorders

Question 1:

On watching the documentary 'The Secret of Our Lives: The Human Genome Project', a student reads up about the international consortium of scientists who fully deciphered the DNA of the most common chromosomal syndrome. What is this syndrome?

- a) Fragile-X- syndrome
- b) Trisomy 18
- c) Trisomy 21
- d) Trisomy 13

Question 2:

A 39-year-old pregnant woman was found to have an abnormal nuchal translucency scan. An amniocentesis done revealed a chromosomal karyotype of 47, XX,+21. This result most commonly occurs due to?

- a) Reciprocal translocation
- b) Non-disjunction in maternal meiosis
- c) Mosaicism
- d) Non-disjunction in paternal meiosis

Question 3:

Your aunt has sent you an image of her 1-year-old son's palm using an instant messaging app. She read online that it is mostly present in genetic disorders and wishes to get your opinion. You identify it as a simian crease and prepare a list of differentials. All of the following diagnoses are correct except:



- a) Down syndrome
- b) Trisomy 13
- c) Cri du chat syndrome
- d) Cystic fibrosis

Question 4:

A girl is born at term after an uneventful pregnancy and delivery. On examination, she is noted to be hypotonic, has a protruding tongue, low set ears, an upward slant to the eyes with epicanthal folds, and a single palmar crease. On cardiac evaluation, what is the most common lesion expected in this syndrome?

- a) Ventricular septal defect
- b) Endocardial cushion defect
- c) Tetralogy of Fallot
- d) Coarctation of aorta

Question 5:

A 41-year-old entrepreneur is excited to start her family. She has been getting a lot of unsolicited advice about the complications of pregnancy in advanced maternal age. On enquiring about such pregnancies, the doctor advised the mother to get prenatal testing done. All are findings associated with the prenatal diagnosis for Down's syndrome except

- a) Reduced femur and humerus length
- b) Nuchal translucency ≥ 3 mm
- c) Increased umbilical blood flow
- d) Ventricular septal defect may be present


Question 6:

An elderly primigravida underwent a nuchal translucency scan at 14 weeks which was highly suggestive of Down's syndrome. The doctor advised her to undergo a triple test and called her back at week 16. On receiving her receipt for the payment of the test, she realises that she has been billed wrongly for a component that is not included in the triple test. What is this component?

- a) β -hCG
- b) Alpha-fetoprotein
- c) Inhibin A
- d) Serum oestriol level

Question 7:

While writing your medical competitive exams, you come across the following image of karyotyping analysis in a question. You recognize this image as being highly suggestive of a particular syndrome. What is the diagnosis?



- a) Patau syndrome

- b) Edward's syndrome
- c) Alagile syndrome
- d) William's syndrome

Question 8:

A 10-month-old boy presented with facial dysmorphism as well as two characteristic findings shown in the picture below. What is the condition affecting this newborn?



- a) Edward's syndrome
- b) Turner's syndrome
- c) Alagile syndrome
- d) William's syndrome

Question 9:

A young girl presents to you with learning disabilities. On examination, you notice a characteristic finding as shown in the picture below. Reports of her investigation scans mention horseshoe kidneys and streak gonads. What is your diagnosis?



- a) Klinefelter's syndrome
- b) Noonan's syndrome
- c) Turner's syndrome
- d) Fragile X- syndrome

Question 10:

Kaveri recently became pregnant following a history of repeated abortions. She was anxious and wanted to undergo karyotyping this time to detect chromosomal disorders pre-natally. Barr body was found to be absent in her report. Which of the following is associated with an absent Barr body?

- a) 46 XX
- b) 45 Xo
- c) 47 XXX
- d) 47 XXY

Question 11:


A 15-year-old adolescent presents with an anti-mongoloid slant, pulmonary stenosis, short stature, webbed neck, and undescended testis. What is the likely diagnosis?

- a) Hypoparathyroidism
- b) Noonan syndrome
- c) Klinefelter syndrome

d) Turner syndrome

Question 12:

A concerned pregnant woman underwent karyotyping which was highly suggestive of a particular syndrome. The image of her results is shown below. Which of the following abnormalities is associated with patients with the given karyotype?



- a) Gynecomastia with long thin limbs
- b) Short stature with webbed neck
- c) High-pitched cry
- d) Round face with protruding tongue

Question 13:

A newborn with previously diagnosed Cri-du-chat syndrome presented to the hospital. Which of the following is not observed in this condition?

- a) Hypertonia
- b) Characteristic shrill cry
- c) Microcephaly
- d) Mental retardation

Question 14:

A senior asks you to read about an autosomal dominant condition known as Treacher- Collins syndrome. While browsing through numerous journals to learn about the topic, you come across a doubt about its features. Which of the following is not associated with this syndrome?

- a) Coloboma of inferior lid
- b) Mandibular hypoplasia
- c) Low intelligence
- d) Cleft palate

Question 15:

All of the following are seen in a 5-month-old child with Pierre- Robin syndrome except?

- a) Hearing defects
- b) Coloboma iris
- c) Respiratory distress
- d) Mandibular hypoplasia


Question 16:

Shantanu, an avid reader, and a first-year resident, recently finished Joseph Heller's famous novel 'Catch-22'. While discussing the book with his friend, he realized that the title was an acronym of a chromosomal disorder they had studied. Which of the following is the least likely feature of this disorder?

- a) Thymic aplasia
- b) Hypocalcemia
- c) Cleft palate
- d) Cranial anomalies

Question 17:

Riya was found to have a height of more than 95 percentile for her age and an arm-span to height ratio of more than 1.05. On further examination, she had flat feet, mild scoliosis, and long slender fingers. An image of her hands showing two characteristic signs is provided below. What other complications can be expected in this child?



- a) Pulmonary hypertension
- b) Mental retardation
- c) Emphysema
- d) Subluxation of lens

Question 18:

Venkatesh decided to present a rare case report of maternal disomy of chromosome 15 at a scientific conference. He collected the following image representing the disorder's features. What is this disorder?



- a) Prader- Willi syndrome

- b) Klinefelter's syndrome
- c) Angelman syndrome
- d) Turner's syndrome

Question 19:

Triangular facies is indicative of which of the following diagnosis?

- a) Russell-Silver syndrome
- b) Down's syndrome
- c) Mobius syndrome
- d) Nephrotic syndrome

Question 20:

During an intersectoral collaborative event, an award-winning geneticist from the National Human Genome Research Institute is invited for an endowment oration. He presents his lecture on an experimental form of treatment that uses the transfer of genetic material to cure diseases known as 'Gene Therapy'. Which of the following techniques does he mention for this therapy?

- a) Electroporation
- b) Electrofocusing
- c) Electrophoresis
- d) Magnetoreception

Question 21:

The following is the finding noted in a 12-month-old baby who is growing well and developing normally since birth. Which of the following is a true statement regarding this finding?



- a) Down's syndrome is the diagnosis
- b) Maternal history of phenytoin intake during pregnancy should be suspected
- c) Reassurance should be provided
- d) Chromosomal analysis should be done to rule out other chromosomal abnormalities

Question 22:

Non-invasive prenatal screening (NIPS) is particularly useful in the modern era of diagnosis in clinical genetics. Which of the following statements is not true regarding this test?

- a) Useful in screening for aneuploidies
- b) Evaluates maternal blood for fetal abnormalities
- c) High positive predictive value
- d) Positive test need confirmation by invasive testing

Answer Key

Question No.	Correct Option
1	c
2	b
3	d
4	b

5	c
6	c
7	a
8	a
9	c
10	b
11	b
12	a
13	a
14	c
15	b
16	d
17	d
18	a
19	a
20	a
21	c
22	c

Detailed Explanations

Solution to Question 1:

Trisomy 21 (Down's syndrome) is the most common chromosomal syndrome. It is the second human chromosome whose DNA has been fully deciphered by the Human Genome Project.

Chromosomal disorders: These may involve autosomal or sex chromosomes.

Note:

- Most common trisomy leading to miscarriage/incompatibility with life: Trisomy 16.
- F body: Y chromosome (quinacrine, a fluorescence dye, binds strongly to the Y chromosome forming a bright fluorescent spot called an F body).

Involving autosomes	Involving sex chromosomes
Down's syndrome (Trisomy 21) Edward syndrome (Trisomy 18) Patau syndrome (Trisomy 13) Cri du chat syndrome (5p deletion)	Klinefelter's syndrome (47, XXY) Turner syndrome (45, Xo)

Solution to Question 2:


The chromosomal karyotype of 47, XX,+21 is suggestive of Down's syndrome which occurs due to nondisjunction in maternal meiosis. Non-disjunction points to the failure of the homologous chromosome to separate during the meiotic division.

Down's syndrome:

- Most common chromosomal disorder
- Incidence: 1/800 live births
- Mechanism:
- Trisomy 21 (meiotic maternal nondisjunction) (95%)
- Robertsonian translocation: 4%
- Mosaic pattern: 1%

Note:

- Robertsonian translocations: involve 2 acrocentric chromosomes (chromosomes 13, 14, 15, 21, and 22) that fuse near the centromeric region, with a subsequent loss of the short arms.



Solution to Question 3:

Simian crease (single transverse palmar crease) is not seen in cystic fibrosis.

Simian crease is seen in the following disorders:

- Down's syndrome
- Fetal alcohol syndrome

- Trisomy 13
- Cri du chat syndrome

Solution to Question 4:

The girl is suffering from Down's syndrome and the most common cardiovascular lesion is an endocardial cushion defect.

Other associated defects are atrial septum defects (ASD), ventricular septum defect (VSD), patent ductus arteriosus (PDA), aberrant subclavian artery, and pulmonary hypertension.

Clinical features: (Mnemonic: I C A PROBLEM Somewhere)

- Intellectual disability/Incurved or short fifth finger with clinodactyly
- Congenital heart disease
- Acute megakaryoblastic leukaemia/Alzheimer's disease/Atlantoaxial instability/Atresia of the duodenum
- Protruding tongue
- Round face
- Occiput flat/Open wide fontanelle/Open mouth
- Brachycephalic/Brushfield's spots
- Low (depressed) nasal bridge
- Epicanthal fold/Ears low set and dysplastic
- Mongoloid slant (oblique palpebral fissure)
- Simian palmar crease and wider gap between first and second toes

Note:

- The most common cause of intestinal obstruction in Down syndrome is duodenal atresia (double-bubble sign).
- Children with Down's syndrome have an increased risk of developing leukaemias, including acute lymphoblastic leukaemia (ALL) and acute myeloid leukaemia (AML).
- AML is more common during the first 3 years of life.
- The most common AML associated with Down Syndrome is acute megakaryoblastic leukemia.
- The most common haematological manifestation in Downs syndrome is transient abnormal myelopoiesis (TAM).
- Brushfield's spots are whitish speckling on iris seen in light-skinned people.

Image: A child having features of Down's syndrome



Image: Brushfield's spots



Solution to Question 5:

Increased umbilical flow is not seen during the prenatal screening for Down's syndrome, instead, the flow is decreased.

Reduced blood flow through the umbilical vessels and abnormal blood flow through ductus venosus (DV) in the first trimester are associated with chromosomal abnormalities, especially Down's syndrome. This decrease in blood flow causes IUGR in infants born with Down syndrome.

Solution to Question 6:

Inhibin A is not included in the triple test:

Note:

- AFP: Maternal serum alpha-fetoprotein
- HCG: Beta-human chorionic gonadotrophin
- PAPP-A: Pregnancy-associated plasma protein A
- UE₃: Unconjugated estriol

Test	Components
Triple test	AFP + hCG + UE ₃
Quadruple test	Triple test + Inhibin A
Combined test(first-trimester screening)	PAPP-A + hCG + nuchal translucency
Integrated test(most sensitive)	Combined test + Quadruple test

Solution to Question 7:

The above karyotype is suggestive of Patau syndrome (Trisomy 13).

Clinical features of Trisomy 13/Patau syndrome (Mnemonic CMC DOCToR OPD)

- Cardiac defects (ventricular septal defect/atrial septal defect/patent ductus arteriosus)
- Microcephaly, Microphthalmia
- Cleft lip/palate
- Developmental delay
- Ocular hypotelorism
- Holoprosencephaly
- Thin posterior ribs (missing ribs)
- Renal abnormalities
- Polydactyly with flexed fingers
- Polycystic kidneys
- Scalp defects (aplasia cutis)

Image: 37-week male with Patau syndrome demonstrating alobar holoprosencephaly with cyclopia. Facial features included a sloping forehead with a nose superior to a single central palpebral fissure.



Solution to Question 8:

The newborn with features of facial dysmorphism, index finger overlapping other fingers, and rocker bottom feet are suggestive of Edward's syndrome (Trisomy 18).

Clinical features of Edward's syndrome: (Remember mnemonic: Hi Mr EDWARDS)

- Hypertonia
- Microcephaly/Micrognathia
- Eighteen trisomy/Elongated skull
- Digit overlapping with a closed fist
- Webbing of 2nd and 3rd toes
- Absent intellect (severe developmental delay)/Absent radii/ Arthrogryposis
- Rocker bottom feet
- Cardiac Disorders (ventricular septal defect, patent ductus arteriosus, atrial septal defect) and kidney, death.
- Small lower jaw (Micrognathia), short sternum, small nipples

Note: P, Patau syndrome: Trisomy 13

E, Edward's syndrome: Trisomy 18

D, Down syndrome: Trisomy 21

Alagille syndrome (AS) is an autosomal-dominant disorder associated with abnormalities of the liver (hypoplasia of hepatic duct), heart (tetralogy of Fallot), skeleton, eye, and kidneys and a characteristic facial appearance.

William's syndrome is a developmental disorder that is characterized by mild to moderate intellectual disability or learning problems, unique personality characteristics, distinctive facial

features, and cardiovascular problems.

Solution to Question 9:

The child shows characteristic webbed neck, learning disabilities, horseshoe kidneys, and streak gonads and is diagnosed with Turner's syndrome.

Turner's syndrome:

- Karyotype, 45; XO, phenotypically females
- Chromosomal abnormality XO is present in 60% of individuals, mosaic pattern in 40%
- Intelligence remains normal

Important features associated with Turner's syndrome: (mnemonic: See A Baby CLOWN Here)

- Short stature, Short fourth metacarpal, Sensorineural hearing loss, Shield chest
- Amenorrhea (primary), infertility
- Barr body absent
- Cardiac defects, Cubitus valgus, Congenital lymphedema
- L: Learning disabilities
- Ovaries underdeveloped (streak ovaries), Gonadoblastoma (increased risk if Y chromosome material is present)
- Webbed neck
- Nipples widely spaced
- Horseshoe kidneys

Note:

The most common cardiac defect in Turner's is bicuspid aortic valve > coarctation of aorta.

Solution to Question 10:

Barr body is absent in female having 45 Xo genome.

Barr body

- Densely staining inactivated condensed X chromosome present in each somatic cell of a female is found in the nucleus.
- Chromatid body (Barr body or sex chromatin) is derived from one of the two X chromosomes which become inactivated.
- The number of Barr bodies is thus one less than the number of X chromosomes.

Turner's syndrome is female without Barr body.

Klinefelter's syndrome is male with Barr body.

Note:

- Lyon hypothesis states that only one of the two X chromosomes in a female is functional, the other becomes inactive during embryogenesis.
- Davidson bodies are small nuclear buds of chromatin shaped like a drumstick found in neutrophils of females.

Solution to Question 11:

The adolescent presenting with antimongoloid slant, pulmonary stenosis, short stature, webbed neck, and undescended testis is suggestive of Noonan syndrome. Although the webbed neck is a feature of Turner syndrome as well, the likely diagnosis would be Noonan syndrome since the patient is male (undescended testis).

Noonan syndrome:

- Autosomal-dominant disorder (12q)
- Male or female with Turner phenotype; normal karyotype.

Clinical features:


- Down-slanting palpebral fissure (antimongoloid slant)
- Pectus carinatum
- Hypertelorism
- Cryptorchidism
- Low levels of clotting factor 11 and 12
- Leukemia (Juvenile chronic myelogenous leukemia)
- Moderate mental retardation in 25%
- Cardiac abnormality: Most common is pulmonary valve stenosis.
- Hypertrophic cardiomyopathy
- Atrial Septal Defect
- Coarctation of aorta
- Short stature, webbed neck, low set ears, cubitus valgus: similarities with Turner syndrome
- Ptosis, micrognathia, clinodactyly, delayed puberty.

Difference between Noonan syndrome and Turner syndrome:

Noonan syndrome	Turner syndrome
Normal karyotype(46XX/46XY)	45, Xo(60%)
Most common heart disease: Valvular pulmonary stenosis	Most common heart disease: Bicuspid aortic valve
25% have mental retardation	Mental retardation rare

Noonan syndrome	Turner syndrome
Genitourinary:Female: Fertile, delayed pubertyMale: Infertile due to cryptorchidism	Genitourinary:Female: Infertile, streak ovariesNo males

Solution to Question 12:



The above karyotype - 47, XXY, is suggestive of Klinefelter's syndrome, in which gynecomastia and long thin limbs are seen.

Klinefelter's syndrome:

- Most common sex aneuploidy in males
- Classically karyotype is 47, XXY. Other variants can have 48 XXXY; rarely, 49 XXXXY; or mosaics with some cells containing normal 46, XY and others 47, XXY
- Results from meiotic nondisjunction of sex chromosomes (mostly during 1st meiotic division)

Clinical features:

- Disproportionately long arms and legs
- Mitral valve prolapse
- Mental retardation; IQ: 60-80
- Minor developmental and learning disabilities
- Lack of secondary sexual characteristics (due to a decrease in androgen production)
- Gynecomastia
- Higher risk of breast carcinoma
- Testicular dysgenesis

- Infertility/ azoospermia

Note:

A variant of Klinefelter's syndrome—49, XXXXY—presents with short stature.

Solution to Question 13:

Hypotonia is seen in cri-du-chat syndrome, not hypertonia.

Cri-du-chat syndrome occurs due to the deletion of chromosome 5p.

Main features:

-
- Hypotonia
- Characteristic shrill cry (cat-like cry) in the first few weeks of life
- Microcephaly
- Hypertelorism
- Bilateral epicanthic folds
- High-arched palate
- Wide and flat nasal bridge
- Intellectual disability

Note:

- Cat-bite infection is caused by *Pasteurella multocida*.
- Tomcat urine odor is seen in multiple carboxylase deficiencies.

Solution to Question 14:

Intelligence is not affected in Treacher–Collins syndrome (like in Turners syndrome).

Treacher–Collins syndrome/Franceschetti syndrome

- Autosomal dominant
- Facial appearance: downward sloping palpebral fissures, colobomas and drooping of the lower eyelids, sunken cheekbones, deformed pinnae, atypical hair growth extending towards cheeks
- Facial clefts, malformed ears, deafness
- Hypoplastic mandible

Image: Treacher–Collins syndrome; note the sunken cheekbones, abnormality of ears, and hypoplastic mandible



Solution to Question 15:

Coloboma iris is not a feature of Pierre–Robin syndrome.

Features of Pierre–Robin syndrome-

- Micrognathia accompanied by a high-arched or cleft palate
- Tongue normal size; the floor of the mouth is foreshortened
- Retrognathia: retracted mandible
- Glossoptosis/Retroglossoptosis: tongue appears to fall into the throat (retraction of the tongue)

Of the children with Pierre–Robin syndrome, 30–50% have Stickler syndrome. Features include prominent joints, arthritis, hypotonia, hypermobile joints, mitral valve prolapse, hearing loss, and ocular problems (myopia, glaucoma, cataracts, retinal detachment).

Pierre–Robin syndrome as shown in the image below:



Solution to Question 16:

DiGeorge syndrome, also known as Catch-22 syndrome, is associated with cardiac anomalies and not cranial anomalies.

DiGeorge/Shprintzen Velocardiofacial syndrome - Remember the mnemonic CATCH 22

- C: Cardiac anomalies
- A: Abnormal facies
- T: Thymic aplasia and agenesis of parathyroid glands.
- C: Cleft palate
- H: Hypocalcemia, hypoplasia of auricle
- 22: 22q11.2 deletion

Solution to Question 17:

The given features are suggestive of a diagnosis of Marfan's syndrome. Subluxation of the lens is a known complication.

Image (a) shows Walker-Murdoch (wrist) sign - the ability to join the thumb and fifth finger around the wrist.

Image (b) shows Steinberg (thumb) sign - thumb tip extends from the palm of the hand in a closed fist

Features of Marfan's syndrome (Mnemonic: MARFAN'S)

- Mitral valve prolapse
- Myopia due to upward dislocation of the lens (bilateral ectopic lentis)

- Aortic root dilatation, Aortic regurgitation
- Retinal detachment
- Fibrillin 1 gene defect: Reduced fibrillin 1 production causing joint laxity
- Fifteen (Chromosome 15 is involved)
- Flat feet
- Arachnodactyly, Astigmatism, Air in lungs (Pneumothorax)
- Nearsightedness
- Subluxation of lens, Scoliosis, Speech disorders, Sleep apnea

The image given below shows the Steinberg (thumb) sign:



Solution to Question 18:

Maternal disomy of chromosome 15 is seen in Prader–Willi syndrome. The image shows the characteristic features of truncal obesity, short stature, and small hands and feet.

Uniparental disomy: Offspring receives both copies of a chromosome from 1 parent and no copies from the other parent.

Chromosome 15 diseases: Chromosome 15 has its own MAP (mnemonic),

- M: Marfan syndrome
- A: Angelman syndrome
- P: Prader Willi syndrome

	Prader-Willi syndrome	Angelman syndrome
Deletion	Paternal chromosome 15	Maternal chromosome 15

	Prader-Willi syndrome	AngelMan syndrome
Uniparental Disomy	Maternal disomy(duplication) seen in 25% of cases	Paternal disomy (duplication) seen in 5% of cases
Clinical features	Elevated Ghrelinlevels: Loss of appetite control, obesitySevere hypotoniaShort statureSmall hands and feetHypogonadotropic hypogonadismCognitive impairment	Short statureSevere mental retardation with lack of speechStiff arm movementsSpastic, uncoordinated walkSeizuresAn inappropriate outburst of laughter (Happy puppets)

Solution to Question 19:

Triangular facies is indicative of Russell–Silver syndrome.

Russell–Silver syndrome:

- Clinical features: Short stature, triangular face (small jaw and pointed chin), microcephaly, limb and facial asymmetry, 5th finger clinodactyly.
- A few cases are due to genomic imprinting disorder involving chromosome 11.

Note: Triangular facies is also seen in Alagille syndrome.

The image given below shows a child with Russell–Silver syndrome.



Solution to Question 20:

Electroporation is used in gene therapy.

Gene therapy is a technique that uses genes to treat or prevent disease through the delivery of DNA into a patient's cell by using recombinant viruses/naked DNA/DNA complexes. The methods

used include intranuclear injection, electroporation, gene-gun, sonoporation, and magnetofection. Gene therapy was first used for severe combined immunodeficiency disorder (SCID), which is due to adenosine deaminase enzyme deficiency.

Solution to Question 21:

The picture depicts a Simian crease – a single transverse palmar crease. It is a normal finding in about 3% of the population. In a child with simian crease and normal growth and development, reassurance should be provided.

- A simian crease is one of the findings of Down's syndrome however presence of only this finding should not lead to a diagnosis of Down's syndrome.
- Maternal history of phenytoin intake during pregnancy does not lead to this finding. However, maternal alcohol intake can be associated with this condition.
- Chromosomal analysis in an otherwise well child is not indicated.

Solution to Question 22:

Non-invasive prenatal screening (NIPS) has a high negative predictive value.

NIPS:

- Useful in screening high-risk pregnancies for aneuploidies.
- Done after 10 weeks of gestation.
- Evaluates maternal blood for screening.
- Positive tests need confirmation by invasive testing (like amniocentesis, chorionic villous sampling).

Metabolic Disorders of Amino Acids

Question 1:

A mother reports to the OPD with the complaint that her child smells funny. On examination, you notice that the child has a global developmental delay and smells mousy. What is your probable diagnosis?

- a) Phenylketonuria
- b) Tyrosinemia
- c) Hawkinsinuria
- d) Hypermethioninemia

Question 2:

A boy who attends school is sent to the principal's office as he often distracts others by rhythmically rocking back and forth on his seat. His mother worried about his behavior brings him to you. She also gives you a history of seizures. On examination, you notice he has blonde hair, fair skin, microcephaly, prominent maxillae with widely spaced teeth, and growth retardation. His urine changes its color transiently to blue-green on the addition of 1ml ferric chloride solution. What is the cause of CNS symptoms in this child?

- a) Hyperphenylalaninemia
- b) Hypotyrosinemia
- c) Phenyl-pyruvate
- d) Phenyl-acetate

Question 3:

An infant with severe vomiting who was misdiagnosed as a case of hypertrophic pyloric stenosis is being screened for metabolic disorders. Screening reveals hyperphenylalaninemia. On ordering an enzyme study, you would include all of the following enzymes except:

- a) Phenylalanine hydroxylase (PAH)
- b) Dihydropteridine reductase
- c) Dihydropteroate synthase
- d) GTP cyclohydrolase

Question 4:

As a physician treating a 2-month-old baby with phenylketonuria what will be your mainstay of treatment?

- a) Phenylalanine-free diet
- b) Low phenylalanine diet
- c) High phenylalanine diet
- d) Tyrosine supplementation

Question 5:

An infant born out of a second-degree consanguineous marriage presents with jaundice, irritability, and vomiting. His mother gives a history of fever for the past 4 days. On examination, petechiae are present, he has hepatomegaly and extensor hypertonia of the neck and trunk. Labs reveal elevated serum transaminases, hypophosphatemia, and elevated levels of succinylacetone in serum and urine. Which of the following enzymes is deficient?

- a) Tyrosine hydroxylase
- b) Tyrosine aminotransferase
- c) 4-HPPD
- d) Fumarylacetoacetate hydrolase

Question 6:

A father while changing his child's diaper observes a boiled cabbage odor emanating from it. Curious about the smell, he calls you. As a pediatrician, you investigate the child thoroughly and conclude that the child is suffering from Type 1 Tyrosinemia. As the child grows all of the following complications can occur except:

- a) Hepatocellular carcinoma
- b) Paralytic ileus
- c) Fanconi-like syndrome
- d) Corneal ulceration

Question 7:

A 2-year-old girl presented with photophobia and speech delay. On examination, she had bilateral herpetiform corneal ulcers and painful hyperkeratotic plaques on the palms and soles. Which among the following is the most likely diagnosis?

- a) HSV type 1
- b) HSV type 2
- c) Hereditary Tyrosinemia Type 1
- d) Hereditary Tyrosinemia Type 2

Question 8:

A child who was being evaluated for hypertyrosinemia is diagnosed with Richner-Hanhart syndrome. Which of the following enzyme deficiencies is the likely cause of hypertyrosinemia in this patient?

- a) Tyrosine hydroxylase
- b) Tyrosine aminotransferase
- c) 4-Hydroxyphenylpyruvate dioxygenase (HPPD)
- d) Fumarylacetoacetate hydrolase

Question 9:

A 15-month-old baby who presented with seizures and self-injurious behavior has increased plasma tyrosine levels. Urine metabolic screening reveals the presence of 4-hydroxyphenylpyruvic acid and its metabolites 4-hydroxyphenyllactic and 4-hydroxyphenyl acetic acids. Which of the following in-born errors of metabolism is the child suffering from?

- a) Hereditary tyrosinemia type 1
- b) Hereditary tyrosinemia type 2
- c) Hereditary tyrosinemia type 3
- d) Hawkinsinuria


Question 10:

A 4-month-old infant who was started on a high-protein diet after weaning from breastfeeding developed severe metabolic acidosis, ketosis, failure to thrive, mild hepatomegaly, and an unusual swimming pool odor. A history of similar illness was noticed in his siblings and his father. What is the likely diagnosis?

- a) Maple syrup urine disease
- b) Isovaleric acidemia
- c) Hawkinsinuria
- d) Phenylketonuria

Question 11:

The following findings in an 18-year-old are characterized by a deficiency of _____



- a) Beta-carotene
- b) Tyrosine aminotransferase
- c) L-ascorbic acid
- d) Homogentisic acid oxidase

Question 12:

A patient presented to the OPD with complaints of pigmentation of the face and severe backache and stiffness. The X-Ray image of the spine is as shown. Her urine color changes to brownish-black on standing. Which of the following drugs can be used in her treatment?



- a) Nitisinone
- b) Thiamine
- c) Folic acid
- d) Pyridoxine

Question 13:

A 10-year-old boy presented with right-sided weakness and deviation of angle of mouth to the left is diagnosed to have a metabolic disorder. Diffusion-weighted MRI reveals diffusion restriction along the left middle cerebral artery. Which of the following defects can be responsible for the boy's condition?

- a) 2,5
- b) 1,3,5
- c) 1,2,3
- d) 1,2,3,4

Question 14:

An 8-month-old boy presented with lethargy, failure to thrive, and developmental delay. On investigation, he had elevated levels of homocysteine in blood and homocystinuria. As the child grows, which of the following features is least likely to develop?

- a) Ectopia lentis
- b) Intellectual disability

- c) Short stature
- d) Myocardial infarction

Question 15:

A 16-year-old boy with frequent episodes of photosensitive dermatitis, diarrhea, and depression is diagnosed to have a metabolic disorder. Metabolism of which of the following amino acids is affected?

- a) Tyrosine
- b) Tryptophan
- c) Threonine
- d) Phenylalanine

Question 16:

Which of the following is false regarding Hartnup disease?

- a) Defect in the transport of monoamino-monocarboxylic amino acids, including tryptophan, is seen
- b) Defect in the BoAT1 protein occurs in the intestine and renal tubules
- c) Pellagra-like rash may develop in symptomatic patients
- d) Treatment is with supplementation of tryptophan

Question 17:

An infant with poor feeding is suspected to have an inborn error of metabolism. Further evaluation reveals hyperammonemia +, acidosis +, ketosis +. What is the most likely diagnosis?

- a) Tyrosinemia
- b) Galactosemia
- c) Phenylketonuria
- d) Organic aciduria

Question 18:

A 10-day-old male infant presented with poor feeding, and recurrent generalized convulsions. On examination, the pediatrician notices repetitive movements of the extremities –appears to be boxing and bicycling and a peculiar burnt sugar odor was present in urine. What is the deficient enzyme in this condition?

- a) Branched-chain α -ketoacid dehydrogenase
- b) Isovaleryl CoA dehydrogenase
- c) Pyruvate carboxylase
- d) Acetyl CoA carboxylase

Question 19:

While doing your morning rounds in the pediatrics department, you notice a peculiar rancid cheese odor emanating from a six-month-old girl who was admitted for respiratory distress. Second-degree consanguinity was documented between the parents. Urine organic acid analysis by gas chromatography-mass spectrometry showed marked excretion of 3-hydroxybutyric acid along with moderate excretion of 3- hydroxy-isovaleric acid. What is the most likely condition is she suffering from?

- a) Tyrosinemia
- b) Multiple carboxylase deficiency
- c) Trimethylaminuria
- d) Isovaleric acidemia

Question 20:

A mother brings her 3-year-old boy with the complaints that her child's urine smells like her tomcat's urine. She also gives a history of generalized seizures and perioral desquamating lesions. On examination, the child has global developmental delay and diffuse thinning of the scalp. Investigations reveal multiple carboxylase deficiency. What is the treatment of choice?

- a) Thiamine
- b) Pyridoxine
- c) Biotin
- d) Methylcobalamin

Question 21:

Which of the following is the most common acquired cause of increased plasma tyrosine levels?

- a) Hepatorenal tyrosinemia
- b) Oculocutaneous tyrosinemia
- c) Alkaptonuria
- d) Transient tyrosinemia of the newborn

Question 22:

Richner-Hanhart syndrome is _____.

- a) Hereditary tyrosinemia type 1
- b) Hereditary tyrosinemia type 2
- c) Hereditary tyrosinemia type 3
- d) Hawkinsinuria

Question 23:

In phenylketonuria, which of the following phenyl ketones is excreted in urine?

- a) Phenylalanine
- b) Phenylpyruvate
- c) Acetoacetate
- d) Both b and c

Question 24:

A 2-month-old infant presented with abdominal distention and increased irritability for 10 days. Physical examination showed jaundice, hepatomegaly, and a boiled cabbage odor. Which among the following is the most likely diagnosis?

- a) Tyrosinemia
- b) Isovaleric acidemia
- c) Hawkinsinuria
- d) Phenylketonuria

Question 25:

A neonate born out of a second-degree consanguineous marriage is suspected to have an inborn error of metabolism. He can present with all of the following except:

- a) Irritability and incessant crying
- b) Poor feeding
- c) Vomiting
- d) Convulsions

Question 26:

A worried mother brings her 2-year-old child with the complaints that he hasn't begun walking or talking. On examination, there is microcephaly, fair hair, light-colored skin, and blue eyes. What is the most probable diagnosis?

- a) Oculocutaneous albinism
- b) Phenylketonuria
- c) Alkaptonuria
- d) Tyrosinemia

Answer Key

Question No.	Correct Option
1	a
2	a
3	c
4	b
5	d
6	d
7	d
8	b
9	c
10	c
11	d
12	a

13	d
14	c
15	b
16	d
17	d
18	a
19	d
20	c
21	d
22	b
23	b
24	a
25	a
26	b

Detailed Explanations

Solution to Question 1:

Mousy odor is characteristically seen in phenylketonuria due to excess phenylacetate (phenyl ketone) formed in the body.

In the absence of phenylalanine hydroxylase, phenylalanine enters into alternate metabolic pathways which result in the formation of phenyl lactate, phenylacetate, and phenylacetylglutamine.

Solution to Question 2:

The clinical vignette describes a case of phenylketonuria. CNS damage in patients with phenylketonuria (PKU) is caused by hyperphenylalaninemia in brain tissue.

High blood levels of phenylalanine saturate the transport system across the blood-brain barrier causing inhibition of the cerebral uptake of other large neutral amino acids such as tyrosine and tryptophan impairing brain protein synthesis.

Microcephaly, prominent maxillae with widely spaced teeth, enamel hypoplasia, and growth retardation are other common findings in untreated children. Low bone mineral density and osteopenia have been reported in affected individuals of all ages.

Excess phenylalanine is metabolized to phenyl ketones (phenylpyruvate and phenylacetate) that are excreted in the urine. However, these metabolites have no role in the pathogenesis of CNS damage; their presence in body fluids simply signifies the severity of the condition.

Hypotyrosinemia can be partially responsible for the signs and symptoms but the primary cause is hyperphenylalaninemia.

Laboratory diagnoses of phenylketonuria include:

- Guthrie's test: growth of bacillus subtilis is seen
- Ferric chloride urine test: transient blue-green color
- Blood phenylalanine levels: elevated
- Enzyme studies
- Genetic mutation studies
- Tandem mass spectrometry: Gold standard screening method for all metabolic disorders.

Solution to Question 3:

Defects of either phenylalanine hydroxylase (PAH) or tetrahydrobiopterin(BH₄) can cause hyperphenylalaninemias.

PAH deficiency is diagnosed as phenylketonuria.

Malignant hyperphenylalaninemias include three enzyme defects, dihydropteridine reductase,pyruvoyl tetrahydropterin synthase, and GTP cyclohydrolase which help in BH₄ synthesis. The deficiency of BH₄ impairs three hydroxylase(phenylalanine, tyrosine, and tryptophan), and severe neurological symptoms develop because of the lack of neurotransmitters.

Dihydropteroate synthase is present only in bacteria and this makes it a useful target for sulfonamide antibiotics, which compete with the PABA precursor.

Solution to Question 4:

A low-phenylalanine diet is the mainstay of treatment of PKU.

The goal of therapy is to reduce phenylalanine levels in the plasma and brain. Because phenylalanine is not synthesized endogenously, small amounts of phenylalanine should be added to the diet to prevent phenylalanine deficiency. Dietary deficiency of this amino acid is manifested by lethargy, failure to thrive, anorexia, anemia, rashes, diarrhea, and even death; moreover, tyrosine becomes an essential amino acid in this disorder and its adequate intake must be ensured.

Sapropterin dihydrochloride , a synthetic form of BH₄, which acts as a cofactor in patients with residual PAH activity, is approved by the U.S. Food and Drug Administration (FDA) to reduce phenylalanine levels in PKU.


Solution to Question 5:

The clinical vignette describes a case of Hereditary tyrosinemia type 1 is caused by the deficiency of fumarylacetoacetate hydrolase (FAAH).

Elevated levels of succinylacetone in serum and urine are diagnostic for tyrosinemia type I.

Liver(jaundice, acute hepatic crisis, elevated serum transaminases), kidney(hypophosphatemia), and nerve damage(extensor hypertonia of neck and trunk) are caused by metabolites of tyrosine degradation, especially fumarylacetoacetate and succinylacetone, and is usually precipitated by minor infections.

A diet low in phenylalanine and tyrosine can slow but does not halt the progression of the condition. The treatment of choice is nitisinone (NTBC), which inhibits 4-HPPD and reduces the flux of tyrosine metabolites to FAH, thus decreasing the production of the offending compounds, fumarylacetoacetate, and succinylacetone.



Solution to Question 6:

Corneal ulceration is not a feature of hereditary tyrosinemia type 1. In fact, ocular manifestations are a feature of hereditary tyrosinemia type 2 which presumably occur due to tyrosine deposition.

Boiled cabbage odor may be present in tyrosinemia resulting from increased methionine metabolites (sulfur-containing compounds).

Features of hereditary tyrosinemia type 1 (hepatorenal tyrosinemia):

- An acute hepatic crisis commonly heralds the onset of disease, which may progress to liver failure, hepatocellular carcinoma, and death.
- Episodes of acute peripheral neuropathy resembling porphyrias often triggered by a minor infection, such crisis can be associated with paralytic ileus
- Renal involvement, which is manifested as a Fanconi-like syndrome with hyperphosphaturia, hypophosphatemia, normal anion gap metabolic acidosis, and vitamin D-resistant rickets

The treatment of choice for HT Type 1 is nitisinone, which blocks early steps in the tyrosine degradation pathway and limits the formation of toxic tyrosine metabolites.

Note:


Boiled cabbage odor can also be found in cases of Hypermethioninemia.

Solution to Question 7:

Palmar and plantar hyperkeratosis, herpetiform corneal ulcers, and intellectual disability are seen in hereditary tyrosinemia type 2 (Richner–Hanhart syndrome/ oculocutaneous tyrosinemia)

Treatment with a diet low in tyrosine and phenylalanine aiming to achieve plasma tyrosine levels $<500 \mu\text{mol/L}$ improves skin and eye manifestations.


The image below shows herpetiform corneal ulcer:



Solution to Question 8:

Oculocutaneous tyrosinemia/Richner-Hanhart syndrome (type II) is caused by the deficiency of tyrosine aminotransferase.

It is a rare autosomal recessive disorder characterized by palmar and plantar hyperkeratosis, herpetiform corneal ulcers, and intellectual disability. Untreated patients have marked hypertyrosinemia. In contrast to tyrosinemia type I, liver and kidney function are normal, as are serum concentrations of other amino acids and succinylacetone.



Solution to Question 9:


Primary deficiency of 4-hydroxyphenylpyruvate dioxygenase (4-HPPD) causes hereditary tyrosinemia type 3. The diagnosis is suspected in children with sustained moderate increases in plasma levels of tyrosine (typically 350-700 $\mu\text{mol/L}$ on a normal diet) and the presence of 4-hydroxyphenylpyruvic acid and its metabolites 4-hydroxyphenyllactic and 4-hydroxyphenylacetic acids in urine.

Asymptomatic infants with 4-HPPD deficiency have been identified by neonatal screening for hypertyrosinemia. In symptomatic patients, developmental delay, seizures, intermittent ataxia, and self-injurious behavior have been reported. Liver and renal abnormalities are absent.

Note:

Transient tyrosinemia of the newborn is the most common acquired cause for increased plasma tyrosine levels. Seen in premature infants receiving a high protein diet and due to the immaturity of 4-HPPD enzyme. Mostly asymptomatic, but sometimes manifested by lethargy, poor feeding, and decreased motor activity. Hypertyrosinemia usually resolves

spontaneously in the first 2 months of life. Treatment includes a low protein diet and vitamin C.



Solution to Question 10:

The clinical vignette and the characteristic swimming pool odor describe Hawkinsinuria.

Hawkinsinuria is an autosomal-dominant condition caused by mutations in the gene for 4-HPPD.

In infants, this disease causes severe metabolic acidosis, ketosis, failure to thrive, mild hepatomegaly, and swimming pool odor. Mental development is usually normal. Affected individuals excrete hawkinsin (4-hydroxyphenylpyruvic acid), and its metabolites.

Treatment consists of a low-protein diet during infancy. Breastfeeding is encouraged. A trial with large doses of vitamin C (up to 1,000 mg/24 hour) is also recommended.

Solution to Question 11:

The following condition is characterized by the deficiency of homogentisic acid oxidase.

The image represents ochronosis: Bluish-black pigmentation of the ear cartilage and/or sclera due to deposits of accumulated homogentisic acid.


This is characteristically seen in alkaptonuria. It is a rare autosomal-recessive disorder caused by the deficiency of homogentisic acid oxidase. Large amounts of homogentisic acid are formed, which are excreted in urine or deposited in tissues. Clinical manifestations of alkaptonuria consist of ochronosis and arthritis in adulthood. Arthritis predominantly involves the spine, hip, and knee. Radiologically, it is characterized by intervertebral calcifications. In children, blackening of the urine on standing is observed; it is caused by the oxidation and polymerization of homogentisic acid.

Solution to Question 12:

The clinical image shows the accumulation of alkapton bodies in connective tissues on the face and urine turning brownish-black on standing are associated with alkaptonuria. X-ray shows calcification of intervertebral discs called ochronotic spondyloarthropathy also seen in alkaptonuria. Nitisinone can be used in the treatment of alkaptonuria.

Nitisinone, formerly known as NTBC, inhibits 4-hydroxyphenylpyruvate dioxygenase (HPPD) (early step in the tyrosine degradation pathway) and efficiently reduces homogentisic acid production in alkaptonuria. Treatment of arthritis is symptomatic. In presymptomatic individuals, treatment using a combination of nitisinone with phenylalanine and a tyrosine restricted diet can be tried.

The patient's urine contains elevated levels of homogentisic acid, which is oxidized to a dark pigment on standing as shown below in the image:




Solution to Question 13:

The clinical vignette describes a thrombotic stroke in the left MCA territory. Elevated levels of serum homocysteine lead to a hypercoagulable state.

Homocystinuria is caused by the following:

- Cystathionine β -Synthase deficiency: This is classic homocystinuria. The most common inborn error of methionine metabolism.
- Defects in methylcobalamin formation: Methylcobalamin is the cofactor for the enzyme methionine synthase, which catalyzes the remethylation of homocysteine to methionine.
- Deficiency of methylenetetrahydrofolate reductase: This enzyme reduces 5,10 methylenetetrahydrofolate to form 5-methyltetrahydrofolate, which provides the methyl group needed for remethylation of homocysteine to methionine.

- Pyridoxine deficiency: B6 is the co-factor for the enzyme cystathionine β -synthase



Metabolic disorders which can cause stroke include:

- Homocystinuria
- MELAS (metabolic encephalopathy, lactic acidosis, and stroke-like episodes)
- Leigh syndrome
- Fabry's disease
- Propionic acidemia
- Glutaric aciduria

Vitamin C deficiency increases the risk of hemorrhagic stroke.

Solution to Question 14:

The above findings support a diagnosis of homocystinuria. Short stature is not a manifestation of homocystinuria.

Manifestations of homocystinuria:

- Ectopia lentis - downward and inward subluxation (inferonasal).
- Intellectual disability and psychiatric and behavioral disorders
- Thromboembolic episodes like stroke and MI
- Skeletal abnormalities resembling those of Marfan syndrome: They are tall and thin, (NOT short) with elongated limbs and arachnodactyly.

Solution to Question 15:

Pellagra-like symptoms- photosensitive dermatitis, diarrhea, and episodic depression are seen in the deficiency of tryptophan.

Niacin is not strictly a vitamin, since it can be synthesized in the body from the essential amino acid tryptophan; deficiency of tryptophan causes pellagra-like symptoms.

Treatment with nicotinic acid or nicotinamide (50-300 mg/24 hr) and a high protein diet results in a favorable response in symptomatic patients with Hartnup disease.

The image below shows Casal's necklace, a lesion seen around the neck which is caused due to the deficiency of niacin or nicotinic acid.

Casal's paint necklace in Pellagra



Solution to Question 16:

The treatment of Hartnup disease is not with the supplementation of tryptophan.

Hartnup disease

- It is caused by a defect in the transport of monoamino-monocarboxylic (neutral) amino acids, including tryptophan, by the intestinal mucosa and renal tubules.
- There is a defect in BOAT1 protein in the intestine/ renal tubules or both, which is caused by a mutation in SLC6A19.
- Tryptophan is an essential amino acid and a precursor for nicotinic acid (niacin) and serotonin. Decreased intestinal absorption of tryptophan in conjunction with its increased renal loss causes reduced availability of tryptophan for niacin synthesis in affected individuals.
- A pellagra-like rash may develop in symptomatic patients, and some patients may have intermittent ataxia.
- Diagnosis is based on aminoaciduria and increased excretion of indole compounds detected by Obermeyer test.
- Treatment is with nicotinic acid or nicotinamide and a high-protein diet.


Solution to Question 17:

In this case, hyperammonemia +, acidosis +, ketosis +. This is seen in organic aciduria.

Organic acidemias most commonly present in infancy with poor feeding, vomiting, hypotonia, high anion gap metabolic acidosis, hepatomegaly, seizures. Organic acid accumulation inhibits gluconeogenesis and inhibits the urea cycle leading to hyperammonemia.

Common organic acidurias:

- Methylmalonic aciduria
- Propionic aciduria
- Isovaleric aciduria
- Maple syrup urine disease
- Glutaric aciduria type I
- Multiple carboxylase deficiency



Solution to Question 18:

The diagnosis is maple syrup urine disease (MSUD) which is due to deficiency of branched-chain α -ketoacid dehydrogenase (BCKDH).

Leucine, isoleucine, and valine are the 3 essential amino acids, which are branched-chain amino acids. Their decarboxylation is accomplished by BCKDH using thiamine (vitamin B1) pyrophosphate as a coenzyme.

Leucine is metabolized by BCAA aminotransferase to α -ketoisocaproic acid, which leads to the disrupted metabolism of neurotransmitters and amino acids (glutamate, GABA, glutamine,

alanine, and aspartate). α -Ketoisocaproic acid can reversibly inhibit oxidative phosphorylation and result in cerebral lactic acidosis. Collectively, these processes are detrimental to the normal function of neurons and glia, clinically manifesting as encephalopathy and brain edema and referred to as leucinoses.

Treatment options include hemodialysis, restricting dietary substances, thiamine for milder forms, and orthotopic liver transplant.

Solution to Question 19:

The clinical vignette suggests a diagnosis of isovaleric acidemia. Sweaty-feet odor or rancid cheese odor may be present.

Isovaleric acidemia is caused by the deficiency of the enzyme isovaleryl CoA dehydrogenase, which affects the breakdown pathway of leucine leading to the accumulation of intermediates.

Solution to Question 20:

Biotin is the treatment of choice for multiple carboxylase deficiency.

Multiple carboxylase deficiency refers to one of two inherited defects of biotin metabolism. Tomcat urine is characteristically seen in multiple carboxylase deficiency.

- Neonatal/infantile form is caused by a deficiency of holocarboxylase synthetase and presents in the first week of life with lethargy, poor muscle tone, and vomiting.
- Juvenile/late form is caused by biotinidase deficiency and is associated with a slow but progressive loss of biotin in the urine, leading to organic aciduria. It is characterized by alopecia, ataxia, ketoacidosis, dermatitis, seizures, myoclonus, and nystagmus.

Both infantile and late-onset multiple carboxylase deficiency can be treated with pharmacological doses of biotin.

It is called multiple carboxylase deficiency because biotin is a water-soluble vitamin that is a cofactor for all 4 carboxylase enzymes in humans:

- Pyruvate carboxylase
- Acetyl CoA carboxylase
- Propionyl CoA carboxylase
- 3-methylcrotonyl CoA carboxylase

Note:

Acquired deficiency of biotin may occur in infants receiving total parenteral nutrition without added biotin, in patients with prolonged use of antiepileptic drugs (phenobarbital, phenytoin, primidone, carbamazepine), and in children with short bowel syndrome or chronic diarrhea who are receiving formulas low in biotin. Excessive ingestion of raw eggs may also cause biotin deficiency because the protein avidin in egg white binds biotin, decreasing its absorption. Infants with a biotin deficiency may develop dermatitis, alopecia, and candidal skin infections. This

condition readily responds to treatment with oral biotin.

Solution to Question 21:

Transient tyrosinemia of the newborn is the most common acquired cause for increased plasma tyrosine levels. Seen in premature infants receiving a high protein diet and due to the immaturity of 4-HPPD enzyme. Mostly asymptomatic, but sometimes manifested by lethargy, poor feeding, and decreased motor activity. Treatment includes low protein diet and vitamin C.

Options A and B, i.e., hepatorenal tyrosinemia and oculocutaneous tyrosinemia, are hereditary causes of increased tyrosine levels.

Option C: Tyrosine levels are normal in alkaptonuria.

Solution to Question 22:

Richner–Hanhart syndrome is hereditary tyrosinemia type 2 (oculocutaneous tyrosinemia).

- It is a rare autosomal-recessive disorder is caused by the deficiency of tyrosine aminotransferase.
- It is characterized by ocular manifestations, skin lesions, and mild to moderate intellectual disability in ~50% of the patients.
- It is diagnosed by an assay of plasma tyrosine concentration.
- Treatment includes a diet low in tyrosine and phenylalanine.

Solution to Question 23:

In phenylketonuria, phenylpyruvate is excreted in the urine.

In phenylketonuria (PKU), excess phenylalanine in the blood is metabolized to phenyl ketones, namely, phenylpyruvate and phenylacetate, which are excreted in the urine, giving rise to the term phenylketonuria (PKU).

Acetoacetate and beta-hydroxybutyrate are keto-acids but are not phenyl ketones. These are not excreted in phenylketonuria.

Solution to Question 24:

The clinical vignette is in favor of tyrosinemia. Boiled cabbage odor may be present in tyrosinemia resulting from increased methionine metabolites (sulfur-containing compounds).

Note:

Boiled cabbage odor can also be found in cases of Hypermethioninemia.

Solution to Question 25:

Irritability and incessant crying are not classical features of inborn errors of metabolism (IEM).

Clinical clues suggesting IEM in a neonate are:

- Unexpected deterioration after a normal initial period.
- Nonspecific, unexplained features such as poor feeding, lethargy, vomiting, hypotonia, failure to thrive, respiratory abnormalities, apnea, bradycardia, and hypothermia.
- Family history of similar illnesses, unexplained sibling deaths, or progressive neurological disease.
- Parental consanguinity: Most acutely presenting metabolic disorders are autosomal recessive.

Solution to Question 26:

The given clinical manifestations point to the diagnosis of Phenylketonuria (PKU).

Characteristic clinical findings in Phenylketonuria:

- Mental retardation
- Failure to walk or talk
- Seizures
- Hyperactivity
- Tremor
- Microcephaly
- Failure to grow
- Deficiency of pigmentation (fair hair, light skin color, and blue eyes)

Oculocutaneous albinism (OCA) is a group of inherited disorders characterized by a reduction or complete lack of melanin pigment in the skin, hair, and eyes.

Its symptoms are albinism with vision problems, which include:

- Strabismus
- Photophobia
- Nystagmus
- Impaired vision or blindness
- Astigmatism

Neuropsychiatric manifestations are not seen.

Metabolic Disorders of Urea Cycle, Complex Molecules, and Carbohydrates

Question 1:

A 3-day-old child is suspected to have an inborn error of metabolism. Further evaluation reveals elevated blood ammonia in the presence of normal blood pH and bicarbonate levels. There are no ketone bodies in the urine. Which of the following is the most likely diagnosis?

- a) Organic aciduria
- b) Disorders of amino acid metabolism
- c) Urea cycle defect
- d) Galactosemia

Question 2:

A young infant started seizing in his cradle. He was immediately rushed to the casualty. On investigation, blood ammonia was markedly elevated. Which of the following is used to lower the ammonia levels?

- a) Sodium benzoate
- b) Sodium phenylacetate
- c) Arginine hydrochloride
- d) All of the above


Question 3:

An infant's mother is distressed that her baby refuses her feeds and starts vomiting immediately when fed and becomes lethargic. After thorough investigations, the baby is diagnosed with HHH syndrome. Which of the following is least likely to be found?

- a) Hyperuricemia
- b) Hyperammonemia
- c) Hyperornithinemia
- d) Homocitrullinemia

Question 4:

As a geneticist studying the familial inheritance of porphyrias, which of the following is likely to show the inheritance pattern given below?



- a) Congenital erythropoietic porphyria
- b) Variegate porphyria
- c) δ -Aminolevulinic acid dehydratase porphyria
- d) X-linked protoporphyria

Question 5:

The most common porphyria which manifests in childhood is _____

- a) Porphyria cutanea tarda
- b) Congenital erythropoietic porphyria
- c) Erythropoietic protoporphyria
- d) Hereditary coproporphyria

Question 6:

Which of the following is also known as plumboporphyria due to its clinical similarity to lead poisoning?

- a) Acute intermittent porphyria (AIP)

- b) δ -Aminolevulinic acid dehydratase porphyria (ALAD)
- c) Porphyria cutanea tarda (PCT)
- d) Erythropoietic protoporphyria (EPP)

Question 7:

A 16-year-old girl was admitted for severe abdominal pain, nausea, vomiting, and diarrhea. She developed generalized seizures in the hospital for which she was given phenytoin. Her symptoms worsened, she developed muscle weakness which progressed to quadriparesis. Urinary porphobilinogen was found to be high. What is the most likely diagnosis?

- a) Congenital erythropoietic porphyria
- b) Porphyria cutanea tarda
- c) Acute intermittent porphyria
- d) X-linked protoporphyria

Question 8:

You are reading an article titled, 'Dark age vampires or our patients', which describes an 18-year-old man who has had an intolerance to sunshine since the age of 6. When playing cricket he suffered intense burning, redness, swelling, and crusting of knees unprotected by shorts, with similar changes on cheeks and bridge of the nose. His labs showed elevated concentrations of total erythrocyte protoporphyrin. Which of the following enzyme deficiencies can explain this boy's condition?

- a) δ -Aminolevulinate synthase 2 (ALAS2)
- b) δ -Aminolevulinic acid dehydratase (ALAD)
- c) Porphobilinogen deaminase (PBGD)
- d) Ferrochelatase (FECH)

Question 9:

A 6-month-old baby boy has coarse facial features, cloudy corneas, and gingival hypertrophy. He has no neck control and is unable to roll over. Laboratory investigations reveal markedly reduced activity of N-acetylglucosaminyl-1-phosphotransferase activity in WBCs. What is the most likely diagnosis?

- a) I-cell disease

- b) Galactosemia
- c) Adrenoleukodystrophy
- d) Hurler's syndrome


Question 10:

An 8-year-old boy is brought with complaints of severe pain in his right thigh which has progressively worsened over the last few months. He is unable to walk because of the same. His mother also says he has had repeated nose bleeds in the last few months and the child has stopped playing with his friends as he gets easily tired for a month. Physical examination reveals hepatosplenomegaly. Lab reports show thrombocytopenia and anemia. An assay of peripheral blood leucocytes shows low levels of beta-glucosidase. What is the inheritance pattern of this condition?

- a) X-linked recessive
- b) X-linked dominant
- c) Autosomal recessive
- d) Autosomal dominant

Question 11:

A 6-month-old infant with progressive failure to thrive and milestone regression is being evaluated for an inborn error of metabolism. On examination, hypotonia, apathy, deafness, hepatosplenomegaly, and generalized lymphadenopathy are present. Fundus examination is shown below. A liver biopsy shows foamy hepatocytes loaded with sphingomyelin. Electron microscopy reveals zebra bodies. What is the likely diagnosis?



- a) Niemann-Pick disease type A
- b) Tay-Sach disease
- c) Niemann-Pick disease type B
- d) Sandhoff's disease

Question 12:

A male child with coarse facies, macroglossia, thick lips with hepatosplenomegaly presents with copious mucous discharge from the nose. The most probable underlying diagnosis is

-
- a) Hurler disease
 - b) Beckwith-Wiedemann syndrome
 - c) Proteus Syndrome
 - d) Hypothyroidism

Question 13:


A case series reviews children with lysosomal storage disorders who usually present with dysmorphic facies, organomegaly, milestones regression, and failure to thrive. Which of the following disorders does not have enzyme replacement therapy as a therapeutic option?

- a) Gaucher's disease
- b) Pompe disease
- c) Sanfilippo syndrome

d) Fabry's disease

Question 14:

A 3-month-old female child was brought to the clinic with a history of failure to thrive. On examination, generalized hypotonia and hepatomegaly were present. A chest X-ray image shows massive cardiomegaly. A muscle biopsy is performed and the findings are attached in the image below. Which of the following is the most likely enzyme that is involved in the suspected condition?



- a) Glucose-6-phosphatase**
- b) Lysosomal acid alpha-glucosidase**
- c) Muscle glycogen phosphorylase**
- d) Glycogen debranching enzyme**

Question 15:

A 6-month-old is brought by his parents with a history of decreased weight gain. On examination, you find that the baby has hepatosplenomegaly, muscle weakness and atrophy, hypotonia, and decreased deep tendon reflexes. Blood tests reveal normal glucose levels. Biopsy of the liver reveals initial stages of cirrhosis due to the accumulation of abnormal glycogen with few branch points, whose structure resembles amylopectin. The clinical and laboratory results are indicative of which glycogen storage disease?

- a) Von Gierke disease**
- b) Andersen disease**

- c) McArdle disease
- d) Tarui disease


Question 16:

A 6-year-old boy develops hypoglycemia and muscle weakness with moderate activity. On examination, he had hepatomegaly and growth restriction. Investigations revealed the accumulation of limit dextrins and increased glycogen deposits in the liver. What is the most likely diagnosis?

- a) Cori's disease
- b) von Gierke's disease
- c) Anderson's disease
- d) Pompe's disease

Question 17:

A 5-day-old neonate is brought to the emergency room with vomiting and poor feeding. Breastfeeding was initiated soon after birth. On examination, the child is lethargic and has hepatomegaly. Labs reveal increased plasma galactose and increased urine galactitol levels. Examination of the mother shows the presence of skin lesions on the breast with the histopathological appearance showing characteristic 'Cowdry Type A' inclusions as shown below. Choose the correct statements from those below:



- a) 2 and 4 are correct

- b) 1, 3, and 5 are correct
- c) 2 and 5 are correct
- d) 1,3, and 4 are correct

Question 18:

A father has noticed that whenever his daughter drinks fruit juice she has multiple episodes of vomiting. He reports to you that his daughter is allergic to fruit juice. Which of the following enzyme deficiency is likely?

- a) Aldolase B
- b) Fructokinase
- c) Glucose 6-phosphatase
- d) Hexokinase

Question 19:

"Tomcat urine" is characteristically seen in _____

- a) Tyrosinemia
- b) Multiple carboxylase deficiency
- c) Phenylketonuria
- d) Trimethylaminuria

Question 20:

The enzyme deficient in hereditary coproporphyria is _____

- a) Coproporphyrinogen oxidase (CPOX)
- b) Protoporphyrinogen oxidase (PPOX)
- c) Porphobilinogen deaminase (PBGD)
- d) Ferrochelatase (FECH)

Question 21:

Deficiency of the enzyme porphobilinogen deaminase causes _____

- a) Porphyria cutanea tarda
- b) Acute intermittent porphyria
- c) Variegate porphyria
- d) X-Linked protoporphyria

Question 22:

A mother who is a hairstylist has noted that her baby's hair looks dull, breaks off easily, and is tufted. She brings the child to you and informs you about the past history of seizures. Investigations reveal elevated plasma ammonia levels and argininosuccinate levels. What is the most likely diagnosis?

- a) Argininemia
- b) Argininosuccinic aciduria
- c) Hyperammonemia type 1
- d) Hyperammonemia type 2

Question 23:

Accumulation of _____ takes place in Farber's disease.

- a) Lecithin
- b) Ceramide
- c) Sphingomyelin
- d) Sulfatides

Answer Key

Question No.	Correct Option
1	c
2	d
3	a
4	d
5	c
6	b


7	c
8	d
9	a
10	c
11	a
12	a
13	c
14	b
15	b
16	a
17	b
18	a
19	b
20	a
21	b
22	b
23	b

Detailed Explanations

Solution to Question 1:

In this case, hyperammonemia is +ve; acidosis is -ve; and ketosis is -ve: This is consistent with the urea cycle defect.

Initial clinical approach to a full-term newborn infant with a suspected genetic metabolic disorder



Solution to Question 2:

All of the above agents are used in the treatment of acute hyperammonemia in an infant.

Treatment of acute hyperammonemia in an infant:

- Provide adequate calories, fluid, and electrolytes intravenously.
- Minimal amounts of protein as a mixture of essential amino acids during the first 24 hours of therapy.
- Give priming doses of the following compounds: Sodium benzoate, Sodium phenylacetate, Arginine hydrochloride.
- Initiate peritoneal dialysis or hemodialysis if the above treatment fails to produce an appreciable decrease in plasma ammonia levels.

Benzoate forms hippurate with endogenous glycine in the liver. Each mole of benzoate removes 1 mole of ammonia as glycine.

Phenylacetate conjugates with glutamine to form phenylacetylglutamine, which is readily excreted in the urine. One mole of phenylacetate removes 2 moles of ammonia as glutamine from the body.

Arginine administration supplies the urea cycle with ornithine. It is effective in all patients, except those with arginase deficiency.

Note:

Phenylbutyrate which is metabolized to phenylacetate in our body can also be used as it scavenges nitrogen.

Solution to Question 3:

Hyperuricemia is not a component of HHH syndrome

HHH syndrome is a rare autosomal-recessive disorder caused by a defect in the transport system of ornithine from the cytosol into the mitochondria (deficiency of ornithine permease/translocase), resulting in the accumulation of ornithine in the cytosol leading to hyperornithinemia and a deficiency of this amino acid in mitochondria results in the disruption of the urea cycle and hyperammonemia. Homocitrulline is formed from the reaction of mitochondrial carbamyl phosphate with lysine, which can become a substrate for the OTC reaction when ornithine is deficient.

Components of HHH syndrome:

- Hyperammonemia
- Hyperornithinemia
- Homocitrullinemia

Solution to Question 4:

The given pedigree chart shows X-linked dominant inheritance. X-linked protoporphyria is inherited in an X-linked dominant pattern.

All porphyrias are autosomal dominant except the following :

- Congenital erythropoietic porphyria (CEP) (autosomal recessive)
- δ -Aminolevulinic acid dehydratase porphyria (ADP) (autosomal recessive)
- Protoporphyria - X-linked protoporphyria (X-linked dominant)
- Erythropoietic protoporphyria (EPP) (autosomal recessive)

Solution to Question 5:

Erythropoietic protoporphyria (autosomal recessive) is the most common porphyria to manifest in childhood.

Porphyria cutanea tarda (option A) manifests in adults.

Congenital erythropoietic porphyria (option B) also known as Günther disease presents itself in utero or in infancy. It is a rare autosomal-recessive disease.

Hereditary coproporphyria (option D) presents itself in the post-pubertal age.

Solution to Question 6:

δ -Aminolevulinic acid dehydratase porphyria (ALAD) is also called plumboporphyria/Doss porphyria due to its clinical similarity to lead poisoning, although it is not related to lead exposure.

ALAD catalyzes the condensation of 2 molecules of ALA to form the pyrrole porphobilinogen (PBG). The enzyme is subject to inhibition by lead because ALAD is the principal lead-binding protein in erythrocytes, and lead can displace the zinc atoms of the enzyme, thus inhibiting it. Therefore, both lead poisoning and ALAD porphyria produce similar signs and symptoms. Symptoms resemble those of other acute porphyrias, including acute attacks of abdominal pain and neuropathy.

Solution to Question 7:

The most likely diagnosis is acute intermittent porphyria which presents as a neurovisceral syndrome with no cutaneous manifestations and no phototoxicity.

Acute Intermittent Porphyria:

- Autosomal dominant
- Deficiency of the enzyme porphobilinogen deaminase

- Presents at post-pubertal age
- Present with hepatic and neurologic manifestations
- Exacerbated by P450 inducing drugs like phenytoin, progesterone and dietary restriction

Solution to Question 8:

Symptoms of cutaneous photosensitivity beginning in childhood and a substantially elevated concentration of total erythrocyte protoporphyrin supports a diagnosis of erythropoietic protoporphyria. The enzyme deficient in erythropoietic protoporphyria is ferrochelatase (FECH).

Symptoms consist of acute pain and itching often occurring within minutes of sunlight exposure and followed by redness and swelling with continued exposure. Swelling may resemble angioneurotic edema and solar urticaria. Symptoms are usually worse in the spring and summer. Chronic changes may include lichenification, leathery pseudovesicles, labial grooving, and nail changes, but changes in pigmentation and pronounced scarring are unusual.

Life-threatening protoporphyric hepatopathy is characterized by greater increases in erythrocyte and plasma protoporphyrin levels, increased photosensitivity, and either chronically abnormal liver function tests or rapidly progressive hepatic failure.

Solution to Question 9:

This clinical vignette is strongly suggestive of I-cell disease or inclusion cell disease. It is also known as type II Mucopolidosis.

It is an autosomal recessive inherited as a lysosomal storage disorder. It is caused due to a defect in N-acetylglucosaminyl-1-phosphotransferase required for the Golgi to phosphorylate mannose residues on glycoproteins. This leads to defective mannose-6-phosphate, an important cell signaling molecule required for trafficking of the lysosomal enzyme from the site of its production in the endoplasmic reticulum to the lysosome. Due to this defect, the lysosomal enzymes are secreted out of the cell and there is an increased serum concentration of the same. Intracellularly, there are abnormal inclusion bodies and vacuolizations, thus earning the disease its name.

Clinically, these patients present with symptoms either at birth or early in life with growth retardation, failure to thrive, delay of developmental milestones, coarse facial features with a cloudy cornea and gingival hypertrophy, and other non-specific findings such as hip dislocations, inguinal hernias, hepatomegaly, joint limitation, and skin changes.

Investigations to establish a diagnosis may include measuring N-acetylglucosaminyl-1-phosphotransferase activity in WBCs or in cultured fibroblasts and estimating serum levels of lysosomal enzymes.

Option B: Galactosemia is caused due to the complete absence of galactose-1-phosphate uridyl transferase. The deficiency of this enzyme leads to an inability to convert galactose-1-phosphate (galactose-1-P) to uridine diphosphate galactose. It is characterized by typical clinical features such as jaundice, vomiting, hepatomegaly, failure to thrive, poor feeding, lethargy, diarrhea, E.Coli sepsis. Infants may also have edema, ascites, full fontanelle, encephalopathy, and excessive

bruising and bleeding. They tend to develop cataracts due to the accumulation of galactitol.

Option C: Adrenoleukodystrophy is an X-linked recessive disorder of β -oxidation of fatty acids. There is a very large chain fatty acids buildup in the adrenal glands, the white matter of the brain, and testes. It is a progressive disease that can lead to adrenal gland crisis, coma, and death.

Option D: Hurler syndrome is an important diagnosis in a patient presenting with clinical features given in this clinical vignette. But it is a mucopolysaccharidosis type 1-H caused due by the defect in the α -L Iduronidase leading to the accumulation of heparin sulfate and dermatan sulfate. It is characterized by skeletal abnormalities, hepatosplenomegaly, and severe intellectual disability. Infants develop characteristic coarse facial features, a wide nasal bridge, and a flattened midface. Features such as hepatosplenomegaly, umbilical or inguinal hernias, and typical skeletal abnormalities, known as dysostosis multiplex are also seen. They also have developmental delays and recurrent sinopulmonary diseases. Failure to thrive and growth restriction are also seen.

Solution to Question 10:

The given clinical scenario of bone pain, epistaxis, hepatosplenomegaly, and low levels of β -glucosidase suggests a diagnosis of Gaucher's disease that is inherited in an autosomal recessive pattern.

Gaucher's disease:

- It is an inborn error of metabolism due to the deficiency of the enzyme glucocerebrosidase (β -glucosidase) that leads to the accumulation of glucocerebroside in the lysosomes of cells.
- It is the most common lysosomal storage disorder.
- The clinical features result from the accumulation of lipid-laden macrophages in the spleen, liver, bone marrow, bone, and other tissues.
- There are 3 clinical subtypes, delineated by the absence or presence and progression of neurologic manifestations: type 1, or the adult, nonneuronopathic form; type 2, the infantile or acute neuronopathic form; and type 3, the juvenile or subacute neuronopathic form
- Typical clinical features include delayed puberty, painless hepatomegaly and splenomegaly, liver cirrhosis, hypersplenism and pancytopenia, bleeding tendencies, bone pain due to avascular necrosis, osteoporosis, pathological fractures.
- Neurological symptoms such as developmental delay, hypertonia, strabismus, supranuclear gaze palsy (seen in type 3).
- Bone marrow examination usually reveals the presence of Gaucher cells (have a characteristic wrinkled-paper appearance)
- Enzyme replacement therapy for Gaucher's disease: imiglucerase, velaglucerase alfa, taliglucerase alfa, eliglustat

Solution to Question 11:

The clinical vignette along with the cherry-red spot on fundus examination suggests a diagnosis of Niemann-Pick disease type A- Lysosomal storage disorder due to an inherited deficiency of sphingomyelinase

Type A is a severe infantile form with extensive neurologic involvement (hypotonia, milestone regression, apathy), marked visceral accumulations of sphingomyelin (hepatosplenomegaly, foamy hepatocytes, zebra bodies on EM), and progressive wasting and early death within the first 3 years of life.

In contrast, patients with Niemann-Pick disease type B have organomegaly but generally no CNS involvement.

Solution to Question 12:


The given clinical scenario of a child with coarse facies, macroglossia, thick lips, hepatosplenomegaly, and copious discharge from the nose is suggestive of Hurler's disease.

It is the severe form of mucopolysaccharidoses type 1 (MPS 1), an autosomal recessive disorder caused by a defect in α -L iduronidase, leading to the accumulation of heparan sulfate and dermatan sulfate. The child usually presents between 6 and 24 months with hepatosplenomegaly, coarse facial features (shown below), corneal clouding, large tongue, prominent forehead, joint stiffness, short stature, and skeletal dysplasia.

Most patients have recurrent upper respiratory tract and ear infections, noisy breathing, and persistent copious nasal discharge. Radiograph shows skeletal dysplasia, called dysostosis multiplex - thick ribs, ovoid vertebral bodies, enlarged and trabeculated diaphyses of the long bones with irregular metaphysis and epiphysis.

Macrocephaly, enlarged J-shaped sella, premature closure of lambdoid sagittal sutures, and shallow orbits are seen with disease progression. Obstructive airway disease, cardiac complications, and respiratory infections are the common causes of death.

The image below shows the radiograph shows enlarged diaphyses of the metacarpal and phalanges.



Other options:

Option B: Beckwith-Wiedemann syndrome is an overgrowth syndrome characterized by macrosomia, macroglossia, hemihypertrophy, omphalocele, and renal anomalies. It is associated with an increased risk of Wilms tumor, hepatoblastoma, rhabdomyosarcoma, neuroblastoma, and adrenal cortical carcinoma.

Option C: Proteus syndrome is a disturbance of cellular growth involving ectodermal and mesodermal tissues, caused by mutation in the AKT1 gene. They present with hemimegalencephaly, asymmetric overgrowth of the extremities, verrucous cutaneous lesions, thickening of the bone, and excessive growth of muscles.

Option D: A child with untreated congenital hypothyroidism may have stunted growth, short extremities, narrow palpebral fissures, swollen eyelids, depressed nasal bridge, thick broad protruded tongue, and short and thick neck, dry and scaly skin.

Solution to Question 13:

Enzyme replacement therapy is not available for Sanfilippo syndrome (MPS type III).

Solution to Question 14:

The muscle biopsy shows the presence of large vacuoles. This along with the clinical vignette is strongly suggestive of Pompe's disease or type II glycogen storage disease. It is caused due to the deficiency of the enzyme lysosomal acid alpha-glucosidase.

Clinically, the child presents with hypotonia, hepatomegaly, hypertrophic cardiomyopathy. The affected child usually dies around the age of 2 years due to heart failure.

Investigations include chest X-ray imaging which shows cardiomegaly. The muscle biopsy showing vacuoles further aids the diagnosis.

Treatment includes enzyme replacement therapy with recombinant enzyme α -glucosidase alfa (Myozyme).

Solution to Question 15:

The clinical and laboratory features indicate Andersen's disease in the baby. Inability to gain weight, hepatosplenomegaly, muscle weakness, and accumulation of abnormal glycogen resembling amylopectin point towards the diagnosis.

Andersen's disease is type IV glycogen storage disease. It is transmitted in an autosomal recessive pattern and occurs due to a deficiency of glycogen branching enzyme because of pathologic variants in the GBE1 gene. This leads to accumulation of an abnormal structure of glycogen known as polyglucosan. It is characterized by

- Growth delay
- Hepatomegaly due to the accumulation of polysaccharides with few branch points
- Myopathy
- There might be fasting intolerance and ketosis in the initial stages. Hypoglycemia usually appears late in the disease when it is associated with cirrhosis, esophageal varices, and ascites.
- Biopsy of the liver will show the presence of abnormal glycogen which resembles amylopectin. Amylopectin is found in plants and has very few branches. Since there is a defect in the branching of the glycogen, the abnormal glycogen appears like amylopectin.

The diagnosis is confirmed by absent branching enzyme activity in skin fibroblasts, muscle, or liver and analysis of the DNA mutation.

Von Gierke's disease (Option A) has features of lactic acidosis, hypoglycemia, hyperuricemia, impaired growth, and bone thinning. It occurs due to a defect in the glucose 6-phosphatase.

McArdle disease (Option C) usually manifests as exercise intolerance, muscle cramping with exercise. It is usually a benign chronic condition. It occurs due to a defect in the muscle glycogen phosphorylase.

Tarui disease (Option D) also presents similar to the McArdle disease. Patients may have developmental delay, and hemolytic anemia in addition to exercise-induced muscle cramps. It occurs due to a defect in the muscle phosphofructokinase.

Solution to Question 16:

The most likely diagnosis is Cori disease.

Glycogen debrancher deficiency (Glycogen Storage Disease III) is also known as Cori disease, Forbes disease, and limit dextrinosis.

Cori's disease may be clinically indistinguishable from von Gierke's disease in infancy and childhood as hypoglycemia, hepatomegaly, hyperlipidemia, and growth retardation is common. Accumulation of limit dextrins is unique to Cori's disease. Definitive diagnosis is through enzyme assays in the liver, muscle, or both.

Solution to Question 17:

The given clinical complex describes a case of classic galactosemia in the infant (increased plasma galactose and urine galactitol levels) and herpes simplex of the breast in the mother (histopathology shown in the image has large multinucleate virus-infected cells with typical eosinophilic Cowdry Type A inclusions).

Galactosemia is an absolute contraindication to breastfeeding and the child is at risk for E.coli sepsis. Herpes lesions on the breast are a relative contraindication to breastfeeding and she is at risk for recurrence.

Deficiency of galactose-1-uridyltransferase leads to the accumulation of galactose-1-phosphate and galactitol causes classic galactosemia.

Infants show vomiting, diarrhea, and poor weight gain within 3-4 days of breastfeeding. Jaundice and liver dysfunction appear in the first or second week of life. Untreated children may show nuclear cataracts (oil-droplet cataract), vitreous hemorrhage, hepatic failure, cirrhosis, ascites, splenomegaly, or intellectual disability. The proximal renal tubular disease may also manifest as metabolic acidosis, galactosuria, glucosuria, and aminoaciduria. The most common cause of death is E.coli sepsis. Acute galactose toxicity is manifested by neurological symptoms

Diagnosis is made by genetic or mutational analysis. There will be the presence of reducing substances in the urine. Elimination of galactose from the diet along with adequate calcium supplementation reverses growth failure and renal and hepatic dysfunction.

Solution to Question 18:

Aldolase B deficiency is likely in this child.

In the given case scenario, the child becomes symptomatic after ingesting fruit juice that contains fructose; this indicates that she has hereditary fructose intolerance, which is caused by the deficiency of aldolase B.

Deficiency of fructose-1,6- biphosphate aldolase (Aldolase B hereditary fructose intolerance): Fructose aldolase B is present in the liver, kidney, and intestine. It catalyzes the hydrolysis of fructose-1,6-bisphosphate into triose phosphate and glyceraldehyde phosphate.

Clinical features: Patients are asymptomatic until fructose or sucrose is ingested (usually from fruit, fruit juice, sweetened cereal, or table sugar). Clinical features include jaundice, hepatomegaly, vomiting, lethargy, irritability, and convulsions. It resembles galactosemia. Acute ingestion causes symptomatic hypoglycemia. Chronic intake results in failure to thrive and hepatic disease. The persistence of fructose intake can lead to liver failure, kidney failure, and death.

Diagnosis: Reducing substance is seen in the urine during an episode (Benedict's test +). An assay of fructaldolase B activity in the liver gives a definitive diagnosis.

Treatment: Complete elimination of all sources of sucrose, fructose, and sorbitol from the diet. With age, symptoms become milder even after fructose ingestion. The long-term prognosis is good.

Note:

The deficiency of fructokinase is not associated with any clinical manifestations. Fructosuria is an accidental finding usually made because the asymptomatic patient's urine contains a reducing substance. No treatment is necessary, and the prognosis is excellent.

Solution to Question 19:

Tomcat urine is characteristically seen in multiple carboxylase deficiency.

It is named so because of its peculiar odour.

Solution to Question 20:

The enzyme deficient in hereditary coproporphyria is coproporphyrinogen oxidase (CPOX).

Hereditary coproporphyria (HCP):

- Autosomal-dominant hepatic porphyria is caused by the deficiency of CPOX.
- Acute attacks of HCP are treated as in acute intermittent porphyria (AIP), which includes intravenous hemin and identifying and avoiding precipitating factors. Cholestyramine may be of some value for photosensitivity occurring with liver dysfunction.
- Gonadotropin-releasing hormone analogs can be effective for the prevention of cyclic attacks.

Solution to Question 21:

Deficiency of the enzyme porphobilinogen deaminase causes acute intermittent porphyria.

Other options:


- Porphyria cutanea tarda: Deficiency of Uroporphyrinogen decarboxylase (UROD)
- Variegate porphyria: Deficiency of Protoporphyrinogen oxidase (PPOX)
- X-Linked protoporphyria: Gain of function mutation of δ -aminolevulinate synthase 2 (ALAS2)

Solution to Question 22:

The most likely diagnosis is argininosuccinic aciduria due to argininosuccinate lyase deficiency.

It is accompanied by high levels of argininosuccinate levels in blood, CSF, and urine. It is also associated with friable, tufted hair known as trichorrhexis nodosa.

Image showing Trichorrhexis nodosa:



Solution to Question 23:

Accumulation of Ceramide is seen in Farber's disease due to ceramidase deficiency.

Clinical features of Farber's disease :

- Painful joint swelling, nodules, progressive joint deformity
- Hoarse cry
- Breathing difficulty
- Failure to thrive

Polio and AIDS

Question 1:

A child is found to have paralysis due to poliovirus infection. Which of the following is false regarding the paralysis seen?

- a) It is an upper motor neuron type
- b) It is an asymmetrical paralysis
- c) Strenuous physical exercise during the onset exaggerates the paralysis
- d) It can be caused by polio vaccination

Question 2:

An outbreak of paralytic polio has occurred in Nigeria. Which of the following is the most likely type to cause such outbreaks?

- a) Type 1 virus
- b) Type 2 virus
- c) Type 3 virus
- d) Type 4 virus

Question 3:

Which of the following signs may be seen in a child with polio?

- a) Tripod sign
- b) Kiss the knee sign
- c) Head Drop Sign
- d) All of the above

Question 4:

You have been appointed as a member of the acute flaccid paralysis (AFP) surveillance team. What age groups would you target for surveillance?

- a) 0-10 years

- b) 0-5 years
- c) 0-15 years
- d) 0-20 years

Question 5:

When was India declared polio-free by WHO?

- a) March 2014
- b) April 2016
- c) January 2013
- d) January 2011

Question 6:

Which of the following is not a component of Polio Endgame Strategy 2019-2023 by WHO?

- a) Stop all circulating vaccine-derived poliovirus (cVDPV) outbreaks within 120 days of detection
- b) Separation of polio surveillance and vaccine preventable disease surveillance
- c) Certify eradication of Wild Poliovirus (WPV)
- d) Interrupt transmission of all wild poliovirus (WPV)

Question 7:

At your vaccination center, you have noticed increased reporting of vaccine-associated paralytic polio (VAPP). Which of the following statement is true regarding this adverse effect of the vaccine?

- a) It occurs mostly after the first dose of oral polio vaccine
- b) It is more likely to occur if the first dose was parenteral inactivated polio vaccine
- c) It is most likely to occur if given on the same day as yellow fever vaccine
- d) It is more likely to occur if multiple parenteral injections have been given at the same time

Question 8:

While volunteering in an orphanage that takes care of disabled kids, you notice the typical hand-knee gait of polio in a child. He walks by putting his right hand on his right lower thigh

while stepping forward. Which of the following muscle weakness is responsible for this finding?

- a) Gluteus medius
- b) Quadriceps
- c) Gastrocnemius-soleus
- d) Gluteus maximus

Question 9:

Serological testing of a neonate born to an HIV-infected mother turned out positive. Which of the following is the most likely route by which the newborn might have acquired the infection?

- a) Intrapartum transmission
- b) Breastfeeding
- c) Intrauterine transmission
- d) Amniocentesis

Question 10:

You are evaluating children infected with HIV in a village, where there is little awareness about HIV and its treatment. Which of the following will be the most common presenting feature in these children?

- a) Gastrointestinal infections
- b) Persistent lymphadenopathy
- c) Oral thrush
- d) Failure to thrive

Question 11:

Match the following HIV cases with their respective stages.

- a) 1-d, 2-a, 3-b, 4-c
- b) 1-c, 2-b, 3-c, 4-a
- c) 1-a, 2-c, 3-d, 4-b
- d) 1-b, 2-d, 3-a, 4-d

Question 12:

A 28-year-old woman in labor comes to the emergency department at 37 weeks gestation. She never had a prenatal checkup. The routine screening revealed she is HIV-positive. Which of the following method is least likely to prevent the perinatal transmission of HIV?

- a) Elective cesarean section
- b) Avoiding breastfeeding
- c) Administration of ART prophylaxis to baby after birth
- d) Vaginal cleansing before delivery

Question 13:

A patient is started on HAART treatment for HIV. Her CD4 count is 150 cells/ μ l. She developed a fever and her symptoms worsened during her first week of treatment. Which of the following is the most likely diagnosis?

- a) AIDS related inflammatory syndrome
- b) Immune reconstitution inflammatory syndrome
- c) HIV related hypereactivity syndrome
- d) CD4 responsive inflammatory syndrome

Question 14:

Which of the following is an indication to start ART in children, as per the WHO guidelines?

- a) <11 months infant of CD4 count <1500 cells/ mm^3
- b) 12-35 months if CD4 count <500 cells/ mm^3
- c) 36-59 months if CD4 count <350 cells/ mm^3
- d) Any child with HIV

Question 15:

A mother brings her 6-week old infant to your PHC for immunization. The baby was not administered the BCG vaccine as he is HIV positive. His recent CD4 counts were 13%. Which of the following vaccines should you not administer in this child?

- a) OPV
- b) Hepatitis B
- c) DPT
- d) Hib

Question 16:

Which of the following is not a part of the 2025 HIV targets of Global AIDS Strategy 2021-2026?

- a) 95% of women of reproductive age have their HIV and sexual and reproductive health service needs met.
- b) 90% of people living with HIV receive preventive treatment for TB
- c) 95% of pregnant and breastfeeding women living with HIV have suppressed viral loads.
- d) 75% of HIV-exposed children are tested by two months of age and again after cessation of breastfeeding

Question 17:

A 26-year-old pregnant labourer who is HIV-positive delivers a healthy male baby. She was advised ART previously but has been non-compliant with her treatment. What is recommended by NACO with regards to feeding the baby?

- a) No breast-feeding
- b) Minimal breast-feeding with complementary feeding
- c) Exclusive breast-feeding
- d) Safe & sustainable alternative formula feeds


Question 18:

You are an intern posted in the postnatal ward of a hospital. You receive an HIV-positive patient on PND-1. Which of the following tests would you use to detect HIV in the baby?

- a) HIV DNA PCR
- b) Cord Blood ELISA
- c) Western Blot
- d) 3rd Generation ELISA

Question 19:

The image given below depicts the mode of infection with which of the following organisms?



- a) *Campylobacter jejuni*
- b) Hepatitis A
- c) Polio
- d) Rota virus

Question 20:

Kenny packs are used in the treatment of _____

- a) Poliomyelitis
- b) Muscular dystrophy
- c) Polyneuropathies
- d) Nerve injury

Answer Key

Question No.	Correct Option
1	a
2	a

3	d
4	c
5	a
6	b
7	a
8	b
9	a
10	d
11	c
12	d
13	b
14	d
15	a
16	d
17	c
18	a
19	c
20	a

Detailed Explanations

Solution to Question 1:

Poliovirus mainly affects the anterior horn cells of the spinal cord and medulla oblongata (cranial nerve nuclei), causing lower motor neuron (LMN) type of paralysis. It damages motor and autonomic nerves; sensory nerves are not involved.

Poliovirus is a non-enveloped, positive-stranded RNA virus belonging to the Picornaviridae family. Three antigenically distinct serotypes (types 1, 2, and 3) are known. Ages 6 months to 5 years are most commonly affected. The virus is transmitted via the fecal-oral route and has an incubation period of 8-12 days. Humans are the only known reservoir for the polioviruses.

The virus colonizes and replicates in the GI mucosa (M cells), followed by entry into the blood via regional lymph nodes. In blood, it causes viremia and gets access to the CNS via the peripheral nerves, finally reaching the anterior horn cells in the spinal cord.

Most cases are subclinical. Some have minor illnesses like fever, sore throat, headache, and malaise. It can present as non-paralytic polio (aseptic meningitis) or paralytic polio (flaccid paralysis with absent deep tendon reflexes). Most frequently, the completely affected muscle is the tibialis anterior. It is asymmetric paralysis which is more severe if strenuous physical exercise occurred during the onset of paralysis. Death usually occurs due to respiratory paralysis.

Vaccination schedule for polio:

- Live attenuated: oral polio vaccine (OPV) (Sabin) at birth (day 0); then boosters at 6, 10, and 14 weeks and at 15-18 months; additional doses till 5 years
- Inactivated polio vaccine (IPV) (Salk) given intradermally; two fractional doses at 6 and 14 weeks as well as 9 months of age.

Solution to Question 2:

Most of the outbreaks of paralytic polio are caused by the type 1 virus.

Three antigenically distinct serotypes (types 1, 2, and 3) of poliovirus are known. Poliovirus type 1 has historically been the predominant cause of poliomyelitis worldwide and continues to be transmitted in endemic areas.

Most infections with poliovirus are asymptomatic. Rarely, patients can have the following presentations:

- Abortive poliomyelitis - a minor illness, manifested by fever, malaise, sore throat, or headache.
- Non-paralytic poliomyelitis - manifested by aseptic meningitis.
- Paralytic poliomyelitis - aseptic meningitis followed by severe back, neck, and muscle pain with rapid/gradual onset of motor weakness. It is the least common presentation of poliovirus infection.

Wild poliovirus type 2 was eradicated in 1999. Until 2015, over 90% of cVDPV (circulating vaccine-derived polioviruses) cases were due to the type 2 component in OPV.

Bivalent oral polio vaccine (OPV) contains serotypes 1 and 3. Serotype 2 has been removed, as it is implicated in the pathogenesis of cVDPV and occasionally vaccine-associated paralytic polio (VAPP).

Note:

On 25 April 2016, India switched from trivalent oral polio vaccine (OPV) to bivalent OPV; this day is called the National switch day.

Solution to Question 3:

All of the above signs may be seen in a child suffering from polio.

- Tripod sign - When the child is asked to sit up unassisted, the child assumes the tripod posture.
- Kiss the knee sign - The child cannot kiss the knee due to spine stiffness.
- Head drop sign - A hand is placed under the patient's shoulder and the trunk is raised. Usually, the head will also raise, but in the case of poliomyelitis, the head drops backward.

Solution to Question 4:

Acute flaccid paralysis (AFP) surveillance is done in children from 0-15 years.

Every case of acute flaccid paralysis in any child under 15 years is to be reported promptly. Fecal samples of all these cases are obtained (2 samples) and transported to the laboratory within 72 hours of collection at 4–8 degree Celcius or frozen at -20 degree Celcius. This process is known as the reverse cold chain.

The differential diagnosis for acute flaccid paralysis include the following:

- Non-polio enteroviruses
- West Nile virus
- Guillan–Barre syndrome
- Other neurotropic viruses: rabies, varicella-zoster, and Japanese encephalitis

Solution to Question 5:

India was certified polio-free on March 27, 2014.

- The last case of polio reported in India was in January 2011.
- Type 2 virus was declared eradicated in September 2015.
- Switch from trivalent to bivalent vaccine was done in India in April 2016.

Solution to Question 6:

Integration, not separation, of polio surveillance with vaccine-preventable disease (VPD) surveillance, is a part of Polio Endgame Strategy 2019-2023.

Goals of the Polio Endgame Strategy 2019–2023

Goal One: Eradication

- Interrupt transmission of all wild poliovirus (WPV)
- Stop all circulating vaccine-derived poliovirus (cVDPV) outbreaks within 120 days of detection and eliminate the risk of emergence of future VDPVs

Goal two: Integration

- Contribute to strengthening immunization and health systems to help achieve and sustain polio eradication
- Ensure sensitive poliovirus surveillance through integration with comprehensive vaccine-preventable disease (VPD) and communicable disease surveillance systems
- Prepare for and respond to future outbreaks and emergencies

Goal Three: Certification & Containment

- Certify eradication of WPV
- Contain all polioviruses

Solution to Question 7:

Vaccine-associated paralytic polio (VAPP) occurs mostly after the first dose of oral polio vaccine (OPV).

VAPP is said to occur if:

- Acute flaccid paralysis occurred 4 to 40 days after receiving OPV, and residual weakness lasted for more than 60 days after the onset of paralysis.
- Isolation of vaccine-related poliovirus in any stool samples and no isolation of wild poliovirus from any stool samples.

Note:

- VAPP: Vaccine-associated paralytic polio
- VDPV: Vaccine-derived polio virus. A very rare strain of polio virus, genetically changed from the original strain contained in OPV.
- cVDPV: circulating vaccine-derived polio virus. In areas of low vaccination coverage, a VDPV strain may revert to a form that can cause cases of paralytic polio, leading to an outbreak. This form is called a cVDPV.

Solution to Question 8:

Polio commonly affects the quadriceps, and weakness of the quadriceps leads to the buckling (bending) of the knee while walking or bearing weight on the leg. So, in order to walk, the person has to prevent the buckling of the knee by supporting with the hand, producing a hand-knee gait.

Examination findings in paralytic polio:

A. In the early stage

- Asymmetric motor paralysis - involvement of affected muscles is haphazard. No associated sensory loss.
- Commonly affects the lower limbs
- Most common muscle affected - Quadriceps, although it is most often partially paralyzed
- Most common muscle to undergo complete paralysis - Tibialis anterior
- The most affected muscle in the hand - Opponens pollicis
- Bulbar/bulbospinal polio - It is a rare but life-threatening disease affecting the motor neurons of the medulla and resulting in the involvement of the respiratory and cardiovascular centers, which may cause death.

B. In the late-stage

- Weakness, wasting, and deformity of the limbs
- At the hip-flexion-abduction-external rotation

- At the knee, flexion deformity is common; in severe cases, there is triple deformity: flexion, posterior subluxation, and external rotation
- At the foot, the commonest deformity is equino-varus. Other deformities - equinovalgus, calcaneo-valgus, and calcaneo-varus
- In the upper limb: affects the shoulder and elbow, hand muscles are spared.

Solution to Question 9:

The most common cause of HIV infection in an infant is the intrapartum transmission.

The mechanism of intrapartum transmission appears to be mucosal exposure to infected blood and cervicovaginal secretions in the birth canal. The intrauterine contractions during active labor/delivery could also increase the risk of late microtransfusions.

Routes of HIV transmission in an infant:

- Intrauterine
- Intrapartum - most common
- Breastfeeding - least common

Risk factors for vertical transmission:

- High maternal viremia (most important)
- Preterm delivery (<34 weeks)
- Vitamin A deficiency
- Prolonged rupture of membranes (>4 hours)
- Low birth weight
- Low maternal CD4 count

Solution to Question 10:

Failure to thrive is the universal finding in children with HIV.

Important features in HIV-infected children include the following:

- Presenting features - failure to thrive, lymphadenopathy, oral thrush, chronic diarrhea, or respiratory symptoms.
- Most common AIDS-defining illness (opportunistic infection) in children - Pneumocystis carinii pneumonia
- Most common fungal infection - oral candidiasis
- Recurrent and chronic bacterial infections (especially, Streptococcus pneumoniae)
- Most common GI disease - chronic/recurrent diarrhea with malabsorption
- Most common renal disease - nephrotic syndrome

- Most common hematological manifestation - anemia
- Most common neoplasms - non-Hodgkin's lymphoma, primary CNS lymphoma, and leiomyosarcomas

Solution to Question 11:

Correct match is: 1-a, 2-c, 3-d, 4-b.

A 5-year old child with persistent diarrhea for 20 days - Stage 3

An asymptomatic 2-year-old with persistent generalized lymphadenopathy - Stage 1

A 7-year-old with recurrent oral ulcers and persistent parotid enlargement - Stage 2

A 4-year-old with HIV-associated cardiomyopathy - Stage 4

Solution to Question 12:

Vaginal cleansing before delivery is least likely to prevent the perinatal transmission of HIV.

Methods to prevent perinatal transmission of HIV include the following:

- Antiretroviral therapy (ART):
- Single-dose nevirapine at the onset of labor followed by a single dose to the newborn within 72 hours of birth.
- Zidovudine can also be used from the 2nd trimester onwards throughout pregnancy and for 6 weeks after birth to infancy.
- Breastfed infants should receive NVP for 6 weeks if the mother is receiving cART, or NVP for the duration of breastfeeding if the mother is not on therapy. Formula-fed infants in resource-limited countries should receive ZDV bid or NVP qd for 6 weeks.
- Elective cesarean section*
- Breastfeeding: Usually contraindicated, except in developing countries where breastfeeding may be continued for 6 months.

*Note:

Maternal cART is documented to decrease the rate of perinatal HIV-1 transmission to $\leq 2\%$, and $\leq 1\%$ if the mother's viral RNA level is $\leq 1,000$ copies/mL at delivery. Thus, elective C-section at 38 weeks of gestation should be considered only for women whose viral load is $\geq 1,000$ copies/mL in late gestation, to further reduce the risk of vertical transmission.

Guidelines for the use of antiretroviral drugs during C-section are as follows:

- For elective (planned) C-section, ART should be given prior to the operation.
- Women on lifelong ART should continue their standard ART regimen.
- In case of emergency C-section in pregnant women who are not on ART, the three-drug regimen should be started before the surgery and continued thereafter.

Solution to Question 13:

Immune reconstitution inflammatory syndrome (IRIS) is characterized by an increased inflammatory response to subclinical opportunistic infections after initiating retroviral therapy.

Features seen are fever, worsening of opportunistic infections, new manifestations like lymphadenopathy, pulmonary infiltrates. It typically occurs within 1st few weeks of starting therapy. It commonly occurs in patients with progressive disease, or severe CD4 + T-lymphocyte depletion.

Solution to Question 14:

Antiretroviral therapy (ART) should be initiated in all children with HIV, regardless of clinical stage or CD4 cell count, as per the latest WHO recommendations 2016.

Also, cotrimoxazole prophylaxis is recommended for infants, children, and adolescents with HIV, irrespective of their clinical and immune conditions.

Solution to Question 15:

The live attenuated oral polio vaccine (OPV) should be avoided in severely immunocompromised infants (CD4 cell percentage $\leq 15\%$) because of an increased risk of paralytic polio.

Vaccine	Recommendation
Oral polio vaccine (OPV)	Yes, if CD4% $\geq 15\%$ No, if CD4% $< 15\%$
Varicella and MMR	Yes, if CD4% $\geq 15\%$ No, if CD4% $< 15\%$
Pentavalent (Hep B, DPT, Hib)	Yes
Inactivated polio vaccine (IPV)	Yes
BCG	No
Live attenuated influenza vaccine (LAIV or nasal spray)	No
Injectable influenza vaccine (flu shot)	Yes

Solution to Question 16:

95%(not 75%) of HIV-exposed children are tested by two months of age and again after cessation of breastfeeding is part of the High level 2025 targets of Global AIDS Strategy 2021-2026

High-level 2025 targets

- 95% of people at risk of HIV infection have access to and use appropriate, prioritized, person-centered, and effective combination prevention options.
- 95% of women of reproductive age have their HIV and sexual and reproductive health service needs met.
- 95% of pregnant and breastfeeding women living with HIV have suppressed viral loads.
- 95% of HIV-exposed children are tested by two months of age and again after cessation of breastfeeding
- 75% of all children living with HIV have suppressed viral loads by 2023 (interim target)
- 95–95–95 testing and treatment targets are achieved within all subpopulations, age groups, and geographic settings, including children living with HIV.
- 90% of people living with HIV receive preventive treatment for TB.
- 90% of people living with HIV and people at risk are linked to people-centered and context-specific integrated services for other communicable diseases, noncommunicable diseases, sexual health and gender-based violence, mental health, drug and substance use, and other services they need for their overall health and wellbeing.

The Strategy keeps people at the center and aims to unite countries, communities, and partners across and beyond the HIV response to take prioritized actions to accelerate progress towards the vision of zero new HIV infections, zero discrimination, and zero AIDS-related deaths.

This Strategy lays out a framework for transformative action to reduce the inequalities by 2025 and to get every country and every community on track to end AIDS by 2030.

Solution to Question 17:

NACO recommends exclusive breastfeeding as the infant feeding choice in the first 6 months. This is irrespective of the fact that the mother is on ART or the infant is provided with ARV prophylaxis for 6 weeks.

Breast-feeding and replacement feed simultaneously (mixed feeds) in the first 6 months should not be done, as it leads to mucosal abrasions in the gut of the baby, thereby facilitating HIV entry.

Breast-feeding should not be stopped abruptly. Only in situations where breastfeeding cannot be done (maternal death, severe maternal illness) or individual mother's choice (at her own risk), then exclusive replacement feeding which is safe, sustainable, affordable and feasible can be started.

In resource-rich settings/developed countries, exclusive alternative formulations are recommended.

Solution to Question 18:

HIV in a baby born to an HIV-infected mother is diagnosed using HIV DNA PCR.


NACO recommends the use of a qualitative HIV-1 DNA PCR (NAT) in children aged <18 months

Antibody assays (e.g., ELISA) that are commonly used for HIV testing in adults and older children are not reliable in infants and young children because of the persistence of transplacentally acquired maternal antibodies until 18 months after birth. For this reason, HIV virologic testing (PCR) must be performed using tests that detect HIV DNA or RNA. Scheduled repeated testing is necessary because test sensitivity increases with time from HIV virus exposure. Also, a very recent infection will not be detectable by PCR.

Cord blood should not be used for testing because of the possibility of contamination of the sample with maternal blood.

Solution to Question 19:

The given image shows, infection through fecal oral route, colonization of the organism, and replication in the GI mucosa (M cells), followed by entry into blood via regional lymph nodes. In blood, it causes viremia and gets access to the CNS via the peripheral nerves, finally reaching the anterior horn cells in the spinal cord. The pathogenesis explained above is that of polio infection.



Option A: Infection with *Campylobacter jejuni* is implicated in the pathogenesis of Guillain–Barre syndrome, where the antibodies against *C. jejuni* cross-react with the gangliosides (GM1) in the peripheral nervous system.

Option B: Hepatitis A infection occurs through the fecal-oral route, and the virus replicates in the GI mucosa and is then transported to the liver.

Option D: Rotavirus causes diarrhea by causing malabsorption.

Solution to Question 20:

Kenny packs are used in the treatment of poliomyelitis.

Treatment of poliomyelitis includes the following:

- Symptomatic and supportive treatment: Analgesics, bed rest, and physiotherapy
- Sister Kenny's treatment: Hot moist packs applied to muscles to relieve pain and spasm
- Optimum positioning of limbs to prevent deformity
- Management of pure bulbar poliomyelitis consists of maintaining the airway and avoiding the risk of aspiration.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Pediatric Bacterial and Parasitic Infections

Question 1:

Which of the following is not commonly caused by *Staphylococcus aureus*?



- a) Bullous impetigo
- b) Folliculitis
- c) Acute pharyngitis
- d) Pyomyositis

Question 2:

Anu, a pre-school student presented with a fever and sore throat. She also had an acute onset generalized erythematous rash with pastia lines over the antecubital fossa. What is the most likely diagnosis?

- a) Ritter's disease
- b) Scarlet fever
- c) Kawasaki disease
- d) Urticaria pigmentosa

Question 3:

A young boy is brought to the OPD with a sore throat and a rash, which appears as shown in the image, for the past 2 days. The rash started around the neck and then spread to the trunk and extremities. On examination, rash blanches on pressure. Tongue examination shows swollen papillae. Which of the following toxins is responsible for this child's condition?



- a) Pyrogenic exotoxin
- b) Exfoliotoxin
- c) Enterotoxin
- d) Panton-Valentine leukocidin

Question 4:

A 7-year-old child abruptly developed symptoms suggestive of obsessive-compulsive disorder and Tourette syndrome. PANDAS, a pediatric neuropsychiatric disorder, is suspected. Which of the following organisms is associated with this condition?

- a) *Staphylococcus aureus*
- b) *Streptococcus pneumoniae*
- c) *Streptococcus pyogenes*
- d) *Staphylococcus epidermidis*

Question 5:

Which of the following is a major criterion in the revised Jones criteria used for the diagnosis of acute rheumatic fever in high-risk populations?

- a) Polyarthralgia
- b) Fever ($\geq 38^{\circ}\text{C}/100.4^{\circ}\text{F}$)
- c) ESR ≥ 30 mm
- d) CRP ≥ 3.0 mg/dl

Question 6:

A 10-year-old girl presents with a 2-day history of fever and joint pain. Further questioning reveals that she had a sore throat 3 weeks back for which she did not seek medical help. Acute rheumatic fever is suspected. Which of the following is not a component of the criteria used for the diagnosis of this child's condition?

- a) Prolonged PR interval
- b) Erythema nodosum
- c) Subcutaneous nodules
- d) CRP ≥ 3.0 mg/dL

Question 7:

Which of the following constitutes Hutchinson's triad?

- a) Interstitial keratitis, 8th nerve deafness, mulberry molars
- b) Interstitial keratitis, 8th nerve deafness, Hutchinson molars
- c) Interstitial keratitis, 8th nerve deafness, mulberry incisors
- d) Interstitial keratitis, 8th nerve deafness, Hutchinson incisors

Question 8:

A 1-year-old baby is brought to the paediatrician by his mother because of jaundice. On examination, the baby has hepatosplenomegaly and lymphadenopathy. He has a saddle nose and persistent rhinitis as seen in the picture. The baby had been 2300g at birth. The delivery was complicated by preterm labour and a single, painless ulcer with a clean base on the mother's labia. Which of the following feature will not be seen in this baby?



- a) Coombs positive hemolytic anemia
- b) Condylomata lata
- c) Painless synovitis
- d) Interstitial keratitis


Question 9:

A neonate was delivered via normal vaginal delivery. His mother was diagnosed with active tuberculosis infection during pregnancy but she refused treatment in fear of side effects to the baby. If the baby had been infected prenatally, which of the following organs is most likely to be involved?

- a) Lung
- b) Liver
- c) Abdomen
- d) Kidney

Question 10:

A 12-year-old boy presented with a persistent cough. There was a contact history with an active tuberculosis case in his neighborhood. His X-ray revealed prominent hilar lymphadenopathy as shown in the image. Which of the following statements is false regarding this patient's condition?



- a) Puhl's lesion at the apex of lung is seen in primary infection
- b) Most cases of pediatric TB are asymptomatic
- c) Sputum smear-negative TB does not rule out pediatric pulmonary TB
- d) The most common site of extrapulmonary TB is the lymph node

Question 11:

A woman with smear-positive tuberculosis is on anti-tubercular therapy (ATT). She delivered a healthy male baby at term. How is the newborn managed in this case?

- a) BCG + Rifampicin + INH + Breastfeeding
- b) Breastfeeding + INH for 2 months then BCG vaccine
- c) Breastfeeding + INH for 6 months then BCG vaccine
- d) BCG + INH for 6 months + withhold Breastfeeding

Question 12:

A 25-year-old lady is detected to have active tuberculosis at 37 weeks of gestation. She delivers a baby girl 2 days later. As per NTEP guidelines, what advice is given immediately after delivery?

- a) INH prophylaxis to be given to the neonate for 6 weeks
- b) Mother is to be started on ATT
- c) Withhold breast feeding

d) Withhold BCG vaccination

Question 13:

A worried couple brought their 4-day-old baby to the pediatrician due to difficulty in feeding and abnormal posture. The baby was delivered at home and the mother never had a prenatal checkup. On examination, the baby's body was rigid, with flexed upper limb and extended back. The umbilical cord was covered with cow dung. The mother says it's their tradition, and that the dirt helps with the falling off of the cord. Which of the following is false regarding this condition?

- a) The toxin responsible affects the excitatory neurons**
- b) Metronidazole is given as a treatment**
- c) It occurs in infants of unimmunized mothers**
- d) Case fatality rate is 70-100%**

Question 14:

A 6-year-old unvaccinated child presents with complaints of sore throat and difficulty in swallowing. The patient also has fever and hoarseness of voice and her facial appearance is as shown below. A provisional diagnosis of Diphtheria is made. Which of the following statements is true regarding this patient's condition?



- a) It can be diagnosed by the demonstration of antibody by ELISA**
- b) Immunization prevents the carrier state**

- c) Treatment of contacts is not indicated
- d) Iron has a critical value in the production of toxins

Question 15:

A diphtheria screening camp is being held at a rural school to find carriers and to provide them adequate treatment. What is the investigation of choice in this scenario?

- a) Throat swab culture
- b) Gram's stain
- c) Albert's stain
- d) Fluorescent antibody


Question 16:

Anuj has been diagnosed with diphtheria and is being treated in the ward. His 3-year-old brother, Ajay who lives in the same house has received vaccination against diphtheria about 16 months back. What is the next best step for the management of the younger sibling, Ajay?

- a) One booster dose
- b) Nothing as the child is already exposed
- c) Erythromycin + diphtheria toxoid
- d) Erythromycin only

Question 17:

A young boy is brought to the clinic with complaints of abdominal pain, diarrhea, and painful straining for 4 months. The symptoms were accompanied by rectal prolapse. Stool examination showed the following finding. Infection with which of the following helminths is most likely?



- a) *Trichinella spiralis*
- b) *Enterobius vermicularis*
- c) *Trichuris trichiura*
- d) *Ascaris lumbricoides*

Question 18:

An anxious mother brought her 4-year-old daughter to the pediatrician. The girl was passing loose, bulky stools for the past 4 days. This was often associated with pain in the abdomen. The pediatrician ordered stool examination, which showed the following organisms. Which vitamin deficiency is the child at the greatest risk for?



- a) Vitamin B1 deficiency
- b) Vitamin A deficiency
- c) Vitamin B6 deficiency
- d) Vitamin C deficiency

Question 19:

In a child, the amoebic liver abscess can be diagnosed by demonstrating which of the following?

- a) Cysts in the sterile pus
- b) Trophozoites in the pus
- c) Cysts in the hepatic venous sample
- d) Trophozoites in the feces

Question 20:

What is the most common route of acquiring *Ancylostoma duodenale* infection?

- a) Ingestion of infected food items
- b) Inhalation of droplets
- c) Penetration of skin by larva
- d) Inoculation into eroded skin

Question 21:


A child from Bihar comes with a fever. Blood examination shows sheathed microfilaria as shown in the image. Identify the organism.



- a) *Brugia malayi*
- b) *Wuchereria bancrofti*
- c) *Loa loa*
- d) *Oncocerca volvulus*

Question 22:

A boy is brought to the emergency with repeated episodes of generalized tonic-clonic seizures. History is significant for a dietary intake of pork. MRI reveals multiple ring-enhancing lesions as shown below. Which of the following organism is the likely cause for seizures in this patient?



- a) Enterobius vermicularis
- b) Taenia solium
- c) Echinococcus
- d) Taenia saginata

Question 23:

Which of the following is false regarding congenital toxoplasmosis?

- a) It occurs through hematogenous dissemination to the placenta
- b) Chorioretinitis is the most common manifestation
- c) Immunoglobulin G (IgG) is diagnostic
- d) It can manifest as hydrops fetalis

Question 24:

The most common manifestation of congenital toxoplasmosis is _____

- a) Hydrocephalus
- b) Chorioretinitis
- c) Hepatosplenomegaly
- d) Thrombocytopenia

Question 25:

Maternal antibodies that do not provide significant immunity in a neonate:

- a) Tetanus
- b) Polio
- c) Diphtheria
- d) Measles

Question 26:

What is the daily dosage of ethambutol for an adolescent of age 14 years?

- a) 20 mg/kg

- b) 10 mg/kg
- c) 40 mg/kg
- d) 30 mg/kg

Question 27:

Which of the following is not a component of Hutchinson's triad in congenital syphilis?

- a) Peg-shaped upper incisor
- b) Interstitial keratitis
- c) Sensorineural deafness
- d) Saber shin

Answer Key

Question No.	Correct Option
1	c
2	b
3	a
4	c
5	a
6	b
7	d
8	a
9	b
10	a
11	c
12	b
13	a
14	d
15	a
16	d
17	c
18	b
19	b

20	c
21	a
22	b
23	c
24	b
25	b
26	a
27	d

Detailed Explanations

Solution to Question 1:

Acute pharyngitis or sore throat is most commonly caused by group A Streptococcus (GAS) - not staphylococcus.

Staphylococcus aureus causes localized infections like furuncle, carbuncle, bullous impetigo, pyomyositis, and folliculitis (skin and soft tissue infections). It may cause bloodstream infections if there is a breach in the skin.


Solution to Question 2:

The given clinical scenario is suggestive of scarlet fever.

Scarlet fever is a delayed type of skin reaction to streptococcal exotoxin and occurs as a complication of streptococcal tonsillopharyngitis. There is a diffuse erythematous maculopapular rash, which blanches on pressure. This gives a sandpaper quality to the skin.

A characteristic physical finding in which red or pink petechial lines appear in skin creases is called Pastia's sign, Pastia lines, or Thompson's sign.

The image below shows Pastia lines in the antecubital fossa.



Solution to Question 3:

The given clinical scenario is suggestive of scarlet fever, which is caused by a pyrogenic exotoxin produced by Group A streptococcus (GAS).

In scarlet fever, the rash appears within 24–48 hours of the onset of upper respiratory infections. It begins around the head and neck and spreads over the trunk and extremities. It usually spares palms, and soles. It is a diffuse, papular, and erythematous eruption, which blanches on pressure. The cheeks are erythematous, with pallor around the mouth. The tongue is usually coated, and the papillae are swollen, giving the tongue a strawberry appearance.

The image below shows a strawberry tongue.



Solution to Question 4:

PANDAS is associated with *Streptococcus pyogenes* infection.

PANDAS stands for Pediatric Autoimmune Neuropsychiatric Disorders Associated with Streptococcal infections.

The symptoms usually occur abruptly and can include the following:

- Obsessive-compulsive disorder (OCD)
- Tic disorder
- Gilles de la Tourette syndrome (coprolalia and tics)

Additionally, children may become moody or irritable, experience anxiety attacks, and/or show separation anxiety. These symptoms are the result of an autoimmune process that leads to inflammation of the basal ganglia and the resultant disruption of cortical-striatal-thalamic axis functioning.

Features:

- Prepubertal onset
- Abrupt and episodic course
- Neurological symptoms
- Disorders associated: Tics, OCD
- Associated with a recent history of sore throat

Note: Non-infectious manifestations of streptococcal infection:

- Rheumatic heart disease
- Glomerulonephritis
- PANDAS
- Guttate psoriasis

Solution to Question 5:

Polyarthralgia is a major criterion in the revised Jones criteria used for the diagnosis of acute rheumatic fever in high-risk populations (ARF incidence ≥ 2 per 100 000 school-aged children or all-age rheumatic heart disease prevalence of ≥ 1 per 1000 population per year).

Strict adherence to the Jones criteria in areas of higher prevalence may result in under-diagnosis. Therefore, the criteria are slightly modified for moderate to high-risk populations as follows:

- The major criterion of arthritis includes polyarthralgia or monoarthritis in addition to polyarthrititis. Polyarthralgia should only be considered as a major manifestation after the exclusion of other causes.
- The minor criterion for joint involvement is monoarthralgia.
- A lower cutoff for fever is used ($\geq 38^{\circ}\text{C}$ [100.4°F] rather than $\geq 38.5^{\circ}\text{C}$ [101.3°F]).

- A lower cutoff for erythrocyte sedimentation rate (ESR) is used (≥ 30 mm/hour rather than ≥ 60 mm/hour), although the C-reactive protein (CRP) cutoff is unchanged (≥ 3 mg/dL).

Solution to Question 6:

Erythema nodosum is not a component of Jones criteria which is used for acute rheumatic fever (ARF) diagnosis. Erythema marginatum is a major manifestation of the Jones criteria. It is an evanescent, non-pruritic rash over the trunk and extremities seen in rheumatic fever.

The image below shows erythema marginatum.



Solution to Question 7:

Hutchinson's triad refers to interstitial keratitis, 8th nerve deafness, and Hutchinson incisors. It is seen in late congenital syphilis.

The image below shows Hutchinson incisors:



Congenital syphilis is due to *Treponema pallidum*. Early features include hepato-splenomegaly, jaundice, diffuse lymphadenopathy, maculopapular or vesiculobullous rash involving hands and feet, persistent rhinitis (snuffles), and condylomatous lesions. Skeletal changes like saber shins, dental abnormalities like Hutchinson's teeth, and saddle nose (depression of the nasal root) are some late findings.

Other clinical features of congenital syphilis:

- Clutton joint - Due to synovitis, with sterile synovial fluid
- Juvenile tabes and paresis
- Rhagades - Linear scars radiating from mucocutaneous fissures of the mouth, anus, and genitalia
- Mulberry molars - Abnormal 1st lower molars with excessive cusps
- Sabre shin - Anterior bowing of the mid-portion of the tibia
- Olympian brow - Bony prominence of the forehead



Solution to Question 8:

The given clinical scenario is suggestive of congenital syphilis. Coombs-negative hemolytic anemia is a characteristic feature of congenital syphilis, not coombs-positive hemolytic anemia.

Congenital syphilis results from transplacental transmission of *Treponema pallidum* or during birth by contact with infectious lesions of the mother.

Clinical features of congenital syphilis:

- Hutchinson's triad - Hutchinson teeth, interstitial keratitis, sensorineural deafness
- Clutton joint - Due to synovitis, with sterile synovial fluid
- Juvenile tabes and paresis
- Rhagades - Linear scars radiating from mucocutaneous fissures of the mouth, anus, and genitalia
- Saddle nose - Depression of nasal bridge
- Mulberry molars - Abnormal 1st lower molars with excessive cusps
- Sabre shin - Anterior bowing of the mid-portion of the tibia
- Olympian brow - Bony prominence of the forehead
- Snuffles - Persistent rhinitis

Solution to Question 9:

The liver is the most commonly involved organ in congenital tuberculosis.

Primary infection in a female just before or during pregnancy is more likely to cause congenital tuberculosis. Congenital transmission usually occurs from a lesion in the placenta through the umbilical vein. The tuberculous bacilli may pass through the umbilical vein and develop a hepatic focus. Tuberculosis in neonates generally presents with respiratory distress, hepatosplenomegaly, and lymphadenopathy.

Note:

The most common route of infection for a neonate is the postnatal airborne transmission from an adult with infectious pulmonary tuberculosis.

Solution to Question 10:

The patient had exposure to an active tuberculosis case and has hilar lymphadenopathy (image), suggestive of primary tuberculosis. Puhl's lesion is chronic tuberculosis (TB) of the apex of the lung, not seen in the primary infection.

The majority of children develop no signs or symptoms at any level.

In primary pulmonary disease, the combination of the parenchymal pulmonary lesion and the corresponding lymph node is called Ghon's complex or the primary complex. Asymptomatic pleural effusion is common and forms part of the primary complex. Erosion of a parenchymal focus of tuberculosis into the blood or lymphatic vessel can result in dissemination of the bacilli and a miliary pattern, with small nodules evenly distributed on the chest radiograph. Sputum smear-negative TB does not rule out pediatric pulmonary TB.

Chronic pulmonary tuberculosis is common in children above 7 years of age. The commonest site is the apex of the lung—Puhl's lesion. Regional lymph nodes are not involved.

The most common site of extrapulmonary tuberculosis in children is the lymph node.

NOTE:

- Ranke complex: Ghon's complex undergoes fibrosis and calcification.
- Simon focus: Hematogenous spread to the apex of the upper lobe, resulting in a post-primary infective lesion, which heals with scarring.
- Simmond's focus: Tuberculous focus in the liver.
- Weigert focus: Caseating metastatic tuberculous focus in the wall of a pulmonary vein.
- Rich focus: Tuberculous granuloma occurring in the cortex of the brain.

Solution to Question 11:

In a neonate born to a mother with smear-positive tuberculosis, breastfeeding along with INH prophylaxis for 3-6 months is recommended. BCG vaccine is recommended at birth or after completion of prophylaxis (3-6 months) in all neonates.

When a child is born to a mother with active TB, the main aim is the TB-free survival of the newborn. It involves detecting active TB lesions (congenital TB) and treating them or prevention of transmission of tubercular infection to the neonate from the mother.

Prophylaxis with isoniazid (3-6 months) is recommended in neonates born to mothers with TB who are infectious (smear-positive). Breastfeeding should be continued in these neonates and isolation is recommended only till the mother is infectious, has multidrug-resistant tuberculosis, or nonadherent to treatment. BCG vaccine is recommended at birth or after completion of prophylaxis (3-6 months) in all neonates. However, if the Mantoux test is positive and evidence of disease is found, ATT is started.

Solution to Question 12:

The mother with TB is advised to start ATT immediately irrespective of time of delivery.

The management guidelines followed in a case of active TB in a mother immediately following delivery include:

- Mother has to be started on ATT.

- Do not separate the baby from the mother.
- Do not withhold breastfeeding.
- Screening of household contacts.
- Screen the neonate for congenital/active TB.

In accordance with the NTEP 2020 guidelines, a newborn whose mother has active TB and is on ATT should be managed with prophylactic isoniazid (INH) for 6 months, BCG vaccination at birth, with the continuation of breastfeeding, after ruling out active TB.

INH prophylaxis is given irrespective of the BCG or nutritional status. 10 mg/kg is administered daily for a minimum of six months. During monthly tablet collection, the child should be closely monitored for TB symptoms to rule out active TB and also assess for any INH-related adverse effects.

BCG vaccination can be given at birth even if INH preventive therapy is planned.

In case the newborn is found to have active TB, ATT is to be initiated along with the administration of BCG vaccine.

Solution to Question 13:

The given clinical scenario is suggestive of neonatal tetanus. It is caused by a neurotoxin produced by *Clostridium tetani*, which inhibits inhibitory neurons, not excitatory neurons.

Clostridium tetani is a spore-forming obligate anaerobe commonly found in soil, dust, and gastrointestinal tracts of various animals. The baby's umbilical stump covered with cow dung serves as the source of infection. It also occurs in infants of unimmunized mothers. Tetanus typically manifests within 3-12 days of birth as progressive difficulty in feeding, rigidity, or spasms.

Tetanus toxin binds at the neuromuscular junction and enters the motor nerve; it is then transported to the alpha-motoneuron via retrograde axonal transport. Subsequently, the toxin enters the adjacent spinal inhibitory interneurons, where it prevents the release of the neurotransmitters glycine and γ -aminobutyric acid. This leads to blockade of normal antagonistic muscle inhibition, while agonist muscles undergo sustained contraction without relaxing.

Neonatal tetanus presents with progressive feeding difficulty (sucking and swallowing), irritability, and crying. Characteristic features include reduced movement or complete muscle paralysis, rigidity, and spasms. These features may or may not be associated with opisthotonos.

Treatment includes supportive care in the form of muscle relaxants (diazepam), along with the administration of human tetanus immunoglobulin (TIG) and metronidazole, which is the antibiotic of choice. Parenteral penicillin G can serve as an alternate antibiotic.

Excising the umbilical stump in a baby with neonatal tetanus is no longer recommended.

Solution to Question 14:

The image shows a bull-neck appearance due to the soft tissue edema, seen in diphtheria. Iron is essential for the production of toxins in diphtheria.

It is caused by *Corynebacterium diphtheriae*. Carriers are the most common causes of infection. Immunization does not prevent the carrier state. It spreads via droplets through the respiratory tract (most important), cutaneous lesions, and fomites.

The regulation of toxin production in the bacterium is partially controlled by iron, thus making it an important prerequisite for toxin production.

Clinical features:

- Respiratory tract diphtheria (most common):
- Thick, adherent pseudomembrane (grey or white) on one/both tonsils and adjacent pharynx. Dislodging of the membrane may cause bleeding.
- Underlying soft tissue edema and enlarged lymph nodes can cause a bull-neck appearance
- Patients with laryngeal diphtheria are at risk for suffocation because of local soft-tissue edema and airway obstruction by the diphtheritic membrane.



- Cutaneous diphtheria is an indolent, non-progressive infection characterized by a superficial, ecthyma-like, non-healing ulcer with a grey-brown membrane.
- Occasionally, it causes mucocutaneous infections at other sites, such as otitis externa, purulent conjunctivitis, and vulvo-vaginitis.

Diagnosis: The culture of the organism from the nose and throat swab specimens is the gold standard. Demonstration of the toxin is done by Elek's gel precipitation test. Gram stain and fluorescent antibody tests are unreliable.

Complications include obstruction due to pseudomembrane, myocarditis, and polyneuropathy.

Solution to Question 15:

The investigation of choice for diphtheria carriers is throat swab culture.

Specimens for culture should be obtained from the nose and throat and any other mucocutaneous lesion. A portion of the membrane should be removed and sent for culture along with the underlying exudate in those with overt disease. Direct smear using Gram stain or specific fluorescent antibody is unreliable.

Solution to Question 16:

In the given clinical scenario, a 3-year-old child is likely to have received the 4th dose (1st booster) of the diphtheria vaccine between 16–24 months of age according to National Immunization Schedule (i.e., 16 months ago). Since he has received a booster dose within 5 years, only erythromycin is indicated in case of close contact with the diphtheria patient.

All household contacts with diphtheria patients should be closely monitored for 7 days. Culture should be taken from both the throat and nose. All contacts should receive antibiotic prophylaxis (benzathine penicillin G or erythromycin), regardless of immunization status.

Children who have not received a booster dose of the diphtheria toxoid vaccine within 5 years should be given the vaccine in an age-appropriate form. Children who have not yet received their 4th dose should be vaccinated accordingly.

If fewer than three doses of diphtheria toxoid have been given, or vaccination history is unknown, an immediate dose of diphtheria toxoid should be given and the primary series completed according to the current schedule.

Asymptomatic carriers of diphtheria are treated with antimicrobial prophylaxis for a period of 10-14 days. If an individual has not received a booster dose of diphtheria toxoid within 1 year, an age-appropriate preparation of the vaccine should be administered promptly.

Droplet precautions (respiratory colonization) or contact precautions (cutaneous colonization) of cases and carriers are maintained until two consecutive negative cultures (collected 24 hours apart) are obtained after cessation of therapy.


Solution to Question 17:

The image shows a barrel-shaped ovum, which along with rectal prolapse is characteristic of trichuris trichiura.

Trichuriasis is caused by *T. trichiura* (whipworm), which inhabits the cecum and ascending colon. They bury their thin heads into the intestinal lining causing edema, loosening the elastic epithelium, and weakening the surrounding muscles. This leads to loss of supports and rectal prolapse. Other clinical features include chronic dysentery, anemia, poor growth, and developmental defects.

Diagnosis is done by fecal smears showing the barrel-shaped ovum of *T. trichiura*.

The image below shows the barrel-shaped ovum of *Trichuris trichiura*.



Albendazole is the drug of choice (400 mg, orally, for 3 days, for all ages).

Solution to Question 18:

The stool examination (image) shows pear-shaped trophozoites having two nuclei and a flagellum, typical of *Giardia lamblia*. Giardiasis is associated with a deficiency of fat-soluble vitamins like vitamin A.

Giardia lamblia is transmitted via contaminated drinking water.

Clinical features include acute infectious diarrhea or chronic diarrhea, failure to thrive, and abdominal pain. Stools initially may be profuse and watery and later become greasy and foul-smelling. Stools do not contain blood, mucus, or leukocytes. It is associated with fat-soluble vitamin deficiencies, iron deficiency, and growth stunting.

Diagnosis is done by stool enzyme immunoassay (EIA) or direct fluorescent antibody tests for *Giardia* antigens; EIA is superior to microscopy.

Treatment includes tinidazole and nitazoxanide.

Solution to Question 19:


The amoebic liver abscess can be diagnosed by demonstrating Trophozoites in the pus

Amebiasis is caused by *Entamoeba histolytica*. Trophozoites are responsible for tissue invasion and destruction.

Amoebic Colitis will present as colicky abdominal pains and frequent bowel movements (6–8/day). Diarrhea is associated with tenesmus, heme-positive stool. Trophozoites invade the intestinal mucosa and produce flask-shaped ulcers.

Amoebic liver abscess occurs in <1% infected individuals. Fever is the hallmark of amoebic liver abscess. It is associated with abdominal pain, abdominal distention and enlargement, and tenderness of the liver. Changes at the base of the right lung, such as elevation of the diaphragm and atelectasis or effusion, can also occur. The most common finding is a single abscess in the right hepatic lobe, detected on USG, CT, or MRI.

The image below shows a CT scan of a right lobe liver abscess.



Invasive amebiasis is treated with nitroimidazoles such as metronidazole or tinidazole. This is followed by a luminal amebicide such as paromomycin.

Solution to Question 20:


Skin penetration is the most common route by which the larva of *Ancylostoma duodenale* (hookworm) enters the human body.

Ancylostoma duodenale (hookworm) larvae enter the human host by skin penetration and undergo extraintestinal migration through the venous circulation to the lungs. They enter alveoli and climb up to the bronchial tree and pharynx, and are swallowed. Larvae reach the small intestine and become adult worms.

Clinical manifestations include iron deficiency anemia, resulting from chronic occult intestinal blood loss. Dermatitis, sometimes referred to as ground itch, may occur when they penetrate human skin. Eosinophilic pneumonia can also be seen.

Infection with a zoonotic hookworm, especially *Ancylostoma braziliense*, can result in the lateral migration of the larvae causing cutaneous tracts of cutaneous larva migrans.

The image below shows the lifecycle of the intestinal hookworm.



The image given below shows cutaneous larva migrans.



Treatment:

- Mebendazole 100 mg bid, orally, for 3 days or single dose of 500 mg
- Albendazole 400 mg, orally, once

Solution to Question 21:

The incidence of *Brugia malayi* is more in children from Bihar and the image showing microfilaria with two nuclei at the tail. Hence the organism is *Brugia malayi*, which causes lymphatic filariasis.

Brugia malayi (Malayan filariasis), *Brugia timori*, and *Wuchereria bancrofti* (bancroftian filariasis) are thread-like nematodes that cause a similar infection called Filariasis. Infective larvae are

introduced into humans during sucking of blood by the mosquito vector (commonly, culex).

Clinical manifestations include episodic fever, lymphangitis of an extremity, lymphadenitis (especially the inguinal and axillary areas), headaches, and myalgias. Hepatosplenomegaly and generalized lymphadenopathy are often seen in children. These symptoms are caused by an acute inflammatory response triggered by the death of adult worms.

Diagnosis is done by demonstration of microfilariae in the blood. Blood samples should be obtained between 10 pm and 2 am because microfilaremia is nocturnal.

Treatment:

- Diethylcarbamazine 1 mg/kg PO as a single dose on day 1
- 1 mg/kg tid on day 2
- 1-2mg/kg tid on day 3
- 6 mg/kg/day divided tid on days 4-14

The image below shows the difference between microfilaria of various filarial nematodes.



Solution to Question 22:

The likely cause of seizures in a child who has multiple ring-enhancing lesions and dietary history of pork intake is an infection with *Taenia solium*.

Children acquire the adult form of the parasite by ingestion of undercooked pork containing larvae cysts. In the intestines, the cyst converts into adult form. The adult form releases eggs in the stool.

If the eggs are ingested (shed by tapeworm carriers or by unhygienic practices), the larvae are released from the eggs. The larva then invades through the intestines and migrate through the bloodstream to the muscles (and other organs), where they form tissue cysts called cysticercosis. When these larvae find their way to the brain, it is termed as neurocysticercosis.

Neurocysticercosis typically presents with seizures (focal or generalized). It can be associated with basilar arachnoiditis that can present with signs of meningeal irritation. Cysticerci in the tissues may present with focal findings from the mass effect. Ocular neurocysticercosis causes decreased visual acuity because of cysticerci in the retina or vitreous, retinal detachment, or iridocyclitis.

Treatment includes management of seizures. albendazole therapy (treatment of choice). Praziquantel can also be used.

Solution to Question 23:

IgG is not diagnostic for congenital toxoplasmosis.

The presence of IgG antibodies in a newborn can be due to an old infection in the mother, as IgG can cross the placenta. IgM cannot cross the placenta, hence the presence of IgM in newborns confirms the diagnosis of congenital toxoplasmosis. IgM immunosorbent agglutination assay (ISAGA) and IgA ELISA are the most useful diagnostic tests.

Toxoplasmosis is caused by *Toxoplasma gondii*. *Toxoplasma* can be transmitted to humans by three principal routes:

- Ingestion of raw or inadequately cooked infected meat
- Ingestion of oocysts through exposure to cat litter or soil
- Newly infected pregnant woman passing the infection to her unborn fetus

It can be acquired orally, transplacentally or rarely, parenterally or by transfusion.

Congenital toxoplasmosis: When a mother acquires the infection during gestation, organisms may disseminate hematogenously to the placenta. Infection may be transmitted to the fetus transplacentally or during vaginal delivery.

The characteristic triad of chorioretinitis, hydrocephalus, and cerebral calcifications may be present. Other clinical features include small size for gestational age, prematurity, persistent jaundice, mild thrombocytopenia, hydrops fetalis, and perinatal death.

Solution to Question 24:

Chorioretinitis is the most common manifestation of congenital toxoplasmosis.

The characteristic triad of congenital toxoplasmosis is chorioretinitis, hydrocephalus, and diffuse intracranial calcifications. Clinical manifestations apart from the triad include hydrops fetalis, intrauterine growth retardation, prematurity, peripheral retinal scars, persistent jaundice, and mild thrombocytopenia.

Congenital toxoplasmosis occurs when a pregnant woman acquires the infection during gestation. The organism may cross to the fetus transplacentally or at the time of vaginal delivery. The severity of the disease in the neonate is greater if the woman gets infected early in her pregnancy. The causative agent is *Toxoplasma gondii*, an obligate intracellular protozoan.

The diagnosis is made by the detection of IgM anti-toxoplasma antibodies. IgG can cross the placenta is hence, cannot confirm infection in the infant. IgM immunosorbent agglutination assay (ISAGA) and IgA ELISA are the most useful diagnostic tests.

Spiramycin can be administered for cases of acute infection in early pregnancy to reduce vertical transmission. It does not cross the placenta and therefore cannot be used to treat an infected fetus. Pyrimethamine-sulfonamide (to be avoided in the first trimester) is used when a fetal infection is confirmed or in maternal infections after 18 weeks.

Solution to Question 25:

Maternal antibodies do not provide significant immunity to polio infection in a neonate.

Administering an oral polio vaccine (OPV) dose at birth can provide early mucosal protection, safeguarding against enteric pathogens before they can hinder the intestinal mucosal immune response. Furthermore, administering OPV during the period when infants are protected by maternal antibodies may help prevent vaccine-associated paralytic polio (VAPP).

All pregnant women should receive a tetanus toxoid, reduced diphtheria toxoid, and acellular pertussis (Tdap) vaccine during each pregnancy, as early in the 27–36-week-of-gestation window as possible. The maternal antibodies formed from Tdap will provide immunity against tetanus, diphtheria, and pertussis in the newborn. Administering the Tdap vaccine earlier, between the 27th and 36th week of gestation, maximizes the transfer of passive antibodies to the infant (options A and C).

Infants less than 6 months are generally protected from measles by maternal antibodies (option D).

Solution to Question 26:

The daily dosage of ethambutol is 20 mg/kg body weight.

First-line antitubercular drugs:

Note:

- All first-line antitubercular drugs are hepatotoxic except ethambutol.
- All first-line antitubercular drugs are bactericidal except ethambutol, which is bacteriostatic.

Drug	Daily dosage(mg/kg)	Adverse effects
Isoniazid	10-15	Peripheral neuritis, hepatic enzyme elevation
Rifampin	15-20	Orange discoloration of secretions and urine
Ethambutol	20	Optic neuritis, decreased red-green color discrimination

Drug	Daily dosage(mg/kg)	Adverse effects
Pyrazinamide	30-40	Hepatotoxicity, hyperuricemia

Solution to Question 27:

Saber shin is not a component of Hutchinson's triad. Saber shin is the anterior bowing of the midportion of the tibia.

Note: Peg-shaped upper central incisor is called as Hutchinson's teeth (erupts during the 6th year of life).

Hutchinson's triad: It is a triad of stigmata seen characteristically in congenital syphilis. The components are as follows:

- Hutchinson's teeth
- Interstitial keratitis
- Sensorineural deafness

Image: Hutchinson teeth (peg-shaped upper incisors)



Measles, Mumps, Rubella and Other Viral Infections

Question 1:

A mother brought her 7-year-old child to the OPD because of fever and facial swelling for the past 2 days. She says that he has been refusing to eat because of the pain near the jaw. He has not been vaccinated since birth because of his parent's fear of side effects. On examination, a tender swelling is noted on the jaw as shown in the picture. Which of the following statement is true regarding this patient's diagnosis?



- a) Meningoencephalitis can precede parotitis
- b) Salivary gland involvement is limited to the parotids
- c) Patient is not infectious prior to clinical parotid enlargement
- d) Orchitis frequently leads to infertility

Question 2:


An outbreak of mumps infection was reported from a local school a few days ago. A boy studying in the 2nd standard of that school presented with painful bilateral swelling near his jaw and cheek for 3 days. His mother is anxious about the possible complications of the disease. Which of the following is the most common complication that can occur in this patient?

- a) Orchitis
- b) Pancreatitis

- c) Meningoencephalitis
- d) Myocarditis

Question 3:

On your pediatrics rotation, you notice a 2-day-old baby with the rash shown in the image below. Which among the following is the most likely cause for this baby's rash?



- a) Rubella
- b) Measles
- c) Erythema infectiosum
- d) Varicella

Question 4:

A woman who came with complaints of fever, myalgia, and rash at 12 weeks of gestation is found to have positive IgM titers for rubella. She is advised MTP. If the pregnancy is taken to full term, which of the following triad of complications would you expect to see in the child?

- a) Chorioretinitis, hydrocephalus, intracranial calcifications
- b) Sensorineural hearing loss, blueberry muffin rash, microcephaly
- c) Cataracts, cicatricial skin lesions, limb defects
- d) Sensorineural hearing loss, cataract, congenital heart defects

Question 5:

Palak, a neonate is being evaluated for the finding shown in the image. The mother's prenatal history is significant for viral exanthem in early pregnancy. On examination, a grade 3 murmur is heard on auscultation. Which of the following is the most likely cause for this patient's murmur?



- a) Patent ductus arteriosus
- b) Ventricular septal defect
- c) Atrial septal defect
- d) Pulmonary stenosis

Question 6:

You are evaluating a baby in NICU with diffuse petechiae. On examination, mild hepatosplenomegaly, bilateral cataract, and a continuous murmur suggestive of patent ductus arteriosus are noted. Which of the following investigations would you do to diagnose this baby's condition?

- a) RT-PCR of CSF sample
- b) Blood spot test
- c) IgM antibody in infant blood
- d) IgG antibody in infant blood

Question 7:


A 2-year-old girl is brought to the clinic with a high fever, cough, red eyes, and rash. The child's parents say that the rash started on the forehead and behind the ears, then subsequently spread to the neck and body. On examination, a confluent maculopapular rash is noted as shown below. Which of the following is false regarding this child's condition?



- a) The period of infectivity is 4 days before and 5 days after the appearance of rash
- b) The most common complication is pneumonia
- c) Warthin-Finkeldey giant cells are present
- d) Koplik spots are pathognomonic

Question 8:

A young child presented with fever and cough in the OPD. He is observed to have the finding shown in the image below on the inner aspect of cheeks. Which of the following statement is false regarding this clinical finding?



- a) It appears 1 to 4 days before the rash
- b) It appears as tiny-rose colored lesions
- c) It is seen at the level of molars
- d) It may also occur in conjunctiva

Question 9:

A 10-year-old boy presents with a history of myoclonic jerks, seizures, and rapid decline in cognitive function. The mother gives a history of fever with a rash at 1.5 years of age. The rash had started behind the ears, progressed to involve the whole body, and resolved on its own. What is the diagnosis?

- a) Subacute sclerosing panencephalitis
- b) Multiple sclerosis
- c) Meningitis
- d) Guillian-Barre syndrome

Question 10:

Akshay developed a fever, headache, and a diffuse rash as shown below. The rash is most prominent on the face and chest. He says the rash is very itchy bothering his everyday life. He is a football player and has an upcoming tournament in the next few days. He wants to know if he'll be able to play by then. Considering the rash infectivity, until when would you advise him to stay isolated?



- a) 4 days after the appearance of lesions
- b) 7 days after the appearance of lesions
- c) 14 days after the appearance of lesions
- d) After crust formation has occurred over lesions

Question 11:

There is an outbreak of chickenpox among the children at a daycare center. What is the most common extracutaneous site of involvement of this disease?

- a) Liver
- b) CNS
- c) Joints
- d) Lungs

Question 12:

Seema is brought to the OPD due to the development of a very itchy rash. Her father says he's worried because she has been constantly scratching her body leading to scratch marks. Before the development of the rash, she had a fever and headache for 2 days. On examination, you note a vesicular rash, which is in different stages of evolution. Which of the following is not a possible complication of this condition?

- a) Subacute sclerosing panencephalitis
- b) Acute cerebellar ataxia

- c) Hepatitis
- d) Herpes Zoster

Question 13:

A child presents with fever, headache, and a diffuse rash as shown below. The rash resolves by crusting and the child recovers in 10 days. Which spinal segments are most commonly involved in a latent reactivation of this infection?



- a) C2-C7
- b) T3-L3
- c) S1-S3
- d) It involves all spinal segments equally

Question 14:

You are a pediatrics resident posted in the labor room. You are examining a baby which was delivered via C-section a few minutes ago. You notice that the baby has zig-zag hypertrophic cutaneous scarring in a dermatomal distribution on the left side of the limbs and face. He also has a small head and small limbs. Antenatal history of the mother will be significant for which of the following infection?

- a) Varicella
- b) Herpes
- c) Cytomegalovirus

d) Syphilis

Question 15:

A baby in the NICU is being evaluated for his features of microcephaly and hepatomegaly. CT scan revealed nodular periventricular calcifications. Congenital CMV infection is suspected. Which of the following samples is the best to detect the organism using PCR?

- a) Liver biopsy
- b) CSF
- c) Blood
- d) Urine

Question 16:

A child is brought to the OPD with complaints of a rash and fever. The worried parents say the rash appeared on his cheeks yesterday. However, this morning, they observed it has spread to the trunk as well. The image of the child is given below. Which of the following is true regarding this condition?



- a) It is also known as sixth disease
- b) It is caused by an RNA virus
- c) It is associated with a transient arrest of erythropoiesis
- d) It is associated with immune fetal hydrops


Question 17:

Niki, a 9-month-old baby is brought to the OPD for a rash. The mother says the baby had a high fever of 103 F for 3 days for which she was given paracetamol. This morning, the baby had no fever but the mother noticed a rash on the trunk. On oral examination, ulcers at the uvulopalatoglossal junction are present. Which of the following statements is false regarding this disease?

- a) It is also called as erythema subitum
- b) Nagayama spots are commonly seen
- c) It is caused by human herpes virus-6
- d) It is a non-pruritic, morbilliform rash on the trunk

Question 18:

A 14-year-old boy came to the OPD with the complaint of fever and sore throat. On examination, you noted generalized lymphadenopathy and hepatosplenomegaly. A peripheral blood smear revealed the following picture. What is the most probable diagnosis of this patient?



- a) Infectious mononucleosis
- b) Kimura disease
- c) Scrofuloderma
- d) Acute lymphoblastic leukemia

Question 19:

Which of the following is the most common cause of infantile diarrhea worldwide?

- a) Rotavirus
- b) Lactose intolerance
- c) Norwalk virus
- d) Vibrio cholerae

Question 20:

Which of the following is not a mode of transmission of hepatitis B from the infected mother to the baby?

- a) Transplacental transmission
- b) Perinatal transmission
- c) Amniocentesis
- d) Breast feeding

Question 21:

Which fungal infection spreads in neonates through the care giver's hand?

- a) Candida albicans
- b) Candida parapsilosis
- c) Candida tropicalis
- d) Candida glabrata

Question 22:

A 16-year-old boy presented with rigidity and difficulty in movement. He has a history of involuntary movements in a single group of muscles of legs suggestive of myoclonic jerks. On examination, choreoathetosis movements and lead pipe rigidity is noted in the limbs. EEG shows suppression-burst episodes which are characteristic for subacute sclerosing panencephalitis. This clinical presentation is a complication of which of the following infection?

- a) Mumps
- b) Chickenpox

- c) Herpes
- d) Measles


Question 23:

Most common heart defect in congenital rubella syndrome is:

- a) ASD
- b) VSD
- c) PDA
- d) PS

Question 24:

A 3-year-old male has the following buccal mucosa finding. After three days, he presents to your OPD with a rash. What is your diagnosis?



- a) Measles
- b) Leukoplakia
- c) Scarlet fever
- d) Kawasaki disease

Question 25:

What is the secondary attack rate of chickenpox?

- a) 60%
- b) 40%
- c) 85%
- d) 50%

Answer Key

Question No.	Correct Option
1	a
2	c
3	a
4	d
5	a
6	c
7	b
8	b
9	a
10	d
11	b
12	a
13	b
14	a
15	d
16	c
17	a
18	a
19	a
20	d
21	b
22	d
23	c
24	a
25	c

Detailed Explanations

Solution to Question 1:

The given clinical scenario is suggestive of mumps. Meningoencephalitis can occur before, along with, or after parotitis. Most commonly, it manifests 5 days after parotitis.

Mumps is caused by RNA paramyxovirus. It is spread from person to person via droplets from clinical and subclinical cases. The incubation period is 12-21 days. The virus appears in the saliva from up to 7 days before to as long as 7 days after the onset of parotid swelling.

The patient presents with prodrome lasting 1-2 days consisting of fever, headache, and vomiting. After the prodrome, unilateral or bilateral parotitis develops, which is tender. It may be associated with ipsilateral ear pain. Ingestion of sour or acidic foods may enhance pain.

It can be prevented by administering live attenuated, MMR vaccine at 9 months, 15 months and 4-6 years.

Option B: Mumps can affect all the salivary glands, but most commonly affects parotid glands.

Option C: The patient is infectious from 7 days before to 7 days after the onset of parotid swelling.

Option D: Orchitis or oophoritis is a common finding in mumps, along with parotitis. But, mumps leading to infertility is very rare.

Solution to Question 2:

Meningoencephalitis is the most common complication that can occur in mumps in children.

Most common complications in mumps:

- Children - Meningoencephalitis
- Adolescents and adult males - Orchitis

Solution to Question 3:

The image shows purple papulonodular lesions, which are suggestive of blueberry muffin rash. It is seen in congenital rubella.

Blueberry muffin rash represents dermal erythropoiesis. In addition to congenital rubella, it can be seen in:

- Congenital viral infections such as cytomegalovirus and parvovirus
- Neoplasms such as neuroblastoma and acute myeloid leukemia
- Rh hemolytic disease

Solution to Question 4:

The patient has an increased risk of congenital rubella if the pregnancy is taken full term. The triad of complications that would be expected in the child include:

- Sensorineural hearing loss: Nerve deafness is the single most common finding among infants with congenital rubella syndrome
- Congenital cataract
- Congenital heart disease: The most common cardiac defect is patent ductus arteriosus

Option A: Chorioretinitis, hydrocephalus, intracranial calcifications - congenital toxoplasmosis.

Option B: Sensorineural hearing loss, blueberry muffin rash, microcephaly - congenital CMV infection.

Option C: Cataracts, cicatricial skin lesions, limb defects - congenital varicella syndrome.

Solution to Question 5:

The image shows a bilateral cataract, which along with the antenatal history of viral exanthem is suggestive of congenital rubella syndrome. Patent ductus arteriosus (PDA) is the most common congenital heart disease associated with rubella, followed by pulmonary artery stenosis (PS).

Frequency of heart disease in congenital rubella: PDA > PS > VSD > ASD.

Solution to Question 6:

The given clinical scenario is suggestive of congenital rubella syndrome. IgM antibody detection in infant blood are done to diagnose congenital rubella syndrome in a newborn.

Congenital rubella syndrome is diagnosed by:

- Presence of IgM rubella antibodies in the infant shortly after birth (since IgM does not cross the placenta), or
- Persistence of IgG antibodies for >6 months (maternally derived antibody would have disappeared)

Solution to Question 7:

The given clinical scenario is suggestive of measles. The most common complication of measles is acute otitis media. The most common cause of death is pneumonia (Hecht's giant cell pneumonia).

Measles (rubeola) is caused by RNA paramyxovirus. The infection is spread via droplets, with a secondary attack rate $\geq$ 90%. The patients are infectious from 4 days before upto 5 days after the appearance of the rash. The rash appears on the 4th day of fever. It starts from the retro auricular region. Koplik spots are pathognomonic.

The most common cause of death is pneumonia. Measles can lead to subacute sclerosing panencephalitis (SSPE) as a late complication.

Measles can be prevented by administering live attenuated, Edmonston–Zagreb strain or combined MMR given at 9 months of age.

Solution to Question 8:

The image shows discrete red lesions with white spots in the center on the inner aspect of the cheeks, suggestive of Koplik spots. They do not appear as tiny rose-colored lesions.

Koplik spots are pathognomonic of measles. These are the spots present at the level of molars. They appear 1-4 days prior to the rash. They may also occur in conjunctival folds and vaginal mucosa.

Forchheimer spots: Tiny red-colored lesions in the oropharynx or petechial hemorrhages on the soft palate, seen in rubella (German measles).

Solution to Question 9:

The given case scenario with a classic history of exanthematous fever suggestive of measles, seizures, myoclonic jerks, and cognitive dysfunction is suggestive of subacute sclerosing panencephalitis (SSPE).

SSPE is a chronic, progressive demyelinating disease that develops years after measles infection, most often in patients between 5-15 years of age. Patients have a history of measles infection at $\leq$ 2 years of age. CSF shows elevated measles antibody levels and MRI shows increased T2 intensity in white matter. No definitive treatment is available.

Other complications of measles:

- Meningoencephalitis
- Giant-cell pneumonitis
- Pneumonia
- Croup
- Secondary infections - otitis media
- Protracted diarrhea

Solution to Question 10:

The above scenario is suggestive of chickenpox. You would advise the boy to stay isolated until after crust formation has occurred over lesions.

People with chickenpox are contagious 24-48 hours before the rash is evident and until the vesicles have crusted, which is usually 3-7 days after the onset of rash.

Solution to Question 11:

The most common extracutaneous site of involvement by the varicella-zoster virus in children is the central nervous system. It is a benign complication of VZV infection in children and generally does not require hospitalization.

The most common infectious complication of varicella is secondary bacterial superinfection of the skin, which is caused by *Streptococcus pyogenes* or *Staphylococcus aureus*.

The most common serious complication of varicella is varicella pneumonia.

Solution to Question 12:

The given clinical scenario is suggestive of chickenpox caused by the varicella-zoster virus (VZV). Subacute sclerosing panencephalitis (SSPE) is not a complication of Varicella-zoster infection. It occurs due to measles.

The characteristic rash of chickenpox is present in different stages of evolution.

Complications of VZV occur with disseminated varicella or with reactivation of infection. This more commonly occurs in immunocompromised patients. Complications include:

- Bacterial infections of the skin and soft tissues
- Pneumonia
- Hepatitis
- Acute cerebellar ataxia
- Encephalitis
- Disseminated intravascular coagulopathy
- Sepsis
- Shingles/Herpes Zoster: vesicular lesions are confined to the unilateral segment of the sensory nerve (dermatome). In children, the rash is mild, with complete resolution within 1-2 week. Unlike in adults, postherpetic neuralgia is unusual in children.

Solution to Question 13:

The clinical scenario suggests a diagnosis of chickenpox. Latent reactivation of the infection or shingles most commonly occurs in the T3 to L3 spinal segments.

Herpes zoster/shingles are characterized by a unilateral vesicular dermatomal eruption associated with severe pain.



Herpes zoster ophthalmicus affects the ophthalmic branch of the trigeminal nerve. It is shown below:



Ramsay–Hunt syndrome (herpes zoster oticus) affects the facial nerve with an eruption on the tympanic membrane, the external auditory canal, and facial palsy.



Solution to Question 14:

The given clinical scenario is suggestive of congenital varicella syndrome. Hence, antenatal history of varicella infection would be significant in the mother of this baby.

Fetal varicella infection occurs if the mother gets infected during the first half of pregnancy.

Clinical features include:

- Cicatrix, a zigzag scarring in a dermatomal distribution
- Microphthalmia
- Cataracts
- Optic atrophy
- Chorioretinitis
- Microcephaly
- Limb hypoplasia
- Motor and sensory deficits.

Solution to Question 15:

Among the following options, urine is the best sample to be sent for the diagnosis of congenital cytomegalovirus (CMV) infection.


In congenital CMV infections, the investigation of choice is PCR or viral culture. It should be done within the first three weeks of life (after this period, it is difficult to distinguish between a congenital and an acquired infection). Since not all affected infants are viremic at birth, urine is a more reliable sample than blood.

Most CMV transmission occurs in the first trimester. The risk of transmission is high in the third trimester. 90% of the infected babies are asymptomatic and have lesser chances of having late sequelae.

The clinical manifestations include hepatosplenomegaly, petechial rashes, jaundice, chorioretinitis, and microcephaly. Neuroimaging shows intracranial periventricular calcifications. One of the long-term sequelae is sensorineural hearing loss.

There is no definitive treatment for congenital CMV infection. Treatment with ganciclovir is indicated only in patients with progressive neurologic disease and deafness.

The image below shows periventricular calcifications:



Solution to Question 16:

The image shows a characteristic slapped cheek appearance of a rash, which is classically present in parvovirus B-19 infection or erythema infectiosum. This is associated with transient aplastic crisis and erythropoiesis is mainly affected.

Erythema infectiosum is also known as the fifth disease. The rash appears first on the cheeks called the slapped cheek appearance then spreads to the trunk. It is caused by parvovirus B-19, which is a single-stranded DNA virus.

Clinical features:

- Febrile, non-toxic child with a nonspecific prodrome
- Transient aplastic anemia, pure red cell aplasia
- Petechial/ purpuric skin eruptions: Papular purpuric "gloves and socks" syndrome (PPGSS)
- Myocarditis
- Primary maternal infection is associated with non-immune fetal hydrops and intrauterine fetal demise

- Arthritis and arthralgia (adults)

Solution to Question 17:

The given clinical scenario is suggestive of roseola infantum. It is also called exanthema subitum, not erythema subitum.

Roseola infantum is also called as sixth disease or exanthema subitum (exanthema=breaking out, subitum=sudden rash). It is caused by a human herpesvirus (HHV) -6A, 6B, and 7.

Clinical features:

- Prodrome of high-grade fever
- Non-pruritic, morbilliform rash, which appears as the fever comes down.
- Leucopenia and relative lymphocytosis
- Nagayama spots: ulcers at uvulo-palatoglossal junction

Image: Exanthem subitum



Solution to Question 18:

The image of the peripheral blood film shows atypical lymphocytes, also known as Downey cells. The given clinical scenario along with characteristic atypical lymphocytes is suggestive of infectious mononucleosis.

Infectious mononucleosis, also called, glandular fever is caused by Epstein–Barr virus (EBV). It is transmitted person-to-person through salivary secretions and blood transfusion. The incubation period is 4 to 5 weeks. It presents with a classic triad of

- Fever

- Pharyngitis
- Lymphadenopathy

Other clinical features include fatigue and hepatosplenomegaly. The patients are advised to avoid contact sports for 3-4 weeks due to the risk of splenic rupture.

The screening test of choice is the monospot test. This is a heterophile antibody test. Atypical lymphocytosis of $>10\%$ is a characteristic laboratory finding. These are CD8 T lymphocytes that develop in response to the infected B lymphocytes.

Other diseases caused by EBV include:

- Hodgkin's lymphoma
- Non-Hodgkin lymphoma
- Burkitt's lymphoma
- Post-transplant lymphoproliferative disease
- Nasopharyngeal carcinoma

Option B: Kimura disease is a chronic inflammatory disorder of unknown etiology that most commonly presents as painless lymphadenopathy.

Option C: Scrofuloderma is the result of cutaneous infection adjacent to a tuberculous focus, which may correspond to peripheral ganglionic tuberculosis (the most common form of extrapulmonary tuberculosis in HIV-positive patients and in children).

Solution to Question 19:

The most common cause of infantile diarrhea worldwide is rotavirus.

Rotavirus causes diarrhea in children of 3-5 years of age, via the fecal-oral route. The incubation period is 1-3 days. Most cases are asymptomatic or with mild symptoms like watery diarrhea, vomiting, low-grade fever, and temporary lactose intolerance. This is presumably because of the protection by maternal antibodies in breast milk. Diagnosis can be made by rapid antigen detection of rotavirus in stool specimens

Lactose intolerance is actually not very common in infants and typically only starts showing up after age 3. All babies are born with lactase in their intestines. As they grow older, the lactase enzyme decreases.

Solution to Question 20:

Breastfeeding is not a mode of transmission of hepatitis B from an infected mother to the baby.

Mother-to-child transmission of HBV can occur via the following three routes:

- Intrauterine transmission
- Transmission during delivery (most frequent method)

- Postpartum transmission

Postpartum transmission: A proportion of babies may acquire the infection after birth due to close contact with the mother. However, breastfeeding is not a risk factor for HBV infection.

Transmission following amniocentesis has been described, but the risk appears low.

Newborns born to seropositive Hepatitis B mothers are given Hep B immunoglobulin as well as Hep B vaccination at birth.

Solution to Question 21:

Candida parapsilosis is increasingly responsible for hospital outbreaks, and the hands of healthcare workers may be predominant environment sources.

Solution to Question 22:

Subacute sclerosing panencephalitis (SSPE) is a rare complication of measles. Hence, prior infection with measles is most likely.

SSPE occurs 3-8 years after a primary measles infection. It is the least common complication of measles but is almost always fatal.

The clinical manifestations occur in 4 stages:

Stage I: Maybe missed due to mild symptoms like irritability and reduced attention span.

Stage II: Hallmark is massive Myoclonus. Involuntary muscle movements and myoclonic jerks start from one group of muscles but gives way to massive spasms and jerks.

Stage III: Involuntary movements disappear and are replaced by choreoathetosis, immobility, dystonia, and lead-pipe rigidity due to destruction of deeper parts of basal ganglia.

Stage IV: Loss of critical centers that lead to coma and then death.

Diagnosis of SSPE is by:

- Detection of measles antibody in CSF
- Characteristic findings on EEG
- Typical histological findings or/and isolation of virus or virus antigen from brain tissue obtained by biopsy or after postmortem.

Solution to Question 23:

Patent ductus arteriosus (PDA) is the most common congenital heart disease associated with rubella, followed by pulmonary artery stenosis (PS).

Cardiac anomalies have been observed in 50% of children infected during the first 8 weeks of gestation. The incidence of heart disease in congenital rubella syndrome is: PDA > PS (right

>left) > ventricular septal defect > atrial septal defect.

Solution to Question 24:

The given image shows small red lesions with bluish-white spots in the center on the inner aspects of the cheek opposite to the lower premolar, indicating the presence of Koplik spots. The appearance of these spots, along with a rash appearing three days later, strongly suggests a diagnosis of measles.

The measles virus is a single-stranded RNA virus. Measles infection has 4 phases: the incubation period, prodromal illness, exanthematous phase (characterized by the rash), and recovery phase.

After an incubation period of 8-12 days, the prodromal phase of measles starts with mild fever and is followed by conjunctivitis, photophobia, coryza, cough, and increasing fever. Koplik spots, appearing 1-4 days prior to the rash, are a characteristic sign of measles.

Symptoms gradually intensify over a period of 2-4 days until the first day of the rash. The rash typically begins on the forehead (around the hairline), behind the ears, and on the upper neck as a red maculopapular eruption. It then spreads downward to the torso and extremities, often becoming confluent on the face and upper trunk.

Complications include acute otitis media (most common), pneumonia (the most common cause of death), croup, bronchiolitis, reactivation of tuberculosis, diarrhea, appendicitis, febrile seizures, and encephalitis. Subacute sclerosing panencephalitis is a rare chronic complication of measles.

Other options:

Option B: Leukoplakia appears as a white patch in the oral cavity that cannot be wiped off.

Option C: Scarlet fever is Group A Streptococcus (GAS) pharyngitis and is characterized by a distinctive sandpaper rash. The rash appears as a diffuse, finely papular, erythematous eruption that turns the skin bright red and blanches on pressure. It starts around the neck and spreads to the trunk and extremities.

Option D: Kawasaki disease (KD) is a systemic inflammatory disorder with arteritis of medium to large-sized vessels. It presents with fever, rash (maculopapular, erythema multiforme urticarial, or micropustular), conjunctivitis, and cervical lymph node enlargement.

Solution to Question 25:

The secondary attack rate of chicken pox is 85% (range 61–100%)

Chicken pox:

- Causative agent: Varicella–zoster virus
- Incubation period: 2 to 3 weeks
- Rash: On 1st day, centripetal, pleomorphic, vesiculopapular lesions with erythematous base (dew drops on rose petal)

Treatment: Acyclovir 20 mg/kg/dose; four doses per day; for 5 days

Complications:

- Cutaneous: Secondary bacterial superinfection of the skin
- Systemic: Acute cerebella ataxia, Guillain–Barre syndrome, transverse myelitis, optic neuritis, facial nerve palsy, pneumonia, Reye's syndrome
- Aspirin is contraindicated due to risk of Reye's syndrome.

Vaccine:

- Oka strain (live attenuated)
- All adults who have never had chickenpox should receive 2 doses 0.5 mL in the deltoid area subcutaneously.
- For <13 years of age, the first dose is administered at 12–15 months and the second dose at age 4–6 years.
- For people older than 13 years, the two doses are administered 4–8 weeks apart.
- For those already immunized, booster doses are not needed if titers are adequate.

Surgical GI Disorders

Question 1:

A young boy came to visit his mother admitted to the antenatal ward for her delivery. You are the paediatrician on rounds in the adjacent post-natal cubical. His speech was very different from a normal child. Out of curiosity, you examine his oral cavity and the following finding is seen. What would be the most likely cause of this condition?



- a) Abnormally short lingual frenulum
- b) Abnormally long lingual frenulum
- c) Fissured tongue
- d) Abnormally large tongue

Question 2:

Sirish, a young talented singer failed to make it through the Music Idol audition. Deeply affected by this, he drank a full bottle of a drain unclogger. He was rushed to the ER with froth in the mouth. Which of the following could be done as the first step in treatment?

- a) Administer mild acid
- b) Administer milk
- c) Administer rice water
- d) Gag the child to induce emesis

Question 3:

Mala, a young student of class 9 accidentally pipettes sulphuric acid while performing volumetric analysis in her chemistry lab. Which type of mucosal damage would you expect in her?

- a) Liquefactive necrosis
- b) Coagulative necrosis
- c) Saponification of fat
- d) Thermic reaction

Question 4:

A 48-hours-old boy baby born at 35 weeks of gestation regurgitates the feeds given to him. Abdominal examination reveals an airless scaphoid abdomen. A contrast study is performed and the findings are as shown below. What is the possible diagnosis?



- a) Esophageal atresia in TEF type A
- b) Esophageal atresia in TEF type D
- c) Esophageal atresia in TEF type C
- d) Esophageal atresia in TEF type B

Question 5:

A full-term infant with a history of polyhydramnios in utero presents with respiratory distress soon after birth. On examination frothing at the mouth is observed. What would be the immediate investigation of choice?

- a) Chest radiography
- b) Lateral neck radiography
- c) Passing a nasogastric tube with subsequent chest radiography
- d) Erect X-ray abdomen

Question 6:

Milind, a young child is brought to you for recurrent respiratory symptoms and dysphagia. After ruling out other causes, you suspect a vascular anomaly that could be compressing the esophagus to be causing the symptoms. A CECT is done and the findings are as shown below. Which of the following vascular anomalies is most likely to cause this condition?



- a) Aberrant left subclavian artery
- b) Aberrant right subclavian artery
- c) Left sided azygous vein
- d) Truncus arteriosus

Question 7:

You are a new intern posted in the department of pediatrics. The lab reports of a newborn baby admitted to the SNCU arrive and the ABG report is as below. Which congenital anomaly

would you suspect in such a setting of electrolyte and acid-base imbalance?

- a) Gastric volvulus
- b) Trachea-esophageal fistula
- c) Congenital hypertrophic pyloric stenosis
- d) Malrotation with volvulus

Question 8:

A 5-week-old infant presented with non-bilious projectile vomiting and lethargy for 2 days. On abdominal examination, an olive-like mass was palpated. Barium study showed an elongated pyloric channel. Which drug is used in the first week of life can cause this presentation?

- a) Misoprostol
- b) Erythromycin
- c) Thalidomide
- d) Phenytoin

Question 9:

Savitri, a 3-week-old baby is having bouts of non-bilious projectile vomiting. On examination, the child seems lethargic and dehydrated. Abdominal examination reveals visible gastric peristalsis. What is the investigation of choice for the diagnosis of this condition?

- a) Barium meal
- b) Ultrasound
- c) CT scan
- d) MRI

Question 10:

Ashwini, the mother of a young baby is concerned after she noticed straw yellow-colored stains in her child's clothing while washing them. She brings the child to you for examination. The child has a mildly elevated temperature and the fluid comes out from the umbilicus, whenever the child strains. What is the diagnosis?

- a) Patent vitellointestinal duct
- b) Urachal fistula

- c) Patent omphalomesenteric duct
- d) Bladder diverticulum

Question 11:

On observing the following finding in the X-ray of a baby, what is the most likely diagnosis you would suspect?



- a) Duodenal atresia
- b) Ileal atresia
- c) Jejunal atresia
- d) Sigmoid volvulus

Question 12:

An infant was brought to the paediatrician with complaints of vomiting and abdominal distention. The parents inform that the baby passes stools that look like currant jelly. On ultrasonography, a target sign was observed. Which of the following is not true about this condition?

- a) Rota virus vaccine is not associated with increased risk
- b) Submucosal lipomas can cause recurrence
- c) The most common cause is idiopathic
- d) The most common location is ileocolic region

Question 13:

A 4-year-old child is admitted to your unit with recurrent pancreatitis. Which of the following developmental anomalies is likely to be associated with this condition?

- a) Pancreas divisum
- b) Annular pancreas
- c) Pancreatic pseudocyst
- d) Small ectopic pancreas

Question 14:

A 7-year-old boy was playing football with his friends during the PET classes. Suddenly he complained of severe abdominal pain and he is rushed to the pediatric ER. An erect X-ray scan of the abdomen showed the following picture. What is the most likely diagnosis?



- a) Intussusception
- b) Gastric volvulus
- c) Sigmoid volvulus
- d) Meckels diverticulum

Question 15:

A 12-year-old boy presents to the emergency department with sudden-onset severe generalised abdominal pain associated with fever, nausea, vomiting, and diarrhoea. On

examination, his abdomen is tense with generalised tenderness, guarding and McBurnys tenderness is present. No bowel sounds are present. What is the best diagnostic investigation in this child?

- a) USG
- b) MRI
- c) CT scan
- d) X-ray

Question 16:

Aaradhana, a 14-year-old girl is feeling embarrassed as she has to often rush to the restroom during school hours. Her friends also make fun of her as she blushes frequently for no reason. She was later diagnosed with an appendicular tumour. Labs revealed low levels of niacin with the tumour being positive for synaptophysin. Which of the following biochemical moieties can be demonstrated in the patient's urine sample?

- a) 5-hydroxy indole acetic acid (HIAA)
- b) vanillylmandelic acid (VMA)
- c) ACTH
- d) catecholamines

Question 17:

A 14-year-old boy presents with recurrent colicky abdominal pain and melena. On examination, you note the following lesions in the buccal mucosa. You suspect Peutz-Jeghers syndrome and advice for the endoscopic investigation to look for polyps. In which part of the intestine would you expect to find maximum polyps?



- a) Jejunum
- b) Appendix
- c) Colon
- d) Gastric mucosa

Question 18:

A full-term neonate was found to have distention of the abdomen. He also passed his meconium late and based on further workup was later diagnosed with Hirschsprung disease. Which of the following conditions is least likely to be found in association with this?

- a) Von Recklinghausen's disease
- b) Joubert syndrome
- c) Edward syndrome
- d) Downs syndrome

Question 19:

A neonate is being evaluated in the nursery for failure to pass meconium in the last 48 hours. Physical examination reveals abdominal distension and an anal dimple. Wangenstein's invertogram shows a low lying defect. Further evaluation reveals the presence of vertebral and renal deformities. Which of the following conditions is least likely to be seen in this child?

- a) Cardiac defects

- b) Tracheal anomalies
- c) Radial aplasia
- d) Lymphedema

Question 20:

You notice a 5-year-old boy walking very differently in an orphanage. On probing further the guardian informs that he has a history of frequently soiling his pants. On examination, you find a mass protruding through his anus and a diagnosis of rectal prolapse is made. Which of the following would be the most likely cause for this presentation?

- a) Pertussis
- b) Constipation
- c) Cystic fibrosis
- d) Trichuriasis

Answer Key

Question No.	Correct Option
1	a
2	b
3	b
4	a
5	c
6	b
7	c
8	b
9	b
10	b
11	a
12	a
13	a
14	c
15	a
16	a

17	a
18	c
19	d
20	d

Detailed Explanations

Solution to Question 1:

The given image shows an abnormally short lingual frenulum called Ankyloglossia.

Ankyloglossia is also called “tongue-tie” due to an abnormally short lingual frenulum that can hinder tongue movement. It can interfere with speech or feeding. In such cases, surgical correction is indicated.

The image below shows tongue tie and the classic W-shaped appearance of the tongue on protrusion:



Fissured tongue (Option C) (scrotal tongue or lingua plicata) is a benign malformation in which numerous small furrows or deep grooves are present on the dorsal surface of the tongue. Food particles and debris may become trapped in the fissures, resulting in irritation, inflammation, and halitosis.

Macroglossia (Option D), or an abnormally large tongue, may be seen in disorders like Down's syndrome, Beckwith–Wiedemann syndrome, Hurler's syndrome, and cretinism.

Solution to Question 2:

The recommended first step in the case of alkali ingestion (most commonly seen with drain uncloggers/ house hold cleaning agents) is dilution through the administration of milk or water.

Contraindications in a case of caustic (strong acid or alkali) ingestion:

- Neutralization
- Induced emesis
- Gastric lavage

Solution to Question 3:

Acid ingestion produces coagulative necrosis.

Acid ingestion produces coagulative necrosis and a somewhat protective thick eschar. The amount of acidic agents usually ingested is low as they are generally bitter.

Alkali ingestion: Most cases (70%) of caustic ingestions are accidental. Drain uncloggers are most common as they are tasteless so more is ingested. Alkali ingestion produces liquefactive necrosis.


Solution to Question 4:

The defect demonstrated in the contrast study depicts oesophageal atresia (EA). Isolated oesophageal atresia falls under tracheoesophageal fistula (TEF) type A.

Oesophageal atresia is the most common congenital anomaly of the oesophagus, >90% have an associated TEF.


Plain radiography in the evaluation of respiratory distress might reveal a coiled feeding tube in the oesophageal pouch and/or an air-distended stomach, indicating the presence of a coexisting TEF. Conversely, pure EA can manifest as an airless scaphoid abdomen.

The following classification was given by Gross:



Solution to Question 5:

Newborn presenting with the above features is suggestive of a case of TEF with possible esophageal atresia. This can be confirmed with a chest X-ray image showing a coiled nasogastric tube in the esophageal pouch, as shown in the image below.



Indicators of esophageal atresia with or without a tracheo-esophageal fistula (TEF) are a newborn presenting with:

- Early-onset respiratory distress
- Frothing at the mouth and nose
- History of polyhydramnios

Radiological findings:

- Passing a nasogastric tube with subsequent chest radiography will reveal a coiled nasogastric tube in the esophageal pouch.
- If an air-distended stomach is also present, it indicates the presence of a coexisting TEF.
- Pure EA can manifest as an airless scaphoid abdomen.

Solution to Question 6:

The given case scenario suggests a diagnosis of dysphagia lusoria, which is caused most likely by the aberrant right subclavian artery. The given image suggests the presence of the right aberrant subclavian artery.

Dysphagia lusoria is dysphagia produced due to extrinsic compression of the oesophagus by a developmental vascular anomaly, often an aberrant right subclavian artery or right-sided or

double aortic arch.

Solution to Question 7:

Hypokalemic hypochloremic metabolic alkalosis, an electrolyte and acid-base imbalance characteristically seen in congenital hypertrophic pyloric stenosis (CHPS).

A pH of 7.55 indicates alkalosis and plasma HCO_3^- is 30 mmol/L, which is in the higher range. Both these findings indicate metabolic alkalosis.

Infants with hypertrophic pyloric stenosis develop a characteristic metabolic abnormality. i.e Hypokalemic hypochloremic metabolic alkalosis with paradoxical aciduria.

- Persistent vomiting results in loss of hydrochloric acid leading to the development of metabolic alkalosis (due to loss of H^+) and hypochloremia (due to loss of Cl^-).
- Kidneys try to compensate for alkalosis by secreting Na^+ and HCO_3^- . However, loss of sodium stimulates the secretion of aldosterone through the RAAS pathway.
- The aldosterone causes active reabsorption of sodium in exchange for K^+ and H^+ ions.
- Loss of potassium ions leads to the development of hypokalemia.
- Loss of H^+ ions in urine lowers the urinary pH resulting in aciduria. It is called paradoxical aciduria, since the body initially lost H^+ ions due to loss of HCl , and instead of compensating for that loss, the body is losing more H^+ ions through urine.

CHPS typically presents with the following:

- Non-bilious vomiting, which is progressive, maybe projectile, occurring immediately after a feed, usually presents after the 3rd week of age.
- As the vomiting continues, a progressive loss of fluid, hydrogen ions, chloride ions, and potassium leads to hypokalemic hypochloremic metabolic alkalosis with paradoxical aciduria.

Hyperbilirubinemia is the most common clinical association of pyloric stenosis, which is also known as the icteropyloric syndrome.

Solution to Question 8:

The given case scenario suggests the diagnosis of Congenital hypertrophic pyloric stenosis. Erythromycin, particularly when given in the first few weeks of life, is found to be associated with hypertrophic pyloric stenosis.

Exposure	Drug	Malformation
Post-natal	Erythromycin	Hypertrophic pyloric stenosis
Pre-natal	Thalidomide	PhocomeliaGrossly underdeveloped or absent limbs

Exposure	Drug	Malformation
Pre-natal	Phenytoin	Fetal hydantoin syndrome Intrauterine growth restriction, microcephaly, cleft palate, hypoplastic phalanges, etc.
Pre-natal	Misoprostol	Moebius syndrome Underdeveloped VI and VII CNs resulting in facial paralysis and lateral rectus palsy

Solution to Question 9:

The presence of non-bilious vomiting and visible gastric peristalsis is highly suggestive of congenital hypertrophic pyloric stenosis. The investigation of choice to confirm the diagnosis is ultrasound.

CHPS is associated with hypertrophy of the circular muscle of the pylorus resulting in constriction and obstruction of the gastric outlet. It is common between 3-6 weeks of age. First-born male infants are at the highest risk.

Infants usually present with projectile, non-bilious vomiting. Clinically, an olive-shaped mass is palpable in the right upper quadrant or epigastrium. When the infant is given a test feed, visible peristalsis is usually observed moving from left to right, followed by projectile vomiting. The characteristic metabolic abnormality is hypochloremic metabolic alkalosis due to recurrent vomiting.


Ultrasound features:

- Double-layered hypertrophic mucosa protrudes into the stomach, creating the so-called 'Nipple sign'
- The sonographic appearance of the hypertrophied pylorus resembles that of the uterine cervix and is sometimes referred to as the 'cervix sign'
- The diagnostic criteria for pyloric stenosis are a muscle thickness of ≥ 4 mm and a pyloric length of ≥ 16 mm

If ultrasound is inconclusive, contrast studies (barium) are performed, which demonstrate

- An elongated pyloric channel (string sign),
- A bulge of the pyloric muscle into the antrum (shoulder sign), and
- Parallel streaks of barium seen in the narrowed channel, producing a "double tract sign."


The image attached below shows the USG findings in CHPS.



Signs in Congenital Pyloric stenosis:

Sign	
String sign	Passing of small streak of barium through pyloric canal
Double/triple track sign	Crowding of muscle folds in pyloric canal
Diamond sign/twining recess	Transient triangular tent like cleft in mid portion of pyloric canal with apex pointing inferiorly.
Beak sign	Mass impression upon antrum with streak of barium pointing towards pyloric channel
Mushroom sign/kirkin sign/umbrella sign	Indentation of base of duodenal bulb by pylorus
Caterpillar sign	Gastric hyperperistaltic waves
Shoulder sign	Round indentation caused by hypertrophied pyloric muscle

Congenital Hypertrophic Pyloric Stenosis



The treatment of choice is Ramstedt - Frankelstedt pyloromyotomy.

Solution to Question 10:

The diagnosis in the above case is a urachal fistula. The clinical vignette suggesting straw-yellow fluid (suggestive of urine) along with secondary infection (mildly elevated temperature) and presence of fluid only on straining are all suggestive of urachal fistula.

The urachus is an embryologic canal connecting the dome of the fetal bladder with the allantois, a structure that contributes to the formation of the umbilical cord. In this condition, there is continuous urinary drainage from the umbilicus on straining. It usually presents as a tender, swollen mass secondary to infection.

Diagnosis is made by ultrasonography or computed tomography (CT).

Definitive management is surgical resection of the tract.

A patent vitellointestinal duct, patent omphalomesenteric duct and bladder diverticulum will cause umbilical discharge even in the absence of straining. The discharge in such a case is composed of the contents of the intestine (curdy white in colour mostly).

Solution to Question 11:

The presence of a "double-bubble" sign on a plain abdominal radiograph is suggestive of a diagnosis of duodenal atresia. The double bubble is caused by the distension of stomach and proximal duodenum.

Duodenal atresia results from failed recanalization of the intestinal lumen during gestation. A history of polyhydramnios is present; half of the infants seen with duodenal atresia are premature. The hallmark of duodenal obstruction is bilious vomiting without abdominal distention, which is usually noted on the 1st day of life.

The initial treatment of infants with duodenal atresia includes nasogastric or orogastric decompression and intravenous fluid replacement. The typical surgical repair for duodenal atresia is duodenoduodenostomy.

Solution to Question 12:

The clinical vignette is suggestive of intussusception and the rotavirus vaccine is associated with an increased risk, albeit only a slightly increased risk

Intussusception occurs when a portion of the alimentary tract is telescoped into an adjacent segment. The most common location of intussusception is ileocolic. Ileoileal location is seen postoperatively and in Henoch–Schönlein Purpura.

90% of cases of intussusception in children are idiopathic in nature. It can be seen in infants during the weaning period following the introduction of new food, vaccination, or upper respiratory tract infection. The currently approved rotavirus vaccines are associated with a slightly increased risk of intussusception.

Lead points are more common in children older than 2 years of age and can cause recurrent intussusception. Some recognizable lead points are as follows: enlarged submucosal Peyer's patch, Meckel's diverticulum, intestinal polyp, neurofibroma, intestinal duplication cysts, leiomyomas, hamartomas, anastomotic suture line, enterostomy tube, hemangioma, or malignant conditions such as lymphoma or Kaposi sarcoma.

Solution to Question 13:

Pancreas divisum is the most common congenital anomaly associated with recurrent pancreatitis. It is also the most common pancreatic developmental anomaly.

It results from the failure of fusion of the dorsal and ventral pancreatic buds, due to which, the tail, body, and part of the head of the pancreas drain through the small accessory duct of Santorini rather than the main duct of Wirsung. It may be associated with recurrent pancreatitis.

Solution to Question 14:

Sigmoid volvulus is the most likely diagnosis. It occurs when the sigmoid colon twists on the sigmoid mesocolon. It is a cause of large bowel obstruction.

Solution to Question 15:


The clinical scenario is suggestive of acute appendicitis. Ultrasound is the best diagnostic investigation in the evaluation of acute appendicitis and has demonstrated >90% sensitivity and specificity.

The advantages of ultrasound include its easy availability, rapid method, low cost, and freedom from ionizing radiation.

The ultrasound criteria for appendicitis include the following:

- Wall thickness ≥ 6 mm
- Luminal distention
- Lack of compressibility
- A complex mass in the right lower quadrant
- Or an appendicolith

Image: Ultrasound showing evidence of appendicitis




Solution to Question 16:

The clinical vignette is suggestive of a carcinoid tumour. Elevated levels of urinary 5-hydroxy indole acetic acid (5-HIAA) can be seen in this condition.

The carcinoid tumours are neuroendocrine tumours arising from enterochromaffin cells that can occur anywhere in the GIT. In children, they are most commonly seen in the appendix. A carcinoid tumour with metastasis to the liver is likely to produce carcinoid syndrome. The tumour elaborates serotonin, 5-HT, and histamine, which produce flushing (frequent blushing), watery diarrhoea, bronchoconstriction, and right heart failure.

Carcinoid syndrome can cause secondary niacin deficiency as they consume and divert the tryptophan NAD⁺ pathway to cause the overproduction of serotonin. Thus, these patients can also have pellagra like features.

The diagnosis can be confirmed by elevated levels of urinary 5-HIAA.



The image attached above demonstrates islands and nests of uniform cells with scant, pink granular cytoplasm and salt and pepper chromatin.

Synaptophysin is typically positive in carcinoid tumors.

Treatment: Essentially surgical. Medical management includes the administration of somatostatin analogues (octreotide).

- Small tumours (<1 cm in size) are treated by appendectomy.
- Larger tumours (>2 cm in size) require further bowel resection.

Solution to Question 17:

The hamartomatous polyps in Peutz-Jeghers syndrome are seen in order of prevalence: Jejunum > ileum > duodenum > colonic or gastric.


Solution to Question 18:

Hirschsprung disease is not associated with Edward syndrome (trisomy 18).

It is associated with:

- Trisomy 21/ Downs syndrome
- Joubert syndrome
- Smith–Lemli–Opitz syndrome
- Shah–Waardenburg syndrome
- Multiple endocrine neoplasia 2 syndrome
- Neurofibromatosis (Von recklinghausen disease)

- Neuroblastoma
- Urogenital or cardiovascular abnormalities.



Hirschsprung disease or congenital aganglionic colon is characterized by the absence of ganglion cells in the submucosal and myenteric plexus. It is more common in males. Risk factors include Down's syndrome and RET oncogene.

The imaging modality of choice in Hirschsprung's disease is a contrast radiograph with barium or gastrograffin enema (image shown above).

The most important film is a lateral or oblique view of the rectum during slow filling. It typically shows a low bowel obstruction. The rectum is seen as a narrow tube and a cone-shaped transition zone, seen proximal to the dilated ganglionated bowel.

A rectal suction biopsy is a gold standard for the diagnosis and operative intervention is the definitive treatment.

Definitive surgeries for Hirschsprung disease are as follows:

- Swenson procedure, where aganglionic bowel is removed and a coloanal anastomosis is performed.
- Duhamel procedure, where the aganglionic rectal stump is left in place and the ganglionated normal colon is pulled posterior.
- Soave procedure: In this procedure, endorectal mucosal dissection is done within the aganglionic distal rectum and then the ganglionated normal colon is pulled through the remnant muscular cuff, and a coloanal anastomosis is performed.

NOTE:

In the fetus, migration of the neuronal crest tissues to form myenteric plexus and ganglion cells is completed by the 24th week of life. If interrupted in migration, it results in Hirschsprung disease.

Solution to Question 19:

The given clinical vignette is suggestive of imperforate anus in a neonate with VACTERL association (presence of vertebral and renal anomalies in addition to the anal atresia). Lymphedema is not a component of VACTERL syndrome.

VACTERL association is a disorder that affects many-body systems. People diagnosed with VACTERL association typically have at least three of the following characteristic features.

Components of VACTERL association:

- Vertebral
- Anorectal
- Cardiac
- Tracheal
- Esophageal
- Renal
- Radial
- Limb anomalies

Anorectal malformations:

- It occurs due to the post-allantoic gut with the proctodaeum.
- Newborns classically present with inability to pass meconium within 48 hours, abdominal distention, intestinal obstruction, improper anal dimple, and sometimes with complaints of passing meconium per urethra.

Wingspread Classification:

- Low - Below the level of the pelvic floor (puborectalis).
- Intermediate - Occurs at the level of the puborectalis with or without fistula.
- High - Anorectal agenesis, rectal atresia.

Investigations:

- Wangenstein's invertogram is done 6-12 hours after birth with a metal coin strapped to the presumed site of the anus. The X-ray taken is used to measure and classify the deformity.
- USG abdomen
- Imaging to evacuate associated abnormalities

Treatment is surgical.

Solution to Question 20:

In this case, the most likely cause of rectal prolapse would be Trichuriasis. It is common in poor rural communities with inadequate sanitary facilities and soil contaminated with human or animal feces. The highest infection rate occurs among children between 5-15 years.

Clinical manifestations include chronic dysentery, anemia, poor growth, and developmental deficits. In heavy infections, adult worms can extend throughout the colon down to the rectum. Over time, inflammation of the rectum causes tenesmus leading to prolapse.

Other options:

Pertussis would present with cough, low-grade fever, and sneezing. Increased abdominal pressure due to the persistent cough could cause a prolapse.

Chronic constipation is unlikely as the child has a history of frequently soiling his pants, suggestive of dysentery.

Cystic fibrosis would manifest much earlier in life with respiratory symptoms like recurrent pneumonia. These children have exocrine pancreatic insufficiency, causing protein and fat malabsorption. This leads to bulky, greasy stools and eventually a prolapse if untreated.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Medical GI disorders

Question 1:

If you were to conduct a death audit on all the cases of under-five paediatric deaths secondary to infective diarrhoea, which of the following pathogens would be responsible for the largest fraction of deaths?

- a) Adeno virus
- b) Rotavirus
- c) Salmonella typhimurium
- d) Shigella dysenteriae

Question 2:

A child from an urban slum area is admitted to your unit with diarrhea. Given below are five important micronutrients. The deficiency of which of the following factors increases the risk of death due to diarrhea in this child?

- a) 2, 4, 5
- b) 1 and 3 only
- c) 2 and 5 only
- d) 2 and 4 only

Question 3:

The Nutritionist in your hospital prepares the diet charts for all the children admitted with diarrhoeal disease. One of the advice papers has fallen off from the case file. The advice sheet mentions that the child has to be put on fasting some times of the day. You would attach the sheet back to the case file with the diagnosis of _____

- a) Motility diarrhea
- b) Infective diarrhea
- c) Osmotic diarrhea
- d) Secretory diarrhea

Question 4:

A 5-year-old child with chronic malnutrition presents with diarrheal episodes lasting for more than 2 weeks. The stool sample is positive for *Aeromonas* and entero-adherent *Escherichia coli*. What is the likely diagnosis?

- a) Acute diarrhea
- b) Persistent diarrhea
- c) Subacute diarrhea
- d) Secondary diarrhea

Question 5:

A young child had milk-based dessert at his friend's birthday party. Later, the same evening he developed complaints of bloating, abdominal discomfort, and flatulence. His medical history is significant for viral gastroenteritis a couple of weeks back. Stool examination reveals an acidic pH and is positive for reducing substances. Which type of diarrhea is most likely to be seen in this child?

- a) Osmotic diarrhoea
- b) Protracted diarrhoea
- c) Secretory diarrhoea
- d) Dysentery

Question 6:

A 4-year-old child was brought to the hospital with a history of loose stools. No history of fever or blood in stools. The mother says he is irritable and eagerly drinks water if given. On examination, eyes are sunken and in the skin pinch test, the skin retracted within two seconds but not immediately. What treatment is advised for the child?

- a) Give oral fluids and zinc supplementation only and ask mother to come back if danger sign develops.
- b) Give oral fluids and ask mother to continue the same and visit again the next day.
- c) Consider severe dehydration start IV fluids, IV antibiotics, and refer to higher centre.
- d) Wait and observe.

Question 7:

Ramya, a 6-year-old child with acute diarrhoea is admitted and started on fluid therapy with reduced osmolarity Oral Rehydration Solution . Which of the following statement is false about reduced osmolarity ORS?

- a) Glucose concentration 75 mmol/L
- b) Sodium concentration 75 mmol/L
- c) Total Osmolarity is 311 mosm/L
- d) Potassium concentration is 20 mmol/L

Question 8:

Sathish, your gym partner and father of a 12-month-old child, calls you for a teleconsult. His child has been passing loose watery stools without fever. He wants to know the medications to be given to his child. What dose of oral zinc would you prescribe to this child?

- a) 40 mg/day for 10-14 days
- b) 10 mg/day for 10-14 days
- c) 30 mg/day for 10-14 days.
- d) 20 mg/day for 10-14 days

Question 9:

In a child admitted with viral enteritis along with signs of dehydration, which type of acid-base imbalance is likely to be seen?

- a) Isotonic dehydration with acidosis
- b) Isotonic dehydration with alkalosis
- c) Hypotonic dehydration with acidosis
- d) Hypotonic dehydration with alkalosis

Question 10:

The intestinal biopsy report of a 7-year-old boy evaluated for chronic watery diarrhoea revealed a flat intestinal mucosa. Which of the following conditions would you suspect in this child?

- a) 1, 2, 3, 4, 5
- b) 3, 4 5

c) 1, 2, 3

d) 1, 2, 4

Question 11:

Suyash, a 10-year-old boy presented to the clinic with a history of malabsorption, diarrhoea, and poor appetite after having his favourite white sauce pasta. His mother adds that he develops such symptoms only when he takes pasta or roti. The duodenal biopsy revealed crypt hyperplasia and CD8+ cells in lamina propria. Which of the following tests is least likely to be used for the diagnosis of the condition?

a) Anti-gliadin antibodies

b) Anti-nuclear antibodies

c) Anti-endomysial antibodies

d) Anti-tissue transglutaminase antibodies

Question 12:

A 9-year-old boy presented with vomiting for 5 days. His sister, who has coeliac disease had similar symptoms. IgA-tissue transglutaminase titre is 5 times the upper limit of normal. Which of the following is not an extraintestinal manifestation of this condition?

a) Polycythemia

b) Osteoporosis

c) Short stature

d) Alopecia

Question 13:

Which of the following is the most common cause of pancreatic insufficiency in childhood?

a) Chronic pancreatitis

b) Cystic fibrosis

c) Shwachman-Diamond syndrome

d) Congenital pancreatic lipase deficiency

Question 14:

A 12-year-old boy presented to the emergency department with fever, severe epigastric pain, nausea and vomiting for 6 hours. He feels comfortable when he bends forward and assumes a knee-chest position. Which of the following tests is not useful in the diagnosis of acute pancreatitis?

- a) Serum lipase
- b) Serum amylase
- c) Serum elastase
- d) CT abdomen

Question 15:

Anwasha, an orator preparing for her international debate finale presents to your clinic with lesions on her lower lip as shown below. She is concerned as these lesions are painful, recurrent and hamper the flow of her speech. Which of the following points about this condition is incorrect?



- a) These are due to stress
- b) These are contagious
- c) They heal without scarring
- d) They are seen in Gluten-sensitive enteropathy

Question 16:

The nanny of a 3-year-old boy brings the child to the ER after he had accidentally ingested button batteries from the mobile phone toy that he was playing with. What would be the next best step in the management of this child?

- a) Urgent endoscopy and removal
- b) Observation and follow up in 24 hours
- c) Observation until the foreign body passes out in faeces
- d) Repeat Xray after 4 hours

Answer Key

Question No.	Correct Option
1	b
2	d
3	c
4	b
5	a
6	a
7	c
8	d
9	a
10	d
11	b
12	a
13	b
14	c
15	b
16	a

Detailed Explanations

Solution to Question 1:

Rotavirus is the most common identifiable viral cause of gastroenteritis in children. It accounts for 29% of deaths caused by diarrhoea in children aged < 5 years.

Solution to Question 2:

The risk of death from diarrhoea is increased in the presence of deficiencies of micronutrients such as vitamin A, and zinc (2 and 4 only).

Both these micronutrients are essential for maintaining mucosal immunity. Vitamin A deficiency increases mortality from diarrhoea, measles, and malaria while Zinc deficiency increases mortality from diarrhoea, pneumonia, and malaria.

Recent studies also indicate the usefulness of folic acid in the prevention of diarrhoea.

Solution to Question 3:

Osmotic diarrhoea results from the presence of osmotically active, poorly absorbed solutes in the bowel lumen that inhibit normal water and electrolyte absorption. This type of diarrhoea ceases with fasting.

Secretory diarrhoea, infective diarrhoea, dysentery and motility diarrhoea occurs independent of dietary intake and do not subside with fasting.

Solution to Question 4:

Persistent diarrhoea(Chronic diarrhoea) is defined as a diarrhoeal episode that lasts for more than 14 days.

Diarrhoea is defined as

- stool volume of more than 10 g/kg/day in toddlers/infants (or)
- stool volume greater than 200 g/day in older children

Persistent diarrhoea is increasingly observed in children with malnutrition. These are due to infections commonly caused due to enteroadherent *Escherichia coli* and *Aeromonas*.

Acute diarrhoeas generally last less than 7 days while chronic diarrhoea lasts more than 14 days and is usually a non-infectious pathology.

Solution to Question 5:

The clinical vignette is suggestive of acquired lactose intolerance which is an example of osmotic diarrhoea.

Lactose is present in milk and dairy products. If it is not absorbed in the small intestine, it reaches the colon, where it is fermented to short-chain organic acids. This releases hydrogen that can be detected in the lactose breath test and also generates an osmotic overload.

The child with lactose intolerance generally has perianal excoriations. The condition is diagnosed by using stool examination which shows acidic stool positive for reducing substances. Treatment is by diet control using a diet devoid of lactose. It can be congenital due to lactase deficiency or acquired after post-viral gastroenteritis. In secretory diarrhoea, the epithelial cells' ion transport processes are turned into a state of active secretion. The most common cause of acute-onset secretory diarrhoea is a bacterial infection of the gut.

Chronic diarrhoea is, in many cases, the result of an intestinal infection that persists longer than expected.


Dysentery/Shigellosis is a type of food poisoning caused by infection with the Shigella species. It is spread by means of faeco-oral transmission.

Solution to Question 6:

In a child with diarrhea, thirst (eagerly drinks water), sunken eyes, restless and irritable behavior, and skin pinch showing slow retraction (<2 seconds) point towards some dehydration. Treatment includes oral rehydration therapy, zinc supplementation, and continued breastfeeding according to WHO integrated management of neonatal and childhood diseases (IMNCI) plan B.

The degree of dehydration is assessed and managed using the WHO IMNCI PLAN given below:

SIGNS	CLASSIFY AS	IDENTIFY TREATMENT
Two of the following signs: - Lethargic or unconscious - Sunken eyes - Not able to drink or drinking poorly - Skin pinch goes back very slowly	SEVERE DEHYDRATION	(Urgent pre-referral treatments are in bold print) ► If child has no other severe classification - Give fluid for severe dehydration (Plan C). OR - If child also has another severe classification: Refer URGENTLY to hospital with mother giving frequent sips of ORS on the way. Advise the mother to continue breastfeeding. ► If child is 2 years or older, and there is cholera in your area, give antibiotic for cholera.
Two of the following signs: - Restless, irritable - Sunken eyes - Drinks eagerly, thirsty - Skin pinch goes back slowly.	SOME DEHYDRATION	► Give fluid, Zinc supplements and food for some dehydration (Plan B) ► If Child also has a severe classification: Refer URGENTLY to hospital with mother giving frequent sips of ORS on the way. Advise the mother to continue breastfeeding. ► Advise mother when to return immediately. ► Follow-up in 5 days if not improving. ► If confirmed/symptomatic HIV, follow-up in 2 days if not improving.
Not enough signs to classify as some or severe dehydration	NO DEHYDRATION	► Give fluid, Zinc supplements and food to treat diarrhoea at home (Plan A) ► Advise mother when to return immediately. ► Follow-up in 5 days if not improving. ► If confirmed/symptomatic HIV, follow-up in 2 days if not improving.



Solution to Question 7:

The total osmolarity of reduced osmolarity ORS is 245 mOsm/L (not 311 mOsm/L)

Reduced osmolarity ORS is now the global standard of care in the management of acute gastroenteritis.

Solution to Question 8:

All children older than 6 months of age with acute diarrhoea should receive oral zinc (20 mg/day) for 10-14 days during and after diarrhoea.

Most of the cases of under-five acute diarrhoea are infectious in aetiology with more than 50% due to the Rotavirus, which is self-limiting. Treatment will include fluid and electrolyte correction and zinc supplementation.

Solution to Question 9:

Isotonic dehydration with acidosis is the most common finding in children with severe viral enteritis.

Viruses that cause human diarrhoea selectively infect and destroy the villus tip cells in the small intestine.

In the small intestine, the upper villus enterocytes are differentiated cells, which have absorptive functions, such as the transport of water and electrolytes. On the other hand, the crypt enterocytes are undifferentiated cells and are net secretors of water and electrolytes.

Selective viral infection of intestinal villus tip cells thus leads to decreased absorption of salt and water (proportionate loss of water and salt—isotonic dehydration) and an imbalance in the ratio of intestinal fluid absorption to secretion.

Metabolic acidosis may be a result of stool bicarbonate losses in children with diarrhoea. These can also be seen due to secondary renal insufficiency or lactic acidosis as a sequelae to diarrhoea.

Solution to Question 10:

Chronic diarrhoea with a flat intestinal mucosa can be seen in conditions Tropical sprue, giardiasis, and HIV enteropathy.

Causes of the flat intestinal mucosa (subtotal or total villous atrophy):

- Celiac disease
- Tropical sprue
- Giardiasis
- Bacterial overgrowth
- HIV enteropathy
- Eosinophilic gastroenteritis
- Cow's milk or soy protein enteropathy
- Graft versus host disease
- Chemo or radiotherapy
- Protein-energy malnutrition
- Tuberculosis or lymphoma

Amoebiasis and neutrophilic gastroenteritis may also rarely have villous atrophy but the presentation is generally acute. Amoebiasis is characteristically associated with dysentery and biopsy would reveal a flask-shaped ulcer.

Solution to Question 11:

The clinical vignette is suggestive of celiac disease and an anti-nuclear antibody test is not used in the diagnosis of celiac disease.

Symptoms are pre-dominantly seen after consuming a diet rich in gluten (Wheat pasta or roti). Diagnosis of celiac disease is made based on lab and histopathological findings.

Celiac disease-specific antibodies are as follows:

- Anti-tissue transglutaminase (Anti-TG2) IgA antibodies
- Anti deaminated - IgG forms of gliadin peptides antibodies
- Endomysial antibodies (EMA)

The duodenal biopsy will show crypt hyperplasia, villous atrophy and CD8+ cells in lamina propria

Solution to Question 12:

The boy is suffering from coeliac disease. Polycythemia is not an extraintestinal manifestation of coeliac disease.

On the contrary, the most common extraintestinal manifestation of celiac disease in the paediatric age group is iron-deficiency anaemia that is refractory to iron therapy, and not polycythemia.

Celiac disease is an immune-mediated systemic disorder triggered by the ingestion of wheat gluten and related prolamines from rye and barley.

The intestinal symptoms are caused by the atrophy of the small bowel mucosa and malabsorption. The symptoms include diarrhoea, vomiting, anorexia, weight loss, failure to thrive, rectal prolapse, and intussusception.

The extraintestinal manifestations are caused due to malnutrition and autoimmunity and they include the following:

System	Manifestation	Possible Cause
Hematologic	Anaemia (Most common extraintestinal manifestation)	Iron malabsorption
Skeletal	RicketsOsteoporosisEnamel hypoplasia of the teeth	Calcium/vitamin D malabsorption
Muscular	Atrophy	Malnutrition
Neurologic	Peripheral neuropathyEpilepsyIrritabilityCerebral calcificationsCerebellar ataxia	Thiamine/vitamin B12 deficiency
Endocrinologic	Short statureSecondary hyperparathyroidism	Malnutrition calcium/vitamin Dmalabsorption
Dermatologic	Dermatitis herpetiformisAlopecia areataErythema nodosum	Autoimmunity
Respiratory	Idiopathic pulmonary hemosiderosis	

Solution to Question 13:

Cystic fibrosis is the most common cause of pancreatic insufficiency in children, which, if untreated, will lead to malnutrition.

Other rare causes include:

- Shwachman–Diamond syndrome

- Isolated enzyme deficiencies
- Chronic pancreatitis
- Protein-energy malnutrition

Solution to Question 14:

Serum elastase is not used for the diagnosis of acute pancreatitis. The measurement of faecal elastase is used as a screening test for pancreatic insufficiency.

The criteria for the diagnosis of pancreatitis in children is defined as 2 of 3 of the following:

- Epigastric abdominal pain;
- Serum amylase and/or lipase activity at least 3 times greater than the upper limit of normal; and
- Imaging findings characteristic of, or compatible with, acute pancreatitis.

Serum lipase is now considered the test of choice for acute pancreatitis as it is more specific than amylase.

CT scan has a major role in the diagnosis and follow-up of children with pancreatitis. Other useful investigations are USG, ERCP, and MRCP.

Solution to Question 15:

The lesions given in the above image are aphthous ulcers, and they are not contagious. All the other points are true.

Aphthous ulcers are painful, recurrent, and non-contagious oral lesions. They are well-circumscribed, ulcerative lesions with a white necrotic base surrounded by a red halo. They heal in 10–14 days without scarring. Their aetiology is unclear but the following factors have been implicated:

- Allergic or immunologic reactions
- Emotional stress
- Genetics
- Injury to soft tissues in the mouth
- Inflammatory bowel disease
- Gluten-sensitive enteropathy
- Behcet disease
- Sweet syndrome
- Human immunodeficiency virus (HIV) infection
- Cyclical neutropenia

Treatment generally includes local anaesthetic to relieve pain and B-complex supplementation to enhance healing.

Solution to Question 16:

Button batteries and sharp objects in the oesophagus and foreign bodies associated with respiratory symptoms all mandate urgent endoscopic removal.

Button batteries, in particular, are very dangerous foreign bodies, which must be expediently removed because they can cause:

- Mucosal injury in 1-hour contact time
- Involve all layers of the oesophagus in 4 hours

Injury is thought to be secondary to electrochemical burns from electrical discharge. Chemical burns from electrolyte alkali leakage and pressure necrosis also may contribute to mucosal injury.

Asymptomatic blunt objects and coins lodged in the oesophagus can be observed for up to 24 hr in anticipation of passage into the stomach.

Disorders of the Liver

Question 1:

A mother observed unusual redness on the palms of her young child while putting him to sleep. She brings him to your OPD and on examination, you find mild abdominal distension, splenomegaly, and pedal edema. You admit the child for further evaluation, but on the night of admission, he suddenly starts vomiting fresh blood. Which is the most likely diagnosis?

- a) Mallory Weiss tear
- b) Esophageal varices
- c) Kala azar
- d) AML

Question 2:

A 10-year-old boy was brought by his mother to the ER after having a sudden episode of vomiting blood. On examination, splenomegaly is detected with engorged superficial veins around his umbilicus. Which is the investigation of choice in this patient?

- a) Abdominal USG
- b) Capsule endoscopy
- c) Triphasic CT
- d) Upper GI endoscopy

Question 3:

Unconjugated hyperbilirubinemia, which did not subside even after 3 weeks of birth, was observed in a neonate. On investigating, liver enzymes, PT/INR and albumin levels were normal. No hemolysis was seen on a peripheral blood smear. A drop in bilirubin level was observed within a week after treating with phenobarbital. What is the most likely diagnosis?

- a) Rotor syndrome
- b) Crigler Najjar type 2
- c) Dubin Johnson syndrome
- d) Crigler Najjar type 1

Question 4:

A 14-year-old boy was brought to the OPD by his father who observed a yellowish tinge to his skin and eyes. He gives a history of recent onset fatigue after playing football for a few minutes and cannot stay active on the playground for a long time like before. Investigations revealed isolated conjugated hyperbilirubinemia but are normal otherwise. Liver histology revealed a black pigment. What is the most likely diagnosis?

- a) Gilbert syndrome
- b) Crigler-Najjar syndrome
- c) Dubin-Johnson syndrome
- d) Rotor syndrome

Question 5:

You are an intern posted in the paediatrics department and a child with Dubin-Johnson syndrome is admitted to your unit. During rounds, the attending paediatrician asks you which molecular defect causes liver disease in this patient. Your answer would be

- a) ATP7A
- b) ATP7B
- c) ABCC2
- d) SERPINA 1

Question 6:

All are true about transient familial neonatal hyperbilirubinemia, except:

- a) Increased unconjugated bilirubin
- b) Occurs due to maternal antibody against fetal UGT1A1
- c) It is also known as Lucey Driscoll syndrome
- d) Stop breast feeding when jaundice sets in

Question 7:

Lately, a teacher observed that an anxious young girl had started drooling saliva from her mouth while listening in the classroom. She had also developed a deterioration in her handwriting while taking down notes. Her mother was informed and she brought her child to

your clinic. On examination, she had tremors and mild hepatomegaly. A slit-lamp examination was done and showed the following picture. The right statement about this disease is -



- a) It is autosomal dominant.
- b) Serum ceruloplasmin levels are high.
- c) Coombs-positive hemolytic anaemia may be present.
- d) Patient can be treated with triethylenetetramine

Question 8:

A mother observed deterioration in her young child's academic performance and loss of interest in everyday activities for the last few weeks. She has also lost the ability to play catch with her brother and complains of muscle weakness. There is mild hepatomegaly noted on examination. Her serum ceruloplasmin levels were reduced. All of the following manifestations can occur in a patient with this disease except?

- a) Sparse depigmented hair
- b) Shrinkage of liver
- c) Ascites and portal hypertension
- d) Renal Fanconi syndrome

Question 9:

In Wilson Disease, hepatic copper content usually exceeds ____ $\mu\text{g/g}$ dry weight.

- a) 150
- b) 250
- c) 350
- d) 450

Question 10:

A male infant was born to a mother with chronic hepatitis B infection. The infant is suspected to have been exposed at the time of delivery. At 4 months, which of the following viral markers signifies ongoing viral replication in the infant?

- a) Anti-HBs
- b) Anti-HBc
- c) HBe Ag
- d) HBs Ag

Question 11:

A 9-year-old boy was brought to the OPD by his mother with a history of fatigue and myalgia for a few days, associated with rashes that started a week before it. On examination, coppery-red coloured skin lesions were seen on the face and limbs including palms and soles. His serology was positive for HBsAg. What is the diagnosis?

- a) Erythema infectiosum
- b) Gianotti-Crosti syndrome
- c) Henoch schonlein purpura
- d) Erythema multiformae

Question 12:

Which of the following drugs are not used in the treatment of hepatitis B?

- a) Interferon $\alpha 2b$
- b) Lamivudine
- c) Adefovir
- d) Ribavirin

Question 13:

A young child developed sudden bouts of vomiting and was brought to the ER. Her condition rapidly worsened as she developed seizures. She had a history of influenza infection one week ago. The patient's liver biopsy will show _____.

- a) Non-alcoholic steatohepatitis (NASH)
- b) Centri zonal hemorrhagic necrosis
- c) Marked microvesicular steatosis
- d) Ring granuloma

Question 14:

A 12-year old girl was rushed to the ER by her parents due to hematemesis. They gave a history of weakness and fatigue for the last few weeks. On examination, the patient had spider telangiectasias and mild palmar erythema. Liver function tests showed the following results- AST-1100IU/L, serum bilirubin- 8mg/dl, albumin- decreased, globulin- elevated, PT- prolonged along with Anti-Nuclear Antibody positive. What is the diagnosis?

- a) Type 1 autoimmune hepatitis
- b) SLE
- c) Type 2 autoimmune hepatitis
- d) Hepatitis C

Question 15:

A 2-week-old low-birth-weight neonate presents with jaundice. Stools are pale/clay coloured. Liver biopsy shows loss of architecture. The most likely diagnosis is _____.

- a) Neonatal hepatitis
- b) Extrahepatic biliary atresia
- c) Wilson disease
- d) Hereditary hemochromatosis

Question 16:

A mother is concerned about her 3-week-old baby having yellowish discolouration of skin and eyes. On examination, the baby is chubby and playful. Stools are pale in color. Further

evaluation with USG shows triangular cord sign and liver biopsy shows bile duct proliferation. The most likely diagnosis is _____.

- a) Neonatal hepatitis
- b) Hallervorden Spatz disease
- c) Wilson disease
- d) Extrahepatic biliary atresia

Question 17:

A mother leaves her 3-year-old son unattended for 20 mins and later finds him holding a blue coloured empty strip of tablets with a few tablets scattered on the floor. A few hours later he is rushed to the ER with nausea and vomiting. As an intern taking the history, you suspect accidental ingestion of paracetamol tablets. The pattern of liver injury you would expect in this case is

- a) Cholestasis
- b) Centrilobular necrosis
- c) Microvesicular steatosis
- d) Fibrosis

Question 18:

You are examining a young child who presented to the ER with a high fever and abdominal pain. The patient winces on abdominal palpation of the right upper quadrant. USG showed hydrops of the gallbladder. All of the following conditions are associated with hydrops of the gallbladder except _____.

- a) Group A streptococcal pharyngitis
- b) Kawasaki disease
- c) Total parenteral nutrition
- d) Cystic fibrosis

Question 19:


The commonest indication for liver transplantation in children is _____

- a) Alagille syndrome
- b) Biliary atresia

- c) Caroli disease
- d) Hepatocellular carcinoma

Question 20:

A young boy was admitted with complaints of fever and abdominal pain for 2 days. ESR and CRP were elevated and CT Abdomen obtained was as shown in the picture. Which of the following is not a common feature in the given condition?



- a) Hepatomegaly
- b) Hypoalbuminemia
- c) Jaundice
- d) Leukocytosis

Answer Key

Question No.	Correct Option
1	b
2	d
3	b
4	c
5	c

6	d
7	d
8	a
9	b
10	c
11	b
12	d
13	c
14	a
15	a
16	d
17	b
18	d
19	b
20	c

Detailed Explanations

Solution to Question 1:

A child presenting with hematemesis and splenomegaly is presumed to have variceal bleeding unless proved otherwise. Bleeding from oesophageal varices is the most common presentation of portal hypertension.

Palmar erythema (redness of the palms) and ascites (swelling of the abdomen) are features of liver failure seen in this patient.

Upper GI endoscopy is the preferred diagnostic test for oesophageal varices.

The management of acute variceal bleeding-

- Attain hemodynamic stability through blood transfusion/fluids
- Vasoactive drugs (e.g., octreotide)
- Endoscopy to perform ligation or sclerotherapy
- Transjugular intrahepatic portosystemic shunt should be considered for variceal bleeding refractory to medical and endoscopic therapy.

Secondary prophylaxis to reduce the recurrence of bleeding includes-

- Nonselective β blockade (propranolol, nadolol)
- Obliteration of varices through serial treatment via ligation or sclerotherapy

Solution to Question 2:

A child presenting with hematemesis and splenomegaly is presumed to have variceal bleeding and hence, upper GI endoscopy is the investigation of choice.

The upper GI endoscopy provides a definitive diagnosis and an option to treat via band ligation or sclerotherapy.

Capsule endoscopy, ultrasound, CT, MRI, and elastography are not recommended because of their suboptimal accuracy compared to endoscopy. Capsule endoscopy is used to investigate an occult/obscure gastrointestinal bleed mostly suspected to be in the small intestine.

Solution to Question 3:

Jaundice persisting 3 weeks after birth is suggestive of pathological jaundice. Among the given options, Crigler Najjar type 2 presents with unconjugated hyperbilirubinemia which responds to phenobarbital therapy.

Dubin Johnson (option C) and Rotor (option A) syndromes lead to conjugated hyperbilirubinemia.

Crigler Najjar type 2 is an autosomal recessive disease with non-hemolytic, unconjugated hyperbilirubinemia caused by homozygous missense mutations in the enzyme UDP UGT1A1 resulting in reduced (partial) enzymatic activity.

The differences between Crigler-Najjar type 1 and type 2 are described in the table below:

Crigler Najjar Type 1	Crigler Najjar Type 2
Autosomal recessive	Autosomal recessive
UDPGT is absent	UDPGT is reduced
Severe unconjugated hyperbilirubinemia	Less severe
Pale stools	Normal-coloured stools
May proceed to kernicterus and does not respond to phenobarbitone	Does not proceed to kernicterus and is responsive to phenobarbitone

Solution to Question 4:

The presence of fatigue, conjugated hyperbilirubinemia and black pigment on liver biopsy, termed as "black liver" is suggestive of Dubin–Johnson syndrome.

Dubin-Johnson syndrome is an autosomal recessive inherited defect in hepatocyte secretion of bilirubin glucuronide. Disease results from absent function of multiple drug-resistant protein 2, an ATP dependent canalicular transporter.

The most commonly reported symptoms are abdominal pain and fatigue, jaundice, dark urine, and slight enlargement of the liver.

Inherited conjugated hyperbilirubinemia-

Dubin–Johnson syndrome	Rotor syndrome
Autosomal recessive	Autosomal recessive
Absent function of multiple drug-resistant protein (MRP-2)	Defect in anion transport
Black liver on biopsy (due to accumulation of epinephrine metabolite)	Black liver is not seen
Normal levels of coproporphyrin in urine	Increased levels of coproporphyrin in urine
Gall bladder is not visualized on oral cholecystography	Gall bladder is visualized on oral cholecystography

Solution to Question 5:

Molecular defect causing liver disease in Dubin–Johnson syndrome is ABCC2 (codes for canalicular protein with ATP-binding cassette), it leads to the mutation of multiple drug-resistant protein 2 (MRP2) receptor. Hence, there is a defect in the transfer of conjugated bilirubin from the liver to bile in Dubin–Johnson syndrome.

Important diseases and their molecular defects:

ATP7A	Menke's disease
ATP7B	Wilson disease
CFTR	Cystic fibrosis
SERPINA 1	α -1 antitrypsin deficiency
ABCC2 (MRP2 protein defect)	Dubin Johnson syndrome

Solution to Question 6:

Lucey Driscoll syndrome or transient familial neonatal hyperbilirubinemia is a self-limiting condition and there is no need to stop breastfeeding.

Lucey Driscoll syndrome is an autosomal recessive metabolic disorder affecting enzymes involved in bilirubin metabolism. It causes transient familial neonatal unconjugated hyperbilirubinemia and jaundice occurs during the first 4 days of life. It is due to maternal antibodies against fetal UGT1A1 (UDP glucuronosyltransferase family 1 member A1).

Solution to Question 7:

The image shows the Kayser-Fleischer ring, which is characteristic of Wilson's disease. It is an autosomal recessive disorder, characterised by low serum ceruloplasmin levels. Coomb's negative hemolytic anemia may be present. The patient can be treated with triethylenetetramine also known as trientine.

Wilson's disease is caused due to a loss-of-function mutation in the ATP7B gene on Chr.13q. The gene encodes a copper transporting ATPase (ATP7B) which has 2 functions:

- Maintains copper in bound form by binding copper to apoceruloplasmin
- Helps excrete free copper in bile.

Thus, an ATP7B deficiency leads to excess deposition of free copper within tissues.

The drugs used in the management of Wilson's disease include:

- Chelating agents as first-line drugs
- D-penicillamine
- Trientine or triethylenetetramine
- Elemental zinc as maintenance therapy.

Lab markers	Change in Wilson's disease	Reason
Serum ceruloplasmin	↓↓	ATP7B deficiency results in reduced bound copper
24h urinary copper	↑↑	Excess free copper excreted in urine
Total serum copper	↓	Due to marked reduction in ceruloplasmin

Solution to Question 8:

The presence of reduced ceruloplasmin levels, motor incoordination, and hepatomegaly suggest the patient is suffering from Wilson's disease. Sparse depigmented hair is not a feature of Wilson's disease. It is a feature of Menke's disease.

Wilson's disease is caused due to a mutation in the ATP7B gene that leads to excess deposition of free copper within tissues.

It has the following clinical features:

Note: Wilson's disease may also present as cirrhosis and hepatic failure, where shrinkage of liver occurs.

System	Clinical features
Hepatic(most common)	JaundiceAbdominal painHepatomegalyFatigueAscitesVariceal bleedCoagulation defects
Neurological	DysarthriaDystoniaParkinsonismSubcortical dementiaDepressionWing beating tremorsDecreased scholastic performance
Ophthalmic	Kayser-Fleischer (KF) ringsSunflower cataract
Others	Hemolytic anemia (Coombs negative)Acute pancreatitisFanconi's syndromeHyperparathyroidismCardiomyopathySpontaneous abortionsInfertility

Solution to Question 9:

In Wilson disease, hepatic copper content usually exceeds 250 µg/g dry weight.

Hepatic copper accumulation is the hallmark of Wilson disease. It is measured using neutron activation analysis/atomic absorption spectrometry.

Solution to Question 10:

HBeAg signifies the ongoing viral replication in case of hepatitis B infection.

Solution to Question 11:

The coppery-red colored papules along with a positive HBsAg are suggestive of Gianotti-Crosti syndrome, also known as papular acrodermatitis.

It is an immunologic reaction to viral infections. The most common associations are Epstein-Barr virus, hepatitis B virus, Coxsackievirus A16, and parainfluenza virus, as well as with many childhood immunizations.

Skin lesions are monomorphic, firm, coppery red papules sized 1-10 mm. They occur in crops and may become profuse and coalesce into plaques, forming a symmetric eruption on the face, ears, buttocks, and limbs, including the palms and soles. The trunk, scalp, and mucous membrane are relatively spared.

The image given below shows the coppery red papules seen in Gianotti-Crosti syndrome.



Solution to Question 12:

Ribavirin is a drug used in the treatment of hepatitis C, not hepatitis B.

Indications for starting antiviral therapy in chronic hepatitis B are:

- Patients without cirrhosis
- HBeAg + with HBV DNA $\geq 20,000$ IU/mL and ALT level ≥ 2 times the upper limit of normal
- HBeAg -ve with HBV DNA ≥ 2000 IU/mL and ALT level ≥ 2 times the upper limit of normal
- Patients with compensated cirrhosis and an HBV DNA ≥ 2000 IU/mL
- Patients developing acute liver failure or decompensated cirrhosis.

Drugs used in the treatment of hepatitis B are:

- First-line treatment: Entecavir or tenofovir
- Other drugs:
 - Pegylated interferon-alpha
 - Telbivudine
 - Lamivudine
 - Adefovir

Solution to Question 13:

Severe vomiting in a 5-year-old child, rapidly worsening to seizures, with a history of viral infection (influenza) is suggestive of Reye's syndrome. Hence, liver biopsy will show marked microvesicular steatosis.

Reye's syndrome is a secondary mitochondrial hepatopathy. It is precipitated due to

- Viral infection (influenza, adenovirus, herpes, coxsackievirus A, Epstein–Barr virus)
- Salicylate or antiemetic use
- Inborn error of metabolism such as single enzyme defects of β oxidation.
- Toxins

There is an acute onset of vomiting and encephalopathy, rapidly progressing to seizures, coma, and death. There is no jaundice.

Investigations

- $\uparrow$ ALT and AST
- $\uparrow$ Ammonia
- normal bilirubin
- Hypoglycemia
- PT prolonged
- INR increased
- Liver biopsy: marked microvesicular steatosis. Necrosis is not seen.

Solution to Question 14:

The symptoms of liver failure and portal hypertension along with a positive anti-nuclear antibody(ANA) is suggestive of Type 1 autoimmune hepatitis. It is associated with-

- Elevated serum aminotransaminase concentrations.
- Liver-associated serum autoantibodies.
- Hypergammaglobulinemia.

Solution to Question 15:

Neonatal jaundice in a low-birth-weight (LBW) neonate with consistently pale/clay coloured stools with liver biopsy finding of loss of architecture is suggestive of neonatal hepatitis.

Wilson disease and hereditary hemochromatosis will not present with jaundice in the neonatal stage. Hence, they can be excluded. Injury to liver cells can present the clinical and histologic picture of "neonatal hepatitis."

Differentiation between Neonatal hepatitis and EHBA

Neonatal hepatitis		Extrahepatic biliary atresia (EHBA)
Definition	It implies intrahepatic cholestasis which includes Hepatic injury due to metabolic, viral, or idiopathic causes Intrahepatic bile duct injury	Obliteration of the entire extrahepatic biliary tree at or above the porta hepatis (85% cases) OR distal segmental bile duct obliteration with patent extrahepatic ducts up to the porta hepatis (uncommon)
Clinical features	Family history is present Pre-mature or LBW baby Clay-coloured stools	Family history absent Normal appearance, without looking sick Persistent acholic stools
Investigations	Liver biopsy: Loss of liver architecture	HIDA scan: screening test USG: triangular cord sign Liver biopsy: bile duct proliferation

Solution to Question 16:

Neonatal jaundice in a neonate without a sick-looking appearance and with acholic stools, USG showing triangular cord sign, and liver biopsy showing bile duct proliferation is suggestive of biliary obstruction, which may be seen in biliary atresia.

Wilson disease and Hallervorden Spatz disease (neurodegeneration with iron accumulation in the brain) do not present with jaundice in the neonatal stage. Hence, they can be excluded.

Differentiation between neonatal hepatitis and EHBA

Liver biopsy:

- Biliary atresia is characterized by proliferation of the bile ductule, presence of bile plugs, and portal or perilobular edema and fibrosis, with the basic hepatic lobular architecture intact.
- In neonatal hepatitis, there is severe, diffuse hepatocellular disease, with distortion of lobular architecture, marked infiltration of inflammatory cells, and focal hepatocellular necrosis.

Neonatal hepatitis	Extrahepatic biliary atresia (EHBA)
It implies intrahepatic cholestasis which includes: Hepatic injury due to metabolic, viral, or idiopathic cause Intrahepatic bile duct injury	Obliteration of the entire extrahepatic biliary tree at or above the porta hepatis (85% cases) OR distal segmental bile duct obliteration with patent extrahepatic ducts up to the porta hepatis (uncommon)
Clinical features	
Family history is present Pre-mature or LBW baby Pigmented stools	Family history absent Normal appearance, without looking sick Persistent acholic stools
Investigations	

Neonatal hepatitis	Extrahepatic biliary atresia (EHBA)
Liver biopsy: Loss of liver architecture	USG: triangular cord sign HIDA scan: screening test Liver biopsy: bile duct proliferation

Solution to Question 17:

The pattern of liver injury caused by paracetamol (acetaminophen) is centrilobular necrosis.

Acetaminophen toxicity results from the formation of a highly reactive intermediate metabolite, N-acetyl-p-benzoquinone imine (NAPQI). In therapeutic use, only about 5% is metabolized by the hepatic cytochrome P450 enzyme CYP2E1 to NAPQI, which is then immediately joined with glutathione to form a nontoxic mercapturic acid conjugate.

In overdose, glutathione stores are overwhelmed, and free NAPQI is able to combine with hepatic macromolecules to produce hepatocellular necrosis. Single acute toxic dose of APAP is generally considered to be >200 mg/kg in children.

Patterns of hepatic drug injury:

Disease	Drug
Centrilobular necrosis	Acetaminophen, Halothane
Microvesicular steatosis	Valproic acid
Acute hepatitis	Isoniazid
Fibrosis	Methotrexate
Cholestasis	Chlorpromazine, Erythromycin, Estrogen
Veno occlusive disease	Busulphan, Cyclophosphamide
Portal and hepatic vein thrombosis	Estrogen, Androgen
Biliary sludge	Ceftriaxone
Hepatic adenoma/hepatocellular carcinoma	Oral contraceptive pills, Anabolic steroids

Solution to Question 18:

Hydrops of the gallbladder is not associated with cystic fibrosis.

Hydrops of gallbladder is the acute acalculous non-inflammatory distension of the gallbladder, which occurs in infants and children. It is characterized by the absence of calculi, bacterial infections or congenital anomalies of the biliary system.

Hydrops is distinguished from acalculous cholecystitis by the absence of a significant inflammatory process and a generally benign prognosis.

Conditions associated with hydrops of GB:

- Infections: Streptococcal pharyngitis, staphylococcal infection, ascariasis, leptospirosis, typhoid fever, and viral hepatitis
- Kawasaki disease
- Total parenteral nutrition, prolonged fasting
- Sick cell crisis, thalassemia
- Sepsis
- Mesenteric adenitis, necrotizing enterocolitis

Note: In cystic fibrosis cholelithiasis, cholecystitis, micro gallbladder is seen.

Solution to Question 19:

Biliary atresia is the most common indication for liver transplantation in children.

Extrahepatic biliary atresia (EHBA) is defined as a partial or total absence of permeable bile duct between the porta hepatis and the duodenum. It is due to inflammatory fibrosis of the biliary tree. It leads to obstruction to bile flow thereby causing the leaking of conjugated bilirubin into the circulation.

Infants have symptoms of jaundice, pale stools, and dark urine right after birth. In advanced disease, infants present with failure to thrive, hepatomegaly, and ascites due to liver cirrhosis.

HIDA scan is the investigation of choice for extrahepatic biliary atresia. An intraoperative cholangiogram is a gold standard investigation for diagnosis.

Children with biliary atresia, who do not achieve successful drainage after hepatoportoenterostomy (Kasai's procedure), require liver transplantation within the first year of life.

Management of extrahepatic biliary atresia		
Type I	Atresia restricted to the common bile duct	Roux-en-Y hepaticojejunostomy
Type II	Atresia of the common hepatic duct	Kasai procedure(Roux-en-Y hepatoportoenterostomy)
Type III	Atresia of the right and left hepatic duct	

Solution to Question 20:

The symptoms of fever and abdominal pain and a hypodense lesion on the CT are suggestive of a Liver abscess in this patient.


Signs and symptoms of a liver abscess include fever, chills, night sweats, malaise, nausea, hepatomegaly, as well as pain and tenderness in the right upper quadrant. They also show elevated inflammatory markers, hypoalbuminemia and leukocytosis. However, jaundice is uncommon.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Neonatal Respiratory Disorders

Question 1:

The following image shows the chest X-ray of a preterm neonate who was admitted to the NICU in view of severe respiratory distress. What does this suggest?



- a) Meconium aspiration syndrome
- b) Hyaline membrane disease
- c) Transient tachypnoea of newborn
- d) Bronchopulmonary dysplasia

Question 2:

Which of the following is the most important risk factor for neonatal respiratory distress syndrome?

- a) Perinatal Asphyxia
- b) Prematurity
- c) Caesarean section
- d) History of affected sibling

Question 3:

A father runs to a PHC with a baby wrapped in a towel. The neonate was delivered a few hours back, at home, by the local dai. On initial examination, the medical officer notes cyanosis and severe tachypnoea. Which of the following features would favor a diagnosis of neonatal respiratory distress syndrome in this child?

- a) Patchy coarse infiltrates on chest X-ray
- b) Air bronchogram on chest X-ray
- c) Emaciated appearance with palmar and plantar creases
- d) Onset of distress after 6 hours of birth

Question 4:

Which of the following statements regarding the treatment of RDS with surfactant is false?

- a) It is given in neonates with RDS who fail CPAP
- b) It is given prophylactically to all neonates at < 31 weeks gestation.
- c) It can be given as rescue treatment when RDS actually develops.
- d) Repeated dosing is given every 6-12 hr for a total of 2 to 4 doses.

Question 5:

A first-year resident receives a call from the casualty regarding an unbooked primigravida, at 32 weeks of gestation. She has presented with regular painful uterine contractions. On examination, her cervix is 2cm dilated. What dose of betamethasone injection should the resident give her?

- a) 6mg q12h; 2 doses
- b) 12mg q24h; 2 doses
- c) 12mg q6h; 4 doses
- d) 6mg q12h; 4 doses

Question 6:

A 30-year-old woman with a history of recurrent late-trimester abortions is in preterm labor, presumed to be due to chorioamnionitis. She has received one dose of betamethasone injection. This is a precious pregnancy and the relatives are concerned about prematurity-related complications in the baby. Which of the following is useful for assessing lung maturity in this fetus?

- a) L/S ratio
- b) Bilirubin content of amniotic fluid
- c) Ultrasound
- d) Amniocentesis


Question 7:

Which of the following is not an indicator of fetal lung maturity?

- a) Lecithin-sphingomyelin ratio > 2.5
- b) Positive shake test
- c) Increased phosphatidyl glycerol
- d) Nile blue test showing $> 50\%$ of blue cells

Question 8:

A female infant was born in a community hospital by spontaneous vaginal delivery at 34 weeks of gestation. 30 minutes later, the baby's breathing became excessively laboured. Despite providing non-invasive ventilation, the baby died. Post-mortem histopathological examination of the lung showed thick pink membranes lining the small, collapsed alveolar spaces and necrotic debris, mainly within the respiratory bronchioles, alveolar ducts, and unopened alveoli. What are these membranes chiefly composed of?



- a) Phosphatidylcholine
- b) Fibrin

- c) Mucoprotein
- d) Complex polysaccharide

Question 9:

A primigravida was taken up for emergency cesarean section in view of placental abruption, at 32 weeks of gestation. Immediately after delivery, you dry the baby, clear secretions from the mouth and nose, and provide tactile stimulation. But even after 30 seconds, you note the baby is still gasping for breath and oxygen saturation has not increased more than 92%. Which is the next best step?

- a) Mechanical ventilation
- b) CPAP
- c) 100% oxygen
- d) Surfactant and ventilation

Question 10:

A 3-hour-old neonate with apnea has been on bag and mask ventilation for the last 30 seconds. She has now started taking spontaneous breaths. The heart rate is still between 100 and 110/min. What is your next step?

- a) Discontinue ventilation
- b) Continue ventilation
- c) Give chest compression
- d) ET intubation

Question 11:

A preterm infant in the NICU was observed to have marked nasal flare, distinct xiphoid retractions, minimal intercostal retractions, and inspiratory lag with an audible grunt. What would be the Silverman Anderson score?

- a) 10
- b) 7
- c) 8
- d) 9

Question 12:

As part of calculating Downe's score for a baby, you would assess all of the following except:

- a) Cyanosis
- b) Nasal flaring
- c) Air entry
- d) Respiratory rate

Question 13:

Your brother-in-law calls you during one of your night duties and tells you that your sister just delivered a baby girl and that both the mother and the child are doing well following the cesarean delivery. The next morning, however, you get a call that the baby started having trouble breathing and has been shifted to the NICU. He also sends you a photo of her chest X-ray via WhatsApp. On the radiograph, you note streaky, perihilar linear densities and fluid in the horizontal fissure. Which of the following is true regarding the given clinical condition?

- a) Only observation and supportive therapy is needed
- b) It can predispose to bronchopulmonary dysplasia
- c) More common in preterm infants
- d) Management involves surfactant administration


Question 14:

A preterm, extremely low birth weight baby, delivered by emergency caesarian section was admitted to the NICU and treated for respiratory distress syndrome. She responded to treatment and was discharged. At 6 months of age, she is still having recurrent respiratory distress and requires supplemental oxygen at home. The baby was diagnosed with bronchopulmonary dysplasia. Which of the following does not cause this clinical complication?

- a) Oxygen therapy in the NICU
- b) Mechanical trauma to the lungs
- c) Inhalation of nitrous oxide
- d) Patent Ductus Arteriosus

Question 15:

A female newborn was delivered at 40 weeks of gestation by emergency C-section, following signs of severe fetal distress. Greenish amniotic fluid was noted. Immediately following delivery, the baby was shifted to the NICU and had to be intubated. Her chest X-Ray is as below. What does it show?



- a) Mediastinal shift
- b) Ground glass opacities
- c) Fluid in the interlobar fissure
- d) Patchy infiltrates and coarse streaking

Question 16:

A newborn infant, delivered at home, at 42 weeks of gestation, became cyanotic and started breathing rapidly. The child was immediately taken to the nearby primary health centre. As the medical officer prepares to intubate, she notes greenish staining on the vocal cords. Which of the following is this baby likely to develop?

- a) Listeriosis
- b) Pathological jaundice
- c) Obstructive emphysema
- d) Meconium ileus

Question 17:

Which of the following is false about apnea of prematurity (AOP)?

- a) Usually resolves by 37 weeks post conceptional age
- b) Gentle tactile stimulation is often adequate for mild to moderate episodes
- c) Recurrent AOP can be treated with caffeine or theophylline
- d) AOP is a risk factor for Sudden infant death syndrome (SIDS)

Question 18:

While monitoring the respiratory rate of the infants in the NICU, you notice one baby which suddenly stops breathing. Which of the following statements is true in this regard?

- a) Primary apnea is characterized by labile BP and tachycardia
- b) Caffeine therapy should be given in primary apnea
- c) Secondary apnea responds to stimulation by establishing normal breathing.
- d) Central asphyxia can cause secondary apnea

Question 19:

A 15-day-old infant had conjunctivitis, which later developed into respiratory distress and pneumonia. Chest X-ray examination revealed bilateral lung infiltrates. Peripheral smear count showed 14,300/dL (N 43, L 54, E3). Which of the following is the most likely pathogenic organism?

- a) Chlamydia trachomatis
- b) Streptococcus pneumonia
- c) Neisseria gonorrhoeae
- d) Mycoplasma pneumonia

Question 20:

The radiologist calls and confirms a Bochdalek hernia in a baby born with multiple congenital anomalies. The baby had been placed under an oxygen hood in the NICU, in view of postnatal respiratory distress. On repeat evaluation, he continues to have sternal retractions and is not maintaining saturation. Which of the following would be your initial clinical intervention in this case?

- a) Intubate and insert a nasogastric tube
- b) Bag and mask ventilation
- c) Insert a central venous pressure line

- d) Ventilate with a high-frequency ventilator


Question 21:

A newborn delivered vaginally, at term, was diagnosed with congenital diaphragmatic hernia. The most important limiting factor for survival in this condition is _____.

- a) Mediastinal shift
- b) Left-sided hernia
- c) Intestinal obstruction
- d) Pulmonary hypoplasia

Question 22:

You are doing a quick pre-op evaluation of a 2-day-old baby shifted for emergency pediatric surgery. Parents had brought him with a history of sudden, noisy breathing. He appeared cyanotic with decreased breath sounds on the left side of the chest. Chest X-ray taken in the casualty is given below. Which of the following is not true about the underlying condition?



- a) Not associated with dextrocardia
- b) Most commonly seen on the left side
- c) Mortality is usually associated with pulmonary hypoplasia
- d) Pulmonary dilators markedly improve the survival

Answer Key

Question No.	Correct Option
1	b
2	b
3	b
4	b
5	b
6	a
7	d
8	b
9	b
10	a
11	c
12	b
13	a
14	c
15	d
16	c
17	d
18	d
19	a
20	a
21	d
22	a

Detailed Explanations


Solution to Question 1:

From the given clinical picture of respiratory distress in a premature neonate and the X-ray findings of diffuse ground-glass (reticulo-granular) opacities with a characteristic white-out appearance, hyaline membrane disease (HMD) is the most likely diagnosis.

HMD, also known as neonatal Respiratory distress syndrome (RDS) is a relatively common condition resulting from insufficient production of surfactant that occurs in preterm neonates. Maternal diabetes, perinatal asphyxia, and multiple gestations are other known risk factors for the condition. Clinical features include tachypnea, chest retractions, expiratory grunting, and nasal flaring. The infant may or may not be cyanosed.

X-ray features of HMD include:

- Air bronchogram sign
- Diffuse ground-glass opacities
- Obliteration of cardiac and diaphragmatic contours giving rise to 'white-out' lung (indicates severe disease)



Chest radiograph (A) shows reticulogranular opacities throughout both lungs with prominent air bronchograms. The cardiac silhouette is preserved and both lungs are hypo-aerated.

Chest radiograph (B) shows more severe changes. There is obliteration of the cardiac and diaphragmatic contours due to marked opacification, giving a complete 'white out' appearance

Solution to Question 2:

Respiratory distress syndrome occurs primarily in premature infants. Its incidence is inversely related to gestational age and birthweight. It occurs in 60–80% of infants delivered at <28 wk of gestational age, in 15–30% of those delivered between 32 and 36 weeks of gestational age, and rarely in those delivered at >37 wk of gestational age.

Other risk factors include the following:

- Maternal diabetes
- Multiple births
- Cesarean delivery
- Asphyxia
- History of previously affected siblings

Solution to Question 3:

Air bronchogram on chest x-ray is suggestive of neonatal respiratory distress syndrome.

Other chest x-ray findings in respiratory distress syndrome (RDS):

- Fine reticulogranular pattern
- Bilateral diffuse ground-glass opacities
- White-out lung (severe disease)

Option A: Patchy coarse infiltrates are associated with meconium aspiration syndrome.

Option C: Reduced fat under the skin resulting in an emaciated appearance as well as prominent creases is a feature of post-term babies. Respiratory distress syndrome is more common among preterm babies

Option D: Onset of respiratory distress syndrome is within 6 hours of birth.

Solution to Question 4:

Surfactant is given prophylactically to all neonates < 28 weeks, not 31 weeks. It is given irrespective of the presence or absence of RDS. The route of administration is intratracheal.

Surfactant is indicated in all infants with moderate to severe RDS. It is also given as a rescue treatment when CPAP fails.

Solution to Question 5:

The woman is currently in preterm labour and the dose of antenatal betamethasone given for avoiding complications of prematurity is 12mg IM every 24 hours (total 2 doses).

Dexamethasone is used, the dose is 6mg IM every 12 hours (total 4 doses).

Corticosteroid administration is recommended for all women in preterm labour (<34 weeks of gestation).

Antenatal steroid administration reduces the risk of the following: (RINN)

- Respiratory distress syndrome (RDS)
- Intraventricular haemorrhage (IVH)
- Necrotizing enterocolitis (NEC)
- Neurodevelopmental delay

Solution to Question 6:

The lecithin–sphingomyelin ratio (L/S ratio) is an amniotic fluid test used to assess fetal lung maturity. L/S ratio of ≥ 2 in amniotic fluid indicates lung immaturity.

Surfactant starts appearing in the fetal lung by 20 weeks of gestation. Levels indicating lung maturity are detected after 35 weeks.

Surfactant is a complex mixture of lipids, proteins, and carbohydrates produced by type II pneumocytes in the lung epithelium. The absence of surfactant in the liquid film lining the alveoli causes an increase in surface tension and alveolar collapse. If not treated, such atelectasis leads to increased work of breathing, intrapulmonary shunting, ventilation-perfusion mismatch, hypoxia, and eventually respiratory failure.

The relative concentration of the phospholipids lecithin (L) and sphingomyelin (S) is an indicator of lung maturity. The outward flow of pulmonary secretions from the fetal lungs into the amniotic fluid maintains the level of lecithin and sphingomyelin equally until 32–33 weeks gestation when the lecithin concentration begins to increase significantly while sphingomyelin remains nearly the same. An L–S ratio of 2.5 or more indicates fetal lung maturity while an L/S ratio of less than 1.5 is associated with a high risk of infant respiratory distress syndrome.

Other tests done to assess fetal lung maturity are:

- Shake test/bubble test
- Phosphatidyl glycerol measurement
- Surfactant:albumin ratio
- Lamellar body counts

Solution to Question 7:

Nile blue test showing $\geq 50\%$ orange cells is a sign of fetal lung maturity, not blue cells.

Tests for the assessment of fetal lung maturity include:

The Nile blue test involves staining desquamated fetal cells from amniotic fluid with 0.1% Nile blue sulfate. The presence of orange cells $\geq 50\%$ suggests lung maturity. Lecithin/sphingomyelin ratio at 31 to 32 weeks is 1, and at 35 weeks ratio is 2. The ratio ≥ 2 indicates lung maturity (option A).

The shake test or bubble test (Clement's test) (option B) involves mixing amniotic fluid with 95% ethanol, shaking it for 15 seconds, and observing it for 15 minutes. A complete ring of bubbles at the meniscus is a positive test and indicates lung maturity. This is a useful bedside test.

The presence of phosphatidylglycerol, (option C) saturated phosphatidylcholine and lamellar body count ($\geq 50,000/\text{mL}$) indicates lung maturity.

Glucocorticoids accelerate lung maturation and lower the incidence of respiratory distress and neonatal mortality in preterms if the birth was delayed for at least 24 hours after initiation of betamethasone. A single course of corticosteroids is recommended for women between 24 and 36 weeks who are at risk of delivery within 7 days. Intramuscular injections of either 2 doses of 12 mg betamethasone are given 24 hourly or 4 doses of 6 mg dexamethasone are given 12 hourly.

Solution to Question 8:

The history of respiratory distress in a premature baby (34 weeks) and a biopsy showing pink (hyaline) membranes in the alveoli is indicative of hyaline membrane disease (HMD). The hyaline membranes are composed of layers of fibrin-rich edema fluid and necrotic cells of lining alveoli. Diffuse ischemic damage to the alveoli causes the transudation of proteins into the alveoli to form the hyaline membrane.

Hyaline membrane disease (HMD), also known as infantile respiratory distress syndrome (IRDS) or respiratory distress syndrome of newborns (RDS) is a syndrome in premature infants caused by developmental insufficiency of pulmonary surfactant production and structural immaturity of the lungs.

Solution to Question 9:

The most common cause of respiratory distress in a preterm infant is neonatal RDS (hyaline membrane disease). The infant has presented with moderate respiratory distress without any signs of respiratory failure so early continuous positive airway pressure (CPAP) is the appropriate treatment.

CPAP started at birth is the choice for the management of a preterm infant at risk for respiratory distress syndrome (RDS). Early nasal CPAP has shown to lower mortality and risk of bronchopulmonary dysplasia. Room air is better than oxygen for the resuscitation of preterm infants, and 100% oxygen is also harmful, causing increased oxidative stress.

The intubate-surfactant-extubate (INSURE) or CPAP technique is recommended for surfactant administration when babies showed signs of RDS and needed more than 30% inspired oxygen to maintain saturation in the normal range.

Solution to Question 10:

As per the NRP algorithm, at any time during resuscitation, if the HR $\geq$ 100 and spontaneous breathing resumes then discontinue ventilation.

The components of NRP resuscitation include: (TABCD)

- Temperature- maintain between 36.5 and 37.5 C
- Airway- "sniffing" position, clear secretions if needed
- Breathing- free flow O₂ (maintain sats 91–95%), PPV (40–60 breaths /min), ETT
- Circulation- chest compressions:ventilation ratio (3:1 = 90:30; lower 1/3rd sternum; two thumb technique preferred)
- Drugs- Epinephrine, volume expanders

Algorithm states:

- HR $<$ 100/min + no respirations → Positive pressure ventilation (PPV)
- No response with PPV → Endotracheal intubation (ET)

- HR < 60/min → Chest compressions
- No response to chest compressions → epinephrine

Solution to Question 11:

The Silverman-Anderson score of the child would be 8.

Calculation of the Silverman-Anderson score of this child:

Total score: 8

A score of more than 7 indicates respiratory failure.

The Silverman-Anderson score is a commonly used score to assess respiratory distress. It helps to assess the progression of distress over time and initiate treatment. It is more suited for preterms with respiratory distress syndrome. Scoring is to be done every half an hour and charting is carried out to monitor progress.

The image below shows Silverman Anderson retraction score.

Criteria	Finding	Score
Expiratory grunt	Audible with the naked ear	2
Nasal flaring	Marked	2
Xiphoid retraction	Marked	2
Intercostal retraction (lower chest)	Minimal	1
Chest movement (upper chest)	Lag	1

Silverman Anderson Retraction Score

	Grade 0	Grade 1	Grade 2
Upper Chest	 Synchronized	 Lag on inspiration	 See-saw
Lower Chest	 No retractions	 Just visible	 Marked
Xiphoid Retractions	 None	 Just visible	 Marked
Nares Dilatation	 None	 Minimal	 Marked
Expiratory grunt	 None	 Heard with stethoscope	 Audible

©MARROW

Downes' Score is also used for assessing the severity of respiratory distress. It is more comprehensive and can be applied to any gestational age and condition. Downe's score of ≥ 7 means impending respiratory failure.

Score	0	1	2
Respiratory rate	<60/min	60–80/min	>80/min
Air entry	Normal	Mild decrease	Marked decrease
Cyanosis	Nil	In-room air	On >40% O ₂
Grunting	None	Only on auscultation	Audible with the naked ear
Retraction	Nil	Mild	Moderate

Solution to Question 12:

Downe's score does not consider nasal flaring.

The severity of respiratory distress is assessed by Downe's score and Silverman–Anderson score. Downe's score can be applied for any gestational age. Downe's score of ≥ 7 means impending respiratory failure. Silverman–Anderson score is more suited for preterms with hyaline membrane disease (HMD).

Downe's scoring


Score	0	1	2
Respiratory rate	<60/min	60–80/min	>80/min
Air entry	Normal	Mild decrease	Marked decrease
Cyanosis	Nil	In room air	On >40% O ₂
Grunting	None	Only on auscultation	Audible with naked ear
Retraction	Nil	Mild	Moderate

Solution to Question 13:

Benign respiratory distress in a newborn with a chest X-ray showing fluid in the interlobar fissure is indicative of transient tachypnoea of newborn (TTN), also known as retained fetal lung liquid syndrome. The management of TTN is supportive oxygen therapy and observation.

It is more common after a term, cesarean delivery and is due to the delayed clearance of the lung fluid from the interstitium into the pulmonary lymphatics.

Other findings of chest x-ray imaging are prominent (perihilar) pulmonary vascular markings, increased interstitial markings, and a “wet silhouette” around the heart.



Chest radiograph (A) on day 1 at 6 hours shows a bilateral diffuse ground-glass appearance and fine granularity due to interstitial opacities.


Chest radiograph (B) of the same patient at the age of 2 days (B) shows that the pulmonary parenchymal opacities have decreased, though perihilar streaky markings are still present.

Treatment is supportive and minimal oxygen treatment (< 40%) is often adequate; infants usually recover within 3 days.

Solution to Question 14:

Inhalation of nitrous oxide (N₂O) is not associated with bronchopulmonary dysplasia (BPD). Inhaled NO is a pulmonary vasodilator approved for the treatment of hypoxic respiratory failure in term and near-term infants with pulmonary hypertension. It may have a role in preventing BPD.

BPD is a pathologic process leading to clinical features of chronic lung disease that originates in the neonatal period and is defined by an oxygen requirement for 28 days postnatally. Inflammatory injury to the developing lungs due to barotrauma and oxygen toxicity resulting from prolonged mechanical ventilation is the key factor driving the pathogenesis of BPD. Together with other antenatal and postnatal factors, it results in aberrant tissue repair, dysregulated vascular and arrested alveolar development, which is the basis of the pathology seen in BPD.



Solution to Question 15:

The image shows patchy infiltrates and coarse streaking, which are characteristic findings in the chest X-ray image of infants with meconium aspiration syndrome.

Solution to Question 16:


Respiratory distress in a post-term neonate with a history of greenish (meconium-stained) amniotic fluid during delivery is suggestive of meconium aspiration syndrome (MAS). The aspirated meconium can block the large and small airways, causing patchy atelectasis and emphysema. This produces a ventilation-perfusion mismatch. Severe overinflation can lead to air leak syndromes such as pneumothorax.

Meconium consists of water, desquamated GIT cells, skin, lanugo hair, vernix, bile salts, pancreatic enzymes, lipids, and mucopolysaccharides. The bile salts and pancreatic enzymes can cause chemical pneumonitis. Meconium can also cause surfactant inactivation by displacing it from the alveolar surface and inhibiting its surface tension-lowering ability. Severe MAS is associated with remodelling and thickening of the muscular walls, resulting in pulmonary hypertension.

Infants who develop MAS typically present in the first 12 hours after birth with varying degrees of respiratory distress. On inspection, many affected infants have 'barrel chests'; crackles (rales) and rhonchi often can be heard on auscultation. In severe cases, the entire lungs can be involved resulting in a total "whiteout" of the lungs on the radiograph.

Listeriosis is an infection that is acquired from the mother and is reported to be rarely associated with meconium-stained amniotic fluid.

Pathogenesis of Meconium aspiration syndrome



Solution to Question 17:

Apnea of prematurity (AOP) is not a risk factor for sudden infant death syndrome (SIDS).

It is defined as cessation of breathing for longer than 20 seconds or for any duration if accompanied by cyanosis or bradycardia. It is a diagnosis of exclusion. It is seen in preterm infants and usually resolves by the 37th week of gestation. Recurrent apnea of prematurity may be treated with caffeine or theophylline. It does not predict the future occurrence of SIDS.

Sudden Infant Death Syndrome (SIDS) is the sudden death of an infant under one year of age, which remains unexplained after a thorough case investigation, including the performance of a complete autopsy, examination of the death scene, and review of the clinical history.

Solution to Question 18:

Secondary apnea can occur as a central CNS mediated asphyxia or due to neuromuscular disorders.

Apnea is the cessation of breathing for >20 seconds or for any duration if accompanied by bradycardia and cyanosis.

Primary Apnea	Secondary Apnea
Failure to initiate respiratory effort at birth	Apnea developing in an infant who has initiated a respiratory effort
There may be bradycardia and decreased tone. BP remains normal	Can originate in CNS due to asphyxia or peripherally due to neuromuscular disorders

Primary Apnea	Secondary Apnea
Respond to tactile stimulation	Require ventilatory assistance and drugs- caffeine or theophylline

Solution to Question 19:

The organism most likely to cause the mentioned infections is *Chlamydia trachomatis*.

Pneumonia caused by *C. trachomatis* can develop in 10–20% of infants born to women with active, untreated chlamydial infection. Only approximately 25% of infants with nasopharyngeal chlamydial infection develop pneumonia.

C. trachomatis pneumonia of infancy has a very characteristic presentation. It usually occurs between 1-3 months of age and is insidious, with a persistent cough, tachypnea, and absence of fever. Auscultation reveals rales; wheezing is uncommon.

The absence of fever and wheezing helps to distinguish *C. trachomatis* pneumonia from respiratory syncytial virus pneumonia.

A distinctive laboratory finding is the presence of peripheral eosinophilia (>400 cells/ μ L).

The most consistent finding on chest radiograph is hyperinflation accompanied by minimal interstitial or alveolar infiltrates.

Solution to Question 20:

Once the diagnosis of congenital diaphragmatic hernia is made or suspected, all patients are intubated. Also a nasogastric tube, connected to continuous suction, is placed in the stomach to reduce lung compression.

Congenital diaphragmatic hernia (CDH) is a birth defect of the diaphragm. The most common type of CDH is a Bochdalek hernia (postero-lateral diaphragmatic hernia); other types include Morgagni hernia (anterior-parasternal), diaphragm eventration, and central tendon defects of the diaphragm. Malformation of the diaphragm allows the abdominal organs to push into the chest cavity, hindering proper lung formation and expansion.

Immediate steps of management:

Endotracheal intubation → Placement of a nasogastric tube → Administration of fluids and inotropic agents → Administration of surfactant therapy to only preterm infants with findings suggestive of respiratory distress syndrome. Other options:

Option B: Bag and mask ventilation is to be avoided as they lead to gastric/abdominal distension and compression of the lung.

Option C: Both arterial and central venous lines are inserted; however, this is not the first line of intervention.

Option D: To support respiratory failure, the first modality to be used is conventional ventilation, and if there is no improvement, then high-frequency ventilation is used.

Solution to Question 21:

Pulmonary hypoplasia is a very poor prognostic factor in congenital diaphragmatic hernia.

Poor prognostic indicators in CDH:

- The presence of liver in the thoracic cavity
- Pulmonary hypoplasia, pulmonary hypertension
- Early respiratory distress, within 24 h after birth
- Need for extracorporeal membrane oxygenation
- Herniation to the contralateral lung (right-sided hernia)

Solution to Question 22:

The given clinical picture of neonatal respiratory distress, absent breath sounds in the left side of the chest, and an X-ray image showing bowel loops in the thoracic cavity is diagnostic of congenital diaphragmatic hernia (CDH). Though less commonly documented, dextrocardia is seen in association with CDH.

A diaphragmatic hernia is defined as a communication between the abdominal and thoracic cavities with or without abdominal contents in the thorax. It is mostly congenital, with a defective formation of the pleuroperitoneal membrane. Females are affected twice as often as males.

Bochdalek hernia accounts for up to 90% of the hernias seen in the newborn period, with 80-90% occurring on the left side. So, the heart will be pushed to the right side (dextroposition).

Predictors of a poor prognosis include an associated major anomaly, symptoms before completion of 24 hours since birth, severe pulmonary hypoplasia, herniation to the contralateral lung, and the need for extracorporeal membrane oxygenation (ECMO).

Nitric oxide is a selective pulmonary vasodilator. It reduces ductal shunting and pulmonary pressures, resulting in improved oxygenation.

Childhood Respiratory Disorders

Question 1:

A mother brought a healthy-looking, 3-month-old infant for routine immunization. As you were giving the shot, you noticed some noisy breathing in the baby, which became worse once he started crying. However, it decreased, as you held him in a prone position. There were no clear signs of distress. According to his mother, he had been feeding well at home and has not had any episodes of apnea or cyanosis. Which is the most likely diagnosis?

- a) Laryngeal stenosis
- b) Laryngeal web
- c) Laryngomalacia
- d) Laryngeal papilloma

Question 2:

A senior casualty nurse informs you that a 5-year-old girl with severe respiratory distress might require intubation. You just finished stabilising and placing an ET tube in Mohan, a 57-year-old man with head trauma and a GCS of 7. Which of the following is not a feature of child's airway?

- a) The larynx being located higher in the neck at the level of C3-C4
- b) The narrowest part being the subglottic region
- c) The tongue being relatively larger
- d) Having a cylindrical shape

Question 3:

During a night shift, you get a call from the ER to come and see a toddler, who presented with complaints of fever, drooling, and a muffled voice since evening. On examination, the child has a stridor and you pick up a "thumb sign" on the lateral x-ray of his neck. What is the most likely diagnosis?

- a) Croup
- b) Acute epiglottitis
- c) Foreign body aspiration

d) Retropharyngeal abscess


Question 4:

Which of the following is the most common etiological agent associated with epiglottitis in children, overall in the world?

- a) H. Influenza type B
- b) Influenza virus
- c) Moraxella
- d) Streptococcus

Question 5:

A 5 year old boy was suffering from low grade fever, inspiratory stridor and barking cough for past 5 days. His chest X-ray is given below. What is the cause of this condition?



- a) Moraxella
- b) Parainfluenza
- c) Hemophilus influenza
- d) Staphylococcus aureus

Question 6:

2-year-old Neha has been having fever since yesterday and is refusing food. Parents became alarmed when she developed an abnormal barking cough today morning and have brought her to the casualty. On examination, the child looked exhausted and had visible chest retractions and stridor. The temperature was 39.6 degrees and SpO₂ was 93% on room air. Chest X-ray reveals a 'steeple' sign. What is the treatment of choice in this child?

- a) Racemic epinephrine nebulization
- b) Humidified oxygen and salbutamol nebulization
- c) Send a throat swab and start IV antibiotics
- d) High dose dexamethasone injection

Question 7:

A child in your unit is admitted with acute bronchiolitis. Which of the following is the most common etiological agent responsible for this condition during infancy?

- a) Influenza virus
- b) Para Influenza virus
- c) Rhinovirus
- d) Respiratory syncytial virus

Question 8:

10-month-old Ishaan was presented to the OPD with a 3-day history of lower respiratory tract symptoms and an expiratory wheeze. Chest X-ray showed mild peribronchial thickening. Which of the following statements is false regarding the condition described?

- a) The risk is higher for infants with young mothers or mothers who smoked during pregnancy.
- b) It is more common in children who have not been breastfed.
- c) The mainstay of treatment is antibiotics.
- d) There is a higher incidence of wheezing and asthma in children with a history of bronchiolitis.

Question 9:

The pediatric intensivist is contemplating whether to start aerosolized ribavirin in a young boy with a severe respiratory infection. There would be a clear indication to start the drug if _____.

- a) The child is immunocompromised

- b) The child has not received any immunization
- c) There is a history of severe atopy
- d) There is a history of recurrent infection

Question 10:

3-year-old Sravya was referred with a one-week history of being unable to look upwards, constant neck flexion, and deviation to the right side. Parents also complained of feeding difficulty. On examination, the child was noted to have poor oral hygiene and the neck was tender on palpation. The X-ray is given below. Which of the following statements are false about the given condition?



- a) It occurs mostly in children younger than 3-4 years
- b) It is more common in boys
- c) Examination shows unilateral bulging of the posterior pharyngeal wall
- d) Trauma to the posterior pharynx is the most common cause

Question 11:

The most common cause of bacterial pneumonia in children 3 weeks to 4 years of age is _____.

- a) Haemophilus influenzae
- b) Streptococcus pneumoniae
- c) Streptococcus pyogenes

d) Staphylococcus aureus

Question 12:

A 9-year-old boy presented with a 4-day history of high-grade fever and a productive cough with yellow sputum. Chest x-ray showed a consolidation in the right middle zone. According to WHO guidelines, which of the following is not an indication for hospital admission?

- a) Chest indrawing**
- b) Persistent vomiting**
- c) Convulsions**
- d) Severe malnutrition**

Question 13:

A 5th standard girl is brought to the OPD with a high fever and cough with expectoration for 3 days. Her chest X-ray shows multiple pneumatoceles. Which of the following is the most common etiological agent implicated in this presentation?

- a) Mycobacterium tuberculosis**
- b) Staphylococcus aureus**
- c) Mycobacterium avium intracellulare**
- d) Pneumocystis carinii**

Question 14:

A one-and-half-year-old child brought to the hospital with a history of breathlessness, fever, cough, and cold. On examination, respiratory rate was 46/min, and there were no chest retractions. Which is the most appropriate management?

- a) Give IV antibiotics and observe**
- b) Give IV antibiotics and refer to higher centre**
- c) Oral antibiotics, warn of danger signs, and send home**
- d) No need of treatment only home remedies**

Question 15:

An adolescent boy with a family history of asthma presented with intermittent wheeze and cough for 2 months. The onset of wheeze is triggered during exercise and while playing at school. Which of the following is preferred as regular therapy for the prevention of his symptoms?

- a) Inhaled short acting beta 2 agonist (SABA)
- b) Inhaled corticosteroids
- c) Inhaled long acting beta 2 agonist (LABA)
- d) Anti – IgE antibody

Question 16:

A 17-year-old boy presented to the clinic with an acute respiratory tract infection. He has a history of asthma associated with recurrent respiratory tract infections and sinusitis. Genetic testing in childhood had revealed CFTR mutations. Which of the following is likely to be decreased in this patient?

- a) Sweat chloride
- b) Serum bicarbonate level
- c) Sperm count
- d) Stool fat

Question 17:

Which of the following has similar inheritance pattern as cystic fibrosis?

- a) Accumulation of glycogen in the lysosome
- b) Trinucleotide repeat expansion of CAG on chromosome 4
- c) Adult onset polycystic kidney disease
- d) RB gene mutation

Question 18:

Which of the following statements about cystic fibrosis is not true?

- a) It is an autosomal-recessive disorder.
- b) Abnormality in CFTR gene leads to defective calcium transport.
- c) Mutation in cystic fibrosis transport regulator is seen.

- d) Males have infertility in adult life

Question 19:

The most common organism causing pneumonia in cystic fibrosis among the pediatric age group is _____.

- a) *Pseudomonas aeruginosa*, non-mucoid
- b) *Staphylococcus aureus*
- c) *Burkholderia cepacia*
- d) *Streptococcus pneumoniae*

Question 20:

A 5-year-old boy was evaluated for cystic fibrosis. Sweat chloride shows value of 38 and 44 mEq/L (normal < 40). Which is the next best test to confirm the diagnosis of cystic fibrosis?

- a) 72-hour fecal fat
- b) CT chest
- c) DNA analysis
- d) Trans epithelial nasal potential difference

Question 21:

A couple planning for pregnancy visits you for genetic counseling. They are both known to be heterozygous carriers of CFTR mutations. Which of the following is the most likely complication in an affected child at birth?

- a) Hyaline membrane disease
- b) Transient tachypnea of newborn
- c) Meconium ileus
- d) Sigmoid volvulus

Question 22:

A 9-year-old boy, diagnosed with cystic fibrosis shortly after birth has been having around 2-3 exacerbations per year, requiring hospital admissions. Infection by which of the following

organisms has increased mortality in such patients?

- a) Staphylococcus aureus
- b) Pseudomonas aerogenes
- c) Burkholderia cepacia
- d) Aeromonas

Question 23:

A 3-year-old boy presents with a history of recurrent pneumonia and chronic diarrhea. His mother states that he has 6-8 foul-smelling stools per day. Physical examination reveals a low-grade fever and scattered rhonchi over both lung fields. Sweat chloride testing is elevated on 2 separate occasions. Which of the following drugs is used as a treatment option for this condition?

- a) Ivacaftor
- b) Ravulizumab
- c) Tecovirimat
- d) Mepolizumab

Question 24:

A 4-year-old girl presented with cough and difficulty breathing following accidental inhalation of a peanut. On examination, there was localised wheezing and decreased breath sounds on the right side. What is the next best step in management?

- a) Chest X-ray
- b) Rigid bronchoscopy
- c) Flexible endoscopy
- d) Direct Laryngoscopy

Question 25:

All the following are true about Macleod syndrome except that _____.

- a) It is also called Swyer-James syndrome.
- b) It is not a true emphysema.
- c) It is mostly linked to adenovirus.

- d) Affected lung is larger in size compared to unaffected lung.

Question 26:

A 15-year-old child has been recurrently hospitalized for infections, since 2 years of age. After a detailed evaluation, a diagnosis of Kartagener's syndrome was suspected. All of the following are components of the Kartagener triad except _____.

- a) Situs inversus totalis
- b) Chronic sinusitis
- c) Male infertility
- d) Bronchiectasis

Question 27:

All the following are components of Young's syndrome except:

- a) Bronchiectasis
- b) Asthma
- c) Sinusitis
- d) Obstructive azoospermia

Question 28:

A term male neonate develops respiratory distress at birth, unresponsive to surfactant. Echocardiography is normal and chest X-ray shows a ground-glass appearance. Parents say that their first child died at 10 days of birth due to similar complaints. What is the most likely diagnosis?

- a) Totally anomalous pulmonary venous congestion
- b) Meconium aspiration
- c) Neonatal pulmonary alveolar proteinosis
- d) Diffuse herpes simplex infection

Question 29:

A 3-month old previously healthy baby was brought to the casualty when the parents found that the baby was not breathing while asleep. Following resuscitative measures, the baby was still apnoeic, pulseless and asystolic. Which of the following is not a risk factor for sudden infant death syndrome?

- a) Infants of higher birth order
- b) Infants born to women aged ≥ 35 years
- c) Infants of mother who smoked during pregnancy
- d) Infants sleeping in prone position

Question 30:

A 6-month-old infant was admitted because of increasing stridor since birth, cough, pulmonary infections, and episodes of acute respiratory distress. On detailed evaluation using bronchoscopy and MRI imaging, a bronchial cyst was found to be the cause. The following are features of the lesion except that ____.

- a) It is mostly mediastinal.
- b) It is usually confined to 1 lobe.
- c) It is rarely uniloculated.
- d) It is never associated with malignancy.

Question 31:

In a child with yellow nail syndrome, you would expect to find all the following clinical features except _____.

- a) Pleural effusion
- b) Lymphedema
- c) Clubbing
- d) Discolored nails

Question 32:

What is the most likely cause of a mediastinal mass in a 2-year-old child?

- a) Neurogenic tumors
- b) Thymoma

c) Germ cell tumors

d) Lymphoma

Answer Key

Question No.	Correct Option
1	c
2	d
3	b
4	d
5	b
6	a
7	d
8	c
9	a
10	d
11	b
12	a
13	b
14	c
15	b
16	c
17	a
18	b
19	b
20	c
21	c
22	c
23	a
24	b
25	d
26	c
27	b
28	c
29	b


30	d
31	c
32	a

Detailed Explanations

Solution to Question 1:

A clinical picture of inspiratory stridor that worsens on crying and relieves in a prone position in a newborn without respiratory distress is suggestive of laryngomalacia.

Laryngomalacia is the most common congenital laryngeal anomaly and the most common cause of stridor in infants and children. Stridor is exacerbated by any exertion: crying, agitation, or feeding. In the prone position, the stridor disappears.



On laryngoscopy, an omega-shaped epiglottis is visualized.

In most children, only expectant observation is required as the condition resolves secondary to nervous system maturation, within 18 to 24 months of birth. However, neonates with severe laryngomalacia may develop marked upper airway obstruction resulting in cor pulmonale, life-threatening apneic episodes, feeding difficulties, obstructive sleep apnea, and failure to thrive. In such cases, surgical resection of supraglottic tissues and supraglottoplasty has been the favoured approach.

Solution to Question 2:

The pediatric airway is funnel shaped. In comparison, the adult airway is cylindrical.

In newborns, the larynx is situated at the level of C3-C4, where it facilitates simultaneous respiration and swallowing during feeding.

Solution to Question 3:

Thumb sign on lateral x-ray of neck is pathognomonic for acute epiglottitis (AE). It is seen due to epiglottic enlargement.

AE is an inflammation of the epiglottis. The arytenoids, false cords, and posterior tongue may also be involved.

In children, *Haemophilus influenzae* type B (Hib) is the most common cause. This has decreased dramatically since the widespread availability of immunization (pentavalent). *Streptococcus pyogenes*, *S. pneumoniae*, and *S. aureus* have also been implicated. Non-infectious causes can be foreign body ingestion and trauma from thermal or caustic agents.

It has a progressive and potentially fulminating course, characterised by high fever, sore throat, dyspnea, and rapidly progressing respiratory obstruction.

AE is a clinical diagnosis and a true emergency, X-ray soft tissue neck is not necessary to confirm it. It may be used to rule out differentials such as laryngotracheobronchitis, foreign body obstruction, retropharyngeal abscess or congenital anomalies. The time taken for X-rays should be avoided in a critically ill child due to the risk of sudden complete airway obstruction.

The definitive diagnosis of AE is made during direct laryngoscopy. However, since manipulation can induce a laryngospasm, it is generally avoided.

The image below shows thumb sign:



Solution to Question 4:

The most common etiological agent, overall in the world, of acute epiglottitis in the pediatric age group, now is *Streptococcus*.

In the past, *Haemophilus influenzae* type b (Hib) used to be the most commonly identified cause of acute epiglottitis. But, since the widespread use of the *H. influenzae* type b vaccine, invasive disease caused by *H. influenzae* type b in pediatric patients has been reduced by 99%.

Therefore, other organisms, such as *Streptococcus pyogenes* (GAS), *Streptococcus pneumoniae*, non-typeable *H. influenzae*, and *Staphylococcus aureus*, have become the common causes of acute epiglottitis in pediatric age group, overall in the world.

Lateral X-ray of the neck shows the thumb sign in acute epiglottitis.

Solution to Question 5:

Barking cough is a feature of laryngotracheobronchitis, which is caused by the parainfluenza virus. The X-ray finding is the 'Steeple sign' due to subglottic narrowing.

The age of presentation of laryngotracheobronchitis ranges from 6 months-6 years. The clinical features include barking cough, inspiratory stridor, hoarseness, low-grade fever.

Management:

- Racemic epinephrine nebulization leads to the rapid improvement of upper airway obstruction. Epinephrine constricts pre-capillary arterioles in the upper airway mucosa and decreases capillary hydrostatic pressure, leading to fluid resorption and improvement in airway edema.
- Dexamethasone-by the least invasive route (Oral>IV>IM). It is indicated in all the grades of croup.
- Oxygen should be administered to children who are hypoxemic ($SpO_2 < 92\%$)

Antibiotics are not indicated in laryngotracheobronchitis.

Solution to Question 6:

Fever, barking cough, inspiratory stridor and a steeple sign on chest radiograph are suggestive of acute laryngotracheobronchitis (croup). The treatment of choice in this child is nebulization with racemic epinephrine.

Epinephrine acts rapidly and provides immediate relief. It constricts pre-capillary arterioles in the upper airway mucosa and decreases capillary hydrostatic pressure. This increases fluid resorption and reduces airway edema.

Westley Croup Severity Score:

Overall score and severity

- 0 to 2- mild
- 3 to 7- moderate
- 8 to 11- severe

- 12 to 17- impending respiratory failure

Treatment of moderate to severe croup :

- Racemic epinephrine nebulization
- Dexamethasone is indicated in all grades of croup and is given by the least invasive route (Oral>IM>IV)
- Oxygen should be administered to children who are hypoxemic (SpO₂ <92 %)
- Antibiotics are not indicated in croup

Feature	0	1	2	3	4	5
Stridor	None	With agitation	At rest			
Air Entry	Normal	Decreased	Markedly decreased			
Retractions	None	Mild	Moderate	Severe		
Cyanosis	None				With agitation	At Rest
Level of Consciousness	Normal,including sleep					Disoriented

Solution to Question 7:

Acute bronchiolitis is predominantly a viral disease in children younger than 2 years. Respiratory syncytial virus (RSV) is the most likely etiological agent and is responsible for more than 50% of cases.

Acute bronchiolitis refers to airway inflammation and obstruction of the lower respiratory tract.

Clinical features of bronchiolitis include fever, cough, fine inspiratory crackles, expiratory wheezes, and breathing difficulty. Cyanosis may be seen in severe cases.

Radiographs may be normal in some patients. In the majority of them, there will be interstitial infiltrates, hyper-expansion of the chest, and peribronchial thickening.

Management is mainly conservative. Hypoxemic children should receive cool, humidified oxygen. Rarely, a child might develop respiratory failure and require hospitalization and invasive ventilation.

Solution to Question 8:

The given clinical scenario is suggestive of acute bronchiolitis. The treatment is mainly supportive. Hypoxemic children should receive cool, humidified oxygen. Rarely, a child might develop respiratory failure and require hospitalization and invasive ventilation.

It is more common in boys, in those who have not been breastfed, and in those who live in crowded conditions. The risk is higher for infants with young mothers or mothers who smoked during pregnancy.

Acute bronchiolitis is predominantly a viral disease caused by respiratory syncytial virus (RSV) and antibiotics are not indicated.

There is a higher incidence of wheezing and asthma in children with a history of bronchiolitis.

Solution to Question 9:

Ribavirin is the only specific antiviral therapy available for the treatment of severe bronchiolitis caused by RSV infections. It is reserved as a last-line agent for life-threatening bronchiolitis in patients who are immunocompromised or who have hemodynamically significant cardiopulmonary disease. In otherwise healthy children, the management of acute RSV bronchiolitis is typically supportive.

Ribavirin is a prodrug, which when metabolized resembles purine RNA nucleotides. In this form, it interferes with RNA metabolism required for viral replication.

MNEMONIC: (for indications of ribavirin)

- C-RIBAvirin:
- Hepatitis C
- RSV
- Influenza B
- Arenaviruses (Lassa, Bolivian, etc.)

Note - Ribavirin is also used in the treatment of hepatitis C and some viral hemorrhagic fevers.

Solution to Question 10:

The history of progressive torticollis, neck pain and dysphagia in a young child along with a neck X-ray demonstrating a prevertebral soft-tissue swelling are all indicative of a retropharyngeal abscess. In young children, an antecedent upper respiratory tract infection is the most common cause of an RP abscess and not trauma.

It involves the suppuration of lymph nodes present in the RP space and is more common in children younger than 3-4 years since these lymph node chains involute after 5 years of age. Boys are affected more often than girls. In older children and adults, the cause is usually trauma to the posterior pharynx, which leads to inoculation of the retropharyngeal space.

Risk factors include poor oral hygiene, diabetes, an immunocompromised state, and low socioeconomic status.

Clinical manifestations are nonspecific and include fever, decreased oral intake, and drooling. Neck stiffness and torticollis may also be present. As the abscess grows in size, it can also lead to upper airway obstruction and asphyxiation. The physical examination can reveal unilateral

bulging of the posterior pharyngeal wall.

Although lateral radiograph is used for initial imaging, CT of the neck with intravenous contrast is the best imaging study to evaluate patients with a retropharyngeal abscess. Treatment ranges from prolonged courses of intravenous antibiotics to surgical incision and drainage.

Solution to Question 11:

Streptococcus pneumoniae (pneumococcus) is the most common bacterial pathogen in children 3 weeks to 4 years of age.

Aetiology of infectious pneumonia grouped by age of the patient:

Age group	Frequent pathogens (in order of frequency)
< 3 weeks	Group B <i>Streptococcus</i> , <i>E. coli</i> , gram-negative bacilli, <i>S. pneumoniae</i> , <i>Hemophilus influenzae</i> (type b)
3 weeks - 4 years	RSV, other respiratory viruses (rhinovirus, influenza, parainfluenza and adenovirus), <i>S. pneumoniae</i> , <i>Hemophilus influenzae</i> (type b), <i>Mycoplasma pneumoniae</i>
> 4 years	<i>Mycoplasma pneumoniae</i> , <i>Chlamydia pneumoniae</i> , <i>S. pneumoniae</i> , <i>Hemophilus influenzae</i> (type b)

Solution to Question 12:


As per the revised 2014 WHO guidelines, it is safe to treat chest indrawing pneumonia at home.

However, a child who is HIV positive and has chest indrawing will need to be referred for inpatient treatment. Only those children who have general danger signs are classified as having “severe pneumonia” and need to be referred to a higher level (hospital admission).

Danger signs:

- Not able to drink
- Persistent vomiting
- Convulsions
- Lethargic or unconscious
- Stridor in a calm child


- Severe malnutrition



Solution to Question 13:

Staphylococcus aureus pneumonia manifests as confluent bronchopneumonia, which is often unilateral and characterized by the presence of hemorrhagic necrosis and irregular areas of cavitation of the lung parenchyma, resulting in pneumatoceles, empyema, or at times, bronchopulmonary fistulas.

The following CT image shows cavitating pneumonia by S. aureus:



Solution to Question 14:

The given clinical scenario is suggestive of acute respiratory illness. According to the IMNCI guidelines 2020, this child can be classified as having 'Pneumonia'. The treatment is oral amoxicillin, vitamin A and home care advice.

SIGNS	CLASSIFY AS	TREATMENT
Any general danger sign	VERY SEVERE PNEUMONIA	- Give diazepam if convulsing now - Quickly complete the assessment - Give any pre-referral treatment immediately - Treat to prevent low blood sugar - Keep the child warm - Refer URGENTLY - Screen for possible TB and HIV
- Any general danger sign OR - Oxygen saturation < 90% - Stridor in calm child - Central Cyanosis - AVPU* = V, P or U	SEVERE PNEUMONIA	- If saturation < 90%, start oxygen therapy and refer or admit - Give the first dose of Benzyl Penicillin & Gentamicin - Treat for and to prevent low blood sugar - Keep the child warm - Treat wheeze if present, admit or refer urgently to hospital - Screen for possible TB and HIV
- Chest indrawing in calm child OR - Fast breathing AND - No signs of severe pneumonia	PNEUMONIA	- Give Amoxicillin Dispersible Tablet and Vitamin A - If wheezing, treat and follow-up in 2 days - Soothe the throat and relieve the cough with a safe remedy - Screen for possible TB and HIV - Review in 2 days. If not possible, admit OR refer children with chest indrawing - Advise mother when to return immediately
No signs of pneumonia or very severe disease.	NO PNEUMONIA: COUGH OR COLD	- If wheezing, treat and follow-up in 2 days - Soothe the throat and relieve the cough with a safe remedy - Follow-up in 5 days if not improving - Screen for possible TB and HIV - Advise mother when to return immediately

*AVPU - Alert, responsive to Voice, responsive to Pain, Unresponsive

Fast breathing is defined as given in the table below:

Age of child	Fast breathing RR
<2 months	60 breaths/min
2-12 months	50 breaths/min
1-5 years	40 breaths/min

Solution to Question 15:

Inhaled corticosteroids are recommended as regular therapy for the prevention of his symptoms of exercise-induced asthma.

According to GINA 2021 guidelines, exercise-induced bronchoconstriction can be prevented with daily treatment (regular therapy) with inhaled corticosteroids. SABA, LABA, or chromones may be used prior to exercise, along with warm-up before exercise.

Those who have mild exercise-induced asthma can use their regular inhaled corticosteroid before exercise and need not take SABA pre-exercise.

Solution to Question 16:

The given clinical scenario is suggestive of cystic fibrosis. In these patients, sperm count is most likely to be decreased.

Male patients with cystic fibrosis usually have infertility due to failure of development of Wolffian duct structures leading to congenital absence of bilateral vas deferens. There is defective sperm transport resulting in obstructive azoospermia.

The sexual function is generally unimpaired.

Option A: Sweat chloride levels are elevated (≥ 60 mEq/L) due to abnormalities in the CFTR protein leading to defective chloride transport.

Option B: Serum bicarbonate levels are increased due to excessive loss of salt in the sweat resulting in hypochloremic alkalosis.

Option D: Patients with cystic fibrosis have fat malabsorption secondary to exocrine pancreatic insufficiency, hence the increased stool fat.

Solution to Question 17:

Cystic fibrosis (CF), is an autosomal-recessive disorder. Accumulation of glycogen in the lysosome is seen in Pompe's disease which is another autosomal-recessive disorder.

Option B: Trinucleotide repeat expansion of CAG on chromosome 4 is seen in Huntington's disease, which is autosomal dominant.

Options C and D: Adult-onset polycystic kidney disease and retinoblastoma (RB gene mutation) are autosomal-dominant diseases.

Solution to Question 18:

In cystic fibrosis, abnormalities in the cystic fibrosis transmembrane conductance regulator (CFTR) gene lead to defective chloride channels and not calcium channels.

Cystic fibrosis (CF) is an autosomal-recessive disorder of children and adults. Dysfunction of the CFTR, the primary defect, leads to a wide and variable array of presenting manifestations and complications.

The most common CFTR mutation is F508 deletion.

More than 95% of males are azoospermic because of failure of development of Wolffian duct structures, but the sexual function is generally unimpaired.

Solution to Question 19:

Staphylococcus aureus is the most common organism causing pneumonia in cystic fibrosis among the given options.

The most common organisms causing pneumonia in CF patients:

- The pediatric age group- *S. aureus*
- The adult population- *P. aeruginosa* (mucoid)

Pseudomonas aeruginosa-mucoid may also cause pneumonia in pediatric CF patients.

In cystic fibrosis, *P. aeruginosa* shows a special feature of developing a mucoid capsule. This complex polysaccharide generates a biofilm that provides a hypoxic environment and thereby protects *Pseudomonas* against antimicrobial agents.

Burkholderia cepacia pneumonia is associated with higher mortality, but it is not the most common. These patients are found to have severe pulmonary hypertension, possibly due to pulmonary inflammatory vascular response.

Solution to Question 20:

DNA analysis is used to confirm the diagnosis of cystic fibrosis in patients with equivocal (one normal and one abnormal value) or frankly normal sweat chloride values.

The diagnosis is confirmed if both of the following criteria are met:

- Presence of
- Clinical symptoms affecting at least one organ system (respiratory, gastrointestinal, or genitourinary) or
- Positive newborn screening or
- A sibling with cystic fibrosis
- Evidence of dysfunction of the CFTR gene by
- Two positive sweat chloride test results obtained on separate days or
- Identification of two CFTR mutations by genetic testing or
- An abnormal measurement of nasal potential difference

Options A and B: 72-hour fecal fat and chest CT are not used in the diagnosis of CF.

Option D: This testing is primarily used in research applications and has never undergone extensive validation as a clinical tool.

Solution to Question 21:

CFTR mutations are associated with cystic fibrosis (CF). Meconium ileus is a common complication associated with CF patients at birth.


Meconium found in the intestine of a neonate consists of succus entericus (bile salts, bile acids, and debris from the intestinal mucosa) and is normally evacuated within 6 hours of birth (or earlier).

Meconium ileus occurs in the first 24-48 hours of life when meconium becomes inspissated and obstructs the distal ileum causing abdominal distension and emesis. Abdominal radiographs show

dilated loops of bowel with air-fluid levels and, frequently, a collection of granular, “ground-glass” material in the lower central abdomen.

Meconium ileus may also be seen with pancreatic atresia or pancreatic duct stenosis.

Radiological image shows contrast enema in a newborn infant with abdominal distention and failure to pass meconium. Several air–fluid levels in the small bowel are visible on the upright lateral view.



Solution to Question 22:

Cystic fibrosis patients with *Burkholderia cepacia* pneumonia are found to have severe pulmonary hypertension, possibly due to pulmonary inflammatory vascular response, and have higher mortality.

Staphylococcus aureus and *Pseudomonas aeruginosa* (mucoid) are the most common pathogens causing pneumonia in CF patients. However, they have lower mortality compared to *B. cepacia*.

Solution to Question 23:

The given clinical scenario is suggestive of cystic fibrosis (CF). Ivacaftor is used in the treatment of cystic fibrosis.

Ivacaftor is a potentiator of the chloride channel and overcomes its defect. It activates the CFTR-G551D mutant protein.

Remember: IvaCaFToR (CFTR mutation in CF)

Solution to Question 24:

In a child with a suspected foreign body in the lung, the next best step would be to do a rigid bronchoscopy as it is both diagnostic and therapeutic.

The rigid bronchoscope has a ventilation mechanism and a suction port making it easy for visualization and handling of the foreign body. However, it is pertinent to note that the initial investigation is usually a chest X-ray.

Solution to Question 25:

Macleod syndrome refers to the inability of the lungs to grow after a severe obstructive bronchiolitis. Children with this condition often have chronic cough, recurrent pneumonia, and wheezing. On CT scan, the affected lung shows small size and decreased vascularity, hyperexpansion, and hyperlucency (false emphysema).

It is also called Swyer–James syndrome. It is mostly linked to adenovirus. Patients respond well to bronchodilators.

Solution to Question 26:

Male infertility is typical in Kartagener syndrome, but not a component of the triad.

Kartagener's syndrome is primary ciliary dyskinesia characterized by the classical triad of:

- Situs inversus
- Chronic sinusitis
- Bronchiectasis

Primary ciliary dyskinesia (also called immotile cilia syndrome) is an autosomal recessive disorder affecting ciliary motility. It occurs due to ultrastructural defects in the dynein arm of cilia. This causes the retention of secretions due to a failed mucociliary clearance mechanism. Clinically, this manifests as chronic sinusitis and recurrent infections that ultimately lead to bronchiectasis.

As ciliary function is necessary to ensure proper rotation of the developing organs in the chest and abdomen. Its absence, will lead to situs inversus or a partial lateralizing abnormality.

Male patients with this condition tend to be infertile, due to asthenospermia (immotile spermatozoa due to impaired sperm flagella function).

Solution to Question 27:

Asthma is not usually seen in Young's syndrome.

Young's syndrome is characterized by:

- Bronchiectasis
- Sinusitis

- Obstructive azoospermia

In contrast, Kartagener's syndrome is characterized by

- Bronchiectasis
- Sinusitis
- Situs inversus


Solution to Question 28:

The history and chest X-ray findings are suggestive of pulmonary alveolar proteinosis (PAP).

PAP is a rare, diffuse lung disease that is characterized by alveolar and interstitial accumulation of a periodic acid-Schiff (PAS) stain-positive phospholipoprotein (derived from surfactant).

Infants and children with PAP present with dyspnea, fatigue, cough, weight loss, chest pain, or hemoptysis. Boys are affected 3 times as often as girls.

Chest X-ray shows ground-glass opacity and HRCT shows crazy-paving pattern (as shown below).



Treatment options include total lung lavage, inhaled or subcutaneous granulocyte-macrophage colony-stimulating factor (GM-CSF).

Solution to Question 29:

Higher maternal age is not a risk factor for sudden infant death syndrome (SIDS).

Risk factors for SIDS include

- Low socioeconomic status
- 3-4 months of age have the highest incidence

- Higher birth order irrespective of maternal age and low birth weight
- Cigarette smoking, alcohol, and drug abuse during pregnancy
- Sleeping prone, even side sleeping increases the risk

Solution to Question 30:

Bronchial cysts are associated with malignancy (pleuropulmonary blastoma).

Bronchial cysts are congenital malformations of the bronchial tree. It mainly presents as a mediastinal mass that may enlarge and cause local compression.

It is usually confined to a single lobe and is mostly multiloculated.

Solution to Question 31:

Yellow nail syndrome is not associated with clubbing.

Yellow nail syndrome is characterized by respiratory manifestation (pleural effusion, bronchiectasis), lymphedema, and discoloured nails.

Yellow nail syndrome almost always occurs after 50 years of age, but a juvenile or familial form has also been observed.

Solution to Question 32:

The most common mediastinal masses in young children are neurogenic tumours (50%-60%). In most cases (< 4 years of age), they are invariably malignant (neuroblastomas).

Anterior mediastinal masses:

- Thymoma - most common
- Malignant lymphoma (Hodgkin's and non-Hodgkin's)
- Teratoma
- Thyroid (retro- or sub-sternal extension)
- Thymic cyst
- Angioma

Middle mediastinal masses:

- Foregut duplication cyst (bronchogenic, enteric, and neurenteric) - most common
- Lymphadenopathy
- Vascular malformations

Posterior mediastinal masses:

- Neurogenic tumour - most common overall
- Foregut duplication cyst
- Vascular malformations
- Extramedullary hematopoiesis

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Fetal Circulation

Question 1:

A junior radiologist performing a fetal doppler follows the direction of the flow of oxygenated blood from the placenta to the fetus to examine the fetal circulatory vessels. The sequence that he would follow is:

- a) Umbilical artery → inferior vena cava → ductus venosus → right atrium → left atrium
- b) Umbilical artery → superior vena cava → right atrium → ductus arteriosus → left atrium
- c) Umbilical vein → ductus venosus → inferior vena cava → right atrium → left atrium
- d) Umbilical vein → ductus arteriosus → superior vena cava → right atrium → left atrium

Question 2:

Oxygenated blood is carried to the fetus by _____.

- a) The umbilical arteries
- b) The inferior vena cava
- c) Both a and b
- d) The umbilical vein

Question 3:

A final year student is asked to give a seminar on 'fetal circulation.' Which among the following statement told by him in the seminar is likely to be false?

- a) Heart receives blood with high oxygen saturation
- b) Pressure in right and left ventricles are equal
- c) Liver receives blood with low oxygen saturation compared to the brain
- d) Blood in inferior venacava has lower oxygen concentration compared to superior venacava

Question 4:

The total fetal cardiac output at term is estimated to be _____ per minute.

- a) 150 mL/kg body weight

- b) 250 mL/kg body weight
- c) 350 mL/kg body weight
- d) 450 mL/kg body weight

Question 5:

During the evaluation of a newborn, you hear a continuous machinery murmur. Where would the structure responsible for the murmur ideally be located?

- a) Just proximal to the left common carotid
- b) Just distal to the left subclavian
- c) Just distal to the left common carotid
- d) Just proximal to the left subclavian

Question 6:

You assist the delivery of a 3.1 kg healthy term neonate by spontaneous vaginal delivery. As the baby transitions from the fetal circulation, what is the sequence of functional closure of the following structures?

- a) Ductus venosus --> Foramen ovale --> Ductus arteriosus
- b) Ductus arteriosus --> Foramen ovale --> Ductus venosus
- c) Ductus venosus --> Ductus arteriosus --> Foramen ovale
- d) Ductus arteriosus --> Ductus venosus --> Foramen ovale

Question 7:

During the examination of a neonate with documented hearing defects and bilateral zonular cataracts, you hear a continuous machinery murmur. The mother gives a history of viral exanthem during the first trimester of the pregnancy. By when should the underlying defective channel that is responsible for the murmur anatomically close?

- a) Birth
- b) 3-4 days
- c) 10-21 days
- d) 10-15 hours

Question 8:

Which of the following is the main factor for the ductal closure postnatally?

- a) Increase in partial pressure of oxygen (paO_2)
- b) Increase in systemic vascular resistance
- c) Increase in circulating prostaglandin levels
- d) Decrease in pulmonary venous resistance

Question 9:

A newborn infant develops respiratory distress 2 days after birth. On examination, she is mottled and has weak upper extremity pulses and absent femoral pulses. Investigations reveal the coarctation of the aorta. Which of the following drugs can you use to keep the ductus open medically before surgical repair?

- a) Trometamol
- b) Indomethacin
- c) Alprostadil
- d) Morphine

Question 10:

A 23-year-old woman with G2P1 and an Rh-negative pregnancy gave birth to a baby with an Rh-positive blood group. The direct Coomb's test is positive. Cord blood bilirubin level is more than 4 mg/dL. The team plans for an isovolumetric and simultaneous double set up exchange transfusion. But the preferred vessel seems obliterated. This obliterated fibrotic vessel would now be called as?

- a) Medial umbilical ligaments
- b) Ligamentum teres hepatis
- c) Ductus venosus
- d) Ductus arteriosus

Answer Key

Question No.	Correct Option
1	c
2	d
3	d
4	d
5	b
6	a
7	c
8	a
9	c
10	b

Detailed Explanations

Solution to Question 1:

The sequence of oxygenated blood flow from the placenta to the fetal heart is umbilical vein → ductus venosus → inferior venacava → right atrium → left atrium.

Solution to Question 2:

Oxygenated blood is carried to the fetus by the umbilical vein.

The umbilical venous blood shunts via the ductus venosus into the inferior vena cava. Thus, the inferior vena cava, which also carries blood from hepatic veins, kidneys, and from lower extremities carries a mixture of deoxygenated and oxygenated blood. So, the highest percentage of oxygen in the fetal circulation is in the umbilical vein, followed by the inferior vena cava.

The umbilical arteries carry deoxygenated blood from the fetus to the placenta for oxygenation.

Note:

In the umbilical cord, there are a pair of umbilical arteries and a single umbilical vein.

Mnemonic: AVA- artery –vein- artery(2 arteries and a single vein).

Solution to Question 3:

Blood in the inferior vena cava has a higher oxygen concentration compared to the superior vena cava.

The superior vena cava carries the deoxygenated blood from the upper part of the body to the right atrium. The highly oxygenated blood is carried to the fetus from the placenta via the umbilical vein with 50% of the umbilical venous blood entering the hepatic circulation and the rest bypassing the liver and joining the inferior vena cava (IVC) via the ductus venosus. Hence, the blood in the inferior vena cava has a higher oxygen concentration compared to the superior vena cava. Also, relatively a larger percentage of oxygen-rich blood is perfused to the brain.

Pressures in both the right and left ventricles are equal in a developing fetus. The upper part of the body is perfused exclusively by the left ventricle while the lower extremities are perfused by the right ventricle.

Solution to Question 4:

The total fetal cardiac output at term is approximately 450 mL/kg/min.


Approximately 65% of descending aortic blood flow returns to the placenta, and hence, only the remaining 35% perfuses the fetal organs and tissues.

The clinical relevance of this 65% blood flow being directed to the placenta comes in the context of delayed cord clamping vs early cord clamping after delivery. Delayed cord clamping entails the umbilical cord is clamped more than one minute after the birth or when cord pulsation has ceased. This delayed clamping allows the blood flow between the placenta and neonate to continue and helps improve the iron status in the infant.

Solution to Question 5:

Continuous machinery murmur in a newborn is characteristic of Patent ductus arteriosus. Most commonly, the ductus arteriosus is just distal to the left subclavian artery and the ductus enters the pulmonary artery at its bifurcation.

The image attached below shows the relations of ductus arteriosus.



Solution to Question 6:

Functional closure of the fetal circulatory adjustments occurs in the following sequence: Ductus venosus --> Foramen ovale --> Ductus arteriosus

Fetal circulatory adjustment	Functional closure	Anatomical closure
Ductus venosus	After removal of the placenta from fetal circulatory unit	7 days
Foramen ovale	At birth - after breathing is initiated	Months to years
Ductus arteriosus	10–15 hours	10–21 days

Solution to Question 7:

The given vignette is suggestive of Patent ductus arteriosus in a child with congenital rubella syndrome. The anatomical closure of the ductus arteriosus occurs at 10-21 days.

Patent ductus arteriosus (PDA) is the most common congenital heart disease associated with rubella, followed by pulmonary artery stenosis (PS). PDA presents with a continuous machinery murmur.


Triad of congenital rubella syndrome

- Sensorineural deafness
- Congenital heart defects – MC is PDA
- Cataracts

Classical rubella syndrome consists of:

- Ocular defects: Congenital cataract, salt and pepper chorioretinopathy, microphthalmos, Glaucoma, and anterior uveitis
- Ear defects: Sensorineural hearing loss due to destruction of organ of Corti
- Heart defects: Patent ductus arteriosus, pulmonary stenosis, and ventricular septal defects

Changes in fetal circulation after birth




©MARROW

Fetal circulatory adjustment	Functional closure	Anatomical closure
Ductus venosus	After removal of the placenta from fetal circulatory unit	7 days
Foramen ovale	At birth - after breathing is initiated	Months to years
Ductus arteriosus	10–15 hours	10–21 days

Solution to Question 8:

Oxygen is the most important factor controlling ductal closure.

Changes in fetal circulation after birth



©MARROW

Gestational age also appears to play an important role; the ductus of a premature infant is less responsive to oxygen, even though its musculature is developed. Hence, patent ductus arteriosus (PDA) is more common in premature infants.

The fetal circulatory adjustments after birth are given below:

The patency of ductus arteriosus may either provide an alternative route for blood to bypass congenital defects such as pulmonary atresia or coarctation of the aorta.

Therapeutic medications can be used to either maintain the patency (prostaglandin E1) or promote closure (indomethacin). Drugs like indomethacin are not recommended during the third trimester of pregnancy.

Functional closure		Anatomical closure
Ductus venosus	After removal of the placenta	7 days
Foramen ovale	At birth - after breathing is initiated	Months to years
Ductus arteriosus	10–15 hours	10–21 days

Solution to Question 9:

Alprostadil, a prostaglandin E1 (PGE1) analogue, can be used to keep the ductus open medically in the coarctation of the aorta.

The ductus arteriosus (the fetal connection between the pulmonary artery and the descending aorta) is functionally closed at 10-15hrs after birth and anatomically closed at 10-21 days after birth.

Survival of newborns with certain heart lesions such as coarctation of the aorta, hypoplastic left heart syndrome and transposition of great vessels depends on the patency of ductus arteriosus. Closure of PDA in such conditions can lead to hypoperfusion, severe hypoxia and cyanosis.

PGE1 is a potent and specific relaxant of ductal smooth muscle. PGE1 analogues like alprostadil, misoprost and rioprostil can be used to keep the ductus open and support systemic blood supply.

In patent ductus arteriosus, the ductus is closed medically by NSAIDs (which inhibit prostaglandin synthesis) like ibuprofen trometamol/ ibuprofen lysine and indomethacin. Morphine can be used to manage respiratory distress but rarely preferred in a neonate.

Solution to Question 10:

The umbilical vein is the preferred vessel for an isovolumetric and simultaneous double set up exchange transfusion. The obliterated remnant of the umbilical vein is the Ligamentum teres.

The vessels that are occluded soon after birth are in due course replaced by fibrous tissue and forms the following ligaments:

The median umbilical ligament or Xander's ligament represents the remnant of the embryonic urachus.

The lateral umbilical ligaments are folds of the peritoneum over the inferior epigastric arteries.

Note:

Cord blood bilirubin level more than 4 mg/dL is an indication for change transfusion in a neonate.

Fetal structure	Adult structure
Foramen ovale	Fossa ovalis
Umbilical vein	Ligamentum teres
Ductus venosus	Ligamentum venosum
Umbilical arteries	Medial umbilical ligaments
Ductus arteriosus	Ligamentum arteriosum

Acyanotic Congenital Heart Diseases

Question 1:

The most common mode of inheritance of congenital heart disease is _____.

- a) Autosomal dominant
- b) Autosomal recessive
- c) X linked recessive
- d) Multifactorial

Question 2:

As a medical officer working under the RBSK scheme, the commonest congenital heart disease you would encounter is _____.

- a) Atrial septal defect
- b) Ventricular septal defect
- c) Patent ductus arteriosus
- d) Tetralogy of Fallot

Question 3:

As an intern posted in the pediatrics department, you notice a neonate with jaundice and a characteristic blueberry muffin rash. The mother gives an antenatal history of fever with a rash in her first trimester. On examination, he has bilateral cataracts, hepatosplenomegaly, and a murmur on auscultation. BERA reveals sensorineural deafness. On cardiac evaluation, which of the following is least likely to be found?

- a) Atrial septal defect
- b) Ventricular septal defect
- c) Patent ductus arteriosus
- d) Pulmonary stenosis

Question 4:

A mother reports to you that her baby is irritable and cries all the time. She has also noticed that her infant sweats excessively while feeding. Which of the following findings noted during the examination would you not include under NADA's major criteria?

- a) Wide and variable S2
- b) Congestive cardiac failure
- c) Grade 2 mid-diastolic murmur at the mitral area
- d) Grade 3 pansystolic murmur at the lower left sternal border

Question 5:

The most common type of ventricular septal defect is _____.

- a) Membranous
- b) Muscular
- c) Supracristal
- d) A and B have equal prevalence

Question 6:

An infant born to a diabetic mother was diagnosed with a ventricular septal defect. On examination of the baby, all of the following features can be found except:

- a) Ejection systolic murmur at the pulmonary area
- b) Holosystolic murmur at the left 4th intercostal space
- c) Reverse splitting of second heart sound (S2)
- d) A mid-diastolic murmur at the mitral area.

Question 7:

A mother is worried about her 4-month-old infant's growth. She says that her infant feeds slowly and often requires periods of rest during feeding. On examination:

- a) Left atrial enlargement
- b) Cardiomegaly with left ventricular apex
- c) Pulmonary edema
- d) Pulmonary oligemia

Question 8:

A 6-week-old infant is brought to the pediatrician for immunization. She has a loud and harsh holosystolic murmur best heard over the lower left sternal border and is accompanied by a thrill. The chest radiographs and EKG are normal. Echocardiography reveals a small muscular VSD. What is the appropriate management?

- a) Fenton's operation at 4 years
- b) Wait and watch till 5 years and then operate
- c) Restrict physical activity and operate at 2 years
- d) Reassure the parents

Question 9:

A good surgeon knows when not to operate. As a pediatric surgeon, on which of the following cases of VSD would you operate?

- a) 5 only
- b) 3,5
- c) 1,2,4
- d) 1,2,3,4


Question 10:

Rahel, a 3-year-old girl who is a known case of congenital heart disease is admitted to your pediatric ward because of low-grade fever with afternoon elevations. On examination, she is febrile and has painless small erythematous lesions on her palms. Her echocardiography showed oscillating intracardiac vegetation. Which one of the following congenital defects is most likely associated with the above findings?

- a) Atrial septal defect
- b) Ventricular septal defect
- c) Patent ductus arteriosus
- d) Pulmonic stenosis

Question 11:


A mother brings her 4-year-old child to the OPD. She says that the child has recurrent chest infections, easy fatiguability, and poor growth. You notice a limb abnormality as shown in the image below and find the given X-ray of the child's arm in his medical history. Based on all the findings, you suspect a congenital heart disease. Which of the following conditions are you most likely to find on screening echocardiography?



- a) Atrial septal defect
- b) Coarctation of aorta
- c) Pulmonic stenosis
- d) Patent ductus arteriosus

Question 12:


A child born to a 35-year-old mother was found to have a round face, flat occiput, and a protruding tongue. Karyotyping of the child is as shown below. Which of the following is commonly associated with this disorder?



- a) Bicuspid aortic valve
- b) Endocardial cushion defect
- c) Ventricular septal defect
- d) Coarctation of aorta

Question 13:

Which among the following is the most common type of atrial septal defect?



- a) 5
- b) 3

- c) 1
- d) 4

Question 14:

A 6-year-old child was brought to her pediatrician for routine well-child care. Her physical examination is normal except for a systolic murmur and a widely split second heart sound that does not vary with respiration. She has no family history of heart disease. Which among the following is the most likely diagnosis?

- a) Total anomalous pulmonary venous connection
- b) Ventricular Septal defect
- c) Atrial septal defect
- d) Pulmonic stenosis

Question 15:

A child who used to play for hours together now comes back home and says he gets tired too easily, his concerned mother brings him to you. On auscultation, you find that there is a wide and fixed splitting of the second heart sound, ECG shows left axis deviation and features of left ventricular hypertrophy. What is your probable diagnosis?

- a) Ostium secundum atrial septal defect
- b) Ventricular septal defect
- c) Ostium primum atrial septal defect
- d) Patent ductus arteriosus

Question 16:

An ECG finding of right ventricular enlargement in a 2-year-old child is most consistent with which of the following disorders?

- a) Atrial septal defect
- b) Patent ductus arteriosus
- c) Ventricular septal defect
- d) Tricuspid atresia

Question 17:

During an outreach program, you incidentally discover a child with an asymptomatic ASD. His parents are anxious as they had read on the web about transcatheter device closure and want to know if their child needs one. Your advice is he does not require it, for now, but needs regular follow-up. During the course of follow-up, you would take up the child for transcatheter device closure immediately in all of the following instances except:

- a) If the child becomes symptomatic
- b) If the size of the defect is $\leq 5\text{mm}$ at 2 years of age.
- c) If the Qp: Qs ratio is $\geq 2:1$
- d) If he develops RV enlargement

Question 18:

As an obstetrician, you discover an atrial septal defect incidentally in a 20-year-old primigravida. You will be watchful about all of the following complications, except:

- a) Infective endocarditis
- b) Stroke
- c) Arrhythmias
- d) Pulmonary hypertension

Question 19:

In a NICU you examine a preterm baby, who has a loud, harsh, continuous, machinery murmur in the Gibson area. All of the following can be found, except:

- a) Narrow pulse pressure
- b) Congestive heart failure
- c) Necrotising enterocolitis
- d) Cardiomegaly

Question 20:

You are posted as an intern in the pediatrics OPD and your professor asks you to examine an interesting case of cyanosis. On examination, you notice the bluish discoloration of the lower limbs but the upper limb and head are pink and on examination of the chest you see a

parasternal heave and a thrill is palpable. These findings can occur in which of the following?

- a) Patent ductus arteriosus
- b) Transposition of great vessels
- c) Tricuspid stenosis
- d) Ventricular septal defect

Question 21:

Where would you place the pulse oximeter to measure pre-ductal oxygen saturation in an infant who was born 3 minutes ago?

- a) Left upper limb
- b) Left lower limb
- c) Right upper limb
- d) Right lower limb

Question 22:

During neonatal resuscitation of a baby delivered via Cesarean section, you hear a continuous murmur. Which of the following would you not include in your differential diagnosis?

- a) Aortopulmonary window
- b) Small ventricular septal defect
- c) Patent ductus arteriosus
- d) Pulmonary arteriovenous malformation

Question 23:

A 10-year-old boy starts seizing while playing catch with his father. He is immediately brought to the casualty. On examination, blood pressure in the upper extremity measured as 200/140 mm Hg and femoral pulses were not palpable. What is the likely diagnosis?

- a) Takayasu aortoarteritis
- b) Renal parenchyma disease
- c) Grand mal seizure
- d) Coarctation of the aorta


Question 24:

A chest X-ray of a patient shows inferior rib notching. Which of the following diseases would you include in your differential diagnosis?

- a) Bidirectional Glen shunt
- b) Modified Blalock-Taussig shunt
- c) Inferior vena caval occlusion
- d) Coarctation of the aorta

Question 25:

What's your diagnosis and which is the most common anomaly associated with reference to the X-ray given below?



- a) Transposition of great vessels , renal dysplasia
- b) Coarctation of the aorta, bicuspid aortic valve
- c) Coarctation of aorta, renal dysplasia
- d) Transposition of great vessels, bicuspid aortic valve

Answer Key

Question No.	Correct Option
--------------	----------------

1	d
2	b
3	a
4	a
5	a
6	c
7	d
8	d
9	d
10	b
11	a
12	b
13	b
14	c
15	c
16	a
17	b
18	a
19	a
20	a
21	c
22	b
23	d
24	d
25	b

Detailed Explanations

Solution to Question 1:

The most common mode of inheritance of congenital heart disease is multifactorial.

Multifactorial inheritance refers to traits that are caused by a combination of inherited, environmental, and stochastic factors.

Solution to Question 2:

Ventricular septal defect(VSD) is the most common congenital heart disease.

The prevalence of congenital heart disease is estimated at 6-8 per 1000 live births.

The relative frequency of major CHD: VSD > ASD = PDA > CoA = ToF

NOTE:

Tetralogy of Fallot is the most common cyanotic heart disease.

Solution to Question 3:

The classical triad of cataract, sensorineural deafness, and congenital heart disease point towards congenital rubella syndrome. Atrial septal defect (ASD) is not found in congenital rubella infection.

Prenatal exposure to rubella increases the risk of congenital heart defects like:

- Patent ductus arteriosus (Most common)
- Peripheral pulmonic stenosis
- Ventricular septal defect (VSD)
- Myocarditis

Nonspecific signs of TORCH infections are hepatosplenomegaly, thrombocytopenia, jaundice, and growth restriction.

Solution to Question 4:

Wide and variable S2 (Abnormal S2) is not included under Nada's major criteria.

Features of congestive cardiac failure, a grade 3 pansystolic murmur with wide and variable S2, and an MDM at the mitral area suggest a diagnosis of large VSD in this baby.

NADA's criteria for clinical diagnosis of congenital heart disease:

The presence of 1 major or 2 minor criteria indicates congenital heart disease.

Pansystolic murmurs are always pathological no matter their intensity.

Short, soft systolic murmurs of intensity <3 are often detected during febrile illnesses, after which they disappear. Such murmurs are called innocent murmurs. Before labeling any murmur as innocent, it should be reassessed after an interval.

Major criteria	Minor criteria
Systolic murmur grade 3 or more	Systolic murmur < grade 3
Any diastolic murmur	Abnormal second sound (S2)
Cyanosis	Abnormal ECG
Congestive cardiac failure	Abnormal chest X-ray findings

Major criteria	Minor criteria
	Abnormal BP

Solution to Question 5:

Membranous is the most common type of ventricular septal defect (VSD) to occur.

VSD is the most common congenital cardiac malformation.

Types of VSD include membranous (most common), muscular, inlet/atrioventricular canal defects, and supracristal defects.

Swiss Cheese septum: a special type of muscular VSD with defects in the mid-portion or apical region of the ventricular septum.

Restrictive VSD -A small VSD where the size of the lesion is usually ≤ 5 mm. It is associated with normal right ventricular pressure.

Non-restrictive VSD -A large VSD where the size of the lesion is usually ≥ 10 mm leading to equalization of pressures in the right and left ventricle.

Solution to Question 6:

Wide, variable splitting of S2 is a feature of VSD and not reverse splitting.

The basic defect in VSD is a left-to-right shunt which occurs at the ventricular level. During systole, the left ventricle empties into the aorta and the right ventricle via the VSD, because of which the total amount of blood in the right side of the heart increases, and that on the left decreases. Due to this increase, the right ventricle takes a longer time to empty causing a delayed P2 and on the other hand, the left ventricle empties sooner causing an early A2. This explains the wide S2. Variations due to respiration still happen and hence, the second heart sound is widely and variably split.

Auscultatory findings of VSD include a pansystolic murmur also known as Roger's murmur/shunt murmur can be best heard in the fourth left intercostal space. Flow murmurs that can be heard include, an MDM at the mitral area and an ejection systolic murmur at the pulmonary area.

A hyperdynamic precordium, a palpable P2, parasternal heave, a laterally displaced apical impulse, and a systolic thrill can be detected on palpation.

Solution to Question 7:

The above findings are suggestive of a large VSD with congestive cardiac failure. On a chest X-Ray of such a patient pulmonary oligemia is not seen.

"Suck- rest-suck cycle ", failure to thrive, high respiratory rate are features suggestive of congestive cardiac failure. Acyanosis, hyperdynamic precordium, palpable P2, pansystolic murmur, and

MDM in the mitral area suggest a large VSD.

In ventricular septal defect (VSD), there is left to right shunting of blood, and this large volume of shunted blood passes through the lungs, which is recognized in the chest X-ray as pulmonary plethora (increased pulmonary vasculature). Other findings seen on a chest X-ray include cardiomegaly with left ventricular apex.

If the defect is large, frank pleural effusion can be seen.



Significant cardiomegaly, prominence of the pulmonary arterial trunk, pulmonary plethora (increased vascular markings seen in the lung fields) can be seen on the X-ray image.

Solution to Question 8:

In patients with small VSDs, parents should be reassured of the relatively benign nature of the lesion, and the child should be encouraged to live a normal life, with no restrictions on physical activity. Surgical repair is not recommended.

Guidelines from the Council on Cardiovascular Disease in the Young of the American Heart Association (AHA) state that an isolated, small, hemodynamically insignificant VSD is not an indication for surgery. VSDs may get smaller or close completely without treatment, but they won't get any bigger.

Echocardiography is used to estimate pulmonary artery pressure, screen for the development of Left ventricular outflow tract pathology (subaortic membrane or aortic regurgitation), and confirm spontaneous closure.

Solution to Question 9:

Severe pulmonary vascular disease unresponsive to pulmonary vasodilators is a contraindication to the closure of a VSD - due to permanent remodeling of the pulmonary vasculature.

Indications for surgical closure of a VSD include:

- Patients at any age with large defects in whom clinical symptoms and failure to thrive cannot be controlled medically;
- Infants between 6 and 12 mo of age with moderate to large defects associated with pulmonary hypertension, even if the symptoms are controlled by medication; and patients older than 24 mo with a Qp: Qs ratio greater than 2: 1.
- Patients with a supracristal VSD of any size are usually referred for surgery because of their higher risk for developing aortic valve regurgitation

A child with large VSD showing transient improvement in symptoms with better exercise tolerance may be the first sign of developing pulmonary hypertension and indicates an urgent need for evaluation and immediate surgery - the increased pulmonary pressure reduces the amount of blood pushed into the lungs and thereby, congestion transiently.

The presence of a mid-diastolic, low-pitched rumble at the apex is caused by increased blood flow across the mitral valve and usually indicates a Qp: Qs ratio $\geq$ 2: 1. This murmur is best appreciated with the bell of the stethoscope.

Surgical risks are somewhat higher for defects in the muscular septum, particularly apical defects and multiple (Swiss cheese-type) VSDs. These patients may require pulmonary arterial banding if symptomatic, with subsequent de-banding and repair of multiple VSDs at an older age.

Solution to Question 10:

The above findings are suggestive of a diagnosis of infective endocarditis according to Modified Duke's criteria. Patients with high-velocity blood flow lesions such as ventricular septal defect (VSD) and aortic stenosis are at high risk for infective endocarditis.

To make a diagnosis of IE, she has evidence of endocardial involvement (1 major) and fever, predisposing heart condition, Janeway lesions (a painless small erythematous lesion on palms) - 3 minor criteria.

In VSD, a high-pressure gradient across the defect $\rightarrow$ turbulent blood flow $\rightarrow$ traumatizes the vascular endothelium $\rightarrow$ predisposition for infective endocarditis. Important complications of VSD include infective endocarditis and Eisenmengerisation,

High-risk cardiac lesions for which endocarditis prophylaxis is advised before dental procedures and invasive procedures like tonsillectomy/adenoidectomy are prosthetic heart valves, prior endocarditis, post-cardiac transplant, congenital heart disease (6 months post repair and unrepaired).

Solution to Question 11:

The X-ray of the upper limb shows radial ray defects (absent radius and thumb) and the family history of heart disease suggest a diagnosis of Holt-Oram syndrome (aka hand-heart syndrome). Out of all the syndromes associated with cardiac defects, HOLT- ORAM has the strongest familial tendency - it is inherited in an Autosomal dominant manner.

Holt–Oram syndrome/cardiac limb syndrome is most commonly associated with atrial septal defect (ASD).

Features of Holt-Oram syndrome include:

- Hypoplastic or absent thumbs, radii
- Triphalangism
- Phocomelia
- 1st-degree heart block
- ASD

Solution to Question 12:

The given clinical findings and the karyotyping suggest a diagnosis of Trisomy 21 or Down syndrome. The most common cardiovascular lesion in Down's syndrome is an endocardial cushion defect.

Other associated defects are atrial septum defects (ASD), ventricular septum defect (VSD), and patent ductus arteriosus (PDA), aberrant subclavian artery, and pulmonary hypertension.


Other congenital defects are seen in some syndromes:

Syndrome	Congenital defect
Trisomy 21 (Down syndrome)	Endocardial cushion defect, VSD, ASD
XO (Turner syndrome)	Bicuspid aortic valve, coarctation of aorta
William's syndrome	Supravalvular aortic stenosis, peripheral pulmonic stenosis
Noonan syndrome	Pulmonic stenosis, ASD
Holt–Oram syndrome	ASD, 1st-degree heart block
Marfan's Syndrome	Mitral or Aortic Regurgitation

Solution to Question 13:

Ostium secundum ASD (labelled as 3) is the most common type of atrial septal defect (ASD), located at the mid-inter-atrial septum, in the region of the fossa ovalis.

Types of Atrial Septal Defects



Based on location

©Marrow

ASD is characterized by an abnormal opening in the atrial septum allowing communication between the right and left atria. There are four types based on their location within the septum:

- Secundum ASD - most common, usually an isolated abnormality
- Primum ASD - associated with abnormal mitral valve leaflet (partial AV septal defect)
- Sinus venous - is located at the junction of the superior vena cava and right atrium. Associated with anomalous right pulmonary venous return to the SVC or RA
- Coronary sinus type ASD - also called unroofed coronary sinus

Ostium primum ASD: is located at lower atrial septum

Endocardial cushion defect/An AV septal defect/ AV canal defect consists of contiguous atrial and VSDs with markedly abnormal AV valves. The lesion is common in children with Down syndrome.

Congenital conditions associated with ASD include:

- Down's syndrome - usually an ostium primum defect
- Holt-Oram syndrome - autosomal dominant disorder with upper limb bony defects and cardiac septal defects, commonly secundum ASD
- Ellis-van Creveld syndrome - chondroectodermal dysplasia associated with postaxial polydactyly and ASD
- Lutembacher syndrome - combined ASD and mitral stenosis due to rheumatic valvulitis
- Mitral valve prolapse

Solution to Question 14:

The age of the patient, normal physical findings except for a systolic murmur, and wide and fixed splitting of S2 are all in favor of a diagnosis of Atrial septal defect.

Most ASDs are asymptomatic and are picked up during routine physical examinations. A soft ejection systolic murmur best heard over the left upper sternal border is also commonly found.

The soft ejection systolic murmur is a flow murmur heard due to increased blood flow through the normal pulmonary valve. A delayed diastolic murmur can also be heard in the tricuspid area. As the right ventricle now receives more blood, the time required for its emptying is increased resulting in prolonged P2. Since the two atria are linked, inspiration does not produce any net pressure changes between them abolishing respiration-related changes in venous return hence causing fixed S2.

Causes of Wide and fixed splitting of S2 include,

- Atrial septal defect
- RBBB
- TAPVC

Other options:

TAPVC - patients usually present with cyanosis and congestive cardiac failure around 4-10 weeks of age - this is when the fetal pulmonary vasculature regresses and there is increased pulmonary blood flow which results in early-onset cardiac failure. The physical findings are identical to ASD.

VSD - is an acyanotic heart disease and physical findings include a pansystolic murmur best heard in the left 4th parasternal area and a wide and variable split S2 which is usually masked by the pansystolic murmur.

PS - is an acyanotic heart disease and physical findings include an ejection systolic murmur and a thrill in the left second intercostal space and a wide and variable split S2 because of delayed closure of P2.

Solution to Question 15:

Wide splitting of the 2nd heart sound with left axis deviation (LAD) and left ventricular hypertrophy (LVH) points to the diagnosis of ostium primum atrial septal defect (ASD). Wide-fixed splitting of the 2nd heart sound is characteristic of ASD.

In ostium primum ASD, the basic abnormality is the combination of left-to-right shunt across the atrial defect and mitral insufficiency due to mitral valve clefts. Mitral valvular insufficiency increases the volume load on the left ventricle (seen as LAD & LVH on ECG). When mitral insufficiency is severe enough to cause clinical symptoms like fatigue on exertion, the left atrium enlarges as a result of the regurgitant flow, and the left ventricle becomes hypertrophied and dilated.

Solution to Question 16:

In atrial septal defect (ASD), there is a left-to-right shunt, resulting in increased blood flow through the right side of the heart, which leads to enlargement of the right atrium and ventricle and dilation of the pulmonary artery, the left atrium may also be enlarged, but the left ventricle and aorta are normal in size.

Patent ductus arteriosus (PDA): There is shunting of blood from the aorta to the pulmonary trunk, which goes into the lungs and then again to the LA and LV. If the ductus is large, left ventricular or biventricular hypertrophy is present.

In ventricular septal defect (VSD), there is shunting of blood from the left ventricle to the right ventricle, but most of the blood is shunted directly to the inflow tract of the pulmonary trunk, so, there is no significant right ventricular hypertrophy. In VSD, the pulmonary artery left atrium, and left ventricle are enlarged.

In Tricuspid atresia, there is right to left shunting of blood through an ASD as the tricuspid valve is dysfunctional due to which the entire systemic venous return enters the left side of the heart using the foramen ovale or through an ASD, and hence, there is left ventricular hypertrophy.

Solution to Question 17:

An ASD defect of size ≤ 5 mm at 2 years of age does not need immediate transcatheter device closure.

Small ASDs (3 to ≤ 6 mm in diameter) may grow smaller or close spontaneously. They do not cause symptoms and usually only need observation.

Transcatheter device or surgical closure is advised for:

- All symptomatic patients
- Asymptomatic patients with a Qp: Qs ratio of at least 2: 1
- Those with RV enlargement.
- ASD size ≥ 8 mm

Closure (if indicated) is recommended before school entry. Repair is preferred during early childhood because surgical mortality and morbidity are significantly greater in adulthood.

Solution to Question 18:

Infective endocarditis is extremely rare in isolated atrial septal defect (ASD) and antibiotic prophylaxis for isolated ASDs is not recommended.

ASD is a low-pressure gradient defect whereas a high-pressure gradient across a defect is required to create a turbulent flow, which traumatizes the vascular endothelium, making the patient susceptible to endocarditis.

Patients with a small or moderate ASD are asymptomatic unless a complication occurs.

Pulmonary hypertension, atrial dysrhythmias, tricuspid or mitral insufficiency, and heart failure may initially appear during the increased volume load of pregnancy.

Solution to Question 19:

A preterm baby with the classical continuous machinery murmur in the Gibson's area is in favor of patent ductus arteriosus as the diagnosis. Preterm babies with patent ductus arteriosus (PDA) have wide pulse pressure, hyperdynamic precordium, and bounding peripheral pulses with a continuous murmur.

If the PDA is large, it can cause overloading of the left ventricle leading to cardiomegaly and congestive heart failure. Retrograde aortic flow, decreased systemic blood flow, and moderate hypotension is common in premature infants with a PDA predisposing them to necrotizing enterocolitis.

Solution to Question 20:

Differential cyanosis can be seen in patent ductus arteriosus (PDA) when there is a reversal of shunt.

Differential cyanosis is manifested as cyanosis of only the lower extremities while upper extremities are unaffected. It is a result of reduced arterial oxygen saturation in the distal aorta compared to that in the aorta proximal to the left subclavian artery.


It is seen with:

- right-to-left shunting across a ductus arteriosus in the presence of coarctation or an interrupted aortic arch
- patent ductus arteriosus (PDA) with severe pulmonary artery hypertension (PAH)

The rest of the options cause cyanosis of all extremities. Among the options, PDA is the only condition that can manifest with differential cyanosis when there is a reversal of shunt.

Image: Right to left shunting of blood flow through a PDA between the aortic arch and pulmonary artery.

Reversal of shunt in Patent ductus arteriosus



©Marrow

Solution to Question 21:

The right upper limb is the best area to check pre-ductal oxygen saturation in a newborn infant as it receives blood supply primarily from the right brachial artery, which arises from the aorta before the ductus arteriosus.

In routine neonatal screening for critical congenital heart diseases (CCHD), saturation probes are placed on the right upper limb (pre-ductal) and the right lower limb (post-ductal) respectively.

If the difference between pre-ductal and post-ductal saturation is more than five percent, it confirms the presence of hypoxemia. Any reading showing an SpO₂ <90% or greater than a three percent difference between the right hand and right foot for three successive readings separated by an hour each would suggest the presence of a congenital cardiac disease.

Screening is done to identify critical conditions before they present symptomatically, and before the neonate leaves the hospital post-delivery. This would allow for early and effective treatment of an underlying disease.

Critical congenital heart diseases include coarctation of the aorta, double outlet right ventricle, Ebstein anomaly, hypoplastic left heart syndrome, interrupted aortic arch, pulmonary atresia, single ventricle, tetralogy of Fallot (TOF), total anomalous pulmonary venous return (TAPVC), d-transposition of the great arteries (TGA), tricuspid atresia, and truncus arteriosus.

Solution to Question 22:

Small ventricular septal defect (VSD) characteristically produces a loud, harsh, or blowing holosystolic murmur (not continuous murmur) and is heard best over the lower left sternal border; it is frequently accompanied by a thrill.

Pansystolic or holosystolic murmurs begin almost simultaneously with the 1st heart sound and continue throughout systole (do not spill into diastole like a continuous murmur).

Continuous murmur is a systolic murmur that continues or “spills” into diastole and indicates continuous flow, such as in the presence of a patent ductus arteriosus or other aortopulmonary communication.

Other options:

Patent ductus arteriosus (PDA): The classic continuous murmur is described as being like machinery in quality.

Aortopulmonary window: The findings can mimic those of a patent ductus arteriosus (PDA) (wide pulse pressure and a continuous murmur).

Pulmonary arteriovenous malformations: Large fistulas are associated with dyspnea, cyanosis, clubbing, a continuous murmur, and polycythemia, commonly associated with Osler Weber Rendu disease.

Solution to Question 23:

Here, the diagnosis is coarctation of the aorta because the classic sign of coarctation of the aorta is a disparity in pulsation and blood pressure in the arms and legs (as seen in this patient). The femoral, popliteal, posterior tibial, and dorsalis pedis pulses are weak (or absent in up to 40% of patients), in contrast to the bounding pulses of the arms and carotid vessels. Seizure is a complication secondary to systemic hypertension since untreated coarctation can increase the risk for intracranial hemorrhage and hypertensive encephalopathy.

The most common site of coarctation of the aorta is distal to the origin of the left subclavian artery near the insertion of ligamentum arteriosum (juxtaductal coarctation).

DDx: Takayasu arteritis is a chronic vasculitis of unknown etiology, predominantly affecting women with an age of onset that is usually between 10 and 40 years. The disease primarily affects the aorta and its primary branches.

Solution to Question 24:

Notching of the inferior border of the ribs (four to eighth ribs) from pressure erosion by enlarged collateral vessels is a characteristic finding in the coarctation of the aorta by late childhood.

Causes of rib notching

Superior rib notching	Inferior rib notching
Poliomyelitis Osteogenesis imperfecta Neurofibromatosis Marfan's syndrome Connective tissue diseases Rheumatoid arthritis Systemic lupus erythematosus Scleroderma Hyperparathyroidism	Coarctation of the aorta Blalock-Taussig shunt Superior vena cava obstruction AV malformation of chest wall Schwanoma



Solution to Question 25:


The given X-ray demonstrates the “3” sign, which is pathognomic of coarctation of aorta. Bicuspid aortic valve is commonly associated with coarctation of the aorta.

3 sign: Indentation of the aorta at the site of coarctation, pre and post-stenotic dilatation along the left paramediastinal shadow

Other Chest X-ray features in coarctation of aorta:

- Dilated left subclavian artery
- Dilated ascending aorta
- Notching of 3rd and 9th ribs due to rib erosion by dilated collateral vessels

10% of individuals with CoA have a circle of Willis aneurysm.



Cyanotic Congenital Heart Diseases

Question 1:

You are called to see a cyanotic neonate in the NICU. You note that the neonate has an oxygen saturation of 60%. Which of the following would not be in your differential diagnosis

- a) Ebstein anomaly
- b) Ventricular septal defect with pulmonic stenosis
- c) Pulmonary AV malformation
- d) Isolated infundibular stenosis


Question 2:

Which of the following is the most common cyanotic congenital heart disease?

- a) Tetralogy of Fallot
- b) Transposition of Great Arteries
- c) Tricuspid Atresia
- d) Total Anomalous Pulmonary Venous Connection

Question 3:

Below is an X-ray of an infant with cyanotic heart disease. Which is not a finding seen here?



- a) Ventricular septal defect
- b) Right ventricular hypertrophy
- c) Atrial septal defect
- d) Pulmonary stenosis

Question 4:

A mother brings her 2-month-old blue baby to your pediatric clinic for the first time. Neonatal records were not available but the mother says that the X-ray of the baby looked like that of a shoe. Which of the following wouldn't you expect in this baby?

- a) Soft P2
- b) Large ascending aortic root
- c) Cardiomegaly
- d) Large "a" waves in jugular venous pulse

Question 5:

A primary school teacher notices a peculiar behavior in one of her pupils. Whenever he plays with his friends, she notices that her fingers and lips turn blue. The child then squats for sometime after which he feels better. Which of the following does not occur during squatting?

- a) It increases the systemic vascular resistance
- b) It increases the magnitude of right-to-left shunt
- c) Leads to trapping of deoxygenated blood in lower limbs
- d) Both A and C

Question 6:

While waiting for corrective open-heart surgery, you decide to manage a case of tet spells medically. Which of the following drugs would worsen the cyanosis?

- a) Sodium bicarbonate
- b) Propranolol
- c) Phenylephrine
- d) Isoprenaline

Question 7:

Blalock-Taussig shunt is done between_____.

- a) Subclavian artery to pulmonary artery
- b) Aorta to pulmonary vein
- c) Aorta to pulmonary artery
- d) Subclavian vein to pulmonary artery

Question 8:

Which of the following is a component of the pentalogy of Fallot?

- a) Atrial septal defect
- b) Patent ductus arteriosus
- c) Coarctation of the aorta
- d) Left ventricular hypertrophy

Question 9:

An adolescent who is a known case of VSD is brought by his parents with complaints of breathlessness on exertion which is gradually progressive, bluish discoloration, and swelling of toes and fingers. Investigations showed features suggestive of pulmonary hypertension. Which of the following is not a part of the condition?

- a) Right ventricular hypertrophy
- b) Bidirectional flow of blood
- c) Obstructive pulmonary hypertension
- d) Predominant left to right shunt

Question 10:

The most common type of total anomalous pulmonary venous connection is _____.

- a) Supracardiac
- b) Cardiac
- c) Infracardiac

d) Mixed

Question 11:

A neonate was noted to have cyanosis of the oral mucosa. Chest x-ray revealed a snowman appearance of the cardiac silhouette. What is the most likely diagnosis?

- a) Transposition of great arteries**
- b) Truncus arteriosus**
- c) Tricuspid atresia**
- d) Total anomalous pulmonary venous connection**

Question 12:

You are examining a week-old male infant who was severely cyanotic at birth in the post-operative ward. He has undergone balloon atrial septostomy following prostaglandin E₁ administration. Post-surgery there was a marked improvement in cyanosis. The most likely diagnosis is_____.

- a) Tetralogy of Fallot**
- b) Transposition of great arteries**
- c) Truncus arteriosus**
- d) Tricuspid atresia**


Question 13:

A 24-year-old woman with bipolar disorder on lithium therapy expresses intent to conceive. She must be advised to stop the drug due to the risk of development of which of the following conditions in the fetus?

- a) Ebstein anomaly**
- b) VACTERAL defects**
- c) Coarctation of aorta**
- d) Supravalvular aortic stenosis**

Question 14:

Akshay, a high schooler presents to you with swelling of fingers and toes. He said he gets palpitations while he is exercising and even when he hangs out with friends. An X-Ray of the patient is given below. ECG showed the presence of delta waves. What is your diagnosis?



- a) Partial anomalous pulmonary venous return
- b) Total anomalous pulmonary venous return
- c) Endocardial cushion defects
- d) Ebstein anomaly

Question 15:

In a 13-year-old child with Ebstein's anomaly, echocardiography would likely reveal:

- a) Mitral regurgitation
- b) Tricuspid regurgitation
- c) Mitral stenosis
- d) Tricuspid stenosis

Question 16:

A cyanotic newborn is suspected of having congenital heart disease. He has an increased left ventricular impulse and a holosystolic murmur along the left sternal border. ECG shows left axis deviation and left ventricular hypertrophy. Which of the following is the most likely diagnosis?

- a) Transposition of Great Arteries
- b) Truncus arteriosus
- c) Tricuspid atresia
- d) Tetralogy of Fallot

Question 17:

You are examining a case of cyanosis in NICU as a resident. Which of the following cardiac features would make the diagnosis of tricuspid atresia unlikely?

- a) Left axis deviation on ECG
- b) Himalayan P waves on ECG
- c) Ventricular septal defect on echo
- d) Left ventricular hypoplasia on echo

Question 18:

Which of the following cases could develop recurrent infections due to pulmonary congestion?

- a) 2 and 3
- b) 1, 3 and 4
- c) 1 and 4
- d) 1, 2, and 3


Question 19:

A cyanotic baby at the 8th hour of life is being evaluated in NICU. Pulse oximetry shows 80 % not improving with 100 % O₂. Records show that continuous machinery murmur was heard at birth but not anymore. Which of the following can be done?

- a) Intubation with 100% O₂
- b) Indomethacin administration
- c) RBC transfusion
- d) Prostaglandin E₁ infusion

Question 20:


The given chest X-ray is suggestive of:



- a) Tetralogy of Fallot
- b) Atrial septal defect
- c) Ventricular septal defect
- d) Ebstein's anomaly

Question 21:

Which among the following options describes the chest x-ray image?



- a) Egg on string appearance

- b) Coeur en sabot
- c) Scimitar sign
- d) Snow man sign

Question 22:

Of the following, all are findings of the condition in the below chest x-ray except:



- a) Increased pulmonary blood flow
- b) Narrow vascular pedicle
- c) Pulmonary venous hypertension
- d) Right atrial enlargement

Question 23:

Below is the chest X-ray image obtained for a neonate on day 7 of life who developed heart failure later. What is the most probable diagnosis?



- a) Tetralogy of Fallot
- b) Ventricular septal defect
- c) Total anomalous pulmonary venous connection
- d) Transposition of great arteries

Answer Key

Question No.	Correct Option
1	d
2	a
3	c
4	c
5	b
6	d
7	a
8	a
9	d
10	a
11	d
12	b
13	a
14	d

15	b
16	c
17	d
18	c
19	d
20	a
21	b
22	d
23	d

Detailed Explanations

Solution to Question 1:

Given clinical scenario is suggestive of cyanotic congenital heart disease. Isolated infundibular stenosis would not be in the differential diagnosis because it is an acyanotic congenital heart disease.

Time of onset of cyanosis in congenital heart disease:

Congenital cyanotic heart disease	Time of onset of cyanosis
Tetralogy of Fallot with pulmonary atresia Pulmonary atresia with intact ventricular septum Tricuspid Atresia Hypoplastic Left Heart Syndrome Dextro - Transposition of Great Arteries (TGA) - with Intact Ventricular Septum Severe Ebstein's anomaly	1st few hours or days of birth
Total anomalous pulmonary venous connection	Neonatal period
Tetralogy of Fallot TGA with VSD and pulmonic stenosis	1st few months of life. (Not at birth)
Truncus arteriosus	Minimal cyanosis at birth Moderate cyanosis in 1-2 months.
Ebstein's anomaly	Mild: teenage or young adult. Severe: at birth

Solution to Question 2:

The most common cyanotic congenital heart disease is tetralogy of Fallot (ToF).

Infants are often asymptomatic until later in life (beyond 1yr of life).

TOF has 4 components:


- Obstruction to right ventricular outflow (pulmonary stenosis)
- Malalignment type of ventricular septal defect (VSD)
- Dextroposition of the aorta so that it overrides the ventricular septum (overriding of the aorta), and
- Right ventricular hypertrophy (RVH)

Solution to Question 3:

Cyanosis and boot-shaped heart (cœur en sabot in French) points towards tetralogy of Fallot (TOF) and the atrial septal defect is not a finding seen in TOF.


Clinical findings in TOF:

- Right ventricular hypertrophy causing left anterior hemithorax bulge.
- Single S2 or soft P2 because of the stenotic pulmonary valve
- Systolic thrill in the left sternal border in 3rd and 4th parasternal spaces.
- The large 'a' wave in the jugular pulse is a reflection of the forceful contraction of the right atrium at the end of diastole as there is reduced distensibility of the hypertrophic right ventricle.
- There is a loud and harsh systolic murmur due to the right ventricular outflow tract (RVOT) obstruction. It becomes louder, longer, and harsher as obstruction increases in severity from mild to moderate.
- However in severe obstruction, especially during tet spell, the murmur becomes less prominent as the blood is shunted towards the aortic valve and not through RVOT.



X-Ray in TOF showing boot shaped heart:

Tetralogy of Fallot - boot-shaped heart



Acyanotic/pink tetralogy of Fallot: This occurs when obstruction to the right ventricular outflow is mild to moderate. The blood would flow more through the mildly obstructed RVOT and less through the VSD. Hence, there is less cyanosis.

Solution to Question 4:

The given history of the shoe (boot) shaped heart in cyanotic baby is suggestive of tetralogy of Fallot and cardiomegaly is not a part of TOF presentation.

In the tetralogy of Fallot (TOF), we see concentric right ventricular hypertrophy without cardiac enlargement. It is because of pulmonary stenosis (PS) and the ventricular septal defect (VSD). Any build-up of pressure in the right ventricle due to PS is released through VSD to the left ventricle.

Option A: P2 is of low intensity and soft because the right ventricular outflow obstruction results in reduced pulmonary artery pressure.

Option B: The aortic root usually appears large because it overrides (straddling) the VSD, allowing deoxygenated blood shunted from the right ventricle mixed with blood from the left ventricle to exit the heart.

Option D: The large 'a' wave in the jugular pulse is a reflection of the forceful contraction of the right atrium at the end of diastole as there is reduced distensibility of the hypertrophic right ventricle.

Solution to Question 5:

Squatting decreases the magnitude of the right-to-left shunt in a child with tetralogy of Fallot (TOF).

The child feels comfortable while squatting due to two compensatory mechanisms:

- Increasing systemic vascular resistance and, thus, decreasing the magnitude of the right-to-left shunt across the ventricular septal defect (VSD). Therefore, more blood goes to the lungs for oxygenation.
- Trapping deoxygenated venous blood in the lower limbs.

Solution to Question 6:

Isoprenaline/isoproterenol should be avoided in tet or hypercyanotic spells. It will cause a decrease in total peripheral resistance (TPR) or systemic vascular resistance (SVR). So, more blood leaves the left ventricle leading to the increased right to left shunt. This results in the worsening of right ventricular outflow and causes worsening of cyanosis and the symptoms.

Treatment of hypercyanotic spell involves placing child in knee-chest position (or squatting), and use of:

- oxygen
- morphine - sedates the patient
- sodium bicarbonate - to counteract metabolic acidosis that develops when arterial $P_{O_2} < 40$ mmHg
- alpha-agonists (phenylephrine)- vasopressor used in severe cases
- beta-blockers (propranolol)- improves resting saturation and decreases the frequency of spells.

Drugs that increase SVR, such as IV phenylephrine (alpha-1 agonist) or propranolol (non-selective beta-antagonist/beta-2 block), can improve right ventricular outflow, decrease the right-to-left shunt, and improve the symptoms. This is same principle as squatting.

Solution to Question 7:

Blalock-Taussig shunt is done between subclavian artery to pulmonary artery

Blalock-Taussig shunt is currently the most common aortopulmonary shunt procedure and consists of a Gore-Tex conduit.

Sometimes, the shunt is brought directly from the ascending aorta to the main pulmonary artery; in this case, it is called the central shunt.

Corrective open-heart surgery performed in critically ill newborns and early infancy is the preferred surgical treatment now.

Name of the surgery	Connects
Blalock-Taussig shunt (B S)	Subclavian artery to pulmonary artery
Waterston's shunt (WA)	Ascending aorta to right pulmonary artery

Name of the surgery	Connects
Pott's shunt (PD)	Descending aorta to pulmonary artery

Solution to Question 8:

Pentology of Fallot consists of

- Tetralogy of Fallot: Ventricular septal defect + Pulmonic stenosis + Overriding of the aorta + Right ventricular hypertrophy
- Atrial septal defect


Solution to Question 9:

Given clinical scenario along with the development of pulmonary hypertension is seen in Eisenmenger's syndrome and predominant right to left shunt (not left to right shunt) is seen here.

Gradual onset of cyanosis, dyspnoea, fatigue, and in later stages, heart failure, chest pain, headaches, syncope, and hemoptysis in a setting of congenital acyanotic heart disease, are signs and symptoms of Eisenmenger's syndrome.

Reversal of the shunt at the level of atria/ventricles/patent ductus arteriosus due to irreversible pulmonary vascular obstructive changes is defined as Eisenmenger's syndrome.

Eisenmenger's syndrome pathophysiology



©Marrow

Features of pulmonary hypertension in Eisenmenger physiology: Right ventricular heave, narrow split S 2 with a loud pulmonic component.

Chest X-ray showing prominent pulmonary artery and right ventricle, peripheral pruning of pulmonary vessels.

ECG shows features of right ventricular hypertrophy, with spiked P waves.

Treatment: Calcium channel blockers, prostacyclin, and its analog treprostinil and lung transplantation.

Solution to Question 10:

The most common type of total anomalous pulmonary venous connection (TAPVC) is supracardiac.

In TAPVC, pulmonary veins instead of draining into the left atrium drain into the right atrium directly or indirectly.

- Supracardiac (most common): Common pulmonary vein drains into superior vena cava or left innominate vein.
- Cardiac: Pulmonary veins drain into right atrium or coronary sinus.
- Infracardiac (always obstructive and most dangerous): Pulmonary veins drain into the hepatic vein or portal vein or inferior vena cava.


Solution to Question 11:

Snowman's appearance on the chest X-ray image is classically seen in total anomalous pulmonary venous connection (TAPVC).

The supracardiac variant can classically depict a snowman appearance/figure of 8/cottage loaf heart on the frontal chest radiograph.

The dilated vertical vein on the left, brachiocephalic vein on the top, and the superior vena cava on the right form the head of the snowman; the body of the snowman is formed by the enlarged right atrium.

The image given above shows a snowman or figure-8 configuration. Cardiomegaly and increased pulmonary vascularity are evident.



Solution to Question 12:


Cyanosis at birth, prostaglandin E1 administration, and septostomy to improve oxygenation are typical features of transposition of great vessels (TGA).

TGA is characterized by the aorta arising from the right ventricle and pulmonary artery from the left ventricle. Since systemic and pulmonary circulations are separate, survival depends on the mixing of oxygenated pulmonary blood and deoxygenated systemic blood which can occur:

- At atrial level: Through atrial septal defect (ASD)
- At ventricle level: Through ventricular septal defect (VSD)
- At great vessel level: Through patent ductus arteriosus

Chest X-ray image shows cardiomegaly, a narrow mediastinum (the classic “egg-on-string appearance”), and normal to increased pulmonary blood flow. The narrow pedicle of the heart is because of the malposition of the great vessels.

Treatment: Here, prostaglandin E1 administration and septostomy help initial stabilization by creating a communication between the right and left atria, thereby facilitating the mixing of systemic and pulmonary circulations.



Solution to Question 13:

She must be asked to stop the intake of lithium as it can cause an Ebstein anomaly in the newborn.

Ebstein anomaly is characterized by the downward displacement of the tricuspid valve into the right ventricle, which results in a thin-walled, low-pressure, "atrialized" segment of the right ventricle due to the anomalous attachment of the tricuspid leaflets.


Lithium can also cause macrocephaly as a teratogenic effect.

Solution to Question 14:

The diagnosis is Ebstein's anomaly.

The presence of clubbing and a box-shaped heart on chest X-Ray in the given case is suggestive of Ebstein's anomaly. Clubbing develops secondary to cyanosis as Ebstein's anomaly is a right to left shunt. Delta wave is observed in Wolff-Parkinson-White (WPW) syndrome. WPW and supraventricular tachycardia causing palpitations can be seen in association with Ebstein's anomaly.

Ebstein's Anomaly - box-shaped heart



Solution to Question 15:

Tricuspid regurgitation would be seen in a child with an Ebstein anomaly.

Mild cases develop cyanosis during adolescence or early adulthood. In severe cases, cyanosis manifests at birth.

Ebstein anomaly is characterized by the downward displacement of the tricuspid valve into the right ventricle, which results in a thin-walled, low-pressure, "atrialized" segment of the right ventricle due to the anomalous attachment of the tricuspid leaflets.

In Ebstein's, tricuspid valve tissue is dysplastic and results in tricuspid regurgitation.

The right ventricle is hypoplastic, which leads to decreased pulmonary outflow and there will be functional pulmonary atresia. As a result, blood shunts through the foramen ovale/atrial septal defect (ASD) results in cyanosis at birth.

Himalayan P wave (amplitude > 5 mm) is associated with tricuspid atresia, Ebstein anomaly, and combined tricuspid and pulmonary stenosis, where they indicate a dilated right atrium. Wolff–Parkinson–White syndrome (WPW) and supraventricular tachycardia may be present in Ebstein anomaly.

Note:

In Ebstein anomaly, the malformed tricuspid valve is usually regurgitant, but may rarely be stenotic.

Solution to Question 16:

Clinical cyanosis in combination with left-axis deviation (LAD) and left ventricular hypertrophy (LVH) noted on the electrocardiogram distinguishes tricuspid atresia from most other cyanotic heart lesions.

Other points in favor:

- Increased left ventricular impulse (in contrast to most other causes of cyanotic heart disease, in which an increased right ventricular impulse).
- Holosystolic murmurs audible along the left sternal border (due to associated ventricular septal defect).

Tricuspid atresia is the congenital absence of the tricuspid valve. So, blood cannot flow directly from the right atrium to the right ventricle, and the following associated defects are necessary for survival:

- Patent foramen ovale or atrial septal defect: for the flow of blood from the right atrium to the left atrium
- Ventricular septal defect: for the flow of blood from the left ventricle to the right ventricle
- There is a hypoplastic right ventricle.

Option B and D: Truncus arteriosus and tetralogy of Fallot and - The ECG demonstrates right axis deviation (RAD) and evidence of RVH.


Option A: Transposition of great arteries- ECG is usually normal, showing the neonatal right-sided dominant pattern.

Solution to Question 17:

Left ventricular hypoplasia would make the diagnosis unlikely as right ventricular hypoplasia is seen in tricuspid atresia.

Other features of tricuspid atresia:

- An increased left ventricular impulse is noted in contrast to an increased right ventricular impulse seen in most other cyanotic diseases.
- A VSD holosystolic murmur is audible along the left sternal border.
- The second heart sound (S2) is usually single.
- Left ventricular hypertrophy and left axis deviation on ECG.
- Tall, peaked P waves on lead II known as Himalayan P waves which are suggestive of an enlarged right atrium.



Solution to Question 18:

Patients with atrial and ventricular septal defects have increased pulmonary blood flow. This results in recurrent pulmonary infection due to congestion in the lungs.

Tetralogy of Fallot (TOF) and tricuspid atresia (TA) are associated with reduced pulmonary blood flow (oligemic lung fields). They do not cause pulmonary congestion and do not predispose to infections.

Reduced pulmonary blood flow is due to

- Right ventricular outflow obstruction in tetralogy of Fallot
- Right ventricular inflow obstruction in tricuspid atresia

Cyanotic congenital heart diseases (CCHD) with decreased and increased pulmonary blood flow:


Decreased pulmonary blood flow	Increased pulmonary blood flow
Tetralogy of Fallot Pulmonary atresia Ebstein's anomaly Tricuspid atresia with an intact ventricular septum Transposition of the great vessels with VSD and pulmonary stenosis	Transposition of the great vessels with VSD or PDA Total anomalous pulmonary venous return (TAPVC) Truncus arteriosus Hypoplastic left heart syndrome (HLHS)

Solution to Question 19:

This baby has a cyanotic congenital heart disease that is dependent on a patent ductus arteriosus. Prostaglandin E1 infusion prevents the closure of PDA in term neonates with duct-dependent

congenital heart disease (CHD).

Cyanosis failing to improve with 100% oxygen suggests an underlying cyanotic CHD.



Functional closure of the PDA occurs soon after birth, usually within first week of life.

The open PDA acted as a conduit between systemic and pulmonary circulations, thereby preventing the onset of cyanosis.

Upon its closure, cyanosis manifested in the baby as the blood could not be pumped into the lungs for oxygenation. Deoxygenated blood is being shunted into the systemic circulation after the closure of PDA. Hence, you should maintain the patency of the duct with PGE1 infusion until surgery can be done.

Duct-dependent lesions: Anomalies, in a newborn with CHD, which are dependent on the patency of ductus arteriosus for survival.

- Right-sided lesions (Tetralogy of Fallot, PA/PS) cannot pump blood to the lungs. These patients are dependent on an open PDA for pulmonary circulation.
- Left-sided lesions (Hypoplastic left heart syndrome, Aortic stenosis, Coarctation of aorta) cannot pump blood into the systemic circulation. These patients are dependent on open PDA for systemic circulation.

Solution to Question 20:

The given chest X-ray shows a boot-shaped cardiac silhouette which is suggestive of the Tetralogy of Fallot.

The boot-shaped (*coeur en sabot*) cardiac silhouette is seen due to the right ventricular hypertrophy and upward shift of the cardiac apex. The hilar areas and lung fields appear relatively clear due to reduced pulmonary blood flow or the small size of the pulmonary arteries, or both.

In Ebsteins's anomaly (Option D) box-shaped heart is seen.


Solution to Question 21:

The given chest x-ray shows 'coeur-en-sabot' or boot-shaped heart, which is seen in tetralogy of Fallot (TOF).

The characteristic silhouette is due to cardiac apex elevation as a result of right ventricular hypertrophy. The hilar areas and lung fields are relatively clear because of diminished pulmonary blood flow or the small size of the pulmonary arteries and the aorta is usually enlarged.

The image below describes the likeness of the cardiac silhouette to a boot:

Tetralogy of Fallot - boot-shaped heart



TOF is cyanotic congenital heart disease with the following four components:

- Obstruction to right ventricular outflow (pulmonary stenosis)
- Malalignment type of ventricular septal defect (VSD)
- Dextroposition of the aorta so that it overrides the ventricular septum (overriding of the aorta)
- Right ventricular hypertrophy (RVH)

The severity of symptoms depends on the degree of the right ventricular outflow obstruction. In mild degrees, the symptoms become pronounced as the baby grows due to RV infundibular hypertrophy and may present with heart failure. In severe degrees, pulmonary blood flow predominantly depends on the ductus arteriosus. As soon as the duct closes, the baby presents with cyanosis and circulatory collapse in the neonatal period.

Clinical features include significant hypercyanotic spells in infants which are characterised by hyperpnea, cyanosis, and gasping followed by syncope. The spells occur most frequently in the morning on initially awakening or after episodes of vigorous crying. These spells are due to a prolonged reduction of pulmonary blood flow resulting in severe systemic hypoxia and metabolic acidosis.

Older children present with exertional dyspnea and assume a squatting position to relieve the dyspneic episode after which the child is able to resume the activity.

Solution to Question 22:

The given chest x-ray shows an egg-on-string appearance seen in the transposition of great arteries (TGA). Enlargement of the left atrium is seen in TGA, not the right atrium.

The reason for left atrial enlargement is due to increased pulmonary blood flow.

TGA is characterized by the aorta arising from the right ventricle and the pulmonary artery from the left ventricle. This abnormal configuration leads to a narrow vascular pedicle.

As pulmonary vascular resistance drops during the first few days of life, blood flow through the pulmonary vasculature is increased. There is also elevated pulmonary venous pressure seen in TGA, especially in infants with an intact interventricular septum.

The globular (egg) appearance of the heart due to abnormal convexity of the right atrial border and left atrial enlargement, and the narrow superior mediastinum (string) result in the classical egg-on-string appearance.

TGA is the most specific heart disease in infants of diabetic mothers.

Early management includes an immediate infusion of prostaglandin E₁ which keeps the ductus arteriosus patent for improving oxygenation. Rashkind balloon atrial septostomy is a palliative catheter-based intervention that can be done to improve the hypoxic state before definitive surgery. The procedure creates an opening in the atrial septum which allows better inter-circulatory mixing of the blood. Jatene arterial switch procedure is the definitive surgery.

Most infants with TGA do not survive till adulthood without corrective surgery.

Solution to Question 23:

The classic "egg on end" appearance on this chest X-ray image is seen in the transposition of great arteries (TGA).

In the chest X-ray image, cardiomegaly with narrow mediastinum gives rise to the classic "egg-shaped heart." The superior mediastinum appears narrow due to the anteroposterior relationship of the transposed great vessels and the radiologic absence of the thymus.

Pediatric Nephrology

Question 1:

Hematuria in children is defined as the presence of at least _____ RBCs per high power field in uncentrifuged urine.

- a) 3
- b) 4
- c) 5
- d) 6

Question 2:

An 11-year-old child was brought to the hospital by his mother with complaints of multiple episodes of passing blood in the urine. Which of the following condition is not associated with this symptom?

- a) Berger's disease
- b) Hemolytic uremic syndrome
- c) Hereditary nephritis
- d) Idiopathic hypercalciuria

Question 3:

A 6-year-old girl with urinary complaints was brought to the hospital by her parents. On further enquiry, they mention a history of skin lesions 5 weeks back. Low C3 levels are found during her investigations. Immunofluorescence microscopy reveals a "lumpy-bumpy" pattern of immunoglobulin and complement deposits on the glomerular basement membrane. Which of the following is not typically associated with this condition?

- a) Cola colored urine
- b) Frothy urine
- c) Oliguria
- d) Hypertension

Question 4:

Akshay, accompanied by his father, presents to the OPD with complaints of cola-coloured urine and facial puffiness for 3 days. He has a history of fever with a sore throat that resolved 2 weeks back. Which of the following findings is consistent with the diagnosis of post-streptococcal glomerulonephritis?

- a) C3 level increases and ASO level increases
- b) C3 level decreases and ASO level increases
- c) C3 level decreases and ASO level decreases
- d) C3 level increases and ASO level decreases

Question 5:

Which of the following is not a characteristic feature of hematuria associated with glomerulonephritis?

- a) Dysmorphic RBC in urine
- b) RBC cast in urine
- c) Painful hematuria
- d) Cola-colored urine

Question 6:

A 7-year-old boy presents with decreased urine output for 3 days. He has a history of sore throat 2 weeks back. On examination, he is hypertensive with pitting pedal oedema. The diagnosis is concluded as post-streptococcal glomerulonephritis. Which of the following is true about this condition?

- a) Serum triglyceride levels are elevated.
- b) Microscopic hematuria resolves in 4 weeks.
- c) Serum C3 levels are normal.
- d) Renal biopsy is indicated in severe impairment of renal function.

Question 7:

A 6-year-old boy with a previously diagnosed renal disorder is brought to your OPD with the following reports for a second opinion. What is the likely diagnosis?

- a) Post-streptococcal glomerulonephritis
- b) Rapidly progressive glomerulonephritis
- c) Focal segmental glomerulonephritis
- d) Minimal change disease

Question 8:

An 8-year-old boy presented with cola-coloured urine, decreased urine output, and facial puffiness for 3 days. He had a history of skin infection 4 weeks back. On examination, he is found to be hypertensive. His serum C3 levels are low and ASO levels are increased. What is the appropriate step in the management of this patient?

- a) Renal biopsy
- b) Supportive therapy
- c) Steroids
- d) Cyclosporine

Question 9:

Sarthak, a 7-year-old boy, was brought to the emergency department by his parents with complaints of blurred vision and severe headache since today morning. They mentioned an episode of fits 2 hours back following which they rushed him to the hospital. On further enquiry, it was elicited that the child had decreased urine output for 5 days with cola coloured urine. He was hypertensive with bilateral pitting pedal oedema. The doctor remembered that Sarthak has a glomerular disorder warranting him multiple prior ward admissions and his existing condition could be a possible complication of the same. All of the following are commonly seen as a complication of this disorder except-

- a) Left ventricular failure
- b) Hypertensive encephalopathy
- c) Hyperkalemia
- d) Bleeding diathesis

Question 10:

A 9-year-old boy presents with gross hematuria 2 days after an episode of diarrhoea. The serum C3 levels are normal. Light microscopy demonstrates focal and segmental mesangial proliferation and increased mesangial matrix in the glomerulus. What is the probable

diagnosis?

- a) Berger's disease
- b) PSGN
- c) Microangiopathic hemolytic anemia
- d) None of the above

Question 11:

A child developed dark coloured urine and flank pain 2 days after an upper respiratory tract infection. The physician suspects IgA nephropathy. What is the location of cellular injury in this condition?

- a) Epithelial cell
- b) Basement membrane
- c) Endothelial cell
- d) Mesangial cell

Question 12:

ANCA-positive , pauci-immune glomerulonephritis include-

- a) Wegener`s granulomatosis
- b) Microscopic polyangiitis
- c) Churg-Strauss syndrome
- d) All of the above

Question 13:

An 8-year-old girl presented with complaints of cough, hemoptysis and swelling of both legs. Urinalysis was notable for hematuria and proteinuria. Serum anti-GBM antibodies were positive. What is the most likely diagnosis?

- a) IgA nephropathy
- b) Minimal change disease
- c) Goodpasture disease
- d) Membranous nephropathy

Question 14:

A 9-year-old boy has swelling of knee joints and non-blanching rashes as seen in the picture. Urine examination reveals hematuria +++ and proteinuria +. Microscopic analysis of his renal biopsy specimen is most likely to show which of the following finding?



- a) Tubular necrosis
- b) Visceral podocyte fusion
- c) Mesangial deposits of IgA
- d) Basement membrane thickening

Question 15:

All are features of Alport syndrome except-

- a) Anterior lenticonus
- b) Conductive hearing loss
- c) Gross hematuria
- d) Proteinuria

Question 16:

A 16-year-old boy presented with dark coloured urine. Urinalysis was notable for macroscopic hematuria and heavy proteinuria. His father is a known case of Alport syndrome. What defect is responsible for this condition?

- a) Fibrillin
- b) Type IV collagen
- c) Amyloid
- d) Tau

Question 17:

All of the following syndromes are associated with renal disease except _____.

- a) Lowe syndrome
- b) Denys-Drash syndrome
- c) Alport syndrome
- d) Sinding-Larsen-Johansson syndrome

Question 18:

Microalbuminuria is defined as a urinary albumin excretion rate of _____.

- a) 30-300 mg/ 24 hrs
- b) $\geq$ 300 mg/ 24 hrs
- c) $\geq$ 150 mg/ 24 hrs
- d) $\geq$ 300 μ g/ 24 hrs

Question 19:

A 5-year-old boy presents with facial puffiness and frothy urine. Upon investigating, his urine was positive for proteins with no hematuria and no RBC casts. To conclude the diagnosis of nephrotic syndrome, the range of proteinuria is defined as $\geq$ _____ mg/m²/hr and urine protein:creatinine ratio of $\geq$ _____.

- a) 40 and 2
- b) 30 and 1.5
- c) 3.5 and 2
- d) 4 and 1.5

Question 20:

It is expected to see all of the following clinical features in a 5-year-old boy with nephrotic syndrome except-

- a) Hypoalbuminemia
- b) Hyperlipidemia
- c) Hematuria
- d) Edema

Question 21:

What is the most common cause of nephrotic syndrome in children?

- a) Minimal change disease
- b) Membranous nephropathy
- c) IgA nephropathy
- d) Alport syndrome

Question 22:

A 5-year-old child was brought to the hospital with complaints of progressive swelling around the eyes, abdomen and feet for 2 weeks. On investigating, his urine was positive for 4+ proteinuria with absent hematuria. A spot urine protein: creatinine ratio was reported to be 2.5. His serum albumin was 2.3 g/dL and serum creatinine was 0.5 mg/dL. Serum cholesterol and triglyceride levels were also elevated. A diagnosis of nephrotic syndrome was concluded. Which of the following correctly mentions the aetiology of its most common variant in children?

- a) Flattening of podocytes
- b) Immune complex deposition
- c) Anti-GBM antibodies
- d) Endothelial cell disease

Question 23:

Which of the following is not a cause of hypercoagulability in nephrotic syndrome?

- a) Vascular stasis due to hemoconcentration and intravascular volume depletion
- b) Increase in hepatic production of fibrinogen

- c) Urinary loss of antithrombin III and protein S
- d) Decreased platelet number and aggregability

Question 24:

A 4-year-old boy presents with facial oedema that has now progressed to total body swelling involving the face, abdomen, scrotum, and feet. Urinalysis revealed protein 4+ and serum cholesterol was also elevated. What is the initial treatment of choice?

- a) Prednisolone
- b) Cyclophosphamide
- c) Levamisole
- d) Cyclosporine

Question 25:

Steroid-resistant nephrotic syndrome is defined as failure to achieve remission after _____ weeks of corticosteroid therapy.

- a) 12
- b) 8
- c) 6
- d) 4

Question 26:

A child suffering from steroid-dependent nephrotic syndrome is brought to the OPD for vaccination. Which of the following is true about vaccination in this child?

- a) All live vaccines under NIS can be given as per schedule
- b) All killed vaccines under NIS can be given
- c) Only pneumococcal vaccine should be given to the child in acute phase
- d) Live vaccines are contraindicated in the child's sibling

Question 27:

Congenital nephrotic syndrome is defined as nephrotic syndrome manifesting _____.

- a) At birth
- b) At birth or within 3 months of life
- c) At birth or within 6 months of life
- d) At birth or within 1 year of life

Question 28:

An 11-year-old boy with a chronic hepatitis B infection develops nephrotic syndrome. What would be the most likely pathology encountered in renal biopsy of this child?

- a) Focal Segmental Glomerular Sclerosis
- b) Membrano proliferative Glomerulonephritis
- c) Anti GBM disease
- d) Membranous nephropathy

Question 29:

Finnish-type congenital nephrotic syndrome is caused by a mutation in the _____ gene.

- a) NPHS 1 (nephrin)
- b) NPHS 2 (podocin)
- c) WT 1 (Wilms tumor suppressor gene)
- d) LMX1B

Question 30:

FSGS is caused by mutations in the _____ gene.

- a) NPHS 1 (nephrin)
- b) NPHS 2 (podocin)
- c) WT 1 (Wilms tumor suppressor gene)
- d) LMX1B

Answer Key

Question No.	Correct Option
1	c
2	b
3	b
4	b
5	c
6	d
7	a
8	b
9	d
10	a
11	d
12	d
13	c
14	c
15	b
16	b
17	d
18	a
19	a
20	c
21	a
22	a
23	d
24	a
25	c
26	b
27	b
28	d
29	a
30	b

Detailed Explanations

Solution to Question 1:

Hematuria in children is defined as the presence of at least 5 RBCs/HPF (high power field) in uncentrifuged urine.

When using commercially available dipsticks (capable of detecting 3-5 RBCs / HPF of urine), the presence of 10-50 RBCs/ μ L suggests an underlying pathology.

False-negative results can occur in the presence of

- Formalin
- High urinary concentrations of ascorbic acid

False-positive results, on the other hand, may be seen

- With alkaline urine (pH $\geq$ 8)
- Contamination with oxidizing agents such as hydrogen peroxide (used to clean the perineum before obtaining a specimen)

Solution to Question 2:

Hemolytic uremic syndrome (HUS) presents with acute-onset microscopic hematuria not recurrent gross hematuria.

Causes of recurrent gross hematuria:

- Alport's syndrome: hereditary nephritis
- Thin glomerular basement membrane disease
- IgA nephropathy: Berger's disease
- Idiopathic hypercalciuria/hyperuricosuria
- Sickle cell nephropathy (papillary necrosis)

Solution to Question 3:

Urinary complaints with a history of skin lesions (pyoderma) point towards a likely diagnosis of glomerulonephritis. Results of complement study and immunofluorescence microscopy also correlate with this diagnosis. Frothy urine is due to proteinuria, which is a feature of nephrotic syndrome.

Classic features of nephritic syndrome/glomerulonephritis:

- Cola coloured urine (hematuria)
- Oliguria
- Hypertension
- Facial or body edema

Solution to Question 4:

C3 level decreases and ASO level increases in acute post-streptococcal glomerulonephritis (PSGN).

Acute PSGN is a classic example of an acute nephritic syndrome characterized by the sudden onset of gross hematuria, oedema, hypertension, and renal insufficiency. It follows infection of the throat or skin by certain nephritogenic strains of group A β -hemolytic streptococci-

- 1 to 2 weeks after an antecedent streptococcal pharyngitis or
- 3 to 6 weeks after streptococcal pyoderma

Solution to Question 5:

Hematuria associated with glomerulonephritis is typically painless.

Hematuria from within the glomerulus is often associated with brown-, cola-, or tea-colored urine. Urine analysis reveals RBC casts and deformed urinary RBCs (particularly acanthocytes).

Solution to Question 6:

The given clinical vignette of oliguria, pedal edema and hypertension point towards glomerulonephritis. Severe impairment of renal functions in a background of post-streptococcal glomerulonephritis is an indication for renal biopsy.

Indications for renal biopsy in glomerulonephritis:

- Presence of acute renal failure or nephrotic syndrome
- Absence of evidence of streptococcal infection
- Presence of normal complement levels
- Persistence of hematuria and proteinuria, diminished renal function, and/or a low C3 level for more than 2 months after the disease onset

Serum triglycerides (Option A) are elevated in nephrotic syndrome, not in nephritic syndrome.

The acute phase generally resolves within 6 to 8 weeks. Although urinary protein excretion and hypertension usually normalize by 4-6 wk after onset, microscopic hematuria (Option B) can persist for 1 to 2 years after the initial presentation.

The serum C3 level (Option C) is significantly reduced in >90% of patients in the acute phase, and returns to normal 6-8 weeks after onset.


Solution to Question 7:

The findings mentioned are characteristic of post-streptococcal glomerulonephritis (PSGN).

Findings in PSGN:

- Light microscopy: Enlarged, pale glomeruli with diffuse mesangial proliferation and an increase in the mesangial matrix. Polymorphonuclear (PMN) infiltration in the early stages.
- Immunofluorescence: “Lumpy-bumpy” deposits of immunoglobulin and complement on the glomerular basement membrane (GBM) and in the mesangium.
- Electron microscopy: Electron-dense deposits, or “humps,” are observed on the epithelial side of the GBM.

The image given below depicts the microscopic picture of PSGN. It shows glomerular hypercellularity due to intracapillary leukocytes. Neutrophilic infiltration is present and is marked with purple hematoxylin stain (black arrows).



Other options:

Idiopathic rapidly progressive glomerulonephritis:

- Light microscopy: Crescents develop in several forms of glomerulonephritis (are a characteristic response to the deposition of fibrin in Bowman's space)
- Electron microscopy: No deposits

Focal segmental glomerulonephritis:

- Light microscopy: Focal sclerotic lesions
- Electron microscopy: Foot process fusion

Minimal change disease:

- Light microscopy: Normal
- Electron microscopy Effacement of foot process

Solution to Question 8:

The features of hematuria, oliguria, facial oedema present with a history of skin infection points towards the diagnosis of post-streptococcal glomerulonephritis (PSGN). Results of complement study and ASO levels also support this diagnosis. Supportive conservative therapy is the best step in the management of PSGN.

Management of PSGN is directed at treating the acute effects of renal insufficiency (sodium restriction and prescribing diuretics - usually with intravenous furosemide) and hypertension.

Although a 10-day course of systemic antibiotic therapy with penicillin is recommended to limit the spread of the nephritogenic organisms, antibiotic therapy does not affect the natural history of acute PSGN.

Note: Steroids and cyclosporine are not used in the management of PSGN.

Solution to Question 9:

The given features of oliguria, hematuria, pedal edema and hypertension are likely of glomerulonephritis. The complication of hypertensive encephalopathy is seen in this case due to symptoms of seizure, blurred vision and severe headache. Hyperkalemia and left ventricular failure are also common complications of glomerulonephritis while bleeding diathesis is not a complication of glomerulonephritis.

Acute complications of glomerulonephritis result from hypertension and acute renal dysfunction-

- Hypertension is seen in 60% of patients and is sometimes associated with hypertensive encephalopathy and intracranial bleeding.
- Acute renal failure can occur, which may require treatment with dialysis.

Other potential complications include the following:

- Heart failure
- Hyperkalemia
- Hyperphosphatemia
- Hypocalcemia
- Hyponatremia
- Acidosis
- Seizures
- Uremia

Solution to Question 10:

Gross hematuria following a gastrointestinal infection, associated with normal C3 levels, in a male patient is suggestive of immunoglobulin A nephropathy (Berger nephropathy).

- IgA nephropathy is the most common chronic glomerular disease in children.

- Focal and segmental mesangial proliferation and increased mesangial matrix are seen in the glomerulus.
- It is characterized by a predominance of IgA within mesangial glomerular deposits in the absence of systemic disease.
- It is seen more often in males.
- Gross hematuria occurs within 1-2 days of onset of an upper respiratory or gastrointestinal infection (in contrast to the long latency period observed in acute post-streptococcal glomerulonephritis).
- It can be associated with loin pain.
- Normal serum levels of C3 in IgA nephropathy (help to distinguish this disorder from PSGN).
- Serum IgA levels have no diagnostic value because they are elevated in only a few paediatric patients.

Solution to Question 11:

The location of cellular injury in IgA nephropathy (Berger nephropathy) is the deposition of IgA antibodies in the mesangial cell.

Mesangial cells are directly exposed to the circulation; therefore, they are affected in diseases characterized by the deposition of immune complexes.

Mesangial cell diseases:

- IgA nephropathy
- IgM nephropathy
- Mesangioproliferative glomerulonephritis
- Class II lupus nephritis
- Diabetic nephropathy

Note: In IgA nephropathy gross hematuria often occurs within 1-2 days of onset of an upper respiratory or gastrointestinal infection. In contrast, a longer latency period of about 1-2 weeks is observed in acute postinfectious glomerulonephritis.

Solution to Question 12:

Anti-neutrophil cytoplasmic antibodies (ANCA)-associated vasculitis is categorized into 3 distinct forms:

- Granulomatosis with polyangiitis (GPA; formerly Wegener granulomatosis)
- Microscopic polyangiitis (MPA)
- Eosinophilic granulomatosis with polyangiitis, formerly called Churg–Strauss syndrome (CSS)

Kidney biopsies typically demonstrate crescentic glomerulonephritis with little or no immune complex deposition.

ANCA-associated vasculitis (AAV) comprises a group of multi-systemic diseases affecting small-to-medium-sized vessels. It is characterized by the presence of ANCA with specificity for either proteinase-3 (PR3) or myeloperoxidase (MPO).

Absence or paucity of immune complex deposits in vessel walls is very characteristic, hence the name "pauci-immune vasculitis."

Solution to Question 13:

The diagnosis is Goodpasture disease (anti-GBM antibody disease). Anti-glomerular basement membrane (anti-GBM) antibodies are diagnostic.

"Goodpasture syndrome" is characterized by pulmonary haemorrhage and glomerulonephritis and "Goodpasture disease" is used for those patients with glomerulonephritis, pulmonary hemorrhage, and anti-GBM antibodies. It is caused due to autoantibodies directed against type IV collagen, which is present in the alveoli and glomerular basement membrane. Currently, the preferred term for both conditions is "anti-GBM antibody disease".

Other options along with the pathogenesis of glomerular diseases:

- IgA nephropathy- mesangial cell disease due to immune complex deposition
- Minimal change disease- epithelial cell disease due to flattening of podocytes

Solution to Question 14:

The boy in this clinical scenario has characteristic features of Henoch–Schonlein purpura (HSP).

The renal biopsy in this patient will show mesangial proliferation with mesangial deposition of IgA

- HSP is a common small-vessel vasculitis in childhood.
- It is characterized by a purpuric rash and commonly accompanied by arthritis and abdominal pain.
- About half the patients with HSP develop renal manifestations, which vary from asymptomatic microscopic hematuria to severe, progressive glomerulonephritis.
- The nephritis associated with HSP usually follows the onset of the rash
- The pathogenesis of HSP nephritis appears to be mediated by the deposition of polymeric immunoglobulin A (IgA) in glomeruli.

Solution to Question 15:

Bilateral sensorineural hearing loss (not conductive hearing loss) is a feature of Alport syndrome (AS).

AS or hereditary nephritis is a genetic disease (most have X-linked inheritance) caused by mutations in the genes coding for type IV collagen, a major component of the basement membranes.

Clinical features of Alport syndrome:

- Hematuria: All patients have asymptomatic microscopic hematuria. Recurrent episodes of gross hematuria may be seen 1-2 days after an upper respiratory infection.
- Bilateral sensorineural hearing loss (not conductive)
- Ocular abnormalities: anterior lenticonus, macular flecks, corneal erosions
- Proteinuria: Progressive proteinuria, often exceeding 1 g/ 24 hrs, is common by the 2nd decade of life.

Solution to Question 16:

A defect in type IV collagen, a major component of the basement membrane, is responsible for Alport syndrome.

Alport syndrome is a genetically heterogeneous disease, 85% is X-linked and the rest may be autosomal recessive or autosomal dominant.

Solution to Question 17:

Sinding–Larsen–Johansson syndrome is an insertional periostitis at the inferior pole of the patella (not associated with Renal disease).

Lowe syndrome	Oculocerebrorenal syndrome: Hypotonia, renal disease
Denys–Drash Syndrome	Associated with Wilm's tumor: Early-onset renal failure with renal mesangial sclerosis, male pseudohermaphroditism
Alport syndrome	Bilateral sensorineural hearing loss, renal disease
Sinding–Larsen–Johansson syndrome	Periostitis at the inferior pole of the patella

Solution to Question 18:

Microalbuminuria is defined as a urinary albumin excretion rate of 30–300 mg/24 hrs.

Here, the presence of albumin in the urine is above the normal level but below the detectable range for the urine-dipstick methods. It is commonly associated with glomerular hyperfiltration and blood pressure elevation.

In adults, persistent microalbuminuria indicates diabetic nephropathy and is also a predictor of cardiovascular and renal disease.

In children, microalbuminuria is associated with obesity and predicts the development of diabetic nephropathy in type 1 diabetes mellitus.

Solution to Question 19:

The currently accepted definitions of “nephrotic range proteinuria” include:

- Urine protein: creatinine ratio ≥ 2
- Urine protein ≥ 40 mg/m²/hr
- Urine protein 3+ or 4+ in dipstick test

As a consequence of the large urinary loss of protein, the patient develops hypoalbuminemia (< 2.5 g/dL) and gross edema.

Solution to Question 20:

Hematuria is not a usual feature of nephrotic syndrome. It is a feature of nephritic syndrome. However, occasionally it is seen in the FSGS variant of nephrotic syndrome as well.

Classic features of nephrotic syndrome:

- Nephrotic range proteinuria (≥ 40 mg/m²/hr urine protein: creatinine ratio ≥ 2)
- Hypoalbuminemia (< 2.5 g/dL)
- Edema
- Hyperlipidemia (cholesterol ≥ 200 mg/dL)
- Hypercoagulability

Solution to Question 21:

The commonest cause of nephrotic syndrome in children is minimal change disease. (85% of cases).

The glomeruli appear almost normal on microscopy, hence the name. Immunofluorescence microscopy are negative, and electron microscopy shows effacement of the epithelial cell foot processes.


Treatment: Steroids; 95% of patients respond.

Other options:

- Membranous nephropathy is the most common cause of nephrotic syndrome in the elderly; it is a rare cause of nephrotic syndrome in children.
- Alport syndrome or hereditary nephritis: All patients have asymptomatic hematuria; progressive proteinuria is common only by the 2nd decade of life and can be severe enough to cause nephrotic syndrome. This nephrotic syndrome is not common in children.
- IgA nephropathy presents more commonly as an acute nephritic syndrome or with asymptomatic hematuria. It presents with nephrotic syndrome quite rarely.

Solution to Question 22:

Nephrotic syndrome is concluded as the diagnosis here because of the symptoms of anasarca and proteinuria. A spot urine protein:creatinine ratio >2 , low serum albumin levels, and hyperlipidemia marked by high cholesterol and triglyceride levels, further add to this diagnosis. Minimal change disease is the commonest cause of nephrotic syndrome in children and constitutes 85% of all paediatric cases. The etiology is due to the flattening of foot processes (podocytes) of epithelial cells (seen on electron microscopy). This finding is marked in the image given below.



- Here, the glomeruli appear normal or show a minimal increase in mesangial cells and matrix.
- It is the most common cause of nephrotic syndrome in children.
- Epithelial cells, in conjunction with the basement membrane, allow the filtration of plasma solutes but block the passage of cells and plasma proteins. Hence, this epithelial cell disease, disrupts the filtration barrier, resulting in proteinuria.
- More than 95% of children with minimal change disease respond to corticosteroid therapy.

Causes of epithelial cell injury:

- Minimal change disease

- Membranous nephropathy
- Focal segmental glomerulosclerosis
- Class V lupus nephritis
- Diabetic nephropathy

Solution to Question 23:

Increased (and not decreased) platelet number and aggregability cause hypercoagulability in nephrotic syndrome.

Hypercoagulability in nephrotic syndrome is due to all of the following factors:

- Vascular stasis due to hemoconcentration and intravascular volume depletion
- Increase in hepatic production of fibrinogen
- Urinary loss of antithrombin III and protein S
- Increased platelet number and aggregability

Solution to Question 24:

The clinical picture and investigations are highly suggestive of minimal change nephrotic syndrome and the initial treatment of choice is corticosteroids such as prednisolone.

Minimal change nephrotic syndrome is the most common cause of nephrotic syndrome. It is seen in ages 2-6 years with a clinical picture of nephrotic syndrome, that is, heavy proteinuria, oedema, hypoalbuminemia and hyperlipidemia. They may be asymptomatic or can present with peripheral oedema. Renal biopsy is not required for confirmation and treatment can be initiated if the child fits the standard clinical picture of nephrotic syndrome.

They show excellent response to steroid therapy with prednisone or prednisolone. A daily dose of 2mg/kg (max 60mg) for 6 weeks followed by an alternate dose of 1.5mg/kg (max 40mg) for the next 6-8 weeks is recommended.

Alternative drugs used in steroid-dependent patients, frequent relapsers, steroid-resistant patients, and those with severe corticosteroid toxicity are as follows:

- Cyclophosphamide
- Calcineurin inhibitors: Cyclosporine and Tacrolimus
- Mycophenolate
- Levamisole
- Rituximab (chimeric monoclonal antibody against CD20)

Solution to Question 25:

Steroid-resistant nephrotic syndrome is defined as failure to achieve remission after 6 weeks of corticosteroid therapy.

According to the latest Indian Society of Pediatric nephrology(ISPN) guidelines, it is defined as the absence of remission despite 6-weeks therapy with prednisolone at daily dose of 60mg/m²

Solution to Question 26:

In a child with steroid-dependent nephrotic syndrome, all killed vaccines can be given.

Option A: Live virus vaccines are administered with caution in children receiving corticosteroid therapy. They can be given to patients in whom the prednisone dose is below 1mg/kg daily or 2mg/kg on alternate days. If the patient is receiving corticosteroid-sparing agents (cyclophosphamide, cyclosporine), live virus vaccines are contraindicated.

Option C: Only live vaccines are contraindicated in this child. Full pneumococcal vaccination (with the 13-valent conjugate vaccine and 23-valent polysaccharide vaccine) and influenza vaccination can be given annually to the child and their household contacts.

Option D: Immunization of siblings/healthy household contacts with live vaccines is necessary to reduce the risk of infection transmission to the immunosuppressed child. Direct exposure to the gastrointestinal/respiratory secretions of the household contacts is avoided for about 3-6 weeks after vaccination.

Minimal change nephrotic syndrome is the most common cause of nephrotic syndrome. It is characterized by heavy proteinuria, edema, hypoalbuminemia, and hyperlipidemia. The diagnosis is confirmed by performing a urinalysis and urine protein: creatinine ratio. Nephrotic range proteinuria is proteinuria greater than 3.5 g per 24 hours (3+ or 4+) or a urine protein : creatinine ratio exceeding 2.

The recommended initial treatment for nephrotic syndrome is a daily dose of 2mg/kg (max 60mg) of prednisolone for 6 weeks followed by an alternate dose of 1.5mg/kg (max 40mg) for the next 6-8 weeks.

Definitions regarding the course of nephrotic syndrome:

- Frequently relapsing nephrotic syndrome: ≥ 2 relapses in 6 months or 3 or more relapses in 12 months.
- Steroid-dependent nephrotic syndrome: Relapse during steroid tapering or within 2 weeks of discontinuation.
- Steroid resistance: Failure to achieve remission despite therapy with daily prednisolone for 6 weeks

Alternative drugs used in steroid-dependent patients, frequent relapsers, steroid-resistant patients, and those with severe corticosteroid toxicity are cyclophosphamide, calcineurin inhibitors: cyclosporine and Tacrolimus, mycophenolate, levamisole, and rituximab (a chimeric monoclonal antibody against CD20).

Solution to Question 27:

Congenital nephrotic syndrome is defined as nephrotic syndrome manifesting at birth or within 3 months of life.

It may be

- Primary: Due to inherited disorders caused by mutations in NPHS 1, NPHS 2, WT1, or LAMB2.
- Secondary: Due to in utero infections like cytomegalovirus infection, toxoplasmosis, syphilis, HIV infection, hepatitis B virus or hepatitis C virus HCV infection, infantile systemic lupus erythematosus (SLE), or mercury exposure.

Solution to Question 28:

The given clinical scenario suggests that the pathology would be membranous nephropathy. Membranous nephropathy can be secondarily caused by Hepatitis B infection.

Membranous nephropathy:

Membranous nephropathy is a rare cause of nephrotic syndrome in children.

- It is classified as the primary, idiopathic form, where there is an isolated renal disease, or secondary, where nephropathy is associated with other identifiable systemic diseases or medications
- In children, secondary form is far more common. The most common etiologies of secondary form are systemic lupus erythematosus or chronic infections. Among the latter, chronic hepatitis B infection and congenital syphilis are the best characterized and recognized causes.

Solution to Question 29:

Finnish type congenital nephrotic syndrome is a steroid-resistant nephrotic syndrome caused by a mutation in the nephrin gene (NPHS 1).

- Affected infants usually present at birth with edema caused by massive proteinuria.
- They are delivered with a large placenta (>25% of the infant's weight).
- There is severe hypoalbuminemia, hyperlipidemia, and hypogammaglobulinemia.
- Prenatal diagnosis can be made by elevated maternal and amniotic α -fetoprotein levels.

Gene	Name	Inheritance	Renal disease
NPHS1	Nephrin	Recessive	Finnish-type congenital nephrotic syndrome
NPHS2	Podocin	Recessive	Focal segmental glomerulosclerosis (FSGS)
WT1	Wilms tumor-suppressor gene	Dominant	Denys–Drash syndrome with diffuse mesangial sclerosis Frasier syndrome with FSGS

Gene	Name	Inheritance	Renal disease
LMX1B	LIM-homeodomain protein	Dominant	Nail–patella syndrome

Solution to Question 30:

- focal segmental glomerulosclerosis (FSGS) is caused by mutation in NPHS2 (podocin) gene.
- Steroid resistant nephrotic syndrome is usually caused by FSGS (in 80% of cases).
- In FSGS, glomeruli show lesions of mesangial cell proliferation and segmental scarring that are both focal (present only in a proportion of glomeruli) and segmental (localized to ≥ 1 intraglomerular tufts).
- FSGS is associated with a 50% risk for end-stage kidney disease.

Gene	Name	Inheritance	Renal disease
NPHS1	Nephrin	Recessive	Finnish-type congenital nephrotic syndrome
NPHS2	Podocin	Recessive	Focal segmental glomerulosclerosis (FSGS)
WT1	Wilms tumor-suppressor gene	Dominant	Denys–Drash syndrome with diffuse mesangial sclerosis Frasier syndrome with FSGS
LMX1B	LIM-homeodomain protein	Dominant	Nail–patella syndrome

Pediatric Urology

Question 1:

During her antenatal ward rounds, an intern noticed a USG report indicating the presence of just one kidney in the fetus. Which of the following association would be expected to be seen in the baby?

- a) Hypogonadism
- b) Presence of single umbilical vein
- c) Polycystic kidney
- d) Presence of single umbilical artery

Question 2:

A 10-month-old female is brought to the clinic with a 102° F fever for 2 days. The mother says that she also cries during urination. The history is significant for similar episodes in the past which were treated with antibiotics. Urine analysis shows nitrite-positivity and gram-negative bacteria are seen on urine microscopy. Which of the following anomaly is most commonly associated with this child's condition?

- a) Posterior urethral valves
- b) Vesicoureteric reflux
- c) Horseshoe kidney
- d) Pelviureteric junction obstruction

Question 3:

A pediatrician suspects vesicoureteric reflux in a girl with a history of recurrent urinary tract infections. What is the investigation of choice to confirm this condition?

- a) Ultrasound abdomen
- b) MRI
- c) Voiding cystourethrogram
- d) DMSA scan

Question 4:

Which of the following is the first-line imaging investigation done in a child presenting with an episode of UTI?

- a) Voiding cystourethrogram
- b) Micturating cystogram
- c) Ultrasonogram
- d) DMSA scan

Question 5:

Which of the following is the most common cause of urinary tract infection in children?

- a) E. coli
- b) Staphylococcus
- c) Proteus
- d) Klebsiella

Question 6:

A mother is concerned that her 1-year-old daughter has been having repeated episodes of UTI since birth. The pediatrician ordered a voiding cystourethrogram, which revealed reflux of urine into the upper collecting systems without any dilation. Under which grade of vesicoureteral reflux will this presentation be classified?

- a) Grade III
- b) Grade II
- c) Grade I
- d) Grade IV

Question 7:

Ashi, a 12-year-old girl presented with features of hematuria, headache, and blurring of vision. Her past history is significant for recurrent UTI. On examination, she was found to have a BP of 150/110 mmHg. Investigations revealed mild renal scarring and urine analysis showed broad casts. Which of the following is the most likely diagnosis?

- a) Acute tubular necrosis

- b) Congenital ureteropelvic junction obstruction
- c) Posterior urethral valve with chronic renal failure
- d) Reflux uropathy with chronic renal failure


Question 8:

Which of the following is the most common cause of abdominal mass in neonates?

- a) Neuroblastoma
- b) Wilms tumor
- c) Multicystic dysplastic kidney
- d) Ectopic kidney

Question 9:

Komal, a high school student developed a fever after she came back from a trip with her friends. She also complained about pain while urinating. She says she has been having these episodes frequently for the past few years. Her father and her grandfather are a known case of cystic kidney disease. Her urine analysis is indicative of UTI with hematuria. Renal ultrasound was done and its findings are shown below. Which of the following chromosomes are associated with this condition?



- a) 14 and 6
- b) 8 and 14
- c) 4 and 16

d) 12 and 16

Question 10:

Which of the following conditions is not associated with the lesion seen in the image?



- a) Vesicoureteric reflux
- b) Posterior urethral valve
- c) Wilms tumor
- d) Multicystic dysplastic kidney

Question 11:

A young girl presents with the complaints of continuous dribbling of urine. However, she also voids normally. Her parents tell you that she had never been totally dry since her birth. What is the most likely diagnosis?

- a) Ectopic ureter
- b) Ureterocele
- c) Congenital megaureter
- d) Vesicoureteric reflux

Question 12:

A 3 yr old male child with left flank pain is diagnosed to have an acute UTI. On USG, the left ureter was found to be duplicated. The most probable site of the opening of ectopic ureter will be?

- a) Prostatic urethra
- b) Seminal vesicle
- c) Ejaculatory duct
- d) Trigone of bladder

Question 13:

A newborn is being evaluated for anuria. Imaging reveals renal agenesis and pulmonary hypoplasia. The delivery was complicated due to oligohydramnios. Which of the following is the most likely diagnosis?

- a) Potter sequence
- b) Pierre Robin syndrome
- c) Congenital diaphragmatic hernia
- d) CHARGE syndrome

Question 14:

Antenatal hydronephrosis due to PUJ obstruction is defined as the anteroposterior diameter of the renal pelvis of more than _____ in the third trimester.

- a) 6 mm
- b) 7 mm
- c) 11 mm
- d) 15 mm

Question 15:

Hepatic fibrosis is associated with _____.

- a) Medullary cystic kidney
- b) Acquired cystic kidney disease
- c) Autosomal dominant polycystic kidney disease
- d) Autosomal recessive polycystic kidney disease

Question 16:

Ask-Upmark kidney refers to _____

- a) Renal cystic dysplasia
- b) Renal segmental hypoplasia
- c) Renal hypodysplasia
- d) Renal medullary hyperplasia

Answer Key

Question No.	Correct Option
1	d
2	b
3	c
4	c
5	a
6	b
7	d
8	c
9	c
10	b
11	a
12	a
13	a
14	b
15	d
16	b

Detailed Explanations

Solution to Question 1:

The presence of a single umbilical artery can be expected to be seen in the case of a baby with unilateral renal agenesis.

Renal agenesis occurs secondary to a defect of the Wolffian duct, ureteric bud, or metanephric blastema. It is often discovered during the course of an evaluation for other congenital anomalies (VACTERL syndrome). Its incidence is increased in newborns with a single umbilical artery. The contralateral kidney undergoes compensatory hypertrophy. Some children also have contralateral vesicoureteral reflux. Most males have an ipsilateral absent vas deferens.

NOTE:


VACTERL syndrome - Vertebral defects, imperforate Anus, Congenital heart disease, Tracheo-Esophageal fistula, Renal and Limb defects

Solution to Question 2:

The given clinical scenario is suggestive of recurrent urinary tract infection. The most common anomaly associated with this condition in children is vesicoureteral reflux (VUR).

VUR usually is congenital and often familial. It is more common in females. It occurs when the submucosal tunnel between the mucosa and detrusor muscle is short or absent.

The image below shows VUR grading



Solution to Question 3:

Voiding cystourethrogram (VCUG) is the investigation of choice for vesicoureteral reflux.

Recurrent febrile urinary tract infections in a child should raise the suspicion of an anatomical defect like vesicoureteric reflux as the cause. Compared to other investigations, a VCUG provides more anatomic information. It can demonstrate:

- Vesicoureteral reflux
- Duplex collecting system
- Ectopic ureter
- Paraureteral (bladder) diverticulum
- Bladder outlet obstruction
- Signs of voiding dysfunction, such as a spinning top urethra

Solution to Question 4:

In a child presenting with an episode of urinary tract infection (UTI), initial imaging should be ultrasonography of the kidneys, ureters, and bladder.

A voiding cystourethrogram is indicated only if the ultrasound study is abnormal, the patient has atypical features or after a recurrent febrile UTI.

DMSA (dimercaptosuccinic acid) scan may be done after 4–6 months of acute infection as it may help to locate and delineate the scarring.

Solution to Question 5:

Urinary tract infections in children are most commonly caused by the following:

- Escherichia coli (Most common)
- Klebsiella species
- Proteus species
- Staphylococcus saprophyticus
- Enterococcus
- Adenovirus and other viral infections (cause cystitis and gross hematuria)

Solution to Question 6:


Vesicoureteral reflux (VUR) severity is graded according to the appearance of the urinary tract on a contrast voiding cystourethrogram.

It is classified into 5 grades (Grade I to V):

Grading of vesicoureteral reflux:

- Grade I- VUR into a non-dilated ureter
- Grade II- VUR into the upper collecting system without dilation
- Grade III- VUR into the dilated ureter and/or blunting of calyceal fornices.


- Grade IV- VUR into a grossly dilated ureter
- Grade V- Massive VUR, with significant ureteral dilation and tortuosity and loss of the papillary impression.



Solution to Question 7:

The given clinical scenario is suggestive of reflux uropathy with chronic renal failure.

The pathogenesis of vesicoureteral reflux is as follows:



Loss of renal parenchyma and renal scarring reduces renal function significantly. This renal insufficiency leads to renin-mediated hypertension and further progresses to end-stage renal disease.

Reflux nephropathy remains a common cause of hypertension in children.

Broad casts or renal failure casts are associated with end-stage renal disease.

Solution to Question 8:

Multicystic dysplastic kidney (MCDK) is the most common cause of an abdominal mass in the newborn but is mostly non-palpable at birth.

In most cases, it is discovered incidentally during prenatal sonography. In some patients, the cysts are identified prenatally, but the cysts regress in utero, and no kidney is identified on imaging at birth. MCDK is usually unilateral and generally not inherited, unlike polycystic kidney disease (PKD). Contralateral hydronephrosis is present in 5-10% of patients.

Solution to Question 9:

The presence of multiple cysts seen on the ultrasound (image) along with a positive family history point towards the diagnosis of autosomal-dominant polycystic kidney disease. Chromosomes 16 and 4 are associated with this condition.

Autosomal-dominant polycystic kidney disease inheritance:

Chromosome involved	Chr16 short arm	Chr 4 long arm
Gene	PKD 1	PKD 2
Protein encoded	Polycystin 1	Polycystin 2
Function	Transmembrane glycoprotein	Non-selective cation channel
Seen in	85 %	10–15 %

Solution to Question 10:

The given image shows a horseshoe kidney. It is not associated with the posterior urethral valve.

Horseshoe kidney is the most common congenital anomaly of the kidneys.

Horseshoe kidney is caused due to the fusion of the lower poles of the kidney. The lower border of the kidneys usually lie in front of L3 vertebra due to arrest of ascent.

The common associations of horseshoe kidney are:

- Vesicoureteral reflux
- Wilms tumor
- Multicystic dysplastic kidney
- Turner's syndrome

Solution to Question 11:

The clinical presentation of a young girl with continuous dribbling of urine, despite normal voiding, strongly suggests the diagnosis of an ectopic ureter.

An ectopic ureter refers to a condition where the ureter drains outside the bladder. It is more common in females and opens either into the urethra below the sphincter or into the vagina. Hence, they present with constant urinary dripping all day, even though the child voids regularly (paradoxical incontinence).

The terminal part of the ureter is frequently narrowed, leading to hydroureteronephrosis. Urinary tract infections (UTIs) are common due to urinary stasis.

In males, ectopic ureters usually enter the posterior urethra above the external sphincter, hence they are usually continent and most patients present with UTI or epididymitis.

Investigations include renal ultrasound (USG), voiding cystourethrogram (VCUG), and renal scan to assess function. USG shows a hydronephrotic kidney or dilated upper pole and ureter. If there is a satisfactory function, the recommended treatment options are ureteral reimplantation into the bladder or ureteroureterostomy. The poor function may require partial or total nephrectomy.

Other options:

Option B: A ureterocele is a cystic dilation of the terminal ureter, typically obstructive due to a small ureteral opening. In childhood, they commonly present with infections. Larger ureteroceles can cause obstruction at the bladder neck.

Option C: Congenital megaureter (dilated ureter) presents UTI, urinary stones, hematuria, and flank pain because of urinary stasis.

Option D: Vesicoureteral reflux (VUR) is characterized by the retrograde flow of urine from the bladder to the ureter and kidney. They usually present with recurrent attacks of UTI.

Solution to Question 12:

Opening of the ectopic ureter in male (most to least common) :


- Posterior urethra (prostatic) above the external sphincter
- Seminal vesicle
- Prostatic utricle
- Ejaculatory duct
- Vas deferens

Solution to Question 13:

A newborn infant with anuria, renal agenesis, and pulmonary hypoplasia, with a history of oligohydramnios, is suggestive of Potter's sequence

Potter's sequence is characterized by the triad of:

- Renal agenesis
- Oligohydramnios
- Pulmonary hypoplasia



Option B: Pierre–Robin syndrome: micrognathia, retrognathia, glossoptosis.

Option C: Congenital diaphragmatic hernia: It can present with pulmonary hypoplasia but other features like renal agenesis, anuria will not be seen.

Option D: CHARGE syndrome: Coloboma of the eye, Heart anomaly, choanal Atresia, Retardation, Genital and Ear anomalies.

Solution to Question 14:

Anteroposterior diameter of the renal pelvis of >7 to 7.5 mm in the third trimester denotes antenatal hydronephrosis due to pelvic ureteric junction (PUJ) obstruction.

Definition of antenatal hydronephrosis by anteroposterior diameter:

Degree of antenatal hydronephrosis	Second trimester	Third trimester
Mild	4 to <7 mm	7 to <9 mm
Moderate	7 to ≤ 10 mm	9 to ≤ 15 mm
Severe	>10 mm	>15 mm

Solution to Question 15:

Hepatic and periportal fibrosis is associated with autosomal-recessive polycystic kidney disease (ARPKD)

Systemic manifestations of cystic kidney diseases

Cystic condition of kidney	Associations
Autosomal dominant polycystic kidney disease	Liver cysts Cerebral aneurysms Hypertension Mitral valve prolapse Kidney stones UTIs
Autosomal recessive polycystic kidney disease	Hepatic fibrosis Pulmonary hypoplasia Hypertension
Acquired cystic kidney disease	-
Nephronophthisis	Retinal degeneration, neurologic malformations, skeletal malformations, hepatic malformations, cardiac malformations
Medullary cystic kidney disease	Hyperuricemia, gout

Solution to Question 16:

The Ask-Upmark kidney is also termed segmental hypoplasia.

Renal segmental hypoplasia:

- Severe segmental renal atrophy with loss of parenchymal elements in small kidneys is commonly known as segmental hypoplasia.
- It refers to small kidneys, usually weighing not more than 35 g.
- The scars are seen as cortical depressions overlying shrunken medullary pyramids and their dilated calyces, and they are characterized histologically by colloid-filled tubular microcysts and a paucity or absence of glomeruli.
- Most patients are 10 years or older at diagnosis and have severe hypertension.
- Nephrectomy usually controls the hypertension.

Disorders of the Nervous System

Question 1:

A 20-year-old primigravida was rushed to the OT for an emergency caesarean section. Her pregnancy was unplanned and she did not take folic acid supplementation prior to conception. Moreover, owing to the pandemic led lockdown and her remote residence, she could not adhere to regular antenatal care. The newborn was born with a neural tube defect and immediately shifted to the NICU. What is the least likely diagnosis?

- a) Myelomeningocele
- b) Anencephaly
- c) Diastematomyelia
- d) Holoprosencephaly

Question 2:

A newborn presented with a tuft of hair over the lumbosacral region. What is the likely diagnosis?

- a) Spina bifida occulta
- b) Spinal lipoma
- c) Dermal sinus
- d) Any of the above

Question 3:

A 27-year-old pregnant woman underwent second-trimester screening and had an abnormal presence of a biochemical marker. Following this, she went to a tertiary care referral centre for further investigations. An anomaly scan revealed cranial lemon and cerebellar banana signs. What is the abnormal marker present in this condition?

- a) Pregnancy-associated plasma protein-A (PAPPA)
- b) Human chorionic gonadotrophin (hCG)
- c) Inhibin A
- d) Acetylcholinesterase

Question 4:

A child is scheduled for myelomeningocele surgery. Until the patient is waiting for surgery, the covering of the sac will be protected by a gauze soaked in_____.

- a) Povidone iodine
- b) Normal saline
- c) Methylene blue
- d) Chlorhexidine

Question 5:

A 28-year-old pregnant woman was rushed to the casualty of a maternity hospital due to imminent delivery. It resulted in a stillborn baby as shown in the picture below. What is the likely diagnosis?



- a) Diastematomyelia
- b) Craniorachischisis
- c) Iniencephaly
- d) Holoprosencephaly

Question 6:

What is the most common cause of ventriculomegaly in newborns?

- a) Arnold-Chiari malformation
- b) Dandy-Walker syndrome
- c) Arachnoid villi malformation
- d) Aqueductal stenosis

Question 7:

An infant is brought to the OPD with complaints of multiple episodes of vomiting that is more in the morning and relieved when sitting. She has an unusually large head with bulging fontanelles. The image of the child showing a characteristic sign is given below. Which of the following is not a cause of the communicating variant of this condition?



- a) Choroid plexus papilloma
- b) Meningeal malignancy
- c) Vein of Galen malformation
- d) Achondroplasia

Question 8:

Shreya, a 2-year-old girl, presented with irritability for 2 days. On examination, macrocephaly, hypotonia and spasticity were present with a positive head transillumination test. She started walking independently at 18 months and became dry by day a month before her 2nd birthday. An MRI-brain revealed cystic expansion of the fourth ventricle and midline cerebellar hypoplasia. What is the most likely diagnosis explaining her obstructive hydrocephalus?

- a) Arnold-Chiari malformation
- b) Aqueductal stenosis
- c) Arachnoid villi malformation
- d) Dandy-Walker syndrome

Question 9:

A 6-month-old baby presented with fever, poor feeding, and frequent vomiting. On examination, the head circumference was 46.5cm with a positive sunset and Macewen sign. Which of the following infections is associated with the most common cause of the underlying neurological disease in this child?

- a) Rubella
- b) Mumps
- c) Bacterial endocarditis
- d) Measles


Question 10:

A 9-month-old baby presented with seizures, developmental delay and failure to thrive. Following an MRI, a diagnosis of a neuronal migration disorder was made. What is the probable diagnosis?

- a) Schizencephaly
- b) Myelomeningocele
- c) Lhermitte–Duclos disease
- d) Plagiocephaly

Question 11:

Which of the following is false regarding the given image?



- a) It represents melanocytic hamartoma
- b) It causes no visual disturbance
- c) It is associated with facial angiofibroma
- d) They are diagnostic of a neurocutaneous syndrome

Question 12:

During a telemedicine consultation, you come across a case of a 9-year-old child with a previously diagnosed autosomal dominant neurocutaneous syndrome known as tuberous sclerosis. Which one of the following is not seen in it?

- a) Astrocytoma
- b) Ependymoma
- c) White matter lesions
- d) Renal angiomyolipoma

Question 13:

During rounds in the neurology ward, an 11-year-old child is observed with the following lesion on his back. Which of the following characteristic skin manifestation of tuberous sclerosis is depicted in this picture?



- a) Adenoma sebaceum
- b) Hypomelanotic macules
- c) Shagreen patch
- d) Confetti skin lesion

Question 14:

A 3-year-old child is rushed to the emergency department because of an episode of fits associated with uprolling of eyes and high fever. Kernig's sign is negative and there are no post-ictal neurological deficits. There is a history of a similar episode when the child had a fever due to gastroenteritis. Which of the following is false regarding this type of seizures?

- a) Likelihood of developing epilepsy is low
- b) Associated with temperature of 38°C or higher
- c) Antipyretics reduce the risk of recurrent febrile convulsions
- d) Absence of having a developmental delay

Question 15:

Your neighbour, Sheila, frantically knocks on your door to seek help as her toddler had suddenly started convulsing with uprolling of eyes. The child had a high fever and this was his third episode of seizures in 6 months. Post-recovery, Sheila is still anxious and wishes to know the risk factors for such recurrences of febrile seizures. You would explain all of the following except-

- a) Duration of fever < 24 hr
- b) Late age of onset
- c) Family history of epilepsy
- d) Lower serum sodium

Question 16:

A child presents with developmental delay and multiple seizure types. The seizures could not be controlled by an array of drugs. Electroencephalogram (EEG) shows 1-2 Hz spike and slow waves with polyspike bursts in sleep. What is the diagnosis?

- a) Juvenile myoclonic epilepsy
- b) Lennox-Gastaut syndrome
- c) Landau-Kleffner syndrome
- d) West syndrome

Question 17:

Which of the following is true for Lennox-Gastaut syndrome?

- a) Constitutes of a single seizure type
- b) Good prognosis and adequate controlled epilepsy
- c) No impaired cognitive function
- d) EEG showing slow (< 3 Hz) spike and wave discharge

Question 18:

A concerned kindergarten teacher calls up the mother of one of her students who was having transient lapses of consciousness. The teacher elaborated that the child was having multiple episodes of staring blankly into space for a few seconds before quickly returning to alertness. Following investigations including an EEG, a diagnosis of atypical absence seizures was made. Which of the following is a feature of it?

- a) Associated with head drop during seizure
- b) Precipitated by hyperventilation
- c) EEG shows 3Hz spike and slow wave pattern
- d) Associated with petit mal seizures

Question 19:

Harish had a panic attack when he was called to recite a poem on the stage. This was followed by a sudden lapse in consciousness and manifested as a blank stare along with fluttering of eyes. He appeared frozen without any change in the body tone. Which of the following is a characteristic feature of this condition on electroencephalography (EEG)?

- a) Continuous stunted spikes
- b) 3 Hz spike and slow wave
- c) Continuous tall spikes
- d) Waves at the rate of 10/ sec

Question 20:

An 8-year-old boy was playing at his crèche centre when he suddenly developed rapid jerky movements of both the upper and lower limbs and was rushed to the emergency department. It had been ongoing for the past 10 minutes and the boy had been unresponsive throughout. Which of the following is not useful in the management of this condition?

- a) Lorazepam
- b) Phenytoin
- c) Phenobarbitone
- d) Carbamazepine

Question 21:

A 15-year-old girl presented with episodes of brief involuntary muscular twitching that causes her to drop her toothbrush or teacup in the morning. Following investigations including an EEG, a diagnosis of juvenile myoclonic epilepsy was made. Which of the following is not a feature of it?

- a) Sleep deprivation provokes seizures.
- b) Prognosis is generally good.
- c) Drug of choice is valproate.
- d) EEG shows 1-2 Hz spike waves.


Question 22:

An adolescent girl complains of dropping objects from hands precipitated during morning hours and exams. There is no history of loss of consciousness. One of her cousins has been diagnosed with epilepsy. EEG was suggestive of epileptic spikes. What is the most probable diagnosis?

- a) Juvenile myoclonic epilepsy
- b) Atypical absence
- c) Chorea-athetoid epilepsy
- d) Centrotemporal spikes

Question 23:

A 7-month-old boy presents with brief episodes of head drops and flexion of the body, followed by tonic extension of arms and legs. These movements last 1 to 2 seconds and occur in several clusters per day, with 15 to 30 spasms per cluster. EEG is given below and is suggestive of infantile spasms. Which of the following is the drug of choice for this condition?



- a) Carbamazepine
- b) Ethosuximide
- c) Valproate
- d) ACTH

Question 24:

An 8-month old male child presented with delayed motor milestones. He was born at 28 weeks of gestation with a birth weight of 1200 grams. A diagnosis of cerebral palsy is made following investigations. An MRI scan of the brain revealed periventricular leukomalacia. Which of the following subtypes of cerebral palsy is most likely in this child?

- a) Spastic diplegia
- b) Spastic quadriplegia
- c) Spastic hemiplegia
- d) Dyskinetic cerebral palsy

Question 25:

Which one of the following is not a feature in cerebral palsy?

- a) Hypotonia
- b) Microcephaly
- c) Ataxia
- d) Flaccid paralysis

Question 26:

A newborn presented with high-grade fever, poor feeding and incessant cry for the last 2 days. Multiple episodes of vomiting are present as well. On examination, hypotonia is present along with weak reflexes. The patient is admitted and during rounds, subtle lip-smacking seizures are also observed along with a vacant stare. CSF examination shows reduced glucose, increased protein, and increased leukocytes with predominant neutrophils. Which of the following organism is unlikely to cause this condition?

- a) Neisseria meningitidis
- b) Escherichia coli
- c) Group B streptococci
- d) Listeria


Question 27:

During your PICU rotation, you are assigned a bedside case of a child with fever, convulsions, and signs of meningeal irritation. On browsing through his file, you find a lumbar puncture report mentioning the following findings. What is the likely diagnosis?

- a) Viral meningitis
- b) Tubercular meningitis
- c) Bacterial meningitis
- d) Partially treated bacterial meningitis

Question 28:

Asif, a young child with previously diagnosed tuberculosis, is brought to your hospital for further management. Examination reveals a characteristic finding as present in the image. What do you expect in the cranial CT of this child?



- a) Ring-enhancing lesions
- b) Focal involvement of the temporal lobe
- c) Subependymal calcifications
- d) Basal exudates

Question 29:

Two girls in the same class are diagnosed with meningococcal meningitis. Their 12-year old friend from the same school is afraid of contracting the disease. What advice should be given to the exposed students?

- a) Two doses of polysaccharide vaccine
- b) Antibiotic prophylaxis
- c) Two doses of conjugate vaccine

- d) Single dose of meningococcal vaccine

Question 30:

How long should a child be isolated after being diagnosed with bacterial meningitis to prevent further transmission?

- a) Till 24 hours after starting antibiotics
- b) Till cultures become negative
- c) Till antibiotic course is complete
- d) Till 12 hours after admission

Question 31:

A 2-year-old child with a history of fall 1 year back with parietal bone fracture, has now presented with a painful and growing parietal swelling. What is the likely diagnosis?

- a) Growing scalp hematoma
- b) Growing fracture
- c) Subdural hygroma
- d) Chronic abscess

Question 32:

You have been called to declare the brain death of a 12-year-old child in the ICU. Which of the following is not seen in brain death?

- a) Normal BP without pharmacological support
- b) Positive spinal reflexes on stimulation
- c) Sweating and tachycardia
- d) Decorticate and decerebrate posturing

Answer Key

Question No.	Correct Option
--------------	----------------

1	d
2	d
3	d
4	b
5	c
6	d
7	c
8	d
9	b
10	a
11	c
12	b
13	c
14	c
15	b
16	b
17	d
18	a
19	b
20	d
21	d
22	a
23	d
24	a
25	d
26	a
27	c
28	d
29	b
30	a
31	b
32	d

Detailed Explanations

Solution to Question 1:


Holoprosencephaly is not a neural tube defect (NTD) and hence, it is the least likely diagnosis. Folic acid deficiency, especially in a young primigravida, is a risk factor for the development of NTDs in children, as seen in the given clinical vignette.

NTDs result from the failure of the neural tube to close spontaneously between the 3rd and 4th week of in utero development. These can be detected in antenatal ultrasound scans and subsequently managed for a better prognosis. The major NTDs include the following:

- Spina bifida occulta
- Meningocele
- Myelomeningocele
- Encephalocele
- Anencephaly
- Syringomyelia
- Diastematomyelia (split spinal cord)
- Iniencephaly

In holoprosencephaly, the forebrain fails to develop into two hemispheres due to mutations in the sonic hedgehog gene at the long arm of chromosome 7. It is a developmental disorder of the brain that results from the defective formation of the prosencephalon and the inadequate induction of the forebrain structures. The three types include, alobar, semilobar, and lobar, apart from the middle interhemispheric fusion variant or syntelencephaly.

The image given below shows alobar holoprosencephaly-



Solution to Question 2:

A tuft of hair over the lumbosacral region in a newborn is suggestive of occult spinal dysraphism or neural tube defect. It comprises a range of conditions, including spina bifida occulta, intradural

lipoma, and dorsal dermal sinus.

Spinal dysraphism is due to the defective fusion of the midline embryonic neural, vertebral, and mesenchymal structures. Associated clinical features are high-arched feet, muscle size and strength differences between legs, and abnormal walking patterns. In newborns and young infants, the neurological examination is usually normal, but older children may experience a lack of sensation in the perineal area and back pain.

Most of the patients will have a cutaneous abnormality overlying the lower spine, including a small dimple, a tuft of hair, or a subcutaneous lipoma.

The image shows a tuft of hair in a case of spinal dysraphism:




Spinal dysraphism also includes meningocele, myelomeningocele, lipomeningocele, diastematomyelia (split spinal cord), thickened filum terminale, and cauda equina tumour.

The initial investigation is usually ultrasound in the neonate and the best investigation is MRI.


Spina bifida is the most common type of spinal dysraphism, in which the spinal column does not close completely, leaving the spinal cord and its protective covering exposed.

Types of spina bifida




Spina bifida occulta (mildest form)

The defects in the vertebrae are so small that the spinal cord does not protrude. They are covered by skin or may have hair growing from them.



Meningocele

The meninges protrude through the defect in the vertebra.



Myelomeningocele (meningomyelocele)

The spinal cord, with the meningeal coverings, protrude through the defect in the vertebra.

Myelomeningocele is the most severe form of spina bifida, where the spinal cord and nerve roots are exposed and may be damaged. A sac-like protrusion, containing spinal cord tissue, meninges, and cerebrospinal fluid, is visible outside the body, often leading to significant neurological impairments and disabilities.

Solution to Question 3:

In the given clinical scenario, the presence of abnormal sonographic markers, like the lemon and banana sign, is highly suggestive of neural tube defects (NTDs). Acetylcholinesterase is used as a biochemical marker of neural tube defects during the screening tests.


NTDs account for the largest proportion of congenital anomalies of the CNS. The failure of closure of neural tube allows excretion of fetal substances such as α -fetoprotein (AFP) and acetylcholinesterase into the amniotic fluid thereby serving as biochemical markers for NTDs.

The lemon and banana sign, noted on antenatal imaging, are one of the many notable fruits' inspired signs. The banana sign refers to the shape of the cerebellum owing to caudal displacement while the lemon sign refers to the lemon-shaped head resulting from scalloping of the frontal bones. It is seen in fetuses with open neural tube defects.

The image below shows the characteristic lemon sign on ultrasound-



The image below shows the characteristic banana sign on ultrasound-



Note:

Pregnancy-associated plasma protein-A (PAPPA), inhibin A, and human chorionic gonadotrophin (hCG) are the biochemical markers of trisomy 21 (Down's syndrome).

Solution to Question 4:

Until the time of surgery, the sac of myelomeningocele has to be covered with a sterile gauze soaked in normal saline. This is to prevent drying and trauma to the sac. As the normal saline is less irritant to the sac, it is preferred.

Note:

Surgery is done as early as possible (within a day of birth).

Solution to Question 5:

The given image shows an occipital bone defect with retroflexion of the neck and trunk, which is typical of iniencephaly. Inion is the tip of the external occipital protuberance and this defect comes under the spectrum of neural tube defects.

Diastematomyelia (Option A) is a congenital disorder in which a part of the spinal cord is split, usually at the level of the upper lumbar vertebra.

Craniorachischisis (Option B) is the anencephaly associated with a continuous bony defect of the spine and exposure of neural tissue.

Holoprosencephaly (Option D) occurs when the procerebrum fails to develop into two hemispheres.

Solution to Question 6:


Aqueductal stenosis is the most common cause of ventriculomegaly and hydrocephalus in newborns.

Ventriculomegaly is the dilation of the ventricles. When ventriculomegaly occurs in association with raised cerebrospinal fluid (CSF) pressure, it is called hydrocephalus.

Pathologic etiologies can be congenital or acquired. Congenital causes include structural abnormalities such as aqueductal stenosis, agenesis of the corpus callosum, Dandy-Walker malformations, Chiari malformations, and neural tube defects. Acquired causes are intracranial infections (cytomegalovirus, toxoplasmosis), intracranial hemorrhage, ischemic injury, and brain parenchymal tumors.

Aqueductal stenosis is the narrowing or obstruction of the cerebral aqueduct, a narrow channel that connects the third and fourth ventricles in the brain. As a result, the fluid accumulates within the ventricles, causing them to enlarge.

Treatment for aqueductal stenosis includes ventriculostomy, diuretics, serial lumbar punctures in specific cases, and CSF diversion. In severe cases, CSF diversion procedures (shunting), may be necessary to redirect the flow of CSF and relieve pressure on the brain.



Solution to Question 7:

The clinical features of frequent vomiting, large head, bulging fontanelles, and a positive sunset sign (as seen in the image) point to a likely diagnosis of hydrocephalus. A vein of Galen malformation can expand to become large and, because of its midline position, obstruct the flow of CSF, thereby causing an obstructive or non-communicating hydrocephalus. Vomiting is more in the morning and becomes better when sitting due to poor absorption of CSF in a lying down position.

Hydrocephalus is not a specific disease; it represents a diverse group of conditions that result from impaired circulation or absorption of CSF.

Causes of hydrocephalus:

Communicating(external/non-obstructive)	Non-communicating(internal /obstructive)
Achondroplasia (raised ICP due to jugular foramen stenosis) Choroid plexus papilloma Meningeal malignancy Meningitis Posthemorrhagic Sarcoidosis	Aqueductal stenosis Arnold-Chiari malformation Dandy-Walker malformation Klippel-Feil syndrome Tumours and neurocutaneous disorders (astrocytoma) Vein of Galen malformation


Solution to Question 8:

The features of hydrocephalus like irritability along with macrocephaly, hypotonia and spasticity indicate a likely diagnosis of Dandy-Walker syndrome. A positive transillumination test is present in this syndrome. The child has developmental delay especially in motor and cognitive milestones

because normally children walk independently at 12-13 months and become dry by day around 18 months.

The Dandy-Walker malformation consists of a cystic expansion of the fourth ventricle in the posterior fossa and midline cerebellar hypoplasia. It results from a developmental failure of the roof of the fourth ventricle during embryogenesis. Most of the affected patients have hydrocephalus, agenesis of the posterior cerebellar vermis, and corpus callosum. Brisk tendon reflexes, spasticity, clonus, Babinski sign, cerebellar ataxia, and delayed motor and cognitive milestones are seen in this. Management is by shunting if hydrocephalus is present.

The image below shows a sagittal section of the magnetic resonance tomography scan of a patient with Dandy-Walker syndrome.



Solution to Question 9:

Symptoms of poor feeding and frequent vomiting with signs of enlarged head circumference and a positive sunset and Macewen sign (crack-pot resonance) points toward a diagnosis of hydrocephalus. Aqueductal stenosis is the most common cause of hydrocephalus. Out of the above options, mumps infection can cause stenosis of the cerebral aqueduct.

Mumps is associated in a child with aqueductal stenosis because-

- Mumps virus is neurotropic and is thought to enter the CNS via the choroid plexus and infect the choroidal epithelium and ependymal cells.
- CNS complications of mumps include meningitis, meningoencephalitis (most commonly manifests 5 days after the parotitis), transverse myelitis, aqueductal stenosis, and facial palsy.

Solution to Question 10:


Among the given options, schizencephaly is a neuronal migration defect. MRI is the study of choice for elucidating schizencephaly and associated malformations.

Neuronal migration is the process during brain development, where the neuronal cells migrate to their proper designated site by various mechanisms, to form the 6-layered cerebral cortex. Any abnormality in the above process leads to neuronal migration defects.

Neuronal migration defect includes the following:


- Schizencephaly: (schiz = split or cleavage) It is the presence of unilateral or bilateral clefts within the cerebral hemispheres. The patient can be severely intellectually challenged, with microcephaly, seizures, and spastic quadriplegia.

The image below shows schizencephaly:



- Lissencephaly (agyria): It is the absence of cerebral convolutions. Rather than the normal 6 layers of the cortex, there may be fewer layers. The child will have developmental delay, microcephaly, and seizures.

The image below shows lissencephaly:



Myelomeningocele (Option B) is a neural tube defect that occurs due to defective closure of the neural tube.

Lhermitte–Duclos disease (Option C) is the cerebellar dysplasia leading to focal enlargement of cerebellum and macrocephaly, cerebellar signs, and seizures.

Plagiocephaly (Option D) is the asymmetry and flattening of the skull of the child, due to extrinsic moulding forces (consistent sleeping on the same side of the head).

Solution to Question 11:

The given image shows discrete tan-colored nodules over the iris called Lisch nodules (pigmented melanocytic hamartomas). They are seen in neurofibromatosis-1 (NF-1). Facial angiofibroma (adenoma sebaceum) is a feature of tuberous sclerosis (TS) and not neurofibromatosis.

The image below shows facial angiofibroma seen in tuberous sclerosis (Bourneville disease).



Lisch nodules are specifically associated with NF-1 and are usually multiple and bilateral. They are best visualized with a slit lamp examination and don't cause any visual disturbances.

Other important ocular manifestations in NF-1 include eyelid plexiform neurofibroma, which causes S-shaped deformity of the upper lid, with a 'bag of worms' like texture, optic nerve glioma, and sphenoidal encephalocele due to dysplasia of greater wing of the sphenoid.

NF-1 (von Recklinghausen's disease) is an autosomal dominant disorder caused by a loss-of-function mutation of the NF-1 gene located on chromosome 17q11.2. NF-1 produces neurofibromin, which acts as a tumor suppressor and a negative regulator of RAS-MAP kinase signaling.

Clinical features of NF-1 include café-au-lait macules, axillary freckling, Lisch nodules, neurofibromas, and optic gliomas.

NF-1 is more common than NF2 and typically diagnosed in childhood. NF-2 is rarer but often presents with bilateral acoustic neuroma, juvenile cataract and schwannoma.

The various neurocutaneous syndromes and their ocular manifestations are as follows:

Neurocutaneous syndrome	Ocular manifestations
Neurofibromatosis type 1	- Eyelid plexiform neurofibroma - Optic nerve glioma - Bilateral Lisch nodules - Congenital ectropion - Uveae - Prominent corneal nerves - Congenital glaucoma - Sphenoidal encephalocele
Neurofibromatosis type 2	- Posterior subcapsular cataract - Hamartomas of the retinal pigment epithelium (RPE) and retina

Neurocutaneous syndrome	Ocular manifestations
Sturge-Weber syndrome	Choroidal hemangioma Iris heterochromia Ipsilateral glaucoma
Tuberous sclerosis	Atypical iris coloboma Iris hypopigmentation Retinal astrocytoma
von Hippel-Lindau syndrome	Retinal hemangiomas

Solution to Question 12:

Ependymomas are not seen in tuberous sclerosis. Neurocutaneous syndromes like tuberous sclerosis are a group of disorders with abnormalities in both the integumentary system and the CNS (defect of primitive ectoderm). They include the following:

- Neurofibromatosis
- Tuberous sclerosis
- Sturge-Weber syndrome
- von Hippel-Lindau disease
- Ataxia telangiectasia

Tuberous sclerosis is caused due to mutation of genes coding for hamartin (chromosome 9) and tuberlin (chromosome 16), which leads to the formation of hamartomas.

Clinical features:

- CNS
- CNS involvement is the hallmark of tuberous sclerosis.
- Glioneuronal hamartomas (cortical tubers).
- Subependymal nodules are lesions found along the wall of the lateral ventricles, where they undergo calcification.
- Subependymal giant cell astrocytomas (SEGAs) are the characteristic brain tumours. These benign, slow-growing tumours arise in the periventricular area.
- White matter lesions are common and include nodules, cysts, and areas of gliosis and hypomyelination.
- Skin
- Facial angiofibroma (adenoma sebaceum)
- Periungual fibroma
- Shagreen patch
- Confetti skin lesions

- Others
- Renal angiomyolipoma
- Cardiac rhabdomyoma

Solution to Question 13:

The lesion seen in the picture depicts a shagreen patch. This is a characteristic skin lesion found in patients with tuberous sclerosis. It consists of a roughened, raised lesion with an orange-peel consistency, located primarily in the lumbosacral region. Shagreen is shark skin used as a decorative material present in the image of the question.

Skin lesions in tuberous sclerosis:

- The typical hypomelanotic ash leaf-type macules on the trunk and extremities are seen in 90% of the patients. It is visualized by the use of a wood ultraviolet lamp.
- Facial angiofibromas (adenoma sebaceum) develop between 4 and 6 years of age; they appear as tiny red nodules over the nose and cheeks and are sometimes confused with acne. Later, they enlarge, coalesce, and assume a fleshy appearance.
- Periungual fibroma are small fibromas or nodules of skin, which form around fingernails or toenails in some adolescent patients.
- Confetti skin lesions are seen less commonly as a minor feature of tuberous sclerosis.

Image below: Facial angiofibromas (Adenoma sebaceum):



Image below: Periungual fibroma:



Images: Confetti skin lesions (R) and confetti (L):



Images: Ash leaf macule (above) and Ash leaves (below):



Images: Shagreen patch (above) and shagreen texture of sting ray skin (below):



Solution to Question 14:

This is a probable case of febrile seizures due to the presence of high fever, no post-ictal deficits and a negative Kernig's sign ruling out meningitis. Antipyretics do not reduce the risk of recurrent febrile seizure, as the seizure often occurs as the temperature is rising or falling abruptly.

Febrile seizures are seizures that-

- Occur between the age of 6 months and 5 years
- Occur at a temperature of 38°C (100.4°F) or higher
- Are not due to any CNS infection or any metabolic imbalance
- Occur without a history of prior afebrile seizures
- Occur with no post-ictal neurological deficit.

Types of febrile seizures:

Investigations:

- Lumbar puncture should be performed in the first episode of seizures with fever, for all infants <6 months age (to exclude meningitis).
- EEG and neuroimaging are advised for children with atypical febrile seizures.

Treatment:

Antipyretics and proper hydration. Midazolam or diazepam (rectal) can be used if seizures present.

Prophylaxis:

- Intermittent prophylaxis: If the febrile seizures are 3 or more in 6 months, or 4 or more seizures in 1 year, oral clobazam (1mg/kg/day) for 3 days during the fever episode.
- Continuous prophylaxis: If intermittent prophylaxis fails or in recurrent atypical febrile seizures, sodium valproate and phenobarbitone can be used.

Simple febrile seizures	Complex (atypical) febrile seizures
Occurs within 24 hours of the onset of fever	Can occur at any time during a fever
Generalized seizures	Focal seizures
Duration is <15 min	Prolonged
No recurrence within 24 hrs of seizures	Recurrent
Chances of going into epilepsy 1%	Chances of going into epilepsy 6%

Solution to Question 15:

Early age of onset (age <1 year) is a risk factor for recurrence of febrile seizures. The other risk factors for such recurrences are as follows:

Major:

- Age <1 yr
- Duration of fever <24 hr
- Fever 38-39°C

Minor:

- Family history of febrile seizures
- Family history of epilepsy
- Complex febrile seizure

- Male gender
- Lower serum sodium at the time of presentation

Solution to Question 16:

A child with features of developmental delay, intractable seizures and an electroencephalogram (EEG) showing 1-2 Hz spike and polyspike bursts in sleep has a likely diagnosis of Lennox–Gastaut syndrome.

Lennox–Gastaut syndrome- Typically starts between the age of 2 and 10 years. Consists of a triad of:

- Developmental delay
- Multiple seizure types (includes atypical absences, myoclonic, astatic, and tonic seizures)
- EEG findings 1-2 Hz spike and slow waves, polyspike bursts in sleep, and a slow background in wakefulness

Patients will have long-term cognitive impairment and intractable seizures despite multiple therapies. The drug of choice for Lennox-Gastaut syndrome is valproate. Other drugs used for this condition are clobazam, lamotrigine, rufinamide and topiramate.

Other options:

Juvenile myoclonic epilepsy (Janz syndrome)- Typically, it starts in early adolescence (not in childhood) and the EEG usually shows generalized 4-5 Hz polyspike and slow-wave discharges

Landau–Kleffner syndrome- A rare condition characterized by loss of language skills in a previously normal child (not a child with developmental delay)

West syndrome- Starts between the ages of 2 and 12 months and typical EEG picture is called hypsarrhythmia (high-voltage, slow with chaotic background)

Disease	Typical EEG features	Drug of choice
Lennox–Gastaut syndrome	1-2 Hz spike and polyspike bursts in sleep	Valproate
Absence seizures	3 Hz spike and slow-wave (spike and dome pattern)	Ethosuximide
Juvenile myoclonic epilepsy	4-5 Hz spike polyspike and slow-wave discharges	Valproate
West syndrome	Hypsarrhythmia (high-voltage waves with chaotic background)	Adrenocorticotrophic hormone (ACTH)

Solution to Question 17:

Lennox–Gastaut syndrome is associated with 1-2 Hz spike and slow-wave discharges.

- Typically starts between the ages of 2 and 10 years
- Consists of a triad of developmental delay, multiple seizure types (that as a rule includes atypical absence seizures, myoclonic, astatic, and tonic seizures) and specific EEG abnormalities.
- Most are left with long-term cognitive impairment and intractable seizures despite multiple therapies.
- Lennox–Gastaut syndrome has a poor prognosis

Solution to Question 18:

Atypical absence seizures have associated myoclonic components and tone changes of the head (head drop) and body.

Absence seizures(typical)	Atypical absence seizures
No change in the body tone	Change in the tone of the head (head drop) and body.
Precipitated by hyperventilation	Precipitated by drowsiness
3 Hz spike and slow wave pattern on EEG	1-2 Hz spike and slow wave pattern on EEG
Usually not associated with other syndromes	Associated with Lennox–Gastaut syndrome

Solution to Question 19:

The features of a lapse of consciousness precipitated by hyperventilation (panic attack), without any change in the tone of the body (head drop), point towards a likely diagnosis of petit mal or typical absence seizures. The electroencephalogram (EEG) in petit mal seizure is associated with a 3 Hz spike and slow-wave discharges.


Features of petit mal seizures (typical absence seizures):

- Starts abruptly in childhood (6 to 8 years)
- Approximately 25% of patients also develop generalized tonic-clonic seizures.
- Characterized by brief lapses of consciousness and sudden discontinuation of activity with staring spell, eye fluttering, or rhythmic movements
- No loss of postural control or incontinence
- No post-ictal confusion.

Hyperventilation for 3-5 min can precipitate the seizures. The drug of choice is ethosuximide.

Image: EEG waveform for an absence seizure- 3 waves per second, with spike and slow-wave pattern

EEG in Absence Seizure - 3 Hz Spike and Wave Pattern




Disease	Typical EEG features	Drug of choice
Lennox–Gastaut syndrome	1-2 Hz spike and polyspike bursts in sleep	Valproate
Absence seizures	3 Hz spike and slow-wave (spike and dome pattern)	Ethosuximide
Juvenile myoclonic epilepsy	4-5 Hz spike polyspike and slow-wave discharges	Valproate
West syndrome	Hypsarrhythmia (high-voltage waves with chaotic background)	Adrenocorticotrophic hormone (ACTH)

Solution to Question 20:

The clinical scenario is that of status epilepticus. This is due to rapid jerky movements causing continuous seizure activity for more than 5 minutes. Carbamazepine is not commonly used in the treatment of status epilepticus.

Status epilepticus is defined as continuous or recurrent seizure activity lasting for more than 5 minutes without regaining consciousness. The image given below depicts the management of this condition in a flowchart-

Management of status epilepticus



Solution to Question 21:

In juvenile myoclonic epilepsy, EEG characteristically shows 4-5 Hz polyspike and slow-wave discharges.

Juvenile myoclonic epilepsy (Janz syndrome):

- It is the most common generalized epilepsy in young adults (12-18 years).
- They are benign generalized myoclonic seizures with a good prognosis.
- They commonly affect the upper limbs more than the lower limbs and sometimes cause the dropping of objects.
- Typically, it starts in early adolescence (12-18 years), with 1 or more of the following manifestations:
 - Myoclonic jerks in the morning ("shock-like" contraction of a muscle, that is often repeated), generalized tonic-clonic or clonic seizures upon awakening; and juvenile absences.
- Sleep deprivation, alcohol (in older patients), and photic stimulation provoke seizures.
- The EEG usually shows generalized 4-5 Hz polyspike and slow-wave discharges.
- The drug of choice for juvenile myoclonic epilepsy is valproate. Lamotrigine is also useful.

Disease	Typical EEG features	Drug of choice
Lennox–Gastaut syndrome	1-2 Hz spike and polyspike bursts in sleep	Valproate
Absence seizures	3 Hz spike and slow-wave (spike and dome pattern)	Ethosuximide
Juvenile myoclonic epilepsy	4-5 Hz spike polyspike and slow-wave discharges	Valproate

Disease	Typical EEG features	Drug of choice
West syndrome	Hypsarrhythmia (high-voltage waves with chaotic background)	Adrenocorticotrophic hormone (ACTH)

Solution to Question 22:

The given clinical scenario describes an adolescent girl with myoclonic jerks (a history of dropping objects) with preserved consciousness. This, along with abnormal EEG findings strongly suggests juvenile myoclonic epilepsy.

Juvenile myoclonic epilepsy (Janz syndrome) is the most common generalized epilepsy in young adults. It starts in early adolescence with one or more of the following features: bilateral myoclonic jerks in the morning, causing patients to drop things, generalized tonic-clonic or clonic-tonic-clonic seizures upon awakening, and juvenile absence seizure.

Consciousness is usually preserved. Precipitating factors include sleep deprivation, alcohol, and photic stimulation. EEG shows 4 to 5 Hz polyspike and slow wave discharges. A family history of epilepsy is often present.

The treatment options for JME include sodium valproate, levetiracetam, and topiramate. Benzodiazepine, lamotrigine, perampanel, phenobarbital, primidone, and zonisamide can be considered as add-on drugs. Complete remission is uncommon and has a high risk of seizure recurrence, hence antiepileptics are continued lifelong in a majority of the patients.

Other options:

Option B: Atypical absence seizures have been associated with myoclonic components and head drop. They are precipitated by drowsiness and EEG shows 1 to 2 Hz spike and slow wave discharges.

Option D: In benign childhood epilepsy with centrotemporal spikes, the child typically wakes at night due to a focal seizure with preserved awareness. Buccal and throat tingling and tonic or clonic contractions on one side of the face are seen, with drooling and preserved consciousness. EEG shows typical wide-based centrotemporal spikes.

Solution to Question 23:

The drug of choice for infantile spasms is adrenocorticotrophic hormone (ACTH). The EEG shows a specific type of abnormal chaotic brainwave pattern called hypsarrhythmia. It is characterized by continuous, high-amplitude, asynchronous delta activities with interspersed independent, multiple spikes. Hypsarrhythmia on EEG can occur even between attacks. Excessive corticotropin-releasing hormone (CRH) production is hypothesized to cause infantile spasms. ACTH administration suppresses CRH secretion and hence controls the seizures.

Infantile spasms (West syndrome)

- Age of onset is 3–8 months of life

- Characterized by salaam spells (sudden dropping of the head or flexion of arms, followed by tonic extension of arms and legs) and hypsarrhythmia on EEG.
- Causes include hypoxic-ischemic encephalopathy, neurocutaneous syndromes, and idiopathic.
- Drug of choice: Adrenocorticotrophic hormone (ACTH), Vigabatrin.

The image given below illustrates infantile spasms in a child-

Disease	Typical EEG features	Drug of choice
Lennox–Gastaut syndrome	1-2 Hz spike and polyspike bursts in sleep	Valproate
Absence seizures	3 Hz spike and slow-wave (spike and dome pattern)	Ethosuximide
Juvenile myoclonic epilepsy	4-5 Hz spike polyspike and slow-wave discharges	Valproate
West syndrome	Hypsarrhythmia (high-voltage waves with chaotic background)	Adrenocorticotrophic hormone (ACTH)

Infantile Spasms



Solution to Question 24:

The most likely subtype in this child with delayed motor milestones and a history of prematurity and low birth weight is spastic diplegia. This is because the most common sequelae of periventricular leukomalacia (PVL) is spastic diplegia. PVL is the ischemia of white matter adjacent to the lateral ventricles of the brain.

Cerebral palsy

It is a group of disorders, affecting movement or posture, due to non-progressive disturbances occurring in the developing fetal brain. These could be due to hypoxic-ischemic encephalopathy.

The most common subtype of cerebral palsy is spastic diplegia.

Cerebral palsy subtype	Pathology
Spastic diplegia	Periventricular leukomalacia
Spastic quadriplegia	Parasagittal brain injury
Spastic hemiplegia	Middle cerebral artery territory infarction
Dyskinetic cerebral palsy	Basal ganglia

Solution to Question 25:

Flaccid paralysis is not a feature of cerebral palsy (CP).

CP is an upper motor neuron disease characterized by rigidity and scissoring gait. Flaccid paralysis would be seen in lower motor neuron disease.

Other options:

Option A: Hypotonia is a characteristic feature of athetoid CP. These infants generally have poor head control.

Option B: Microcephaly can be seen in CP due to perinatal insults to the developing brain.

Option C: Ataxia is a common feature of extrapyramidal CP.

The subtypes of cerebral palsy are given in the table below:

UL - upper limb; LL - lower limb; GDD - global developmental delay

Types	Etiology	Presentation	MRI findings	
Spastic(m/c type)	Diplegia(second m/c in India)	PrematurityIschemiaInfection	Bilateral spasticity (LL>UL)Scissoring gaitCommando crawlingLearning disabilities	Periventricular leukomalacia
Hemiplegia	Vascular insultInfectionGenetic	Early hand preference(UL>LL)Walking delayed up to 18-24mEquinovarusdeformity of footSeizures (33%)	Porencephalic cystFocal infarct changes	

Types	Etiology	Presentation	MRI findings	
Quadriplegia(most severe)(m/c type in India)	IschemiaInfection	Marked motor impairmentFlexioncontracturesStrong association withseizuresSevereintellectual impairment andGDDDifficulty swallowing (supranuclear palsy)	Multi-cystic encephalomalaciaSevere periventricular leukomalacia	
Dyskinetic(15-20%)	ExtrapyramidalAthetoidChorioathetoid	AsphyxiaKernicterusMitochondrialGenetic	Rigidity/hypotonia, dystonia, athetosis, choreaCognitive disabilityHip displacementSeizures	Basal ganglia scarringThalamic scarring



Solution to Question 26:

The acute features of fever, poor feeding, incessant cry, vomiting and seizures points toward a likely diagnosis of neonatal meningitis. The CSF findings correlate with that of bacterial meningitis due to purulent appearance, increased proteins, decreased glucose and a predominance of polymorphs. Neonatal meningitis is not caused by *Neisseria meningitidis* because, in neonates, bacteria are often acquired from the maternal vaginal flora.

The common causes of acute bacterial meningitis:

Age group	Organism
Neonates	Group B streptococci, <i>Escherichia coli</i> , <i>Listeria</i>

Age group	Organism
Infants and children	Neisseria meningitidis, Streptococcus pneumoniae, and Haemophilus influenzae
Adolescents and young adults	S. pneumoniae, N. meningitidis

Solution to Question 27:

Fever, convulsions, and signs of meningeal irritation (nuchal rigidity, Kernig's sign, Brudzinski's signs) in a child, with CSF showing reduced glucose, increased protein, and increased leukocytes levels is a typical presentation of bacterial meningitis.

CSF findings in different types of meningitis

Note:

In partially treated bacterial meningitis, CSF can be clear, with normal glucose levels and lymphocyte predominance. Culture is usually sterile, and antigen can be detected by agglutination test.

Parameter/Meningitis	Bacterial	Tubercular	Viral
CSF pressure	Raised	Raised	Raised
Appearance	Turbid	Clear (cob-web formation)	Clear
Glucose	Decreased	Decreased	Normal
Protein	High	Very high	slightly high
Chloride	Low	Very low	Normal
Cells	Neutrophils	Lymphocytes	Lymphocytes

Solution to Question 28:

A history of tuberculosis along with a positive Kernig's sign (as seen in the image) is suggestive of tubercular meningitis. Cranial computed tomography (CT) in a case of tubercular meningitis shows basal exudates and hydrocephalus. Intracranial tuberculosis can also manifest as ring-enhancing lesions (tuberculoma), but the basal exudates are the most common and characteristic feature of tubercular meningitis.

Tubercular meningitis

It is common between 6 months and 4 years of age and usually spread by the hematogenous route. The gelatinous exudate infiltrates the corticomeningeal blood vessels leading to hydrocephalus.

The three stages are-

- Prodromal stage
- Stage of meningitis
- Stage of coma

CSF study shows reduced sugar level, increased protein level, and lymphocytosis. Treatment includes antitubercular drugs for 12 months and corticosteroids (to reduce neurologic sequelae). TB meningitis is the only indication of steroids in TB.

Other options:

Ring-enhancing lesions are seen in cerebral abscess, neurocysticercosis and tuberculoma (rich focus). Focal involvement of the temporal lobe is seen in herpes simplex encephalitis. Subependymal calcifications are seen in tuberous sclerosis.

Solution to Question 29:

Antibiotic prophylaxis is provided to the close contacts of persons with confirmed meningococcal disease.

Close contacts of individuals with confirmed meningococcal disease are at an increased risk of developing the illness. Prophylaxis is also offered to individuals with direct and prolonged contact with a case of meningococcal disease. Antibiotics are effective in preventing additional cases by eradicating the carriage of the invasive strain.

Treatment is provided to the contacts within 24 hours of identification of the index case. The antibiotics effective include rifampicin, ciprofloxacin, ceftriaxone, and azithromycin.

Prophylaxis is not generally recommended for medical personnel, except in cases where there is exposure to aerosols of respiratory secretions, such as during mouth-to-mouth resuscitation, intubation, or suctioning. This recommendation applies within 24 hours before or after the initiation of antibiotic therapy in the index case.

For the treatment of cases, penicillin is the preferred drug, but ceftriaxone or other third-generation cephalosporins are recommended for patients with penicillin allergy. Droplet infection control precautions should be followed for hospitalized patients for 24 hours after effective therapy has been initiated.

Offering quadrivalent meningococcal A, C, W, Y conjugate vaccines in addition to antimicrobial prophylaxis can further reduce the risk of disease among close contacts of cases caused by vaccine capsular groups.

Meningococcal vaccines are available in two forms: polysaccharide vaccines and polysaccharide-protein conjugate vaccines. Polysaccharide vaccines can be given as a single dose to individuals aged 2 years and older. They are offered in bivalent (A, C), trivalent (A, C, W135), and quadrivalent (A, C, W135, Y) formulations.

Meningococcal conjugate vaccines are available as monovalent (A or C) or quadrivalent (A, C, W135, Y) formulations. Conjugate vaccines, which offer increased immunogenicity and potential for herd protection, are preferred over polysaccharide vaccines, especially in children under 2 years of age. Both types of vaccines are safe to use during pregnancy.

Solution to Question 30:

Isolation of the child is recommended until 24 hrs after the initiation of effective therapy after being diagnosed with acute bacterial meningitis to prevent further transmission.

Droplet precautions are recommended for patients with *Neisseria meningitidis* and *Haemophilus influenzae* type b (Hib) meningitis until they have received 24 hours of effective therapy. Patients should be in private rooms, and hospital personnel should wear a face mask when they are within 3 feet (1 meter) of the patient.

The child with meningitis shows nonspecific symptoms like fever, anorexia, poor feeding, headache, myalgia, hypotension, petechia, purpura, or a rash. Meningial irritation is indicated by nuchal rigidity, back pain, and Kernig, and Brudzinski signs (usually not seen in infants). Seizures and altered mental status may be present.

The recommended empirical antibiotic regimen in a suspected case of meningitis outside the neonatal period is vancomycin combined with a third-generation cephalosporin. Patients allergic to penicillin or cephalosporin can be treated with meropenem, fluoroquinolones, or chloramphenicol.

Chemoprophylaxis with rifampicin/ceftriaxone/ciprofloxacin is recommended for all close contacts with meningococcal meningitis regardless of age or immunization status. Rifampicin prophylaxis should be given to all household contacts of patients with invasive disease caused by *Haemophilus influenzae* type b, if an immunocompromised child of any age resides in the household or close family member younger than 48 months has not been fully immunized.

Solution to Question 31:

The clinical presentation of a child with a painful and growing parietal swelling, coupled with a history of a prior parietal bone fracture one year ago, strongly suggests the diagnosis of a growing fracture, also known as traumatic encephalocele or leptomeningeal cysts.

Skull fractures associated with a dural tear may fail to heal properly. The herniated brain, dilated ventricles, or a leptomeningeal cyst may cause an enlargement of this tear. These fractures are called growing skull fractures. They are usually seen in children < 3 years of age.

They present as a localized swelling or palpable skull defect that increases in size, most commonly seen in the parietal region. However, they may go undetected for years and present with delayed onset of neurologic symptoms (headache, seizure, or neurologic deficit).

Other options:

Option A: The extended duration of one year makes scalp hematoma unlikely, as it usually resolves within a few months.

Option C: Subdural hygromas are characterized by fluid accumulation in the subdural space. Most patients with subdural hygromas are asymptomatic, but uncommon symptoms may include headaches, changes in mental status, nausea and vomiting, focal neurological deficits, and seizures. They don't present with scalp swelling.

Option D: Chronic abscess may show inflammatory changes like erythema, pus, sinus, and systemic features like fever.

Solution to Question 32:

Decerebrate or decorticate posturing is not seen in brain death as they are motor responses to noxious stimuli that originate from the midbrain and cortex.

Brain death, also known as death by neurologic criteria refers to the irreversible and total cessation of all functions of the entire brain, including the brainstem.

The three components for diagnosing brain death are a demonstration of coexisting irreversible coma with a known cause, absence of brain stem reflexes, and apnea. Coma requires the patient to be unresponsive, even to noxious stimuli. The diagnosis of brain death can still be made despite positive spinal reflexes. (option B)

Brainstem reflexes (pupillary responses, oculoccephalic reflex, oculovestibular responses, corneal reflex, jaw reflex, facial grimacing, pharyngeal gag reflex, coughing) are absent. Apnea is the absence of respiratory drive at PaCO₂ of ≥ 60 mmHg and > 20 mmHg above baseline, which is consistent with brain death.

The following manifestations are occasionally seen and should not be misinterpreted as evidence of brainstem function:

- Normal systolic blood pressure - SBP > 100 mmHg (option A)
- Movements originating from the spinal cord or peripheral nerve may occur
- Sweating, flushing, tachycardia (option C)
- Deep tendon reflexes; superficial abdominal reflexes; triple flexion response
- Babinski reflex
- Spontaneous movements of limbs other than pathologic flexion or extension response
- Respiratory-like movements (shoulder elevation and adduction, back-arching, intercostal expansion without significant tidal volumes)

Disorders of Thyroid

Question 1:

Which of the following is the most common, preventable cause of mental retardation?

- a) Congenital Hypothyroidism
- b) Inborn errors of metabolism
- c) Congenital hyperthyroidism
- d) Down's Syndrome

Question 2:

In a 4-month old, term infant with open posterior fontanelle and hypotonia, you suspect congenital hypothyroidism and send thyroid studies. Which of the following is the most common cause of this condition?

- a) Maternal hypothyroidism
- b) Thyroid dysgenesis
- c) Iodine deficiency
- d) Thyroid dyshormonogenesis

Question 3:

You are called to see a 3-day-old, term neonate, who has not passed meconium yet. On clinical examination, there is generalized hypotonia and the abdomen is slightly distended without any evidence of ascites or hepatosplenomegaly. Incidentally, you also note a palpable goitre in this baby. Laboratory investigations reveal low T4 and elevated TSH levels. Which of the following is the most likely cause for this child's condition?

- a) Thyroid dyshormonogenesis
- b) Thyroid dysgenesis
- c) Central hypothyroidism
- d) Antithyroid antibodies

Question 4:

At how many weeks of gestation does thyroxine secretion begin in a fetus?

- a) 28 weeks
- b) 20 weeks
- c) 12 weeks
- d) 6 weeks

Question 5:

An anxious mother has brought a 4-month-old baby with complaints that the baby is not feeding well and is excessively sleepy all the time. Clinical examination reveals jaundice and hypotonia. His picture is provided below. What is the probable diagnosis?



- a) Congenital adrenal hyperplasia
- b) Growth hormone deficiency
- c) Congenital adrenal insufficiency
- d) Congenital hypothyroidism

Question 6:

A newborn with abnormal neonatal screening tests was diagnosed with Pendred syndrome. Which of the following is not a feature of this syndrome?

- a) Goitre
- b) Autosomal recessive disorder

- c) Conductive hearing loss
- d) Defective gene on chromosome 7

Question 7:

A 9-day-old infant has elevated TSH level and low T4 level on newborn screening. If this condition is left untreated, which of the following is most likely to be seen in this infant in the first few months of life?

- a) Hyperreflexia
- b) Hyperirritability
- c) Diarrhea
- d) Prolonged jaundice

Question 8:

During a rural health camp, a father brings a dull-looking baby with macroglossia and an umbilical hernia. On enquiry, you find out that the parents were told about a thyroid problem in the child at birth, for which they did not pursue any treatment. Which of the following is least likely to be found in children with the given condition?

- a) Abnormal facies due to craniosynostosis
- b) Palpable goitre
- c) Open posterior fontanelle
- d) Epiphyseal dysgenesis

Question 9:

As part of a quality improvement project undertaken by the maternal and child health services, you are evaluating the current practice of newborn screening and detection rates of hypothyroidism in various community health centres. Which of the following statements is false?

- a) Simultaneous measurement of T4 and TSH levels is the ideal method.
- b) The sample is obtained by heel-prick between 1-5 days of life using dried blood spot.
- c) Newborn screening always detects the affected twin in same-sex twins.
- d) Primary TSH, with back up T4 is the most common strategy used.

Question 10:

As an intern posted in paediatrics, you are asked to interpret the results of a heel prick test that you performed on a newborn. Which of the following results is suggestive of primary hypothyroidism?

- a) Low T4 level and low TSH level
- b) Low T4 level and high TSH level
- c) High T4 level and low TSH level
- d) High T4 level and high TSH level

Question 11:

The image given below depicts an x-ray of the knee in a term baby. Which among the following conditions is associated with it?



- a) Congenital hypothyroidism
- b) Graves disease
- c) Congenital rickets
- d) Osteogenesis imperfecta

Question 12:

Ayesha is a 9th standard girl who presented with a neck swelling, which began more than a year ago. Her mother says she is embarrassed about it and always covers her neck with a

shawl when she goes to school. Her menstrual cycles have also become irregular as of late. Blood tests reveal slightly elevated TSH levels and the presence of thyroid peroxidase antibodies. Which of the following is the most likely diagnosis?

- a) Iodine deficiency
- b) Congenital hypothyroidism
- c) Graves disease
- d) Chronic lymphocytic thyroiditis

Question 13:

A lady, with a history of a thyroid disorder and three prior abortions, delivered a female neonate prematurely at 34 weeks of gestation. In the newborn infant, tachycardia, irritability, and prominent eyes were noted and neonatal hyperthyroidism was suspected. Which of the following statements is false about the condition?

- a) Maternal Graves' disease is the most common cause of neonatal hyperthyroidism
- b) It results from the transplacental passage of maternal stimulatory TSH receptor antibody
- c) Microcephaly and frontal bossing with triangular facies are common clinical findings
- d) Neonatal hyperthyroidism occurs in most infants born to mothers with a history of Graves' disease

Question 14:

A pregnant woman with a past history of Grave's disease has currently completed 36 weeks of gestation. She underwent partial thyroidectomy 10 years back and has remained euthyroid since then. The previous child had craniosynostosis. Which of the following antibodies is most likely to be detected, if a cord blood sampling is done at delivery?

- a) Anti-TPO antibody
- b) TSH receptor stimulating antibody
- c) TSH receptor blocking antibody
- d) Anti-Thyroglobulin antibody

Question 15:

A 4-year-old girl, a known case of celiac disease, presented to the pediatric endocrine clinic with a history of neck swelling, tremor, and weight loss. On evaluation, she was found to have

Graves' disease. Other conditions that need to be screened in this patient include all of the following except _____.

- a) Vitiligo
- b) Pernicious anemia
- c) Diabetes mellitus type 2
- d) Addison disease

Question 16:

An 11-year-old girl was brought by her parents with complaints of weight loss, increased appetite, and a decline in scholastic performance. Following investigations, a diagnosis of Grave's disease is made. Which of the following is the drug of choice for initial therapy?

- a) Propylthiouracil
- b) Methimazole
- c) Radioactive iodine
- d) Levothyroxine

Question 17:

Your unit chief has asked you to present the case of a teenager, who was admitted for evaluation of mental retardation, deaf-mutism and a gait disturbance. He also has a significant patellar clonus. On general exam, you note a goitre but according to lab reports, he is apparently euthyroid. His pubertal development has also been normal. The most likely diagnosis is _____.

- a) Neurological endemic cretinism
- b) Grave's disease
- c) Pendred syndrome
- d) Myxedematous endemic cretinism

Question 18:

During a health outreach program in Sikkim, you come across 3 children from the same family, with goiter. Endemic cretinism was suspected. Which of the following can be a manifestation of this condition?

- a) Facial nerve involvement

- b) Blindness and hypothyroidism
- c) Strabismus and spastic diplegias
- d) Reversible mental retardation

Answer Key

Question No.	Correct Option
1	a
2	b
3	a
4	c
5	d
6	c
7	d
8	a
9	c
10	b
11	a
12	d
13	d
14	b
15	c
16	b
17	a
18	c

Detailed Explanations

Solution to Question 1:

Congenital hypothyroidism is the most common preventable cause of mental retardation which results from a deficient production of thyroid hormone. It is more common among girls and the prevalence is about 1:3000 to 1:4000 births.

Down's syndrome is the most common overall cause of mental retardation, followed by Fragile X syndrome, which mainly affects boys.

Causes of mental retardation:	
Prenatal	Congenital hypothyroidism Infection (toxoplasma, rubella, CMV, HSV) Genetic abnormalities Trisomy 21 Fragile X syndrome other point mutations and chromosomal abnormalities Toxins and teratogens Alcohol Drugs Radiation Inborn errors of metabolism
Perinatal	Hypoxia Prematurity Intracranial haemorrhage Perinatal CNS infection
Postnatal	Acquired brain injury CNS infection, haemorrhage, malignancy Severe malnutrition and deprivation Environmental toxins (lead, mercury)

Solution to Question 2:

Thyroid dysgenesis is the most common cause of permanent congenital hypothyroidism (80–85% of cases) & dyshormonogenesis (15%).

Dysgenesis may involve ectopy, agenesis (absence) or hypoplasia (underdevelopment) of the thyroid gland. Dyshormonogenesis results from a defect in any of the steps in thyroid hormone biosynthesis.

*TSH - thyroid-stimulating hormone; TRH - thyroid-releasing hormone

Acquired hypothyroidism can be caused by iodine deficiency, thyroiditis, radiation, histiocytosis, and drugs. The most common cause of acquired hypothyroidism is autoimmune thyroiditis (Hashimoto's).

Note: Mutation in PAX-8 gene is associated with congenital hypothyroidism and genitourinary abnormalities such as renal agenesis.

Etiology of congenital hypothyroidism:

Primary hypothyroidism: Thyroid dysgenesis, Thyroid dysmorphogenesis, Iodine deficiency (endemic goiter), Defect in thyroid hormone transport- Mutation in MCT8 (monocarboxylate transporter 8) gene, Maternal antibodies- thyrotropin receptor-blocking antibody (TRBAbs), Maternal medications- iodides, amiodarone, propylthiouracil, methimazole, radioiodine

Central (hypopituitary) hypothyroidism: TSH deficiency- mutation in TSH β -subunit gene, TRH deficiency- mutation in TRH gene, TRH unresponsiveness- mutation in TRH receptor gene, Multiple congenital pituitary hormone deficiencies

Solution to Question 3:

A history of decreased activity and late passage of meconium, associated with low T₄ and high TSH is suggestive of primary hypothyroidism. In an infant with a palpable goitre, the likely diagnosis is congenital hypothyroidism secondary to thyroid dysmorphogenesis.

The most common cause of congenital hypothyroidism is thyroid dysgenesis but it is not associated with goitre. Thyroid dysmorphogenesis results from the deficiency of one or more enzymes involved in thyroid hormone synthesis/secretion.

Congenital hypothyroidism can be:

Goitrous (Palpable goitre)	Non-goitrous (No palpable goitre)
Thyroid dysmorphogenesis, Iodine deficiency, Placental passage of antithyroid drugs	Thyroid dysgenesis, Thyroid receptor blocking antibodies, Central (hypopituitary) hypothyroidism

Solution to Question 4:

Thyroxine secretion begins in the fetus at 12 weeks of gestation.

The fetal thyroid arises from an outpouching of the foregut at the base of the tongue (foramen cecum). It migrates to its normal location over the thyroid cartilage by 8-10 wk of gestation. The bilobed shape of the thyroid is recognized by 7 weeks of gestation.

Solution to Question 5:

The above picture shows a child with a protuberant abdomen, umbilical hernia, macroglossia, and edematous facies. Also, the infant is sluggish, has jaundice, feeding difficulties, and hypotonia, all of which suggests a diagnosis of congenital hypothyroidism.

Clinical features of congenital hypothyroidism:

- Prolongation of physiologic hyperbilirubinemia- early sign (it is due to delayed maturation of glucuronide conjugation)
- Constipation- usually does not respond to treatment
- Wide open anterior and posterior fontanelles (>0.5 cm) and cranial sutures
- Developmental delay, delayed bone growth, delayed dentition
- Cardiovascular- slow pulse rate, cardiomegaly, asymptomatic pericardial effusion, edema
- Respiratory- macroglossia, feeding difficulties, choking spells, respiratory difficulties
- Haematological- macrocytic anaemia refractory to treatment with haematinics
- Nervous system- hypothermia, hypotonia, slow relaxation of deep tendon reflexes, sluggishness, and somnolence

Solution to Question 6:

Sensorineural hearing loss is present in Pendred syndrome; not conductive hearing loss.

Pendred syndrome is an autosomal-recessive disorder, caused by a mutation in the gene SLC26A4 on chromosome 7, which codes for pendrin, a chloride–iodide transport protein common to the thyroid gland and the cochlea.

Clinical features include sensorineural hearing loss and goitre with hypothyroidism. It is the most common cause of syndromic deafness. The biochemical reason for hypothyroidism in patients with Pendred syndrome is impaired iodide organification (resulting from a mutation in the Pendrin gene).

Solution to Question 7:

Elevated TSH levels and low T4 levels in a 9-day-old infant are suggestive of congenital hypothyroidism. Prolongation of physiologic hyperbilirubinemia is an early sign of congenital hypothyroidism. It is due to delayed maturation of the hepatic glucuronide conjugation system.

Hyperreflexia, hyperirritability and diarrhoea are seen in hyperthyroidism; not hypothyroidism.

Note:

Kocher–Debré–Sémélaigne syndrome: The muscles are usually hypotonic in congenital hypothyroidism. Rarely, hypothyroidism in infancy can be characterized by generalized muscular pseudohypertrophy, myxoedema, short stature and cretinism. Affected older children can have an athletic appearance. Boys are more prone to the development of the syndrome.

Solution to Question 8:

A dull-looking face with macroglossia, and umbilical hernia in a child with a thyroid disorder are all indicative of congenital hypothyroidism. Craniosynostosis is not seen in neonatal hypothyroidism. It is seen in hyperthyroidism.

Craniosynostosis refers to the premature fusion of sutures and closure of fontanelles and often results in abnormal, triangular facies. In congenital hypothyroidism, there is delayed closure of the fontanelles. Palpable goitre, though not common, can be seen in cases of untreated thyroid dysmorphogenesis.

Solution to Question 9:

Newborn screening can fail to detect the affected twin with hypothyroidism.

Discordance for thyroid dysgenesis is the rule in monozygotic twins. The transfusion of euthyroid blood from the unaffected twin normalizes the serum levels of T₄ and TSH in the affected twin, resulting in delayed or missed diagnoses. Therefore, a second sample for congenital hypothyroidism screening at 14 days of age should be considered in monozygotic twins.

Neonatal thyroid screening strategies:

- Primarily TSH, backup T₄; T₄ is measured if TSH level is >20 mU/L
- Primarily T₄, backup TSH: TSH is measured only if T₄ level is low
- Concomitant T₄ and TSH; it is the most sensitive method

Note: the sample is obtained after 24-48 hours, to minimize the false elevation of TSH values due to normal post-delivery neonatal TSH surge.

Solution to Question 10:

Low T₄ and high TSH levels indicate primary hypothyroidism.

Blood specimen for neonatal thyroid screening is obtained by heel prick between 1 and 5 days of life.

Levothyroxine therapy is started immediately after diagnosis, at 10 to 15 µg/kg per day.

Condition	Goitre	T4	TSH
Thyroid dysgenesis	Absent	Low	Elevated
Thyroid dyshormonogenesis	Present	Low	Elevated
Thyroid receptor blocking antibodies	Absent	Low	Elevated
Central hypothyroidism	Absent	Low	Normal/Low
Hyperthyroidism	Present	High	Low

Solution to Question 11:

The deformity shown in the X-ray image is the absence of distal femoral epiphysis, which is a typical radiological feature of congenital hypothyroidism.

X-ray findings in hypothyroidism

- Absence of distal femoral and proximal tibial epiphyses at birth.
- Punctate epiphyseal dysgenesis (multiple foci of ossification)
- Deformity or beaking of the 12th thoracic or 1st or 2nd lumbar vertebra
- X-ray image of the skull shows large fontanelles and wide sutures; wormian (intrasutural) bones; and, often, an enlarged and round sella turcica.


Image below: The infant on the left with congenital hypothyroidism demonstrates the absence of the distal femoral and proximal tibial epiphyses, while in the normal infant on the right the distal femoral epiphysis is present.



The image below shows small upper femoral epiphyses:



The image below shows enlarged sella turcica and wormian bones (intrasutural):



Solution to Question 12:

The presence of thyroid peroxidase antibodies (TPO-Abs) and elevated thyroid-stimulating hormone (TSH) level in a child with a goitre is suggestive of chronic lymphocytic (Hashimoto) thyroiditis.

Chronic lymphocytic (Hashimoto) thyroiditis is an autoimmune disease. Clinical presentation can include goitre, growth retardation, and other features of hypothyroidism. The thyroid is diffusely enlarged, firm, and non-tender.

TSH is moderately elevated and TPO-Abs and anti-thyroglobulin antibodies (anti-Tg Abs) can be detected.

The image given below shows goitre due to Hashimoto's thyroiditis.



Solution to Question 13:

Neonatal hyperthyroidism occurs in only about 2% of infants born to mothers with a history of Graves' disease. But maternal Graves' disease is by far the most common cause of neonatal hyperthyroidism.

Neonatal hyperthyroidism (due to Grave's) results from the transplacental passage of maternal stimulatory thyrotropin/TSH receptor antibody (TSHR-Ab).

Clinical features are as follows:

- Low birth weight (IUGR)
- Premature birth
- Microcephaly
- Frontal bossing and triangular facies
- Warm, moist skin
- Irritability, hyperactivity, restlessness, poor sleep
- Tachycardia

Note:

TSHR-Ab (TSH-R-stimulating antibodies) is another name for TSH-R-stimulating immunoglobulins (TSI).

Solution to Question 14:

The given clinical scenario with a history of Graves' disease in the mother and craniosynostosis (neonatal Graves') in the previous child suggests that the fetus is at increased risk of neonatal Grave's disease. Thus, the antibody likely to be found at birth in the cord blood sampling is the TSH receptor stimulating antibody (TRSAbs).

Neonatal Graves is caused by the transplacental passage of TRSAbs from a mother with a history of Graves' disease. The mothers can have

- Active Graves' disease
- Graves' in remission
- Prior history of Graves' treated with radioactive ablation or thyroidectomy

TRSAbs bind to the receptor for TSH and stimulate cyclic AMP, leading to thyroid hyperplasia and unregulated overproduction of thyroid hormone.

Solution to Question 15:

Type 2 Diabetes mellitus is not associated with Grave's disease.

Conditions associated with Grave's disease (autoimmune diseases):

- Diabetes mellitus type 1
- Pernicious anaemia
- Celiac disease
- Vitiligo
- Addison disease
- Alopecia areata
- Myasthenia gravis

Solution to Question 16:

Methimazole is the initial drug of choice for treating Grave's disease (hyperthyroidism) in children.

Option A: Propylthiouracil is not recommended in children because it is associated with fatal hepatotoxicity.

Option C: Radioactive iodine is not preferred as the first-line treatment in hyperthyroidism for children.

Option D: Levothyroxine is used to treat hypothyroidism.

Solution to Question 17:

The clinical presentation of a euthyroid patient with goitre as well as mental retardation, deaf-mutism, and normal pubertal development points towards neurological endemic cretinism.

Endemic cretinism includes 2 different, but overlapping, syndromes: neurologic type and myxedematous type. It is essentially the same disorder, phenotypically modified by the length and severity of postnatal hypothyroidism. Neurological manifestations are interpreted as reflecting the effects of maternal and fetal hypothyroxinemia, secondary to severe iodine deficiency, on the developing nervous system. Both neurologic and myxedematous type can manifest neurologic symptoms.

Although neurological cretins are euthyroid, linear growth arrest lines in the long bones are often demonstrated radiologically, suggesting previous hypothyroidism.

Option B: Graves disease includes a triad of hyperthyroidism with goitre, ophthalmopathy, and dermopathy (no mental retardation, deafness, or pyramidal signs).

Option C: Pendred Syndrome is caused by a mutation in chloride-iodide transporter in thyroid and cochlea resulting in euthyroid/hypothyroid goitre and sensorineural hearing loss (pyramidal signs like clonus are not seen).

Solution to Question 18:

Strabismus and spastic diplegias are among the manifestations of the neurological type of endemic cretinism.

Cretinism is a condition characterized by irreversible abnormalities in intellectual development. Endemic cretinism occurs due to the nutritional deficiency of iodine. It can be neurological cretinism or myxedematous cretinism. It is associated with severe, irreversible mental retardation.

Congenital Adrenal Hyperplasia and Related Disorders

Question 1:

The parents of a neonate diagnosed with congenital adrenal hyperplasia come with queries about the disease. They have been told that it's an inherited condition and ask you how is it possible for the baby to be affected when neither of the parents are. What is the mode of inheritance of the disease?

- a) Autosomal recessive
- b) Autosomal dominant
- c) X-linked recessive
- d) X-linked dominant

Question 2:

A baby delivered at home was born with ambiguous genitalia. The parents could not identify the sex of the child. The baby was diagnosed with congenital adrenal hyperplasia. Which of the following enzyme deficiencies is this child most likely to have?

- a) 3β -Hydroxysteroid dehydrogenase deficiency
- b) 17-Hydroxylase deficiency
- c) 11β -Hydroxylase deficiency
- d) 21-Hydroxylase deficiency

Question 3:

Which of the following types of congenital adrenal hyperplasia is not due to 21-hydroxylase deficiency?

- a) Salt-wasting type
- b) Simple virilizing type
- c) Non-classic type
- d) Under virilizing type

Question 4:

A concerned mother brings her 15-day old baby to you with complaints of decreased activity and vomiting. On examination, the baby is hypotensive, has depressed fontanelles and reduced skin turgor. The image of the external genitalia of the baby is shown below. Which electrolyte abnormality can you expect to see in this baby?



- a) Hypokalemia
- b) Hyperkalemia
- c) Hypocalcemia
- d) Hypercalcemia

Question 5:

Investigation of choice to diagnose congenital adrenal hyperplasia due to 21-hydroxylase deficiency is estimation of _____ levels.

- a) 21-Hydroxylase
- b) 17-Hydroxyprogesterone
- c) 3β -Hydroxysteroid dehydrogenase
- d) 11-Deoxycortisol


Question 6:

You see Sanjay who is a known case of congenital adrenal hyperplasia, on steroid therapy in the ER with vomiting and fever. You administer double the regular dose of steroids. Which drug is being referred to here?

- a) Dexamethasone
- b) Betamethasone
- c) Prednisolone
- d) Hydrocortisone

Question 7:

The autopsy image of a baby who died during the neonatal period is shown below. Parents of the child undergo genetic counseling regarding future pregnancies. When should the mother be started on steroids in a future pregnancy?



- a) As soon as pregnancy is diagnosed
- b) At 28 weeks of pregnancy
- c) Before conception
- d) After karyotyping and determination of sex of the baby

Question 8:

You see Sathya, an adolescent in the endocrinology clinic referred by his family physician. He was told by his parents that he was a boy, but he felt differently. As he grew up, he felt increasingly uncomfortable with his identity and this caused him a great deal of sadness and

isolation. On examination, he is hypertensive and has genitalia resembling a female phenotype. After karyotyping, you diagnose him with 46XY DSD. Which of the following enzyme deficiency would be in your differential?

- a) 21-Hydroxylase
- b) 11 β -Hydroxylase
- c) 17-Hydroxylase
- d) 3 β -Hydroxysteroid dehydrogenase

Question 9:

An infant with hypospadias, bifid scrotum presents with signs and symptoms of salt wasting. Lab reports reveal elevated levels of 17-hydroxy pregnenolone and dehydroepiandrosterone. Which of the following deficiency would you suspect in this patient?

- a) 3 β -Hydroxysteroid dehydrogenase
- b) 11 β -Hydroxylase
- c) 17-Hydroxylase
- d) 21-Hydroxylase

Question 10:

The most common enzyme deficiency leading to childhood hypertension among the following is _____.

- a) 17-Hydroxylase deficiency
- b) 21-Hydroxylase deficiency
- c) 3 β -Hydroxysteroid dehydrogenase
- d) 11 β -Hydroxylase deficiency

Question 11:

A boy presents with precocious puberty and hypertension to your OPD. The estimation of _____ will help in the diagnosis of CAH.

- a) 17-Hydroxy progesterone
- b) Cortisol
- c) Aldosterone

d) 11 Deoxycortisol

Question 12:

On a vacation trip to Hawaii, a father was helping his 7-year-old son change into swimming trunks. He noticed the presence of pubic and axillary hair. He knew it was abnormal and brought the child to the pediatrician. Which of the following is the least likely diagnosis?

- a) 17-Hydroxylase deficiency
- b) 21-Hydroxylase deficiency
- c) 11 β -Hydroxylase Deficiency
- d) None of the above

Question 13:

Congenital adrenal hyperplasia most commonly presents as _____.

- a) Male pseudohermaphroditism
- b) Female pseudohermaphroditism
- c) True hermaphroditism
- d) 46, XY intersex

Question 14:

A postgraduate examines a female baby in late infancy with failure to thrive. The genitalia appears normal. Reports reveal serum sodium of 125 MEq/L and serum K⁺ of 6.5 MEq/L. Estimation of which of the following hormones will be helpful in the diagnosis?

- a) Aldosterone
- b) DHEA
- c) 17-hydroxyprogesterone
- d) Cortisol

Question 15:

A 12-year-old boy presents with obesity and short stature, with signs of abnormal masculinization as shown below. Which statement regarding this condition is false?



- a) Most common cause is adrenocortical tumor
- b) Hypertension seen here can occasionally lead to heart failure
- c) Hyperglycemia is a finding.
- d) Generalized obesity is a common finding in younger children.

Question 16:

A high school girl Meenu presents to you with anorexia, vomiting, and muscle weakness for the past few months. On examination, she had the finding shown in the image. Which of the following will not be seen in this condition?



- a) Hyponatremia

- b) Hypoglycemia
- c) Hypotension
- d) Hypertension

Question 17:

You have been called to see a 10-year-old child in the emergency room with complaints of fever and vomiting for the past three days. On examination the child had tachycardia and there was significant postural hypotension. Upon investigation, hyponatremia and hyperkalemia were present. You suspect adrenal insufficiency. Which of the following is the most likely organism grown in blood culture?

- a) Haemophilus influenzae
- b) Neisseria meningitidis
- c) Mycobacterium tuberculosis
- d) Corynebacterium diphtheriae

Question 18:

In the pediatric ward, a resident is evaluating a case of primary adrenal insufficiency who has failure to thrive, recurrent emesis, and hepatosplenomegaly. Bilateral adrenal calcification is noted radiologically. The most likely diagnosis is _____.

- a) Pheochromocytoma
- b) Nelson's disease
- c) Wolman's disease
- d) Adrenal hemorrhage

Question 19:

You are evaluating a case of hyperpigmentation. The patient reports having undergone bilateral adrenalectomy a few years back for Cushing's disease. MRI shows an enlargement of the pituitary adenoma. What is the diagnosis?

- a) Nelson syndrome
- b) Steele-Richardson-Olszewski syndrome
- c) Hamman-Rich syndrome
- d) Job's syndrome

Question 20:

Using novel chemiluminescence immunoassay techniques, you measure the hormonal profile of the adrenal gland of a normal fetus. Which of the following would be present in higher concentrations?

- a) DHEA
- b) Cortisol
- c) Corticosterone
- d) Progesterone

Question 21:

A woman has a previous child with congenital adrenal hyperplasia. In the present pregnancy, steroid therapy should be started _____

- a) As soon as pregnancy is diagnosed
- b) At 28 weeks of pregnancy
- c) Before conception
- d) After karyotyping and determination of sex of the baby

Question 22:

You see a newborn with an enlarged clitoris and partially fused labia. You also note features such as frontal bossing, brachycephaly, midface hypoplasia, slender fingers, medial bowing of ulna, and heart abnormalities. Investigation reveals a deficiency of P450 oxidoreductase. What is your diagnosis?

- a) Antley Bixler syndrome
- b) Waterhouse-Friderichsen syndrome
- c) Nelson syndrome
- d) Lipoid adrenal hyperplasia

Answer Key

Question No.	Correct Option
1	a
2	d
3	d
4	b
5	b
6	d
7	a
8	c
9	a
10	d
11	d
12	a
13	b
14	a
15	a
16	d
17	b
18	c
19	a
20	a
21	a
22	a

Detailed Explanations


Solution to Question 1:

Congenital adrenal hyperplasia / adrenogenital syndrome follows an autosomal recessive inheritance pattern.

Depending on the enzyme that is deficient, patients will have :

- Mineralocorticoid deficiency (hyponatremia, hyperkalemia) or excess (hypernatremia, hypokalemia)
- Incomplete virilization or precocious puberty in males
- Virilization or sexual infantilism in females

Pathophysiology of congenital adrenal hyperplasia




©Marrow

Solution to Question 2:

This child is most likely to have a 21-hydroxylase deficiency as it is the most common enzyme deficiency implicated in congenital adrenal hyperplasia. The second most common is 11 beta-hydroxylase deficiency.

21-Hydroxylase deficiency



©Marrow

21-hydroxylase enzyme hydroxylates progesterone and 17-hydroxyprogesterone to yield 11-deoxycorticosterone and 11-deoxycortisol, respectively. They help in the synthesis of aldosterone and cortisol. Therefore, the deficiency of this enzyme leads to the deficiencies of aldosterone and cortisol.

Also, the accumulated steroid precursors are diverted into the pathway for androgen (testosterone) production.

Clinical profile in 21-hydroxylase deficiency:

- Aldosterone deficiency: Salt-wasting crises (hyponatremia, hyperkalemia, hypotension, and metabolic acidosis)
- Cortisol deficiency: Hypoglycemia, hyperpigmentation (due to increased ACTH production), weight loss, weakness, anorexia, vomiting, and shock
- Androgens excess: Virilization of females and precocious puberty in males

Solution to Question 3:

Under virilizing type of congenital adrenal hyperplasia (CAH) is not caused by 21-hydroxylase deficiency.

Depending on the extent of enzyme deficiency, CAH caused by 21-hydroxylase deficiency is broadly classified into classic and non-classic forms.

Classic forms can be of two types - salt-wasting and simple virilizing types.

- Salt-wasting type is the severe form in which both cortisol and aldosterone are deficient, with excess androgen production.
- In the simple virilizing type, there is an impaired synthesis of cortisol and adequate amounts of aldosterone. Here, clinical features are due to the overproduction of androgens.

The non-classic form is characterized by partial 21-hydroxylase deficiency. Cortisol and aldosterone are produced normally, but at the expense of producing excess androgens.

Solution to Question 4:

The image shows ambiguous genitalia. This along with signs of dehydration and hypotension suggest congenital adrenal hyperplasia due to 21-hydroxylase deficiency. Hyperkalemia and hyponatremia (due to mineralocorticoid deficiency) can be seen in this condition.

These babies present at 10–14 days of age with progressive weight loss, weakness, anorexia, vomiting, dehydration, hypoglycemia, and hypotension.

Without treatment, shock, cardiac arrhythmias, and death may occur within days or weeks.


Solution to Question 5:

17-hydroxyprogesterone levels estimation is the investigation of choice to diagnose CAH due to 21-hydroxylase deficiency. It will be markedly elevated.

Other findings include:

- Elevated androstenedione and testosterone (in females).
- Low cortisol and aldosterone.

- Elevated ACTH levels.



Solution to Question 6:

Hydrocortisone, which is the mainstay treatment for commonest type of CAH, is the drug referred to in the given clinical scenario.

Double or triple doses are indicated during periods of stress, such as infection or surgery.

Glucocorticoid replacement therapy with hydrocortisone, 15-20 mg/m²/24 h, orally, in 3 divided doses is continued lifelong.

Mineralocorticoid replacement with fludrocortisone is used for patients with the salt-wasting disease. The daily dose of fludrocortisone in infants is higher than that in older children.


Solution to Question 7:

The autopsy image shows a bilaterally enlarged adrenal gland (white arrows) with a history of death during the neonatal period (probably due to shock, cardiac arrhythmias) which is suggestive of CAH in the baby. Mothers with a previously affected child of CAH should be started with steroid therapy as soon as pregnancy is diagnosed. Since it is an autosomal recessive disorder, there is a risk of the subsequent children being affected.

Dexamethasone at 20 µg/kg is started to suppress the secretion of steroids by the fetal adrenals (including adrenal androgens) and prevent the virilization of the external genitals in affected females. Then, chorionic villus biopsy is performed to determine the sex and genotype of the fetus, and therapy is continued only if the fetus is an affected female.

Solution to Question 8:

46XY DSD - a genotypical male with female phenotype- along with hypertension makes the diagnosis of congenital adrenal hyperplasia due to 17-hydroxylase deficiency more likely.



Clinical profile in 17-hydroxylase deficiency:

Aldosterone excess: Salt retention, hypernatremia, hypokalemia, and hypertension.


Sex steroid deficiency: In contrast to 11 β -hydroxylase deficiency, patients with 17-hydroxylase deficiency are unable to synthesize sex hormones. Affected males are incompletely virilized (male pseudo-hermaphroditism/ ambiguous genitalia 46XY) and there is a failure of sexual development in affected females.

Cortisol deficiency is not manifested, as elevated corticosterone is an active glucocorticoid.

Intersex or DSD means a condition in which development of chromosomal, gonadal, or anatomical sex is atypical. There is a greater incidence of gender dissatisfaction, gender non-conformity, gender fluidity in such individuals.

Solution to Question 9:

The given clinical scenario of incomplete virilization (hypospadias with or without bifid scrotum and cryptorchidism), salt wasting due to cortisol and aldosterone deficiency, and elevated dehydroepiandrosterone (DHEA) and 17 hydroxy pregnenolone levels are suggestive of 3 beta-hydroxysteroid dehydrogenase deficiency.



Clinical profile in 3 β -hydroxysteroid dehydrogenase (3 β -HSD):

- Aldosterone deficiency- Salt-wasting crises (hyponatremia, hyperkalemia, hypotension, and acidosis)
- Cortisol deficiency- Hypoglycemia, hyperpigmentation (due to increased ACTH production), and shock
- Androgen deficiency- Males are incompletely virilized (male pseudo-hermaphroditism/ambiguous genitalia 46XY/hypogonadism, cryptorchidism)
- Excess DHEA (weak androgen)- Females are mildly virilized (acne, hirsutism).

Solution to Question 10:

The most common enzyme deficiency leading to childhood hypertension is 11 β -hydroxylase deficiency causing congenital adrenal hyperplasia (CAH).

Hypertension in CAH is due to 11 β -hydroxylase deficiency followed by 17-hydroxylase deficiency.

Cortisol and aldosterone deficiency cannot be manifested because some corticosterone is synthesized from progesterone, and hence, there are no signs of adrenal insufficiency.


21- hydroxylase deficiency causes salt-wasting resulting in hypotension.

Solution to Question 11:

Precocious puberty in a male with hypertension is suggestive of CAH due to 11 β -hydroxylase deficiency. An increase in 11-deoxycortisol levels is diagnostic of the enzyme deficiency.

11 β -hydroxylase catalyzes the conversion of deoxycortisol to cortisol. Therefore, 11 β -hydroxylase deficiency causes an increase in 11 deoxycortisol levels.

11 β -Hydroxylase deficiency




Solution to Question 12:

The presence of pubic and axillary hair in a seven-year-old boy is suggestive of precocious puberty and 17-Hydroxylase deficiency is least likely to cause it.

In 17-hydroxylase deficiency, patients are unable to synthesize sex hormones, and hence the affected males are incompletely virilized and present as phenotypic females (but gonads are usually palpable in the inguinal region or the labia).

In both 21-hydroxylase and 11 β -hydroxylase deficiencies, the respective precursors to the hormones are shunted to the androgen synthesis pathway, resulting in all the signs and symptoms of androgen excess, including virilization of female and precocious puberty in males.

17 α -Hydroxylase deficiency



Solution to Question 13:


Congenital adrenal hyperplasia (CAH) most commonly presents as female pseudohermaphroditism as the most common enzyme deficiency is 21-hydroxylase deficiency.

A female pseudohermaphrodite is a genotypically normal female (46, XX), whose external genitalia resembles a male's.

The image below shows ambiguous genitalia in a female with CAH:



The mechanism underlying the endocrine defects in 21-hydroxylase deficiency is depicted in the image below:



Babies with 21-hydroxylase deficiency typically present around 10-14 days of age with the subset of symptoms due to aldosterone deficiency, cortisol deficiency and androgen excess.

Aldosterone deficiency leads to salt-wasting crises marked by hyponatremia, hyperkalemia, hypotension, and metabolic acidosis.

Cortisol deficiency causes manifestations like hypoglycemia, hyperpigmentation, weight loss, weakness, anorexia, vomiting, and shock.

Androgen excess in 21-hydroxylase deficiency results in virilization of females, leading to female pseudohermaphroditism, and precocious puberty in males. Virilization is usually apparent at birth in females and within 2 to 3 years of life in males.

Without treatment, shock, cardiac arrhythmias, and death may occur within days or weeks.

Other options:

Option A: Male pseudohermaphroditism is seen in CAH due to 17α -hydroxylase deficiency resulting in incomplete virilization. 17α -hydroxylase deficiency is 2nd most common enzyme affected in CAH.

Option C: True hermaphroditism is presently referred to as an ovotesticular disorder of sex development(DSD). Here, both ovarian and testicular tissue is present. It is not due to CAH.


Option D: 46 XY intersex (now called 46 XY DSD) has under virilization of genitalia. Adrenal causes include lipid adrenal hyperplasia, 3β -hydroxysteroid dehydrogenase deficiency, 17-hydroxylase/17,20-lyase deficiency, and not 21 hydroxylase deficiency.

Solution to Question 14:

The given clinical scenario of failure to thrive in late infancy, normal genitalia, hyponatremia, and hyperkalemia suggests aldosterone synthase deficiency. Estimation of aldosterone levels will help in the diagnosis.

Normal genitalia in female children rules out congenital adrenal hyperplasia (CAH) caused by 21-hydroxylase and 3β -hydroxysteroid dehydrogenase deficiency. Hyponatremia and hyperkalemia rule out CAH due to 17-hydroxylase deficiency and 11β -hydroxylase deficiency.

Aldosterone synthase deficiency vs other forms of CAH



©MARROW

Infants with aldosterone synthase deficiency may have severe electrolyte abnormalities with hyponatremia, hyperkalemia, and metabolic

Since cortisol synthesis is unaffected, infants rarely become as ill as untreated infants with salt-losing forms of CAH. So some infants escape diagnosis at birth because of normal cortisol levels.

Adults often are asymptomatic, although they may develop electrolyte abnormalities when depleted of sodium through procedures such as bowel preparation for a barium enema.



The given clinical scenario along with the image showing round face (moon facies), obesity, abnormal masculinization, and purple skin striae are suggestive of Cushing's syndrome. The most common cause of Cushing's syndrome is prolonged exogenous administration of glucocorticoids, not an adrenocortical tumor.

- Prolonged exogenous administration of glucocorticoids
- Excess endogenous cortisol secretion:
- In children <7 years: Most often by a functioning adrenocortical tumor
- In children >7 years: Most often by an ACTH-producing pituitary adenoma (Cushing's disease)

- Obesity or weight gain (most common)
- Moon facies
- Purple skin striae

- Abnormal masculinization (acne, bruising, enlargement of the clitoris in girls)
- Pubertal developmental delay and amenorrhea in girls
- Proximal muscle weakness
- Osteoporosis
- Hypertension
- Hyperpigmentation
- Hypokalemic alkalosis
- Hyperglycemia
- Sepsis

Solution to Question 16:

Given clinical scenario is suggestive of Addison's disease and hypertension is not seen here.

Addison's disease is a primary adrenal insufficiency due to acquired lesions of the adrenal cortex, which prevent the production of cortisol and, often, aldosterone. Hence, due to mineralocorticoid deficiency, there will be hyponatremia, hyperkalemia, and hypotension.

The deficiency of cortisol stimulates the production of adrenocorticotrophic hormone (ACTH). Increased ACTH leads to diffuse hyperpigmentation, including areas not exposed to the sun, such as buccal mucosa and gums.

Addison's disease is characterized by the following:

- Hyponatremia
- Hyperkalemia
- Hypotension
- Hypoglycemia
- Increased ACTH levels
- Metabolic acidosis
- Hyperpigmentation

ACTH stimulation test (cosyntropin) - Low cortisol levels before and after administration of ACTH are suggestive of primary adrenal insufficiency, Addison's disease.

Solution to Question 17:

The most common infectious etiology for adrenal insufficiency in children is meningococemia caused by *Neisseria meningitidis*.

Waterhouse–Friderichsen syndrome:

Adrenal apoplexy due to fulminant bacterial infection (most common *Meningococcus*).

Clinical features: Non-specific like fever, rigors, vomiting, headache, disseminated intravascular coagulation (large petechial rashes in the skin and mucous membrane), hypotension, and septic shock

Petechial rashes in meningococcal infection:



Most common in adrenal insufficiency:

- The most common infectious cause of Addison's disease in India is tuberculosis.
- The most common cause of adrenal insufficiency is the sudden withdrawal of steroids in a patient on long-term steroid therapy.
- The most common cause of primary adrenal insufficiency in children is congenital adrenal hyperplasia (CAH).
- The most common antigen against which antibodies are found in Addison's disease is CYP 21 (21-hydroxylase).

Solution to Question 18:

The given clinical scenario is suggestive of Wolman's disease.

It is a rare lipid disorder caused by a deficiency of lysosomal acid lipase (chromosome 10) that leads to the accumulation of cholesterol esters and triglycerides in most of the visceral organs.

Patients present in the first few weeks of life with failure to thrive, vomiting, abdominal distention, steatorrhea, hepatosplenomegaly, and hyperlipidemia with extensive bilateral calcifications of the adrenal glands.

Solution to Question 19:

The clinical scenario of hyperpigmentation, history of bilateral adrenalectomy and MRI showing enlarged pituitary adenoma points to the diagnosis of Nelson's syndrome.

Nelson's syndrome is a rare complication of total bilateral adrenalectomy (TBA). TBA is a procedure used to control hypercortisolism in Cushing's disease (due to ACTH-secreting pituitary adenoma) when resection of the primary tumor is not possible.

The loss of feedback inhibition of the hypothalamic-pituitary-adrenal axis following TBA increases CRH. This leads to stimulation of ACTH-secreting corticotroph cells, causing enlargement of the original pituitary tumor or formation of a new one.

Nelson's syndrome is associated with a clinical triad of:

- Hyperpigmentation
- Excessive ACTH secretion
- Corticotroph adenoma

Screening is done with fasting plasma ACTH levels and MRI brain performed at regular intervals. Prophylactic pituitary irradiation may be recommended to prevent its incidence. The primary treatment is transsphenoidal surgery. Patients will also require life-long glucocorticoid replacement, as they do not produce cortisol.

The use of steroidogenic inhibitors in the treatment of Cushing's disease has decreased the need for bilateral adrenalectomy these days.

Option B: Steele-Richardson-Olszewski syndrome, also known as progressive supranuclear palsy is a neurodegenerative disease. It affects the brainstem and basal ganglia.

Option C: Hamman-Rich syndrome also known as acute interstitial pneumonia, is a rare and severe form of idiopathic interstitial lung disease.

Option D: Job syndrome, also known as hyper-IgE syndrome is an autosomal dominant immunodeficiency disorder. It occurs as a result of STAT3 mutation leading to deficiency in the IL-17 producing T cells. This results in impaired recruitment of neutrophils to sites of infection.

Solution to Question 20:

The fetal adrenal gland produces a maximum amount of DHEA (dehydroepiandrosterone).

DHEA and DHEA-S serve as estrogen precursors. The maternal adrenal gland does not produce a sufficient amount of DHEA-S. The fetal adrenal glands are quantitatively the most important source of placental estrogen precursors in pregnancy. During most of gestation, the fetal adrenal gland is almost solely dedicated to the production of dehydroepiandrosterone sulfate DHEAS.

Solution to Question 21:

Mothers with pregnancies at risk (previously affected child) of congenital adrenal hyperplasia (CAH) should be started with steroid therapy as soon as pregnancy is diagnosed.

Dexamethasone 20 µg/kg is started to suppress the secretion of steroids by the fetal adrenals (including adrenal androgens) and prevent the virilization of the external genitals in affected females.

Then, chorionic villus biopsy is performed to determine the sex and genotype of the fetus and therapy is continued only if the fetus is an affected female.

Solution to Question 22:

Deficiency of P450 oxidoreductase (POR) along with the given findings are suggestive of Antley-Bixler syndrome.

Features of Antley-Bixler syndrome:

These include deficiencies of 17-hydroxylase and 21-hydroxylase along with other congenital anomalies.

It leads to under virilization of males and virilisation of females via an alternate pathway. Virilization of mother is seen because of decreased placental aromatase activity leading to excess androgens in fetus and mother.

Congenital anomalies seen are as follows:

- Head and face: Craniosynostosis, brachycephaly, frontal bossing, severe midface hypoplasia, with proptosis and choanal atresia.
- Limbs: Humero-radial synostosis; medial bowing of the ulnas; long, slender fingers, with camptodactyly and anterior bowing of the femurs.
- Malformations of the heart and kidneys.



Disorders of the Pituitary Gland

Question 1:

Which of the following is the most common tumor associated with growth hormone deficiency in children?

- a) Prolactinoma
- b) Craniopharyngioma
- c) Pinealoma
- d) Somatotropinoma

Question 2:

Asha was being bullied about her height in school. The concerned parents bring her to the pediatrician who plots the growth chart as shown below. On evaluation, growth hormone levels are found to be high and insulin-like growth factor levels are low. Which among the following could be a cause?



- a) Turner's Syndrome
- b) Pituitary Dwarfism
- c) Laron Syndrome
- d) Maternal Deprivation Dwarfism

Question 3:

Which of the following is the most common genetic cause for multiple pituitary hormone deficiency?

- a) Mutations of GNAS1 gene
- b) Mutations of c-MYC gene
- c) Mutations of WT-1 gene
- d) Mutations of PROP1 gene

Question 4:

You come across a child with growth hormone deficiency during your ward rounds. Which of the following features would you not expect in this child?

- a) History of short stature at birth
- b) High-pitched voice
- c) Normal intelligence
- d) Proportionate short stature

Question 5:

You are evaluating children with short stature on whether they are eligible to receive growth hormone therapy as per FDA guidelines. In which of the following children would you prescribe it?

- a) An 8 year old with Laron dwarfism
- b) A neonate diagnosed with congenital hypothyroidism
- c) A 7 year old with Chronic kidney disease
- d) A 6 year old diagnosed with Maternal deprivation dwarfism

Question 6:

Chotu, a 5-year-old boy is on regular outpatient follow-up for growth hormone deficiency diagnosed 2 weeks back. He is planned for recombinant growth hormone therapy. All of the following are possible side effects except-

- a) Type 1 diabetes mellitus
- b) Pseudotumor cerebri

- c) Slipped capital femoral epiphysis
- d) Gynecomastia

Question 7:

The most common pituitary adenoma diagnosed in childhood is _____.

- a) Prolactinoma
- b) Corticotropinoma
- c) Somatotropinoma
- d) Gonadotropinoma

Question 8:

Akash is the tallest kid in his 5th-grade class. While at a routine visit for a booster immunization, his pediatrician plots his height on a growth chart and finds him to be above 3-SD for his age. Which of the following can be an endocrine cause for his height?

- a) Sotos syndrome
- b) McCune-Albright syndrome
- c) Fragile X syndrome
- d) Beckwith-Wiedemann syndrome

Question 9:

"Weak giants" are produced by _____.

- a) Hypothalamic tumors
- b) Craniopharyngiomas
- c) Pituitary adenomas
- d) Pinealomas

Question 10:

A young boy is diagnosed to have central diabetes insipidus. A lesion in which of the following nuclei is likely to cause this condition?

- a) Dorsomedial nuclei

- b) Supraoptic and Paraventricular nuclei
- c) Median preoptic nuclei
- d) Ventromedial nuclei

Question 11:

A 4-year-old boy was brought by his mother with new-onset nocturnal bedwetting. She also reported, he has been drinking more water for the last 3-4 months. 24-hour urine volume was 4 liters and the urine osmolality was low. ADH assays recorded a value was 0.7 pg/mL. T1 weighted MRI brain indicated an absence of a pituitary bright spot. What is the treatment of choice for his condition?

- a) Leuprolide
- b) Thiazide diuretics
- c) Desmopressin
- d) Terlipressin

Question 12:

A 6-year-old boy diagnosed with Wolfram syndrome is admitted to the pediatric ward of your hospital. Which of the following features would you expect to find in this child?

- a) Diabetes mellitus type 2
- b) Syndrome of inappropriate ADH secretion
- c) Conductive hearing loss
- d) Optic atrophy

Question 13:

While working in a tertiary care center, a 5-year-old girl presents with polyuria, polydipsia, and nocturnal enuresis for the past 1 year. A water deprivation test is done and shows low urine osmolality on stopping the intake of water. On desmopressin administration, there is a 7% rise in urine osmolality. The treatment of this condition uses all of the following except?

- a) Thiazide diuretics
- b) Increased salt intake
- c) Amiloride
- d) Indomethacin

Answer Key

Question No.	Correct Option
1	b
2	c
3	d
4	a
5	c
6	a
7	a
8	b
9	c
10	b
11	c
12	d
13	b

Detailed Explanations

Solution to Question 1:

The most common tumor associated with growth hormone (GH) deficiency in children is craniopharyngioma.

Growth hormone deficiency can be caused by congenital CNS malformations, genetic defects or acquired neurological insults.

Note:

- Most common cause of GH deficiency: Idiopathic
- Most common infection causing GH deficiency in India: Tuberculosis & Histoplasmosis
- Most common infiltration causing GH deficiency in India: Histiocytosis & Sarcoidosis.
- Trophic hormone failure with pituitary compression or destruction: GH& FSH& LH& TSH& ACTH.
- Most common tumors associated with central diabetes insipidus: Pinealomas, Germinomas

Solution to Question 2:

The presence of high GH levels and low insulin-like growth factor (IGF-1) levels point towards Laron syndrome/dwarfism. It is caused by end-organ resistance to growth hormone (GH insensitivity) due to the GH receptor mutation. It is an autosomal-recessive disorder with clinical features as those seen in pituitary dwarfism.

Here, growth hormone is produced but is unable to act on the target tissues. Thus, the resting and stimulated GH levels are high and IGF-1 levels are low.

In Turner's syndrome (Option A), loss of the short arm of the X chromosome is implicated in dwarfism. The short stature occurs due to a state of GH resistance, relative deficiency, and decreased utilization.

Pituitary dwarfism (Option B) is a condition with GH deficiency.

Maternal deprivation dwarfism (Option D) is due to psychosocial causes such as emotional deprivation and neglect. There is an inadequate response to GH stimulation tests. The mechanism is not fully understood. No history suggestive of disturbed maternal or family relationships rules this out.

Solution to Question 3:

The most common genetic cause for multiple pituitary hormone deficiencies (MPHD)/panhypopituitarism is the mutation of the PROP1 gene.

LHX3 and GLI2 are other genes involved in MPHD.

GNAS1 mutations cause McCune Albright Syndrome.

c-MYC gene is mutated in Burkitt's lymphoma

WT-1 gene is mutated in Wilms tumors.

Solution to Question 4:

Children with growth hormone deficiency usually have normal growth at birth. Growth retardation (decline in growth velocity) becomes apparent around 1 year of age.

Growth in-utero is dependant on genetic, environmental, nutrition and growth factors. GH levels peak at mid-gestation, increase at birth, and then reduce by 2 weeks of life. At around 6 months of age, GH levels reach pre-pubertal levels and from there on have a major influence on growth. Thus the deficiency becomes apparent after a significant period of GH-dependent growth has occurred.

Characteristic features of growth hormone deficiency (Pituitary Dwarfism):

- Normal growth at birth with growth retardation becoming apparent around 1 year of age
- Proportionate short stature
- Hypoglycemic seizures in newborn

- Small larynx with a high-pitched voice
- Small genitalia (micropenis)
- Delayed skeletal maturation
- Delayed dentition
- Delayed puberty (sexual infantilism)
- Doll-like facial appearance: small, round facies, frontal prominence, depressed nasal bridge (saddle nose), eyes may bulge, underdeveloped mandible and chin, normal head circumference but craniofacial disproportion due to small facies
- Short neck
- Normal intelligence.

Solution to Question 5:

Chronic kidney disease is an FDA-approved indication for growth hormone (GH) therapy.

Renal failure is a state of GH resistance. Moderate-to-severe growth retardation in chronic kidney disease is associated with increased morbidity and mortality. GH therapy with recombinant growth hormone has been shown to improve catch-up growth and to allow the child to achieve normal adult height; the final adult height is still significantly below the genetic target.

FDA-approved indications for growth hormone (GH) treatment-

- GH deficiency
- Turner's Syndrome
- Chronic Renal Failure before Transplantation
- Small for Gestational Age / Short Stature
- Idiopathic Short Stature
- Noonan Syndrome
- Prader-Willi Syndrome
- AIDS-related wasting
- Malabsorption associated with the short-bowel syndrome

Solution to Question 6:

The use of recombinant growth hormone is associated with the risk of type 2 diabetes mellitus (not type 1 diabetes mellitus).

Other adverse effects:

- Pseudotumor cerebri
- Slipped capital femoral epiphysis

- Gynecomastia
- Worsening of scoliosis

Children who have been treated with rhGH have a higher risk of developing hemorrhagic strokes in adulthood.

Some studies show an increased risk of new neoplasms in cancer survivors.

Solution to Question 7:

The most commonly diagnosed adenomas during childhood are prolactinoma > corticotropinoma > somatotropinoma.

Note:

Hypothalamic hamartomas that secrete excess gonadotropin-releasing hormone can cause precocious puberty.

Solution to Question 8:

McCune–Albright syndrome (endocrine dysfunction affecting multiple glands) is an endocrine cause of tall stature in childhood. The triad of McCune–Albright syndrome is precocious puberty, polyostotic fibrous dysplasia, and café au lait macules

Causes of childhood tall stature:

Non-endocrine causes	Endocrine causes
Familial (constitutional) tall stature	Excess GH secretion(pituitary gigantism)
Cerebral gigantism (Sotos syndrome)	McCune–Albright syndrome
Marfan syndrome	Precocious puberty
Fragile X syndrome	Hyperthyroidism
Homocystinuria	
Klinefelter syndrome(XXY)	
XYY syndrome	
Beckwith–Wiedemann syndrome	

Solution to Question 9:

"Weak giants" are due to hyperpituitarism that is most commonly produced by pituitary adenomas.

Hyperpituitarism causes growth hormone (GH) excess, leading to tall stature (giants) with weak muscles. Overproduction of GH results in gigantism in children (before fusion of bone growth plate) and acromegaly in adults.

Clinical features include:

- Gigantism, prognathism (protrusion of jaw), hypertrophy of frontal bones
- Proximal myopathy (though there is muscle hypertrophy, it is weakened due to atrophy of type II muscle fibers)
- Acral enlargement in hands and feet (due to thickening of soft tissues)
- Visceromegaly: Enlargement of most organs especially the tongue, thyroid, salivary gland, liver, spleen, kidney, and prostate
- Local effects of the adenoma: Headache, visual field defects, raised ICT

Note: Although hypothalamic hamartomas can also produce gigantism, the most common cause of gigantism is pituitary adenomas.

Solution to Question 10:

Lesions in the supraoptic and paraventricular nuclei of the hypothalamus can cause central diabetes insipidus.

These nuclei synthesize vasopressin, a 9-amino-acid peptide with antidiuretic and vascular pressor activity. It is transported to the posterior pituitary via axonal projections, where it is stored for release into the systemic circulation.

Diabetes insipidus (DI) is characterized by pathologic polyuria and polydipsia ($>2 \text{ L/m}^2/24 \text{ hr}$ or 5 mL/kg/hr). It can result from either:

- Vasopressin deficiency (decreased anti-diuretic hormone production) causes central DI.
- Vasopressin insensitivity at the level of the kidney causes nephrogenic DI.

Solution to Question 11:

The given clinical scenario is suggestive of central diabetes insipidus. The treatment of choice is desmopressin. It is a vasopressin analog (V2 receptor agonist). It can be administered intranasally, or subcutaneously.

Option A: Leuprolide is a GnRH (gonadotropin-releasing hormone) analog used in the treatment of prostate cancer, breast cancer, and endometriosis.

Option B: Thiazides paradoxically exert an antidiuretic effect in DI but are not the drug of choice. They are generally used in nephrogenic DI.

Option D: Terlipressin is a long-acting analog on vasopressin that acts on V1 receptors of vascular smooth muscle. It is used as a vasopressor in shock and acute variceal bleeding.

Solution to Question 12:

Optic Atrophy is seen in Wolfram syndrome. The components of the syndrome are represented by the acronym DIDMOAD:

- Diabetes Insipidus
- Diabetes Mellitus (type 1)
- Optic Atrophy
- Neural Deafness

It is caused by mutations of the WFS1 gene.

Solution to Question 13:

The given clinical scenario is suggestive of nephrogenic diabetes insipidus. The treatment includes pharmacological treatment combined with salt restriction and not increased salt intake.

Treatment of nephrogenic diabetes insipidus:

- Adequate water intake along with a low sodium diet
- Thiazide diuretics: Acts by
- Volume contraction and increases the responsiveness of other parts of the tubule
- Increase the AQP2 channels independent of vasopressin
- Potassium-sparing diuretics (triamterene and amiloride): Amiloride is the drug of choice for lithium-induced DI
- Prostaglandin synthesis inhibitor (indomethacin)

Disorders of Puberty

Question 1:

You are suspecting a case of delayed puberty in a 15-year-old boy. To assess puberty in this patient, what is the first sign you would look for?

- a) Increase in testicular volume
- b) Penile growth
- c) Pigmentation of the scrotum
- d) Pubarche

Question 2:

As an intern, you are currently posted in the Paediatric endocrinology OPD where you see multiple cases of girls having precocious puberty. The condition is defined by the onset of secondary sexual characteristics before the age of _____.

- a) 5 years
- b) 7 years
- c) 8 years
- d) 9 years


Question 3:

A 6-year-old girl was brought to the OPD by her mother who is concerned about her significant increase in height compared to her classmates in the last 2 years. On examination, the girl shows breast development corresponding to B2 Tanner stage. What is the most common brain lesion causing this condition?

- a) Pituitary adenoma
- b) Tuberous sclerosis
- c) Astrocytoma
- d) Hypothalamic hamartoma

Question 4:

A young girl was referred to the paediatrician with stage IV breast development and growth acceleration seen at the age of 5 years. Her estradiol concentrations were elevated. The sagittal MRI of the brain is as shown below. What is the treatment of choice for this condition?



- a) Cyproterone
- b) Danazol
- c) Medroxyprogesterone
- d) GnRH agonists

Question 5:

A primary school teacher notices bloodstains on the skirt of her second-grade student. She immediately informs her parents and accompanies the child to the doctor on campus. On examination, the 7-year-old girl had blood-stained vaginal discharge, enlarged breasts and areola. Which of the following is not a cause for this condition?

- a) Hypothalamic hamartoma
- b) Hypothyroidism
- c) McCune-Albright syndrome
- d) Prader-Willi syndrome

Question 6:

An incessantly crying 5-year-old girl is brought to the OPD by her father. You observe that she needs support to walk and her trunk is not symmetrically aligned. There is a history of recurrent fractures in her limbs. She also has blood-stained vaginal discharge from a few days. On examination, the girl appears as shown in the image below. What is the most probable diagnosis?



- a) Prader-Willi Syndrome
- b) Reifenstein Syndrome
- c) McCune-Albright syndrome
- d) Polyostotic fibrous dysplasia

Question 7:

A 15-year-old boy on examination is found to have genital development corresponding to Tanner stage 1 and his height is below the 3rd Percentile on growth charting. What is the most common cause for this condition?

- a) Kallman syndrome
- b) Klinefelter syndrome
- c) Constitutional delay
- d) Familial short stature

Question 8:

Breast and papillae appear as a small mound on examination of a young girl. If you are suspecting a case of delayed puberty, it is defined as a lack of development of secondary sexual characters beyond the age of _____.

- a) 15 years
- b) 14 years
- c) 13 years
- d) 12 years

Question 9:

A child presents with features of delayed puberty. On investigating, he has decreased levels of LH, FSH, and testosterone. Which of the following is the most likely diagnosis?

- a) Klinefelter's syndrome
- b) Kallman's syndrome
- c) Androgen insensitivity syndrome
- d) Testicular infection

Question 10:

All of the following are features of Laurence-Moon-Biedl syndrome except _____.

- a) Obesity
- b) Mental retardation
- c) Polydactyly
- d) Hypotension

Question 11:

A female child presents with bilateral inguinal masses, which were thought to be hernias but were found to be testes in the inguinal canals. Which karyotype would you expect to find in the child?

- a) 46,XX
- b) 46,XY
- c) 47,XXY
- d) 45,XO

Question 12:

An infant was found to have elevated levels of LH. The baby has a 46, XY chromosome pattern on karyotyping and the vagina is short and ends blindly in a pouch. Which of the following is false in the condition suffered by the infant?

- a) Internal female genitalia is absent
- b) Internal female genitalia is normal
- c) Affected individuals have female phenotype
- d) High chance of testicular malignancy

Question 13:

A 31-year-old female delivers a term baby. On examination, it has a micropenis and unpalpable testis. She gives a history of acne from 16 weeks of gestation which worsened throughout pregnancy and was treated after delivery. The baby was diagnosed to have female pseudohermaphroditism. Which of the following statements are right about the given condition?

- a) XY genotype, female external genitalia
- b) Female external genitalia and male internal genitalia
- c) XX genotype, male external genitalia
- d) XY genotype, male external genitalia

Question 14:

What is true about ovotesticular disorders of sex development?

- a) Karyotype is 46 XY.
- b) Ovotestis is seen in the gonadal biopsy
- c) The response of testosterone to hCG is increased.
- d) Uterus is usually absent.

Question 15:

A 3-year-old girl was brought by her mother with complaints of enlarged clitoris. Karyotype revealed 46XX. What is the likely cause?

- a) 21 hydroxylase deficiency
- b) 11 hydroxylase deficiency
- c) 3 beta dehydrogenase deficiency
- d) 2, 3 lyase deficiency

Answer Key

Question No.	Correct Option
1	a
2	c
3	d
4	d
5	d
6	c
7	c
8	c
9	b
10	d
11	b
12	b
13	c
14	b
15	a

Detailed Explanations

Solution to Question 1:

The first sign of puberty in males is the increase in testicular volume.

In males at 11 to 12 years, the growth of the testis (>4 mL in volume) and thinning of the scrotum occurs, which are the first signs of puberty. This is followed by the pigmentation of the scrotum, growth of the penis, and pubarche.

Growth of testis > pubarche > adrenarche > growth spurt (TPAG).

In females at 10 to 11 years, the breast bud (thelarche) appears, which is the first sign of puberty followed by the appearance of pubic hair (pubarche), growth spurt, and menarche.

Thelarche & pubarche & growth spurt & menarche (TPGM).

Solution to Question 2:

Precocious puberty is defined as the onset of secondary sexual characteristics before the age of 8 years in girls and 9 years in boys.

Depending on the primary source of hormonal production, precocious puberty can be classified in the following manner:

Note:

Central Precocious Puberty(more common) & Peripheral Precocious Puberty

Central Precocious Puberty(Gonadotropin-dependent puberty)	Peripheral Precocious Puberty(Gonadotropin independent)
Due to the excess secretion of gonadotropins (FSH and LH) from pituitary	Due to excess secretion of testosterone or estrogen from the gonads/adrenals.
Always isosexual (appearance of phenotypically appropriate secondary sexual characteristics before age 8 years in girls and before 9 years in boys)	Can be isosexual or heterosexual (appearance of secondary sex characters of the opposite sex).
GirlsIdiopathic (most common)BoysCNS pathology (most common)	GirlsEstrogenic ovarian cysts (isosexual)Congenital adrenal hyperplasia (CAH) (testosterone increases; heterosexual)BoysCAH (isosexual; most common)

Solution to Question 3:

Hypothalamic hamartoma, a neuronal migration defect, is the most common cause of organic central precocious puberty. The onset of breast development before the age of 8 years in females and onset of testicular development (>4 mL) before the age of 9 years in males is suggestive of central precocious puberty.

It is a congenital malformation that consists of ectopically located neural tissue characterised by-

- Early-onset of puberty
- Seizures associated with unnatural crying or laughing episodes are known as gelastic seizures.

Tuberculous meningitis, tuberous sclerosis, and astrocytomas can also cause central precocious puberty.

Solution to Question 4:

A pedunculated mass in the floor of the 3rd ventricle as shown in the MRI, with elevated levels of estrogen is suggestive of a Hypothalamic Hamartoma, which is the most common brain lesion causing central precocious puberty and is treated with GnRH agonists.

Treatment of Hypothalamic Hamartoma-

- GnRH agonists (depot-forms or implant) are the treatment of choice of tumour-induced Central precocious puberty
- In patients with hypothalamic hamartoma and associated intractable gelastic or psychomotor seizures, stereotactic radiation therapy (Gamma Knife surgery) is more effective and has fewer risks than neurosurgical intervention.

Long-acting formulations of gonadotropin-releasing hormone (GnRH) agonists are the preparations of choice for the treatment of central precocious puberty as they maintain fairly constant serum concentrations of the drug for weeks or months.

Solution to Question 5:

Vaginal bleeding with well-developed breasts before the age of 8 years is suggestive of precocious puberty. Prader–Willi syndrome is one of the causes of delayed puberty in girls (not precocious puberty), where there is hypogonadotropic hypogonadism.

Hypothalamic hamartoma, McCune–Albright syndrome and hypothyroidism are causes of precocious puberty.

Note:

In prolonged hypothyroidism, adolescents typically have delayed puberty. Younger children might present with pseudo-precocious puberty, characterized by breast development and vaginal bleeding in girls and macroorchidism in boys.

Elevated levels of TSH appear to interact with the FSH receptor (specificity spillover), thus inducing FSH-like effects in the absence of LH effects on the gonads. The FSH-like effect suffices to induce estradiol secretion by the ovaries, whereas in males, testicular enlargement occurs without substantial testosterone secretion.

Solution to Question 6:

Precocious puberty(vaginal discharge), bony lesions (polyostotic fibrous dysplasia), and cafe au lait spots, as shown in the picture, are features of McCune–Albright syndrome. There is pain due to the multiple bony lesions and repeated fractures may result in incessant crying in the children and may require regular visits to the orthopaedics department.

McCune–Albright syndrome is a classical cause of peripheral precocious puberty caused by a mutation of GNAS1 gene. It affects the bones, skin, and several endocrine tissues.

Fibrous dysplasia of multiple bones (polyostotic) represents a major cause of morbidity in this syndrome. The base of the skull and the proximal femurs are most commonly involved, but any bone can be affected. A CT scan is usually recommended even in absence of deformities. The prognosis is favourable for longevity, but deformities, repeated fractures, pain, and occasional cranial nerve compression may result from the bony lesions.

Cardinal features:

- Polyostotic fibrous dysplasia
- Cafe au lait macules
- Endocrine abnormalities: hyperthyroidism, acromegaly, adrenal adenomas, and autonomous ovarian hyperfunction, leading to precocious puberty
- Predominantly affects girls

Solution to Question 7:

Failure of development of pubertal features by the age of 14 years in male is suggestive of Delayed Puberty.

The most common cause of delayed puberty in males and females is a constitutional delay in growth and puberty. Lack of pubertal changes in boys by the age of 14 years is suggestive of delayed puberty. Delayed puberty is more common in boys than in girls.

Solution to Question 8:

Delayed puberty in girls is defined as lack of secondary sexual characteristics by the age of 13 years or absence of menarche by the age of 16 years or within 3 years of pubertal onset.

Solution to Question 9:

This patient has low levels of luteinizing hormone (LH), follicle-stimulating hormone (FSH), and testosterone pointing towards hypogonadotropic hypogonadism and Kallman's syndrome is the only condition in the given choices causing hypogonadotropic hypogonadism.

Causes of delayed puberty:

Hypogonadotropic hypogonadism (Secondary/central)


Hypogonadism is due to the impaired secretion of gonadotropins (FSH and LH) by the pituitary gland and the resultant lack of sex steroid production

- Malnutrition, systemic diseases
- Kallmann (anosmia), Prader–Willi syndrome (mental retardation, obesity, low intelligence)

Hypergonadotropic hypogonadism (Primary gonadal failure)

Hypogonadism is due to an impaired response of the gonads to the gonadotropins (FSH and LH), which in turn leads to a lack of sex steroid production and elevated gonadotropin levels (as an attempt of compensation by the body)

- Sex chromosome anomalies (Turner's, Klinefelter's syndrome)
- Androgen insensitivity syndrome
- Ovotesticular dysgenesis



Solution to Question 10:

Hypertension is seen in Laurence–Moon–Biedl syndrome.

Laurence–Moon–Bardet–Biedl is an autosomal-recessive disorder of hypogonadotropic hypogonadism.

Characteristic features:

- Polydactyly
- Obesity
- Retinitis pigmentosa
- Male hypogonadism
- Hypertension
- Mental retardation

Solution to Question 11:

Here, this baby girl with bilateral inguinal masses (testis) probably has androgen-insensitivity syndrome, where karyotype is 46,XY.

Androgen-insensitivity syndrome is an X-linked recessive disorder caused by mutations in the androgen receptor gene that makes XY fetuses insensitive to androgens. The patients phenotypically resemble females because a part of the testosterone is converted to estrogen, which leads to the development of female external genitalia, normal development of breast, and female habitus.

Patients with Klinefelter's syndrome have a karyotype of 47, XXY and can have undescended testis, but they have male external genitalia where the phallus tends to be smaller than average.

Characteristic features:

- Most common forms of male disorders of sexual differentiation (DSD).
- 46 XY male child with female phenotype, present with primary amenorrhoea.
- Normal male gonads, mostly undescended ones, can present as inguinal masses or hernia.
- Here, as testosterone receptors are defective, they do not act on tissues, leading to the absence of secondary sexual characters (absence of pubic hair and axillary hair).

Solution to Question 12:

An infant with a 46, XY chromosome pattern and female external genitalia are suggestive of Androgen-insensitivity syndrome(AIS) where there is absent internal female genitalia.

These patients have absent internal female genitalia with normal female external genitalia with the vagina ending blindly, in a pouch and normal male gonads in the form of undescended testis (mostly) which can present as inguinal masses.

In a male child (46, XY), testicular hormones are resistant to receptors, and a part of the testosterone is converted to estrogen leading to the development of female external genitalia and eventually normal development of breast, and female habitus.

Features of Androgen insensitivity syndrome at different age groups-

- In infants, elevated LH levels should suggest the diagnosis of AIS
- Prepubertal females with this disorder are often detected when inguinal masses which prove to be testes
- In older children and adults, amenorrhea is the usual presenting symptom.

Note:

Reifen-Stein syndrome: Males with various forms of ambiguous genitalia and undervirilization (hypogonadism, severe hypospadias, and gynecomastia). They are phenotypically normal-appearing males with infertility.

Solution to Question 13:

Female pseudohermaphrodites are genetic females (XX genotype) with male external genitalia.

Causes:

- Enzyme deficiencies: 21 hydroxylase, 11 hydroxylase, and 3-beta-hydroxysteroid dehydrogenase (HSD) deficiencies
- Maternal androgen excess

Male pseudohermaphrodites are genetic males (XY genotype) with female external genitalia.

Causes:

- Enzyme deficiencies: 17 hydroxylase and 3-beta HSD deficiencies
- Defect in androgen action

True hermaphroditism/ovotesticular dysgenesis: An intersex condition in which an individual is born with ovarian and testicular tissue.

Solution to Question 14:

Ovotesticular disorder of sex development (DSD) is characterized by the presence of both ovarian and testicular tissue in the same individual. An ovotestis is present in approximately 2/3 of affected individuals.

Most affected individuals have a 46, XX chromosomal karyotype, which normally results in female sexual development. In about 10% of patients, testicular tissue in an individual with a 46, XX karyotype is present as a result of a translocation of the SRY gene on the Y chromosome to the X chromosome or another chromosome. There is a defect in R-spondin1 protein encoded by the RSPO1 gene, described in 46, XX ovotesticular DSD.

An abnormal vagina and an underdeveloped uterus are usually present. The testes are usually undescended, and the penis may present with hypospadias. Upon reaching puberty, breast development, feminization, and menstruation may occur. Most affected individuals are infertile, but ovulation or spermatogenesis is possible. The response of testosterone to hCG is normal or reduced.

Solution to Question 15:

The given clinical scenario of a girl (46XX) with an enlarged clitoris is highly suggestive of congenital adrenal hyperplasia (CAH) due to 21-hydroxylase deficiency.

Congenital adrenal hyperplasia/adrenogenital syndrome is a family of autosomal-recessive disorders caused by the deficiency of enzymes involved in the synthesis of steroid hormones in the adrenal cortex. The majority of cases of CAH are caused by 21-hydroxylase deficiency. This deficiency leads to a lack of synthesis of cortisol and aldosterone, resulting in the salt-wasting form of the disease, which is the most severe form.

Severe 21-hydroxylase deficiency presents with symptoms including weight loss, dehydration, weakness, hypotension, and electrolyte imbalances. Accumulated steroid precursors like

17-hydroxyprogesterone, are shunted into the pathway of androgen synthesis leading to elevated levels of androstenedione and testosterone. This can result in abnormal genital development in affected female fetuses during early gestation.

Females may have masculinized external genitalia. This can include enlargement of the clitoris, partial or complete labial fusion, and a shared opening of the vagina and urethra (urogenital sinus).

Untreated individuals develop signs of androgen excess after birth. These signs include accelerated skeletal maturation, premature epiphyseal closure leading to stunted adult stature, excessive muscular development, the appearance of pubic and axillary hair, acne, and deepening of the voice. In males, enlargement of the penis, scrotum, and prostate. In females, the clitoris may become further enlarged.

Diagnosis is confirmed by elevated 17-hydroxyprogesterone levels. Glucocorticoid replacement with hydrocortisone is done. In a salt-wasting crisis, fludrocortisone supplementation is given. Vaginoplasty and clitoral recession are done for females with ambiguous genitalia.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Paediatric Hemato-oncology

Question 1:

Cancer-'The emperor of all maladies spares no one'. Which of the following is the most common childhood malignancy?

- a) Neuroblastoma
- b) Astrocytoma
- c) Acute lymphoblastic leukemia
- d) Acute myeloid leukemia

Question 2:

You are asked to review an article titled, 'Childhood cancers- a menace'. Which of the following statements is wrong and needs to be edited out?

- a) Acute lymphoblastic leukemia is the most common type of leukemia in children with Down syndrome.
- b) The most common intra-abdominal malignancy is Wilm's tumor.
- c) The most common extracranial solid malignancy is neuroblastoma.
- d) The most common brain tumors in children are infratentorial.

Question 3:

What is the most common leukemia you would encounter in a child with the following findings?



- a) Acute myeloid leukemia
- b) Acute lymphoblastic leukemia
- c) Chronic myeloid leukemia
- d) Chronic lymphoid leukemia

Question 4:

Jeevit, a 7-year-old boy brought to you by his mother, complains of bone pain so severe that he cannot sleep. Investigations reveal:

- a) TdT+, HLA DR+, CD20+, Cy Ig +
- b) CD19+, CD20+, CD24+, Surface Ig+
- c) CD3+, CD4+, CD5+, CD34+
- d) CD2+, CD7+, TdT+

Question 5:

You are posted in the pediatric hematology department as a first-year postgraduate. All of the following cases you encounter are at an increased risk of developing leukemias except:

- a) 1,4
- b) 2,5
- c) 1,3,4
- d) 5 only

Question 6:

To diagnose a 5-year-old child with acute lymphoblastic leukemia, a pathologist has to see at least ____ percentage of blast cells in a bone marrow biopsy.

- a) 15
- b) 10
- c) 25
- d) 30

Question 7:

A 4-year-old child who was being evaluated for pancytopenia is diagnosed with acute lymphoblastic leukemia. Which of the following findings would indicate a poor prognosis?

- a) Minimal residual disease $> 0.01\%$ at the end of induction therapy
- b) $t(12,21)$ on cytogenetic study
- c) Early B- cell immunophenotype.
- d) Initial WBC count of 20,000 cells/mm³

Question 8:

A 14-year-old boy is rushed to the casualty due to difficulty in breathing and swelling of the face and neck. Clinically, he has features suggestive of superior vena cava obstruction. Chest X-ray image shows superior mediastinal widening. Which of the following is the most likely diagnosis?

- a) B-cell acute lymphoblastic leukemia
- b) T-cell acute lymphoblastic leukemia
- c) Chronic myeloid leukemia
- d) Chronic lymphocytic leukemia

Question 9:

A 2-month-old infant with seizures and failure to thrive has cutaneous nodules and hepatosplenomegaly. On investigations, CBC reveals hyperleukocytosis with a leukocyte count of 1,10,000 cells/mm³. Peripheral smear and immunophenotyping studies reveal large,

irregular lymphoblasts, with a phenotype negative for the CD10 (common ALL antigen) marker. On a cytogenetic study which of the following is most likely to be seen?

- a) t(11, 14)
- b) t (4,11)
- c) t (12,21)
- d) t (15,21)


Question 10:

A mother has noticed recently that her child refuses to be touched, seems to be in distress, and cries inconsolably. She brings her child to you who was investigated thoroughly and was diagnosed to have acute lymphoblastic leukemia. Which of the following drugs can be used to treat her?

- a) Imatinib
- b) L-asparaginase
- c) Tretinoin
- d) Cladribine

Question 11:

You are posted in the hematology department as a pathologist. You are asked to report the bone marrow biopsy slide given below, which belongs to a 3-year-old child who is being evaluated for pancytopenia. To make a diagnosis of acute myeloid leukemia, what percentage of such cells should you visualize at the least?



- a) 10
- b) 20
- c) 25
- d) 30

Question 12:

A 3-year-old boy was brought to the pediatric OPD with complaints of persistent fever for the past 1 week. His CBC is given below:

- a) t(8;21)
- b) t(15;17)
- c) Inv 16
- d) Monosomy 7

Question 13:

A child with a violaceous, raised nodular skin lesion which on biopsy is found to be infiltrated by myeloblasts is likely suffering from which type of Acute myeloid leukemia (AML)?

- a) AML M2
- b) AML M3
- c) AML M7
- d) AML M1

Question 14:

Erythemic myelosis is _____.

- a) AML M2
- b) AML M5
- c) AML M7
- d) AML M6

Question 15:

A neonate with trisomy 21 has anemia, petechiae, hepatosplenomegaly, and no life-threatening complications on examination. CBC reveals high leucocyte count and peripheral smear shows numerous myeloblasts. The genetic makeup of blast cells when studied reveals GATA-1 mutations. At 3 months of age, his CBC counts come back to normal and there are no blast cells in his peripheral smear. As his treating physician, what would you recommend?

- a) Chemotherapy
- b) Radiotherapy
- c) Bone marrow transplant
- d) A close follow-up

Question 16:

An 18-month-old child presents to the pediatric OPD with complaints of fever and rashes for the past 1 week after extensive workup gets diagnosed as Juvenile chronic myeloid leukemia. Which of the following findings is not consistent with this baby's diagnosis?

- a) The genetic study may reveal NF-1 mutation
- b) The presence of Philadelphia chromosome on FISH
- c) The bone marrow shows a myelodysplastic pattern with $\geq 20\%$ blasts.
- d) The peripheral smear reveals nucleated RBCs

Question 17:

The most common non-Hodgkin's lymphoma in children is _____.

- a) B-cell acute lymphoblastic leukemia (B-ALL)
- b) Mantle cell lymphoma
- c) Diffuse large B-cell lymphoma (DLBCL)
- d) Burkitt lymphoma

Question 18:

Which of the following statements best describe the behavior of Non-Hodgkin lymphoma in a 10-year-old child?

- a) Is dominantly nodal
- b) Is of high grade and aggressive
- c) Has variable growth fraction
- d) Has poor outcome


Question 19:

A 6-year-old boy is brought with complaints of abdominal distension, fatigue, and night sweats for the past 4 months. He also has a 2-month history of postprandial fullness and recurrent non-bilious vomiting. On examination, he is pale, has hepatosplenomegaly and shifting dullness. A CT scan shows mesenteric and retroperitoneal lymph node enlargement. A biopsy specimen of the mesenteric lymph node reveals tumor cells with high mitotic index and starry sky pattern. The cytogenetic study revealed t(8;14). To stage the disease, which of the following systems would you use?

- a) Ann-Arbor staging
- b) Binet staging
- c) St Jude staging
- d) RAI staging

Question 20:

A lymph node biopsy of a 9-year-old child with cervical lymphadenopathy showed the following picture. The diagnosis is_____.



- a) Burkitt Lymphoma
- b) Hodgkin's disease
- c) Acute lymphoblastic leukemia
- d) Chloroma

Question 21:

A 12- year-old female comes to you with a flu-like illness and left supraclavicular swelling of uncertain nature. Her chest X-ray shows left mediastinal enlargement. CT scan of the chest and abdomen reveals the presence of enlarged lymph nodes in the neck, mediastinum, and liver, and an FDG-PET/CT scan shows multiple scattered consolidation lesions involving also the bones. The diagnosis of classical Hodgkin's lymphoma, nodular sclerosis subtype, is made on cervical lymph node biopsy. According to Ann-Arbor's staging system, how will you stage her disease?

- a) 2
- b) 1
- c) 4
- d) 3

Question 22:

The following parameter in ALL indicate a poor prognosis _____.

- a) Age $\geq$ 10 years

- b) Leukocyte count $< 50,000/\text{mm}^3$
- c) Hyperdiploidy
- d) Trisomy of chromosomes 4, 10, and 17

Question 23:

Sanctuary sites of acute lymphoblastic leukemia are_____.

- a) Central nervous system
- b) Testis
- c) Both A and B
- d) None of the above

Question 24:

A 4-year-old boy presents with pancytopenia and hepatosplenomegaly. Bony tenderness is also present. He is most likely suffering from the following type of ALL_____.

- a) B cell
- b) T cell
- c) Both B and T cell
- d) None of the above

Answer Key

Question No.	Correct Option
1	c
2	b
3	b
4	a
5	b
6	c
7	a
8	b
9	b

10	b
11	b
12	d
13	a
14	d
15	d
16	b
17	d
18	b
19	c
20	b
21	c
22	a
23	c
24	a

Detailed Explanations

Solution to Question 1:

Leukemias are the most common childhood malignancies. Among these, acute lymphoblastic leukemia (ALL) is the most common, accounting for about 77% of childhood leukemias.

Most commons in childhood malignancies:

- Second most common malignancy in childhood: Brain tumor
- Third most common malignancy in children: Lymphoma
- Most common solid tumor in children: Brain tumor

Solution to Question 2:

The most common intraabdominal malignancy in children is neuroblastoma.

- The most common leukemias seen in Down's syndrome: acute lymphoblastic leukemia (ALL) > acute myeloid leukemia (AML)
- The most common site of a brain tumor in children: Infratentorium
- The most common extra-cranial solid neoplasm in children: Neuroblastoma
- The most common intra-abdominal neoplasm in children: Neuroblastoma
- The most common primary neoplasm of renal tissue in children: Wilm's tumor

Solution to Question 3:

The image shows a typical Down's facies (flattened nose, flat occiput, low set ears, protruding tongue). The most common leukemia in children with Down syndrome is acute lymphoblastic leukemia (ALL).

Children with Down syndrome are at increased risk of both acute lymphoid leukemia (ALL) and acute myeloid leukemia (AML).

10% of neonates with Down's syndrome develop transient leukemia or myeloproliferative disorder. These features usually resolve within the first 3 months of life. However, they require close follow-up, because 20–30% will develop typical leukemia (often acute megakaryocytic leukemia) by 3 years of life.

Note:

- Most common leukemia in Down's syndrome: B-cell Acute lymphoblastic leukemia (overall)
- The most common leukemia in children <3 years of age with Down syndrome is acute myeloid leukemia (AML).
- Most common AML in Down's syndrome: M7/megakaryoblastic

Solution to Question 4:

B-lymphoblastic leukemia(pre-B-ALL) is the most common immunophenotype with TdT+, HLA DR+, CD10+, CD19+, CD20+, CD24+, Cy Ig + , with onset at 1-10 years of age.

The most common type of acute lymphoblastic leukemia (ALL) in children is B-lymphoblastic leukemia (previously termed pre-B-ALL)

Based on current WHO classification: The most common subtype of ALL is B lymphoblastic leukemia & T lymphoblastic leukemia & Mature B-cell Leukemia (Burkitt leukemia)

The production of symptoms and signs of ALL is a result of the replacement of bone marrow by malignant cells and caused by secondary bone marrow failure. On physical examination, findings of pallor, listlessness, purpuric/petechial skin lesions, or mucous membrane hemorrhage can reflect bone marrow failure. The proliferative nature of the disease may be manifested as lymphadenopathy, splenomegaly, or, less often, hepatomegaly. At presentation 75% of patients have leukocyte counts of <20,000/ μ L; thrombocytopenia is seen in 75% of patients and hepatosplenomegaly in 30–40% of patients. Patients with joint pain may have evidence of joint swelling and effusion. Deep bone pain may be present with marrow involvement, but tenderness will not be elicited.

Solution to Question 5:

Fanconi's anemia (not syndrome) predisposes to the development of childhood leukemias eg: B-ALL due to defective DNA repair. Fanconi syndrome is a syndrome of inadequate reabsorption in the proximal renal tubules. Congenital pernicious anemia does not predispose to leukemias.

Factors predisposing to childhood leukemias


- Genetic conditions:
- Fanconi's anemia
- Bloom syndrome
- Ataxia telangiectasia
- Li-Fraumeni syndrome
- Down syndrome
- Severe combined immunodeficiency
- Neurofibromatosis 1
- Paroxysmal nocturnal hemoglobinuria
- Diamond-Blackfan anemia
- Schwachman diamond syndrome
- Kostmann syndrome
- Environmental factors:
- Ionizing radiation
- Alkylating agents
- Nitrosoureas
- Epipodophyllotoxin
- Benzene exposure
- Advanced maternal age

Solution to Question 6:

The lower cut-off percentage for diagnosing acute lymphoblastic leukemia on bone marrow biopsy is 25%.

Unlike with myeloid leukemias, there is no agreed-upon lower limit for the proportion of blasts required to establish a diagnosis of lymphoblastic leukemia (ALL). In general, the diagnosis should be avoided when there are < 20% blasts.

The image below shows numerous lymphoblasts seen on a peripheral smear.



Work-up of ALL:

It is important that all studies necessary to confirm a diagnosis and adequately classify the type of leukemia be performed, including bone marrow aspiration and biopsy, flow cytometry, cytogenetics, and molecular studies. The initial evaluation should also include a CSF examination. If lymphoblasts are found and the CSF leukocyte count is elevated, overt CNS or meningeal leukemia is present. This finding reflects a poorer stage and indicates the need for additional CNS and systemic therapies.

Solution to Question 7:

MRD (Minimal residual disease) is a quantitative estimate of the burden of leukemic cells present in marrow. MRD $> 0.01\%$ at the end of induction suggests a poorer prognosis.

Initial WBC count of $< 50,000$ cells/mm³, B-cell immunophenotype and t(12,21) indicate a standard risk (Favourable prognosis)

Risk-directed therapy is the standard of current ALL treatment and accounts for:

- Age at diagnosis
- Initial WBC count
- Immunophenotypic and cytogenetic characteristics of blast populations
- The rapidity of early treatment response (i.e., how quickly the leukemic cells can be cleared from the marrow or peripheral blood)
- Assessment of MRD at the end of induction therapy.

Solution to Question 8:

T-cell acute lymphoblastic leukemia (T-ALL) is the most likely diagnosis since the chest X-ray shows superior mediastinal widening- indicating an involvement of the thymus gland.

T-ALL accounts for about 15% of childhood ALL cases; it is more common in adolescents - this is the time when the thymus reaches maximum size.

60% of patients with T-ALL present with a mediastinal mass. The mediastinal mass may be asymptomatic and detected only on chest x-ray or may cause cough and dyspnea, particularly on lying flat or with facial swelling and plethora due to vascular compression (superior vena cava syndrome).

B-ALL presents usually between 1-10 years of age perhaps because the number of normal bone marrow pre-B-cells (the cell of origin) is greatest very early in life.

Solution to Question 9:

The above findings are diagnostic of Infantile acute lymphoblastic leukemia. Translocation (4;11) involving the MLL gene is most commonly associated with Infantile leukemia accounts for the very high relapse rate.

2% of cases of leukemia during childhood occur before the age of 1 year and are referred to as infantile leukemia.

Patients often present with hyperleukocytosis and extensive tissue infiltration producing organomegaly. Subcutaneous nodules, known as leukemia cutis, caused by infiltration by leukemic cells are observed more often in infants than in older children.

The leukemic cell morphology is usually that of large, irregular lymphoblasts, with a phenotype negative for the CD10 (common ALL antigen) marker (pro-B), unlike most older children with B-ALL, who are CD10 +. Has a very poor prognosis.

Note:

Infants with AML present with CNS and skin involvement and most commonly have a subtype known as acute myelomonocytic leukemia.

Solution to Question 10:

L-asparaginase is used in the treatment of acute lymphoid leukaemia (ALL). ALL can present with hepatosplenomegaly, severe joint and bone pains that the kid refuses to be handled.

It is observed that some lymphoid malignancies are unable to synthesize asparagine on their own and must derive it from the bloodstream. L-asparaginase catalyses circulating L-asparagine, thereby inhibiting protein synthesis in the cancer cells.

Treatment of ALL

- Induction: 4 weeks
- L-asparaginase: single dose (side effects: pancreatitis, thrombosis)
- Vincristine: weekly (side effects: neuropathy, SIADH)

- Prednisolone
- Anthracycline
- CNS prophylaxis- intrathecal methotrexate
- Consolidation:
- Methotrexate
- Etoposide
- Cyclophosphamide
- Cytosine-Arabinoside
- Maintenance: 24-36 months
- 6-Mercaptopurine
- Methotrexate

Other options:

- Imatinib is a tyrosine kinase inhibitor used to treat chronic myeloid leukemia and acute lymphoblastic leukemia with t(9;22) BCR-ABL fusion gene.
- Retinoin is used in acute promyelocytic leukaemia with t(15;17) PML-RARA fusion gene.
- Cladribine is used in hairy cell leukaemia.

Solution to Question 11:

Diagnosis of acute myeloid leukemia (AML) is confirmed by the presence of more than 20% myeloblast cells (as shown in the image) in the bone marrow.

Flow cytometry and special stains can be used to identify myeloperoxidase-containing cells, which confirms the myelogenous origin of leukemia.

Note:

The diagnosis of acute lymphoid leukemia (ALL) requires the presence of more than 25% blasts in the bone marrow.

Solution to Question 12:

Monosomy 7 is associated with a poor prognosis in patients with acute myeloid leukemia (AML).

The production of symptoms and signs of AML is a result of the replacement of bone marrow by malignant cells and caused by secondary bone marrow failure-pallor, fatigue, exercise intolerance, bruising, oral mucosal bleeding, or epistaxis, as well as fever, which may be caused by infection or the disease.

In addition, patients with AML present with signs and symptoms that are uncommon in Acute lymphoblastic leukemia, including subcutaneous nodules or "blueberry muffin" lesions (especially

in infants), infiltration of the gingiva (especially in monocytic subtypes), signs and laboratory findings of disseminated intravascular coagulation (especially indicative of acute promyelocytic leukemia), and discrete masses, known as chloromas or granulocytic sarcomas.

Prognostic implications of chromosomal abnormalities in AML are given below:

Favorable prognosis
t(8;21); RUNX1-RUNX1T1Inv (16) or t(16;16); CBFβNPM1m utationCEBPAmutation
Poor prognosis
inv(3)t(6;9)5q-del(7q)comple x karyotype (≥3 abnormalitie s)Monosomy7

Solution to Question 13:

Nodular, violaceous skin lesion infiltrated with myeloblasts is descriptive of Chloroma otherwise known as extramedullary myeloblastoma or granulocytic(myeloid) sarcoma. Myeloid sarcomas are seen in AML subtypes M2, M4, M5a, and M5b.

“Chloroma” was coined due to the peculiar green color of the tumor caused by myeloperoxidase present in tumor cells.

These masses can occur in the absence of apparent bone marrow involvement and typically are associated with t(8;21) translocation. Chloromas also may be seen in the orbit and epidural space.

Solution to Question 14:

Erythemic myelosis is AML M6.

Constitute about 1% of AML cases

Two subtypes of AML M6:

- Erythroid or myeloid leukemia: M6a (more common)
- Pure erythroid leukemia: M6b, also called erythemic myelosis (Di Guglielmo disease).

Solution to Question 15:

This infant has a transient myeloproliferative disease, which usually resolves spontaneously in the first 3 months of life, and they require close-follow up because 20–30% will develop typical leukemia (acute megakaryocytic leukemia) by 3 years of life.

Approximately 10% of neonates with Down syndrome develop transient leukemia or myeloproliferative disorder characterized by high leukocyte counts, blast cells in the peripheral blood, and associated anemia, thrombocytopenia, and hepatosplenomegaly. These features usually resolve within the first 3 months of life. Although these neonates can require temporary transfusion support, they do not require chemotherapy unless there is evidence of life-threatening complications.

As only 20-30% of such children develop acute megakaryocytic leukemia a bone marrow transplant would be a drastic step in our case.

Solution to Question 16:

Patients with juvenile myelomonocytic leukemia do not have the Philadelphia chromosome.

Formerly termed juvenile chronic myeloid leukemia (CML), it is a clonal proliferation of hematopoietic stem cells that typically affects children younger than 2 years of age. Patients with JMML present with rashes, lymphadenopathy, splenomegaly, and hemorrhagic manifestations.

Patients with this disease do not have the Philadelphia chromosome (the Philadelphia chromosome is characteristic of CML). The peripheral blood smear shows an elevated leukocyte count with increased monocytes, thrombocytopenia, and anemia with the presence of erythroblasts. Bone marrow shows a myelodysplastic pattern, with blasts accounting for $\leq 20\%$ of cells. Patients with neurofibromatosis type 1 and Noonan syndrome have a predilection for this type of leukemia.

JMML in the setting of Noonan syndrome is unique, with most patients having a spontaneous resolution.

Solution to Question 17:

Burkitt lymphoma is the most common non-Hodgkin's lymphoma in children under the age of 15 years.

Diffuse large B-cell lymphoma (DLBCL) is the most common form of non-Hodgkin lymphoma (NHL) in adults and adolescents (15-19) years, worldwide.

B-cell acute lymphoblastic leukemia (B-ALL) is the most common childhood malignancy and the most common leukemia in children. (It is not a non-Hodgkin's lymphoma).

Solution to Question 18:

In contrast to adult non-Hodgkin's lymphoma (NHL), which is typically indolent, pediatric NHL is usually high-grade and aggressive.

NHL in the pediatric age group has distinct clinical and behavior properties when compared to adults. These include the following:

NHL in adults	NHL in children
Low/intermediate grade	High grade
Dominantly nodal	Extra-nodal
Variable growth fraction	High growth fraction
Poor long-term outcome	Good outcome

Solution to Question 19:

Retroperitoneal and mesenteric lymphadenopathy with hepatosplenomegaly, anemia, and short duration of disease, point towards a diagnosis of lymphoid neoplasms. The classic translocation of t(8;14) and starry sky appearance with a high mitotic index on HPE point toward a diagnosis of Burkitt lymphoma. Burkitt's lymphoma is the most common type of Non-Hodgkin lymphoma worldwide. St Jude system of staging is used for staging childhood non-Hodgkin lymphoma.

The current revised staging system used for NHL is the International Pediatric Non-Hodgkin Lymphoma Staging System (IPNHLSS) which reflects our increasing ability to diagnose lower levels of organ involvement with disease. For instance, the older staging system (St. Jude/Murphy classification) did not account for molecular or flow cytometry involvement of bone marrow, which is now reflected in the new system.

Disease	System of staging
Hodgkin lymphoma	Ann–Arbor staging
Non-Hodgkin Lymphoma	St Jude(Murphy) staging
Chronic lymphoid leukemia	Rai staging Binet staging

Solution to Question 20:

In the picture given above, Reed–Sternberg (RS) cells are seen, which is a pathognomonic feature of Hodgkin's lymphoma (HL).


Reed–Sternberg cells are large and either multinucleated or have a bilobed nucleus (thus resembling an "owl's eye" appearance) with prominent eosinophilic inclusion-like nucleoli. The RS cell is clonal in origin and arises from the germinal center B cells but typically has lost most B-cell gene expression and function.

Hodgkin lymphoma encompasses a group of lymphoid neoplasms that typically arise in a signal node or chain of nodes and spread first to anatomically contiguous sites.

WHO classification of Hodgkin lymphoma includes 5 subtypes namely;

- Nodular sclerosis
- Mixed cellularity
- Lymphocyte-rich
- Lymphocyte depletion
- Nodular lymphocyte predominance

Histology of lymph node showing
Reed-Sternberg cell (Owl's eye appearance)



©MARROW

Solution to Question 21:

The clinical vignette describes Hodgkin's lymphoma with the involvement of liver and bone marrow, which is classified as Stage 4 according to Ann-Arbor staging.

Ann Arbor Staging of Hodgkin's Lymphoma

Stage I	Involvement of a single lymph node (I) or Lymphoid structure (eg: spleen, thymus, Waldeyer's ring)
Stage II	Involvement of 2 or more lymph node regions on the same side of the diaphragm
Stage III	Involvement of lymph node regions or lymphoid structures on both sides of the diaphragm (III)

Stage I	Involvement of a single lymph node (I) or Lymphoid structure (eg: spleen, thymus, Waldeyer's ring)
Stage II	Involvement of extranodal site beyond that designated as 'E' More than one extranodal deposit at any location Any involvement of liver or bone marrow
A	No symptoms
B	The presence of fever $>38^{\circ}\text{C}$ (100.4°F) for 3 consecutive days Recurrent drenching night sweats Unexplained loss of $>10\%$ of body weight in the 6 months
E	Localized, solitary involvement of extra lymphatic tissue, excluding liver and bone marrow

Solution to Question 22:

Age >10 years indicates a poor prognosis in ALL.

Leukocyte count $<50,000/\text{mm}^3$, hyperdiploidy, and trisomy of chromosomes 4,10,17 indicate a good prognosis.

Solution to Question 23:

Both CNS and testis are considered as sanctuary sites of acute lymphoblastic leukemia (ALL).

- A sanctuary site is an area of the body that is poorly penetrated by pharmacological agents and, therefore, serves as a place in which tumor cells or infectious organisms can escape the effects of drug therapy.
- Early institution of CNS prophylaxis (intrathecal methotrexate) is essential to eradicate these escaped leukemic cells.
- Testicular involvement is rarely evident at diagnosis, testicular involvement can present as painless testicular enlargement.

Solution to Question 24:

B-cell lymphoblastic leukemia (L1)

- This is the most common immunophenotype of acute lymphocytic leukemia (ALL) in a child between the ages of 1-10 years.
- They present with features of bone marrow failure and organ infiltration such as pancytopenia and hepatosplenomegaly, respectively, so B-cell ALL is the most likely diagnosis.
- Bony tenderness may be present due to sub-periosteal infiltration of lymphoblastic cells.
- T-cell ALL (L2) typically presents in adolescent boys with a large anterior mediastinal mass (thymus or mediastinal lymph nodes), causing respiratory distress or superior vena cava obstruction.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Solid Neoplasms of Childhood

Question 1:

Which of the following is the second most common malignancy in childhood?

- a) Leukemias
- b) Brain tumor
- c) Neuroblastoma
- d) Wilms tumor

Question 2:

The MRI scan of an 11-year-old girl revealed a brain tumor. In which of the following conditions would you do a lumbar puncture for cytological analysis?

- a) Infratentorial tumours
- b) Tumors causing obstructive hydrocephalus
- c) Supratentorial tumors with midline shift
- d) Germ-cell tumours

Question 3:

A mother brought her 10-month-old baby to the hospital with sudden onset vomiting. She has also been noticing an alarming increase in the head size of the baby. On examination, significant macrocephaly is seen. MRI reveals a supratentorial tumor. Which of the following tumors is the most likely cause of this child's presentation?

- a) Medulloblastoma
- b) Pilocytic astrocytoma
- c) Ependymoma
- d) Choroid plexus carcinoma


Question 4:

Sanam, a 7-year-old kid is suspected to have an ependymoma. Which of the following features on histopathology would favor this diagnosis?

- a) Rosenthal fibres
- b) Perivascular pseudorosettes
- c) Flexner-Wintersteiner rosettes
- d) Homer-Wright rosettes

Question 5:

A young child was evaluated for increased ICP. A brain tumor was found on an MRI scan. It was resected and sent for histopathological examination. The slide is shown below. This tumor is found most commonly in the _____.




- a) Filum terminale
- b) Pia mater
- c) Floor of 4th ventricle
- d) Cerebellopontine angle

Question 6:

A concerned mother brought her daughter to the clinic for evaluation of symptoms that have worsened in the last few weeks. The daughter frequently wakes up at night with severe headaches and vomiting. She has also been having some difficulty walking. On examination, ataxia and multiple cranial nerve palsies are noted. Imaging is suggestive of a brainstem


tumor. Which of the following is the most likely diagnosis?



- a) Pontine glioma
- b) Medulloblastoma
- c) Ependymoma
- d) Acoustic neuroma

Question 7:

On your neurology rotation, you come across Ashu, a 5-year-old child who presented with a 6-month history of ataxia, headaches, and vomiting. He was admitted to the ward for further evaluation. MRI scan showed a brain tumor with a combination of solid and cystic components. Biopsy findings of the tumor are shown in the image. Which of the following is the most likely diagnosis?



- a) Medulloblastoma
- b) Ependymoma
- c) Brainstem glioma
- d) Pilocytic astrocytoma


Question 8:

What is the WHO grading for Pilocytic astrocytoma?

- a) Grade I
- b) Grade II
- c) Grade III
- d) Grade IV

Question 9:


You are a pathology resident who is looking at a slide, showing small round blue cells and Homer-Wright rosettes as shown below. The history you received from the pediatric surgery resident was that the specimen was resected from a 5-year-old child with ataxia. The tumor was present in the midline vermis. Which of the following is the most likely diagnosis?



- a) Medulloblastoma
- b) Ependymoma
- c) Pilocytic astrocytoma
- d) Glioblastoma multiforme

Question 10:

A child is brought to the emergency department by his teacher when he started having abnormal jerking movements during math class. The parents give a history of the child having similar episodes in the past. On the way to the CT room, the child had an episode of seizure and was stabilized with lorazepam. The imaging done is shown below. What is the most likely diagnosis?



- a) Astrocytoma
- b) Medulloblastoma
- c) Oligodendroglioma
- d) Craniopharyngioma

Question 11:

An erythropoietin-secreting brain tumor is detected in a child. His father passed away one year back from end-stage renal cell carcinoma. Family history is significant for VHL mutation. On examination, a fiery red retina is seen. Which of the following is the most likely tumor present in this child?

- a) Pilocytic astrocytoma
- b) Cerebellar hemangioblastoma
- c) Cavernous hemangioma
- d) Neuroblastoma

Question 12:

Akhil, a 15-year-old boy with sun-setting eyes, has a tumor in the pineal region as seen in the MRI. Parinaud syndrome is suspected. Which of the following findings will not be seen in this boy?




- a) Pseudo-Argyll–Robertson pupil

- b) Paresis of upward gaze
- c) Convergence retraction nystagmus on attempted upgaze
- d) Infranuclear vertical gaze disorder

Question 13:

An 8-year-old child is brought to the casualty after a minor accident that occurred when he failed to see a bike coming from sideways. His mother says he is a clumsy child, who sometimes knocks things over accidentally even at home. Visual field examination findings are given below. On imaging, an intracranial mass is seen. The resected specimen shows cholesterol-rich machinery-oil-like fluid. What is the most likely diagnosis?



- a) Prolactinoma
- b) Adamantinomatous craniopharyngioma
- c) Ependymoma
- d) Meningioma

Question 14:

Which of the following is the most common extracranial solid tumor in the pediatric age group?

- a) Malignant bone tumors
- b) Wilms tumor
- c) Retinoblastoma

d) Neuroblastoma

Question 15:

A 5-year-old child presents with rapid dancing eye movements, rhythmic jerking of limbs and poor coordination. You suspect a paraneoplastic syndrome. Which of the following conditions should you look for in this child?

- a) Cerebellar hemangioblastoma**
- b) Neuroblastoma**
- c) Medulloblastoma**
- d) Rhabdomyosarcoma**

Question 16:

In the ophthalmology OPD, you see a 3-year-old child with the following finding. There is no history of trauma. There is no significant family history. Which of the following is the most likely diagnosis?



- a) Retinoblastoma**
- b) Neuroblastoma**
- c) Orbital rhabdomyosarcoma**
- d) Basilar skull fracture**


Question 17:

A 10-month-old infant is diagnosed with neuroblastoma. Which of the following factors is associated with a good prognosis?

- a) Hyperdiploidy
- b) Loss of heterozygosity of 11 q
- c) Gain of 17q
- d) N-myc amplification

Question 18:

A 3-year-old boy presents with a 2-month history of an enlarging scalp lesion. There is no history of trauma to the head or scalp. The child is otherwise healthy and investigations are normal. X-ray of the head is given below. What is the most likely diagnosis?



- a) Multiple myeloma
- b) Langerhans cell histiocytosis
- c) Hairy cell leukaemia
- d) Neuroblastoma

Question 19:

A 12-year-old child is brought with 4 small, purplish, rubbery lesions on his back and arms. On examination, you note inguinal freckling and nodules on the iris as shown below. Which

chromosome is involved in the syndrome described in this child?



- a) 7
- b) 17
- c) 22
- d) 11

Question 20:

Vestibular schwannomas, meningiomas, and spinal cord ependymomas are the CNS manifestations of _____.

- a) Neurofibromatosis type 1
- b) Neurofibromatosis type 2
- c) von Hippel-Lindau syndrome
- d) Turcot syndrome

Question 21:

A 12-year-old boy presented with focal seizures involving the right side of his body. On investigations, a subependymal giant cell astrocytoma is found. Which of the following syndromes is most likely to be associated with this?

- a) Neurofibromatosis type 1
- b) Neurofibromatosis type 2

- c) Tuberous sclerosis
- d) Turcot syndrome

Question 22:

What is the most common intraocular tumor diagnosed in the pediatric age group?

- a) Retinoblastoma
- b) Rhabdomyosarcoma
- c) Malignant melanoma of choroid
- d) Basal cell carcinoma of eyelid

Question 23:

James was brought by his mother for evaluation of a squint she noticed. On examination, the given finding was observed. Investigations revealed retinoblastoma with a germline RB1 gene mutation. The secondary malignancy most likely to develop in him is:



- a) Osteosarcoma
- b) Pancreatic tumour
- c) Thyroid malignancy
- d) Hepatoblastoma

Question 24:

A 5-year-old boy presented with an abdominal mass. On examination, the mass is firm, smooth, non-tender, and does not cross the mid-line. The boy is also found to have moderate hypertension. Urinalysis reveals microscopic haematuria. Wilms tumor is suspected. This condition can be associated with all of the following except _____.

- a) Aniridia
- b) Beckwith syndrome
- c) Polycystic kidney
- d) Hemi-hypertrophy

Question 25:

You are assisting on a left partial nephrectomy in a child with Wilms tumor. Intraoperatively, a small mass is noted on the right kidney as well. Frozen section reveals it to be a Wilms tumor. What is the stage of Wilm's tumor in this child?

- a) Stage 2
- b) Stage 3
- c) Stage 4
- d) Stage 5

Question 26:

While posted in the pediatric oncology ward, you come across a case of metastatic triphasic nephroblastoma in a 4-year-old. What is the most common site for metastasis of this tumor?

- a) Lung
- b) Brain
- c) Bone
- d) Adrenal

Question 27:


What is the most important determinant of prognosis in a child diagnosed with Wilm's tumor?

- a) Stage of disease
- b) Loss of heterozygosity of chromosome 1 p

- c) Histology
- d) Age < 1 year at presentation

Question 28:

A 5-year-old child comes with abdominal distension, a palpable mass, and gross hematuria. A biopsy of the mass shows the following picture. Which of the following paraneoplastic syndromes is not associated with this tumor?



- a) Hypertrophic primary osteoarthropathy
- b) Polycythemia
- c) Hypercalcemia
- d) Bleeding


Question 29:

The most common soft tissue sarcoma in the pediatric age group is_____.

- a) Retinoblastoma
- b) Rhabdomyosarcoma
- c) Malignant melanoma
- d) Cavernous hemangioma

Question 30:

A 4-year-old girl presents with the lesion shown in this picture. The diagnosis is_____.



- a) Fibrosarcoma
- b) Sarcoma botryoides
- c) Leiomyosarcoma
- d) Inflammatory polyp

Question 31:

The precursor lesion of Wilm's tumor is _____.

- a) Nephroblasts
- b) Embryonic nephrogenic rests
- c) Medullary cysts
- d) Cortical cysts

Question 32:

The term "trilateral retinoblastoma" refers to bilateral retinoblastoma + _____.

- a) Osteosarcoma
- b) Pineoblastoma
- c) Rhabdomyosarcoma
- d) Malignant melanoma

Question 33:

Tumor which may undergo spontaneous regression is _____.

- a) Medulloblastoma
- b) Hepatoblastoma
- c) Neuroblastoma
- d) Nephroblastoma

Question 34:

Which of the following pediatric tumors is called a "triphasic" blastoma?

- a) Hepatoblastoma
- b) Neuroblastoma
- c) Medulloblastoma
- d) Nephroblastoma

Answer Key

Question No.	Correct Option
1	b
2	d
3	d
4	b
5	c
6	a
7	d
8	b
9	a
10	c
11	b
12	d
13	b
14	d
15	b

16	b
17	a
18	b
19	b
20	b
21	c
22	a
23	a
24	c
25	d
26	a
27	c
28	a
29	b
30	b
31	b
32	b
33	c
34	d

Detailed Explanations

Solution to Question 1:

The second most common malignancy in childhood is a brain tumor.

The most common malignancy in childhood is Leukemias.

Brain tumors in childhood:

- Most common solid tumor in children- Brain tumor
- Most common site of a brain tumor in children- Infratentorium
- Most common brain tumor in childhood- Pilocytic astrocytoma

Solution to Question 2:

Germ-cell tumor is an indication for lumbar puncture for cytological analysis of CSF.

Usually, lumbar punctures are contraindicated in intracranial space-occupying lesions with high intracranial pressure as it can lead to brain herniation and death. They are thus delayed until

surgery or shunt placement. However, in tumors with a risk of leptomeningeal metastases, lumbar puncture is indicated for cytology.

Tumors where LP is indicated	Tumors where LP is contraindicated
Medulloblastoma or primitive neuroectodermal tumors	Infratentorial tumors
Ependymoma	Supratentorial tumors with midline shift
Germ cell tumors	Tumors with hydrocephalus secondary to CSF obstruction

Solution to Question 3:

Choroid plexus carcinoma is a pediatric supratentorial tumor presenting with macrocephaly and focal neurologic deficits.

Choroid plexus tumors are the most common CNS tumors in children aged less than 1 year. These tumors are intraventricular epithelial neoplasms arising from the choroid plexus.

Children present with signs and symptoms of increased intracranial pressure (ICP), whereas infants may present with macrocephaly and focal neurologic deficits.

Medulloblastoma, pilocytic astrocytoma, and ependymoma, are all predominantly infratentorial tumors.

The age-related differences in the primary location of the tumor are given below:

Age	Predominant Site	Examples
< 1 year	Supratentorial	Choroid plexus tumor and teratoma
1 - 10 years	Infratentorial	Juvenile pilocytic astrocytoma and medulloblastoma
> 10 years	Supratentorial	Diffuse astrocytoma

Solution to Question 4:

Perivascular rosettes will be seen on histopathological examination of ependymoma. Other features include ependymal rosettes and monomorphic nuclear morphology.

Ependymomas are tumors derived from the ependymal lining of the ventricular system occurring predominantly in childhood. They are usually non-invasive, extending into the ventricular lumen and/or displacing normal structures, sometimes leading to significant obstructive hydrocephalus.

Solution to Question 5:

The slide shows perivascular pseudorosettes and ependymal rosettes, characteristic of ependymoma. The most common site of ependymoma in children is the floor of the 4th ventricle. Monomorphic nuclear morphology can also be noted.

Ependymomas are tumors derived from the ependymal lining of the ventricular system, occurring predominantly in childhood. These tumors are usually non-invasive, extending into the ventricular lumen and/or displacing normal structures, sometimes leading to significant obstructive hydrocephalus.

Solution to Question 6:

The MRI given shows a diffuse, infiltrating pontine lesion. Multiple cranial nerve dysfunctions, cerebellar signs, motor weakness, and features of raised intracranial pressure (ICP) are all strongly suggestive of a pontine glioma.

These are not amenable for surgical resection. The standard approach is radiation therapy. The prognosis is very poor.

Solution to Question 7:

The given slide shows condensed masses of glial filaments occurring in compact areas suggestive of Rosenthal fibers. The clinical features, MRI findings, and histopathology point towards a diagnosis of pilocytic astrocytoma.

Pilocytic astrocytoma is a low-grade astrocytoma tumor. It is the most common type of astrocytoma in children. These are classically found in the cerebellum (infratentorial) and the optic pathway regions.

There is usually a 3-6 month history of limb ataxia, secondary headaches, and vomiting. Mural nodule, a contrast medium enhancing nodule within the wall of a cystic mass is seen on MRI.

The tumor has a low metastatic potential and is rarely invasive.

Solution to Question 8:

Piloxyoid astrocytoma falls under WHO Grade II.

Piloxyoid astrocytoma:

Occurs most commonly in the hypothalamic/optic chiasmic region and carries a high risk of local as well as cerebrospinal spread. This astrocytoma affects young children and infants. It is classified as a WHO grade II tumor.

WHO Grades of brain tumor:

WHO grade	Example
Grade I	Pilocytic astrocytoma Meningeal Ependymoma/subependymoma
Grade II	Diffuse astrocytoma Pilo-myxoid astrocytoma Papillary/clear-cell ependymoma
Grade III	Anaplastic astrocytoma Anaplastic ependymoma
Grade IV	Glioblastoma multiforme Malignant pineoblastoma and medulloblastoma

Solution to Question 9:

The given clinical scenario and slide are suggestive of Medulloblastoma.

Medulloblastoma is an embryonal CNS tumor, occurring predominantly in the cerebellar vermis that causes ataxia, poor balance, and dysmetria. It may cause obstructive hydrocephalus and often metastasizes extraneurally.

Histopathology shows a tumor composed of small, round, blue cells and is characterized by Homer–Wright rosettes and an immunopositivity for synaptophysin.

Standard treatment for medulloblastomas is surgical resection, with adjuvant craniospinal radiotherapy for those over 3 years of age and chemotherapy.

Prognosis is strongly influenced by surgical resection, the presence of CSF metastases at the time of diagnosis, and the expression of the c-erbB-2 (HER2/neu) oncogene.

Note: Small, round, blue-cell tumors are malignant neoplasms that have a characteristic appearance under the microscope. It consists of sheets of small round cells that stain blue on routine hematoxylin and eosin (H&E) staining. Examples are:

- Medulloblastoma
- Neuroblastoma
- Retinoblastoma
- Ewing's sarcoma
- Rhabdomyosarcoma


Solution to Question 10:

Calcified cortical mass with a seizure is suggestive of oligodendroglioma.

Oligodendrogliomas are uncommon tumors of childhood. These infiltrating tumors occur predominantly in the cerebral cortex and originate in white matter. Treatment involves surgical resection followed by involved field radiation therapy.

The image below shows histology of oligodendroglioma showing characteristic fried egg appearance and chicken wire capillaries.

Oligodendroglioma



©MARROW

Craniopharyngioma is associated with suprasellar calcification. Also, the child may often present with a history of endocrinological and visual complaints.


Solution to Question 11:

The given clinical scenario is suggestive of a cerebellar hemangioblastoma.

Von Hippel-Lindau (VHL) syndrome is an autosomal dominant condition. The major neurological features of VHL include cerebellar hemangioblastomas and retinal angiomas.

Cerebellar hemangioblastomas are erythropoietin-secreting tumors, which can lead to polycythemia. It typically appears as a cystic cerebellar lesion with a vascular mural nodule on CT or MRI scan.

Retinal angiomas are small masses of thin-walled capillaries, usually located in the peripheral retina that is fed by large and tortuous arterioles and venules appearing as a fiery red vascular mass. It can lead to retinal detachment and visual loss.



Associations of VHL syndrome include:

- Cystic lesions of kidneys, pancreas, liver, and epididymis
- Renal cell carcinoma (most common cause of death)
- Pheochromocytoma
- Cerebellar hemangioblastoma
- Retinal angioma

Solution to Question 12:

Parinaud's syndrome refers to supranuclear vertical gaze disorder.

Parinaud's syndrome/dorsal midbrain syndrome is caused by damage to the posterior commissure. Classically, signs of hydrocephalus are seen due to aqueductal stenosis.

Causes:

- Pineal region or midbrain tumors (Most common cause)
- Neurocysticercosis
- Stroke
- Multiple sclerosis

Features include the following:

- Loss of up-gaze (downward gaze is usually preserved)
- Convergence–retraction nystagmus on attempted upgaze
- Downward ocular deviation- Setting-sun sign
- Lid retraction- Collier's sign

- Pseudo-Argyll–Robertson pupil- Accommodative paresis ensues and pupils become mid-dilated and show light-near dissociation
- Bilateral papilledema
- Skew deviation
- Pseudoabducens palsy

Solution to Question 13:

Patients with adamantinomatous craniopharyngioma present with bitemporal hemianopia (image), and the cystic component of this tumor contains a cholesterol-rich, thick, brownish-yellow fluid that is comparable to "machine oil."

Craniopharyngiomas develop from remnants of the Rathke pouch. Adamantinomatous variant of craniopharyngioma predominates in childhood.

These tumors are usually large and heterogeneous, displaying both solid and cystic components, and occur in the suprasellar region. Patients usually present with headache and visual field abnormalities. These children also have endocrinological abnormalities such as growth failure and delayed sexual maturation due to hypopituitarism (GH deficiency).

These tumors are minimally invasive, adhere to the adjacent brain parenchyma, and engulf normal brain structures. CT may show calcifications associated with the solid and cystic components.

Solution to Question 14:

Neuroblastoma is the most common extracranial solid tumor in children and the most commonly diagnosed malignancy in infants. It is an embryonal cancer of the peripheral nervous system.

The signs and symptoms depend on the location of the tumor and the extent of the disease. Neuroblastomas originating from the superior cervical ganglion can present with Horner syndrome.

Metastatic disease can cause a variety of signs and symptoms, including the following

- Fever
- Irritability
- Failure to thrive
- Bone pain
- Cytopenia
- Bluish subcutaneous nodules
- Orbital proptosis, and periorbital ecchymoses

Rapid dancing eye movements (opsoclonus), rhythmic jerking of limbs/trunk (myoclonus), and ataxia are characteristic of the opsoclonus-myoclonus-ataxia syndrome (OMA). It is

a paraneoplastic syndrome of autoimmune origin with a strong association with neuroblastomas. Some neuroblastomas may undergo spontaneous regression, whereas some may be unresponsive to therapy and run a very aggressive course.

	Neuroblastoma	Wilm's tumor (nephroblastoma)
Age group	<2 years of age	Peaks at 3-4 years of age
Calcifications	Very common (90%)	Uncommon (10-15%)
Margins	Poorly marginated	Well circumscribed
Vascular invasion	Encases vascular structures but does not invade them	Extends into the IVC and renal vein
Relation with surrounding structures	Elevates the aorta away from the vertebral columnCommonly crosses the midline, especially behind the aorta	Displaces adjacent structures without insinuating between themForms the claw sign with the kidney

Solution to Question 15:

Rapid dancing eye movements (opsoclonus), rhythmic jerking of limbs/trunk (myoclonus) and ataxia is characteristic of opsoclonus-myoclonus-ataxia syndrome (OMA). It is a paraneoplastic syndrome of autoimmune origin with a strong association with neuroblastoma.

All children with OMA should undergo investigations for neuroblastoma such as ultrasound followed by CT or MRI of chest and abdomen, bone scans and marrow biopsies. The definitive diagnosis of neuroblastoma is through biopsy which shows small round blue cells.

Other small round blue cell tumours are medulloblastoma, retinoblastoma, Ewing's sarcoma and rhabdomyosarcoma.

Solution to Question 16:

This image shows periorbital ecchymoses and proptosis suggestive of metastases of neuroblastoma (a sign of orbital bone involvement).

Neuroblastomas may develop at any site in the sympathetic nervous system. Metastatic spread is common in children who are diagnosed after 1 year of age. It occurs via local invasion, hematogenous, or lymphatic routes.

Metastatic disease can cause a variety of signs and symptoms, including the following:

- Fever
- Irritability
- Failure to thrive
- Bone pain

- Cytopenia
- Bluish subcutaneous nodules
- Orbital proptosis, and periorbital ecchymoses

Solution to Question 17:

Hyperdiploidy is associated with a good prognosis in neuroblastoma if the child is younger than 1 year of age at diagnosis.

Bad prognosis in neuroblastoma is seen in cases of:

- Amplification of N-MYC
- Loss of Heterozygosity of 1p, 11q, 14q
- Gain of 17q
- TrkB expression

Solution to Question 18:

The given clinical scenario with the x-ray image showing a punched-out lytic lesion is suggestive of Langerhans cell histiocytosis (LCH).

LCH is a rare histiocytic disorder characterized by single or multiple osteolytic bone lesions that can infiltrate nearly every organ, most notably the skin (of the scalp), lungs, thymus, liver, spleen, bone marrow, or the central nervous system.

LCH is so named because of its resemblance to the Langerhans's cells, which are specialized dendritic cells found in the skin and mucosa.

Solution to Question 19:

The given clinical scenario is suggestive of neurofibromatosis type 1. The NF1 gene on chromosome 17q11 is involved. It is an autosomal dominant neurocutaneous syndrome.

It is diagnosed when at least 2 of the following 7 criteria are met:

- 6 or more café-au-lait macules
- ≥ 5 mm prepuberty
- ≥ 15 mm post-puberty
- Skin-fold freckling- axillary and inguinal
- 2 or more iris lisch nodules (image)
- 2 or more neurofibromas or 1 plexiform neurofibroma (purplish, rubbery lesions)

- Osseous lesions
- Osseous dysplasia
- Cortical thinning of long bones without a pseudoarthrosis
- Low-grade astrocytomas like optic gliomas
- 1st degree relative with diagnosed NF-1

Solution to Question 20:

Vestibular schwannomas, meningiomas, and spinal cord ependymomas are the CNS manifestations of neurofibromatosis type 2

It is an autosomal-dominant neurocutaneous syndrome, which occurs due to a mutation in the NF2 gene on chromosome 22q12.

von Hippel–Lindau syndrome is an autosomal dominant neurocutaneous syndrome, which occurs due to a mutation in the VHL gene on chromosome 3p25-26. The CNS manifestation is usually a hemangioblastoma.

Turcot syndrome is an autosomal dominant neurocutaneous syndrome, which occurs due to a mutation in the APC gene on chromosome 5q21. The CNS manifestation is a medulloblastoma.

Solution to Question 21:

Subependymal giant-cell astrocytoma is a characteristic CNS finding in tuberous sclerosis.

Syndrome	CNS Manifestations	Gene
Neurofibromatosis type 1 (autosomal dominant)	Optic pathway gliomas Astrocytoma Malignant peripheral nerve sheath tumours Neurofibromas	NF1
Neurofibromatosis type 2 (autosomal dominant)	Vestibular schwannomas Meningiomas Spinal cord ependymomas	NF2
Von Hippel–Lindau (autosomal dominant)	Cerebellar hemangioblastoma	VHL
Tuberous Sclerosis (autosomal dominant)	Subependymal giant-cell astrocytoma Cortical tubers	TSC1 TSC2
Turcot syndrome (autosomal dominant)	Medulloblastoma Glioblastoma	APC

Solution to Question 22:

Retinoblastoma is the most common intraocular tumor of the pediatric age group. It is an embryonal malignancy of the retina.

Solution to Question 23:

Children with germline RB1 mutations are at significant risk for the development of secondary malignancies, especially

- Osteosarcoma (Most common)
- Soft tissue sarcomas
- Malignant melanoma

The risk of secondary malignancies is further increased by the use of radiation therapy.

A pineoblastoma occurring in a child with bilateral hereditary retinoblastoma is called 'trilateral retinoblastoma'.

The image shows leukocoria, a white pupillary reflex. It may be seen in the following conditions:

- Congenital cataract
- Retinoblastoma
- Retinopathy of prematurity (retrolental fibroplasia)
- Persistent hyperplastic primary vitreous
- Coats disease
- Toxocara endophthalmitis

Solution to Question 24:

Polycystic kidneys are not associated with Wilm's tumor.

Wilms tumor (WT), also known as nephroblastoma, is the most common primary malignant renal tumor of childhood.

Polycystic kidney disease can also present with an abdominal mass, hypertension, and hematuria. However, there will be multiple irregular cysts on ultrasound and is not associated with Wilms tumor.

Solution to Question 25:

Wilms tumor with involvement of both kidneys is stage 5.

Children's Oncology Group (COG) Staging System

Stage I	Tumor confined to the kidney and completely resected with negative lymph nodes.
Stage I I	Tumor extends beyond the kidney but is completely resected with negative margins and lymph nodes.
Stage I II	Residual tumor present following surgery confined to the abdomen, including gross or microscopic tumor.
Stage I V	Hematogenous metastases (lung, liver, bone, brain, etc.) or lymph node metastases outside the abdominopelvic region
Stage V	Metastasis to opposite kidney (bilateral renal involvement)

Solution to Question 26:


The most common sites for metastases of nephroblastoma or Wilms tumor are the following:

- Lungs (Most common)
- Regional lymph nodes
- Liver

The triphasic components visualized on histopathology are-

- Malignant small round (blue) cells (blastemal component).
- Tubular structures/rosettes (epithelial component).
- Loose pauci-cellular stroma with spindle cells (stromal component)

Wilms Tumor/Triphasic Nephroblastoma



©Marrow

Solution to Question 27:

Histology is the most important determinant of prognosis in Wilms Tumor. The absence of anaplasia is considered a favorable histological finding.

Other prognostic factors for risk-adapted therapy include

- Age
- Stage
- Tumor weight
- Loss of heterozygosity at chromosomes 1p and 16q.

Solution to Question 28:

The histopathological image shows the characteristic triphasic components of Wilms tumor.

Hypertrophic primary osteoarthropathy is not a paraneoplastic syndrome of Wilms tumor.

Several distinct paraneoplastic syndromes have been reported in Wilms tumor. They are as follows:


- Hypertension
- Erythrocytosis/Polycythemia
- Hypercalcemia
- Cushing syndrome
- Acquired von Willebrand disease

Bleeding may occur as a paraneoplastic syndrome in Wilm's tumor. It is due to acquired deficiency of von Willebrand factor or factor VII deficiency.

The triphasic components visualized on histopathology are-

- Malignant small round (blue) cells (blastemal component).
- Tubular structures/rosettes (epithelial component).
- Loose pauci-cellular stroma with spindle cells (stromal component)

Wilms Tumor/Triphasic Nephroblastoma



©MARROW

Solution to Question 29:

Rhabdomyosarcoma, a tumor of striated skeletal muscle, is the most common pediatric soft tissue sarcoma.

The most common sites are- head and neck (25%) > genitourinary tract (24%) > extremities (19%) > orbit (9%).

Solution to Question 30:

The lesion seen in the picture is sarcoma botryoides

Vaginal rhabdomyosarcoma may manifest as a grape-like mass of tumor tissue bulging through the vaginal orifice, known as sarcoma botryoides.

Solution to Question 31:

- Wilms tumor is derived from foci of incompletely differentiated/undifferentiated renal mesenchyme called nephrogenic rests.

- Nephrogenic rests usually regress or differentiate, but those that persist can become malignant.
- Nephrogenic rests that persist abnormally in the kidneys into postnatal life are seen in 1% of children in the general population, but are seen in up to 90% of children who develop or have a family history of Wilms tumor.

Solution to Question 32:

A pineoblastoma occurring in a child with bilateral hereditary retinoblastoma is called “trilateral retinoblastoma.”

In hereditary retinoblastoma (having spontaneous germline RB1 mutation), second malignancies are a major concern. The most common second malignancy is osteosarcoma; other second neoplasms include rhabdomyosarcoma and melanoma.

Solution to Question 33:

Neuroblastoma is an embryonal cancer of the peripheral sympathetic nervous system with heterogeneous presentation and prognosis; some tumors may undergo spontaneous regression, whereas some may be unresponsive to therapy and run a very aggressive course.

Solution to Question 34:

- The classic nephroblastoma, aka Wilms tumor, is a mixed tumor (more than one neoplastic cell type) derived from one germ-cell layer (examples: Wilms tumor and pleomorphic adenoma of salivary gland).
- Histologically, it is made up of the classic "triphasic" combination (recapitulating stages of normal renal development) of
 - Stromal
 - Epithelial, and
 - Blast cells

Musculoskeletal Disorders

Question 1:


A 3-year-old child is brought to the clinic because of walking difficulties. His mother says he has always walked on his toes and walks by tilting side-to-side. He also has progressive muscle weakness. There is no history of trauma, fever, or rash. On examination, calf hypertrophy with atrophied thigh muscles is noted as shown below. There is no tenderness on palpation. What is the most probable diagnosis?



- a) Duchenne muscular dystrophy
- b) Myotonic dystrophy
- c) Spinal muscular atrophy
- d) Emery-Dreifuss muscular dystrophy

Question 2:

The image below shows the model of a skeletal muscle fiber. Which of the following structures is affected in Duchenne muscular dystrophy?



- a) A
- b) B
- c) C
- d) D


Question 3:

Abhi, a young boy is being evaluated in the ward for proximal muscle weakness. Waddling gait and Gowers sign are observed in this child. Genetic analysis confirms the diagnosis as Duchenne muscular dystrophy. Which of the following is false regarding this patient's condition?

- a) It is less severe than Becker's dystrophy
- b) Tendon reflexes are lost after age 6
- c) Gene affected is the largest gene in humans
- d) Intellectual impairment is common

Question 4:

You observe a six-year-old child in the ward and notice the following. Which of the following are true regarding this sign?



- a) 3, 5
- b) 2, 4
- c) 1, 3, 5
- d) 1, 4, 5

Question 5:

A concerned mother brought her son to the clinic because of difficulty writing in school. She also says that his eyes remain partially open when he sleeps. There is no history of trauma or surgery. On examination, he has asymmetric wrist and finger weakness. The mouth appears puckered and rounded, deltoid muscles are flattened and the following finding is observed. What is the likely diagnosis?



- a) Facioscapulohumeral dystrophy
- b) Long thoracic nerve palsy
- c) Klippel-Feil syndrome
- d) Myotonic dystrophy

Question 6:

A 7-year-old child presents with leg pain. She had fever and cough 3 days prior. Currently, there is pain on palpation of the calf muscles. Reflexes and other clinical examinations are normal. Lab test reveals CK levels of 2000 IU. What is the likely diagnosis?

- a) Viral Myositis
- b) Duchenne
- c) GBS
- d) Dermatomyositis

Question 7:

Rekha brought her 5-day-old infant to the clinic because of feeding difficulties. She says he does not latch properly. Her antenatal history is significant for polyhydramnios and decreased fetal movements. On examination, hypotonia and hyporeflexia are noted in the baby. which of the following is the most likely diagnosis?

- a) Spinal muscular atrophy
- b) Classical myotonic dystrophy
- c) Duchenne muscular dystrophy
- d) Emery-Dreifuss syndrome

Question 8:

Pinky, a high school girl, has difficulty in combing her hair and climbing up stairs since 6 months. On examination, Gower sign is positive and maculopapular rash present over MCP joints. What is the next appropriate investigation to be done?



- a) Nerve conduction studies
- b) Serum creatine kinase
- c) Electromyography
- d) Muscle biopsy

Question 9:

A 1-day-old girl is being evaluated for decreased muscle tone. On examination, thin cheeks, high arched palate, inverted V-shaped upper lip, and a narrow head are observed. Facial muscle wasting is prominent. She has a club foot deformity. Her mother has a history of delayed relaxation of hand muscles after opening doors. What is the most likely diagnosis?

- a) Myotonic dystrophy
- b) Spinal muscular atrophy
- c) Duchenne muscular dystrophy
- d) Emery-Dreifuss dystrophy

Question 10:

A small baby who was being evaluated for multiple bone fractures with minor trauma is found to have brittle bone disease. Which of the following is false regarding this disease?

- a) It is caused by defects in type 2 collagen.
- b) Predominantly autosomal-dominant inheritance is seen.
- c) Blue sclerae and deafness may be associated.

d) Biphosphonates are useful in the treatment.

Question 11:

All of the following are radiological signs of scurvy except _____.

- a) White line of Frankel
- b) Cupping and fraying
- c) Ground glass appearance
- d) Zone of rarefaction

Question 12:

A 6-month-old infant presents with anemia and hepatosplenomegaly. X-ray imaging done as a part of the skeletal survey showed the following image. What is the diagnosis?



- a) Achondroplasia
- b) Scurvy
- c) Osteogenesis imperfecta
- d) Osteopetrosis

Answer Key

Question No.	Correct Option
1	a
2	b
3	a
4	a
5	a
6	a
7	a
8	b
9	a
10	a
11	b
12	d

Detailed Explanations

Solution to Question 1:

The finding of enlarged calf muscles and atrophied thigh muscles with a history of muscle weakness and waddling gait are suggestive of Duchenne muscular dystrophy (DMD).

Duchenne muscular dystrophy is the most common hereditary neuromuscular disease. It is an X-linked recessive disorder. In DMD, pseudohypertrophy of the calf muscles which occurs due to muscle enlargement by the infiltration of fat. It is paradoxically weak, unlike the muscle that enlarges due to hypertrophy. Pseudohypertrophy of calf muscles is the characteristic feature of DMD, whereas the other 3 conditions mentioned in the options do not have this feature.

Clinical features:

- Progressive weakness with developmental delay
- Pseudohypertrophy of calves
- Wasting of thigh muscles
- Characteristic Trendelenburg gait or hip waddle.
- Gower's sign
- Dilated cardiomyopathy

Diagnosis:

- Elevated serum creatine kinase levels
- Polymerase chain reaction (PCR) analysis for dystrophin gene

- If PCR is normal, but clinical suspicion is high, muscle biopsy with dystrophin immunocytochemistry is done.

Treatment includes supportive management for cardiac decompensation, nutritional management, physiotherapy, and glucocorticoids. Death usually occurs by 18 to 20 years of age due to respiratory failure, intractable heart failure, pneumonia, or aspiration.

Note- The FDA has approved Vamorolone, a novel corticosteroid for the treatment of DMD. The drug offers enhanced anti-inflammatory properties and mineralocorticoid receptor antagonism as well. It is potentially a safer and more effective alternative to glucocorticoids.

Other options:

Option B: Myotonic dystrophy presents with distal muscle weakness, proximal muscle involvement is very late.


Option C: Spinal muscular atrophy (SMA), type 1 (which is most common) presents in infancy. Death occurs early.

Option D: Emery–Dreifuss muscular dystrophy presents with predominant involvement of scapular and humeral muscles. The given question describes the involvement of lower limb muscles. Hence, Emery–Dreifuss dystrophy, in this case, is less likely.

Solution to Question 2:

The structure marked as B in the image represents dystrophin protein (a sarcolemmal protein), which is absent in Duchenne muscular dystrophy (DMD) due to a mutation in the dystrophin gene.

This gene is located on the short arm of chromosome X near the p21 locus (Xp21). Lack of dystrophin leads to cellular instability, which leads to progressive leakage of intracellular components.



Solution to Question 3:

Duchenne muscular dystrophy is more severe than Becker's dystrophy. In DMD, tendon reflexes are lost after age 6.

Becker's dystrophy is similar to Duchenne muscular dystrophy with a genetic defect at the same locus (Xp21) but clinically less severe. Gene affected in both is dystrophin, which is the largest gene in humans.

Important differences between Duchenne and Becker's muscular dystrophy:


	Duchenne's dystrophy	Becker's dystrophy
Onset	Age: 2–3 years	Age: 5–15 years
Weakness	Progressive	Variable
Intellectual impairment	Common	Uncommon
Prognosis	Wheelchair dependant by adolescence	Ambulatory till early adult life
Death	By the age of 18–20 years	By 40s or 50s

Solution to Question 4:

The image shows the Gowers sign. Patients use their hands to “climb up” their legs when asked to rise from a prone position to compensate for proximal lower-extremity weakness. An early Gowers sign is often evident by age of 3 years and is fully expressed by age of 5 or 6 years.

This maneuver can be seen in patients with:

- Duchenne muscular dystrophy
- Becker's muscular dystrophy
- Limb-girdle muscular dystrophy
- Facioscapulohumeral dystrophy
- Polymyositis
- Dermatomyositis



Solution to Question 5:

The image shows winging of the scapula, which when combined with this patient's features are suggestive of facioscapulohumeral muscular dystrophy.

Facioscapulohumeral dystrophy is an autosomal dominant condition that generally presents with asymmetric weakness in facial and shoulder girdle muscles. Occasionally, finger and wrist weakness is the first symptom.

The mouth is rounded and appears puckered because of the protruding lips. Upper facial weakness will manifest as an inability to close the eyes completely in sleep. Scapular winging is prominent, even in infants. Flattening or even concavity of the deltoid contour is seen, and the biceps and triceps brachii muscles are wasted and weak. Eventually, proximal muscles get involved and undergo atrophy, and Gowers sign & Trendelenburg gait appear. Lumbar lordosis, foot drop, and kyphoscoliosis are common complications.

Treatment includes orthopedic measures for scoliosis and foot drop, surgical intervention for scapular deformity. Physiotherapy has no value.

Other options:

Long thoracic nerve palsy: No history of trauma or surgery rules out this diagnosis.

Klippel-Feil syndrome: Classic triad of Klippel-Feil syndrome (congenital musculoskeletal condition)

- low posterior hairline
- short neck
- decreased cervical range of motion

Myotonic dystrophy patients have a characteristic V-shaped upper lip, a history of myotonia, and progressive weakness.

Solution to Question 6:

The clinical picture showing pain in the calves occurring within a week of an upper respiratory tract infection (URI), with elevated creatine kinase levels suggests the diagnosis of viral myositis. The specific history of URI suggests influenza myositis.

Infectious myositis has a male predominance and is typically seen in young adults. Childhood form presents with fever, malaise, and rhinorrhea and is usually followed by severe pain, especially in the calves. The muscle pain is usually worse with movement and the symptoms last for about a week. In children, toe walking and wide-based gait may be seen because of the involvement of the gastrocnemius-soleus muscles.

Other viral causes of myositis include HIV-1 (one of the most common causes of myositis), group B coxsackievirus (epidemic myalgia), HTLV-1 (Human T-lymphotropic virus), and cytomegalovirus (CMV).

Lab features of influenza myositis include elevated CK which may be as high as 500 times normal the value. Urine myoglobin is usually positive. Detection of the virus with PCR testing of nasopharyngeal specimens can be carried out but is not of much clinical significance. The histopathological examination would reveal muscle fiber necrosis without much inflammatory change.

Treatment includes bed rest, IV fluids, and symptomatic management with analgesics and antipyretics. Amantadine and other antiviral agents may be considered in adults. Complications include myocarditis and respiratory dysfunction. Rhabdomyolysis is an uncommon complication.

Solution to Question 7:

Among the given options, only Spinal muscular atrophy presents in infancy. The rest present in late childhood.

Spinal muscular atrophy (SMA) is a degenerative disease of motor neurons, that is inherited in an autosomal recessive manner. It usually starts in infancy with hypotonia, hyporeflexia, and thin muscle mass (atrophy). Polyhydramnios may be present prenatally due to weakness in swallowing.

Diagnosis is done by genetic testing for SMN gene mutation. Muscle biopsy is done for cases with negative genetic testing and strong clinical suspicion. The muscle biopsy in infancy shows a characteristic pattern of perinatal denervation that is unlike that of a mature muscle. Groups of giant type I fibers are mixed with fascicles of severely atrophic fibers.

Treatment is symptomatic and supportive.

Recent updates:

- Nusinersen is an antisense oligonucleotide approved by the FDA for treating SMA. It is administered intrathecally.
- Risdiplam (first oral drug) has been approved by FDA to treat infants >2 months of age with SMA. It is an oral splicing modifier that corrects the splicing defect of SMN2, which leads to

increased levels of full-length SMN protein.

Solution to Question 8:

A maculopapular rash over the metacarpophalangeal joints is indicative of Gottron papules in a child with proximal muscle weakness is suggestive of juvenile dermatomyositis. As it is an inflammatory myopathy, the next investigation is serum creatine kinase.

Though the gold standard investigation is a muscle biopsy, it is indicated only when the diagnosis is in doubt or when grading of the disease severity is required. Nerve conduction studies are usually normal unless associated with severe muscle necrosis and atrophy. Electromyography would show the characteristic signs of myopathy and muscle fiber necrosis but is preferably performed in a specialized center.

Solution to Question 9:

The patient's clinical features along with the history of myotonia in the mother are suggestive of Myotonic dystrophy. Facial weakness, inverted V-shaped upper lip, and loss of muscle mass in the temporal fossae are characteristic of myotonic muscular dystrophy.

A severe congenital form of myotonic dystrophy appears in infants born to mothers with symptomatic myotonic dystrophy. It is inherited as an autosomal-dominant trait. Clinical features include:

- Skeletal deformities (clubfoot deformities alone or congenital contractures involving many joints and can even include the cervical spine).
- Generalized hypotonia and weakness at birth
- Facial wasting and weakness
- Inverted V-shaped upper lip
- Loss of muscle mass in the temporal fossa

Infants can require gavage feeding or ventilator support for respiratory muscle weakness or apnea.

Other Options:

Spinal muscular atrophy (SMA), type 1 (which is most common) presents in infancy with hypotonia but has no facial features that are present in this child.

In Duchenne muscular dystrophy, the baby appears normal at birth and the facial features present in this child are not seen.

Emery–Dreifuss muscular dystrophy presents with predominant involvement of scapular and humeral muscles. The given scenario makes this diagnosis less likely.

Solution to Question 10:

Brittle bone disease is another name for osteogenesis imperfecta (OI). It is due to defects in type 1 collagen, not type 2 collagen.


There are 12 types of OI, with type 1 being the most common. Triad of fragile bones, blue sclera, and deafness is known as van der Hoeve syndrome. Treatment includes bisphosphonate therapy and orthopedic rehabilitation.

Solution to Question 11:

White line of Frankel, ground-glass appearance, and zone of rarefaction are the radiological signs seen in scurvy.

Radiological signs in scurvy includes:

- White line of Frankel: calcified cartilage at metaphyses
- Trummerfeld zone: lucent metaphyseal band underlying Frankel line
- Pelkan spur: lateral prolongation of the white line, which is present at the cortical ends
- Wimberger ring sign: circular, opaque radiologic shadow surrounding epiphyseal centers of ossification



Solution to Question 12:

Anemia and hepatosplenomegaly in an infant with the X-ray image showing dense bones, and bone within the bone appearance is suggestive of osteopetrosis.

Osteopetrosis/Marble bone disease/Albers-Schönberg disease

It is a disorder involving defective bone resorption due to mutation of CLCN7 in osteoclasts.

Clinical features include macrocephaly, developmental delay, anemia (reduced bone marrow cavity), and hepatosplenomegaly (compensatory extramedullary hematopoiesis). Low serum calcium and phosphorus levels with elevated parathyroid hormone (PTH) and normal vitamin D levels are observed. Radiological features include diffuse bone sclerosis with clubbing of metaphyses. Bone within the bone appearance and "sandwich" appearance of the vertebral bodies can be seen (image). Treatment is symptomatic. Hematopoietic stem cell transplantation can be done in severe cases.

Achondroplasia usually presents with pulmonary symptoms like dyspnea, obstructive sleep apnea, and recurrent infections.

In both scurvy and osteogenesis imperfecta (OI), the child cries on touching. On radiography, certain typical features are seen in scurvy, while multiple fractures are seen in OI.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Paediatric Rheumatology

Question 1:

Most common rheumatic disease in children is _____.

- a) Juvenile idiopathic arthritis
- b) Juvenile ankylosing spondylitis
- c) Systemic lupus erythromatosus
- d) Rheumatic fever


Question 2:

You are evaluating JIA patients in a pediatric rheumatology clinic. As per the ILAR criteria, which of the following patients would you not include under JIA?

- a) A 3-year-old girl with Psoriatic arthritis
- b) A 10-year-old boy with Enthesitis-related arthritis
- c) A 18-year-old with RF negative Polyarthritis first diagnosed 3 years back
- d) A 14-year-old female with Reactive arthritis

Question 3:

Mohit is brought to the pediatrician by his mother because of red-eye. He complains of blurring of vision and on examination, the pupil appears as shown in the picture. Which of the following subtype of JIA is associated with the highest risk for the ophthalmic complication noted in this patient?



- a) RF-positive polyarthritis
- b) Systemic JIA
- c) ANA-positive oligoarthritis
- d) Enthesitis-related arthritis

Question 4:

A worried mother brings her 4-year-old girl who has been complaining about pain in the joints of arms and legs for 2 months. On examination, multiple nodules over the bony prominences suggestive of rheumatoid nodules are noted. On investigation, RA-factor was found positive pointing to the diagnosis of RF-positive polyarthritis. Which of the following complication is not likely to occur in this patient?

- a) Hip joint erosion
- b) Atlantoaxial subluxation
- c) Micrognathia
- d) Ankylosis

Question 5:

Which of the following is false regarding systemic Juvenile Idiopathic Arthritis?

- a) Positive RA factor
- b) Fever with evanescent rash
- c) Hepatosplenomegaly

d) Generalized Lymphadenopathy

Question 6:

Anuj, a school-going kid has been brought to the pediatric OPD by his mother with complaints of fever for 3 weeks. She says her son has been skipping school because of the fever but in the evening he gets better and plays with his siblings. She also noticed a rash that appears along with fever and disappears spontaneously. He has a history of joint pain on and off since pre-school. On examination, generalized lymphadenopathy and hepatosplenomegaly are noted along with an evanescent erythematous rash on the trunk. Which of the following characteristically defines the fever associated with this child's condition?

- a) Quartan fever
- b) Tertian fever
- c) Quotidian fever
- d) Continuous fever

Question 7:

Aarav is a known case of juvenile idiopathic arthritis (JIA). He is referred to a Rheumatology specialist center from a local hospital, when his condition started deteriorating on steroid treatment. On examination, the patient has lymphadenopathy, hepatosplenomegaly, and purpura along with the acute onset of high spiking fevers. Vitals show persistent hypotension (85/45 mmHg), tachycardia (160/min), and oliguria. He was urgently admitted into the PICU. The resident physician is confused between macrophage activation syndrome (MAS) and JIA flare-up. Which of the following is best suited to distinguish these two?

- a) Hyperferritinemia
- b) Hypofibrinogenemia
- c) Hypertriglyceridemia
- d) Hemophagocytosis

Question 8:

A mother brought her 12-day old baby to the OPD due to the rash shown in the picture for the last 5 days. The mother says the rash gets worse in the sun. The infant was born at home and she never had a prenatal check-up. She herself has a history of joint pains and a facial rash which she says is "no big deal". On examination, Vitals are normal for the baby. A subsequent ECG shows P wave rate 120/min and R wave rate at 70/min with no relation between the two.

Which of the following antibodies will be used to diagnose this baby's condition?



- a) Anti-Jo antibodies
- b) Anti- dsDNA antibodies
- c) Anti-Ro antibodies
- d) Antihistone antibodies

Question 9:

Vikas, a high school student is observed to have an indented 'en coup de sabre' lesion as shown in the image below. What is the most likely diagnosis?



- a) Neonatal Lupus
- b) Systemic JIA
- c) Juvenile Dermatomyositis
- d) Juvenile Scleroderma

Question 10:

A 9-year-old boy has swelling of knee joints and non-blanching rashes as seen in the picture. Urine exam reveals haematuria +++ and proteinuria +. Platelet levels are normal. What is the probable diagnosis?



- a) Systemic lupus erythematosus (SLE)
- b) Henoch Schonlein purpura (HSP)
- c) Immune thrombocytopenic purpura (ITP)
- d) Polyarteritis Nodosa (PAN)

Question 11:

Pinky, a small child with Kawasaki disease is admitted to the ward for treatment. Which of the following feature would you not expect to see in her?

- a) Purulent conjunctivitis
- b) Cervical lymphadenopathy
- c) Perineal desquamation

d) Polymorphous rash

Question 12:

A worried father brought her 5-year-old son to the pediatrician with a history of fever, rash, irritability, and poor feeding. A few days back, he developed perianal desquamation and non-purulent conjunctivitis with edema of the hands and feet. He has been up to date with his immunization. Tongue findings are shown in the image. Which of the following is not likely to occur in the child?



- a) Elevated ESR**
- b) Thrombocytopenia**
- c) Myocarditis**
- d) Coronary artery aneurysm**

Question 13:

Nisha presents to the clinic with clinical features suggestive of Kawasaki disease. She was admitted to the ward for treatment. Which of the following is the treatment of choice for this disease?

- a) IV Immunoglobulins**
- b) Steroids**
- c) Cyclophosphamide**
- d) Mycophenolate mofetil**

Question 14:

Who among the following children with Kawasaki's disease is at greatest risk for coronary aneurysms?

- a) 8 year-old girl with 6 days of high fever and strawberry tongue
- b) 5-year-old boy with 8 days of fever and CSF pleocytosis
- c) 2-year-old girl with IVIG treatment at 7th day of fever
- d) 11-year-old boy with 15 days of fever and low PCV

Question 15:

A 15-year-old boy comes to the clinic complaining about recurrent painful oral ulcers as shown in the picture. He also experiences repeated episodes of eye redness and pain which subsides with medication. On examination, similar ulcers are seen on his genitals. Behçet disease is suspected. Which of the following feature would you not expect to see in this patient?



- a) Dural sinus thrombosis
- b) Pulmonary aneurysms
- c) Nailfold capillary abnormalities
- d) Tender red nodules on the legs

Answer Key

Question No.	Correct Option
1	a
2	d
3	c
4	d
5	a
6	c
7	d
8	c
9	d
10	b
11	a
12	b
13	a
14	d
15	c

Detailed Explanations

Solution to Question 1:

The most common rheumatic disease in children is juvenile idiopathic arthritis (JIA).

Juvenile idiopathic arthritis represents a heterogeneous group of chronic disorders characterized by

- Arthritis* in ≥ 1 joint
- Age at onset: < 16 years
- Duration of disease: ≥ 6 weeks

*Arthritis clinically defined as swelling or effusion, or the presence of 2 or more of the following signs:

- limitation of range of motion
- tenderness or pain on motion
- increased heat

Solution to Question 2:

Reactive arthritis is not a subtype of Juvenile Idiopathic Arthritis (JIA) so you would not include this subset of patients under your evaluation.

ILAR (International League of Associations for Rheumatology) defines JIA as arthritis of one or more joints with onset below 16 years of age and persisting for at least 6 weeks.

The subtypes of Juvenile Idiopathic Arthritis (JIA):-

- Oligoarthritis (4 or lesser joints) (Most common type)
- Polyarthritis (5 or more joints)
- RF Positive
- RF Negative
- Systemic JIA
- Psoriatic arthritis
- Enthesitis-related arthritis
- Undifferentiated

Solution to Question 3:

The patient has red-eye and the blurring of vision along with the finding of the irregular pupil, which is suggestive of uveitis. ANA-positive oligoarthritis is associated with the highest risk for uveitis in patients with Juvenile Idiopathic Arthritis.

The presence of a positive antinuclear antibody (ANA) confers increased risk for uveitis in children with JIA and requires periodic slit-lamp examination every 3 months.

Solution to Question 4:

Ankylosis is not seen in patients with RF-positive Polyarthritis.

Polyarthritis is characterized by inflammation of ≥ 5 joints in both upper and lower extremities. Rheumatoid factor (RF)-positive polyarthritis resembles the characteristic symmetric presentation of adult rheumatoid arthritis. Rheumatoid nodules on the extensor surfaces of the elbows, spine, and over the Achilles tendons are associated with a more severe course and almost exclusively occur in RF-positive individuals.

Complications may include:

- Micrognathia (due to chronic temporomandibular joint disease)
- Cervical spine involvement, manifesting as decreased neck extension, occurs with a risk of atlantoaxial subluxation and neurologic sequelae.
- Hip disease may be subtle, with findings of erosion.

Solution to Question 5:

Rheumatoid arthritis (RA) factor is negative in systemic JIA. It is positive in a subtype of polyarthritic type of JIA.

Systemic JIA is characterized by a fever of at least 2 weeks and Arthritis in 1 or more joints + at least one of

- Evanescent erythematous rash
- Prominent visceral involvement, including hepatosplenomegaly
- Generalized lymphadenopathy
- Serositis

Solution to Question 6:

The patient has clinical features of systemic juvenile idiopathic arthritis (sJIA), which is associated with quotidian fever. Quotidian fever - fever that rises to 39°C (102.2°F) once a day and returns to 37°C (98.6°F) between fever peaks

Systemic JIA is characterized by a fever of at least 2 weeks and Arthritis in 1 or more joints + at least one of

- Evanescent erythematous rash that is non-pruritic and migratory with lesions lasting <1 hr.
- Prominent visceral involvement, including hepatosplenomegaly
- Generalized lymphadenopathy
- Serositis

Solution to Question 7:

Macrophage activation syndrome (MAS) is characterized by all of the listed features. However, the diagnosis is confirmed by bone marrow biopsy demonstrating hemophagocytosis and is the best test to distinguish MAS from a flare-up of JIA.

MAS is a rare but potentially fatal complication of systemic JIA. It is characterized by an uncontrolled activation and proliferation of T lymphocytes and macrophages. It is the result of a cytokine storm that leads to a dysregulated inflammatory activation of the immune system, with rapid progression to multiorgan failure if not promptly diagnosed and properly treated. It is characterized by the following:

- Macrophage hemophagocytosis in the bone marrow
- Hypofibrinogenemia (paradoxical fall in erythrocyte sedimentation rate [ESR])
- Hypertriglyceridemia

- Extreme hyperferritinemia
- Fever $\geq 38.5^{\circ}\text{C}$
- Splenomegaly
- Cytopenia affecting at least two of three cell lineages

Treatment usually includes high-dose corticosteroids and immunosuppressive agents in refractory cases.

Solution to Question 8:

The baby has raccoon eyes (periorbital rash) with complete heart block (P-R dissociation) suggestive of neonatal lupus erythematosus (NLE). Anti-Ro antibodies are used to diagnose neonatal lupus.

Neonatal Lupus Erythematosus(NLE):

NLE is not an autoimmune condition. It occurs when maternal autoantibodies enter the fetal circulation (passive transfer). The antibodies responsible for this condition are anti-Ro antibody (anti-SSA) and anti-La antibody (anti-SSB).

The periorbital rash typically appears within 6 weeks of life and lasts 3-4 months. However, it can be present at birth. Infants may also have cytopenias and hepatitis, but the most feared complication is congenital heart block. Cardiomyopathy is seen in the most severe cases. The noncardiac manifestations of neonatal lupus are usually reversible, whereas congenital heart block is permanent. Conduction system abnormalities can be detected in utero by fetal echocardiogram beginning at 16 weeks of gestational age.

Fluorinated corticosteroids seem to improve cases of hydrops. Significant conduction system abnormalities after birth are treated with cardiac pacing and occasionally intravenous immunoglobulin and steroids, while severe cardiomyopathy may require cardiac transplantation. Transient, noncardiac manifestations are conservatively managed, with topical steroids used occasionally to treat the rash.

Other options:

Anti-Jo antibodies are a type of myositis-specific antibodies. These are not related to neonatal lupus.

Anti-dsDNA antibodies are specific for SLE (Systemic Lupus erythematosus).

Anti-histone antibodies are present in Drug-induced SLE.

Note:

SS in SSA and SSB stands for Sjögren's syndrome-related. Anti-SSA and anti SSB are antinuclear antibodies and are mainly found in patients with SLE and Sjögren's syndrome.

Solution to Question 9:

En coup de sabre lesion is seen in juvenile localized scleroderma (morphea).

It appears as an indented, vertical, colourless, line of skin on the forehead. The indentation looks as though the person may have been struck (coup) by a sword (sabre). This lesion involves the scalp and/or face and often extend into the central nervous system, resulting in neurologic sequelae such as seizures and headaches.

Solution to Question 10:

The given clinical scenario showing swelling of knee joints (arthritis), non-blanching rash in the lower limbs (palpable purpura), along with hematuria and proteinuria (renal involvement) in a child is suggestive of Henoch Schonlein purpura (HSP).

HSP (IgA vasculitis) is a small vessel vasculitis most commonly seen in the age group of 3-8 years. It generally follows an upper respiratory tract infection and there is a history of atopy in one-third of the cases.

The pathology of HSP involves leukocytoclastic vasculitis and the deposition of IgA in the affected blood vessels. IgA is deposited in the glomerular mesangium in a distribution similar to that of IgA nephropathy.

Clinical manifestations include palpable purpura, the hallmark feature of HSP, which characteristically involves the extensor surfaces of arms and legs. Musculoskeletal symptoms such as arthritis and arthralgia are common. Gastrointestinal symptoms can include abdominal pain, vomiting, diarrhea, and bleeding. Renal involvement may lead to gross or microscopic hematuria, nephritic syndrome, or nephrotic syndrome. Renal manifestations are more severe in adults. A rapidly progressive form of glomerulonephritis is seen in some cases. Nephritis usually follows the appearance of the rash.

Diagnosis of HSP is primarily based on clinical findings, including the palpable purpura, along with the presence of other symptoms. Laboratory tests are not specific to HSP but may show nonspecific markers of inflammation and sometimes elevated levels of IgA. Imaging studies and biopsies of affected tissues can provide additional diagnostic information

Treatment of HSP is generally supportive, focusing on hydration, pain relief, and nutrition. Corticosteroids may be used in severe cases or for significant gastrointestinal involvement. Intravenous immune globulin (IVIG) and plasma exchange are occasionally employed for severe disease. Nephritis due to IgA vasculitis should be treated with immunosuppressants like prednisolone and azathioprine. Complications of HSP can include serious gastrointestinal problems, kidney disease, and recurrences, but overall, the prognosis for childhood HSP is favorable. Regular monitoring of blood pressure and urine is recommended to detect and manage potential complications.

Solution to Question 11:

Purulent conjunctivitis is not a feature of Kawasaki disease. Bilateral nonexudative conjunctival infection with limbal sparing is characteristically seen in Kawasaki disease.

Kawasaki disease is also known as mucocutaneous lymph node syndrome and is commonly seen in children between 3 months to 5 years of age.

Solution to Question 12:

The above clinical features are suggestive of Kawasaki disease. Thrombocytosis is a characteristic finding in Kawasaki disease and commonly occurs in the second week.

Important laboratory findings of Kawasaki disease are anemia, elevated ESR, and CRP. Major complications of Kawasaki disease are arteritis and aneurysms of coronary vessels, myocarditis, myocardial infarction, and pericarditis.

Solution to Question 13:

Treatment of choice in Kawasaki disease is IV immunoglobulins. IV immunoglobulins along with aspirin have proven beneficial in preventing coronary artery abnormalities.

Solution to Question 14:

An 11-year-old boy with 15 days of fever and low PCV is at the greatest risk for coronary aneurysms.

Risk factors for coronary aneurysms in Kawasaki's disease:

- Age younger than 1 year or older than 9 years
- Male sex
- Fever ≥ 14 days
- Hematocrit(PCV) < 35 percent
- Serum sodium concentration < 135 mEq/L
- White blood cell count $> 12,000/\text{mm}^3$
- Elevated ESR and CRP

Administration of IVIG during the first 10 days of illness reduces the prevalence of coronary aneurysms fivefold.

Solution to Question 15:

Behcet disease is not associated with nail fold capillary abnormalities which is typically seen in systemic sclerosis.

Behcet disease is a neutrophilic vasculitis affecting small- to medium-sized vessels that classically presents as a clinical triad of recurrent oral aphthous ulcers, genital ulcers, and uveitis. Other skin findings include erythema nodosa (red tender nodules on legs) and positive pathergy test (Pustular reaction, 24-48 hrs after sterile needle puncture or normal saline injection).

There can also be gastrointestinal and pulmonary manifestations, with disease mortality related to severe neurologic involvement or rupture of vascular aneurysms. Dural sinus thrombosis is the most common CNS manifestation in children. Pulmonary aneurysms are the most severe feature of pediatric BD.

It is associated with HLA subtype B51. T helper 17(Th17) cells play an important contribution in the recruitment of neutrophils, which will later infiltrate into the vessel wall.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Pediatric Hematology - Introduction, Bleeding, and Clotting Disorders

Question 1:

Which of the following is the predominant site of hematopoiesis in a 20-weeks old fetus?

- a) Yolk sac
- b) Liver
- c) Spleen
- d) Bone marrow

Question 2:

Electrophoresis and globin gene sequencing are performed on a normal embryo's blood sample as a part of hematology research. Which of the following type of hemoglobin is likely to be present?

- a) HbM Iwate
- b) Hb Portland
- c) Hb Auckland
- d) HbM Boston

Question 3:

When does hemoglobin synthesis switch from HbF to HbA?

- a) At 8 weeks of intrauterine life
- b) At 24 weeks of intrauterine life
- c) At 38 weeks of intrauterine life
- d) At 6 months of postnatal life

Question 4:

Hemoglobin fractions were analyzed in a neonate with a family history of sickle cell disease. The test shows elevated levels of fetal hemoglobin (HbF). Which of the following is true about

HbF?

- a) HbF accounts for 100% of the total hemoglobin at term.
- b) HbF is resistant to alkali denaturation.
- c) HbF has lower affinity for oxygen than HbA.
- d) HbF has higher affinity for 2,3 bisphosphoglycerate than HbA.

Question 5:

You are measuring the life span of RBCs in a normal neonate using chromium-labeled erythrocyte studies as a part of a research project under the clinical hematology department. What would be the result?

- a) 90-120 days
- b) 60-90 days
- c) 45-60 days
- d) 30-45 days

Question 6:

A concerned mother brings her preschool child to you. She states that her child was fine until yesterday night and now is covered with bruises and purple dots, which she had noticed while changing his clothes in the morning. She informs you that one month back, her child had a cough and cold. Findings on physical examination are normal, other than petechiae and purpura. Which of the following is most likely to be seen while investigating this child?

- a) Normal platelet count
- b) Prolonged prothrombin time
- c) Prolonged activated partial thromboplastin time
- d) Prolonged bleeding time

Question 7:

A pediatrician sees a child in the OPD. He has bruising, petechiae, and occasional minor epistaxis, which has very little interference with his daily activities. Based on further workup, he diagnoses the child as a case of mild ITP. How will he manage this child?

- a) Short course corticosteroid therapy
- b) Intravenous anti-D administration

- c) A single dose of intravenous immunoglobulin
- d) Education, counseling and watchful expectancy

Question 8:

Veena, an adolescent girl has had recurrent episodes of bleeding from her nose and gums. She also gives a history of heavy menstrual bleeding and repeated blood transfusions. Her bleeding time is deranged and flow cytometry reveals markedly decreased CD41 and CD61 expression. A defect in which of the following is most likely seen in her?

- a) Platelet adhesion
- b) Platelet activation
- c) Platelet aggregation
- d) Platelet count

Question 9:

Which of the following is the most common congenital bleeding disorder?

- a) Hemophilia A
- b) Hemophilia B
- c) Von Willebrand disease
- d) Factor VII deficiency

Question 10:

Shortly after birth through normal vaginal delivery, a neonate develops petechiae and purpura. You suspect bleeding disorder and send samples for further workup. A prolonged bleeding time, normal platelet count, normal aPTT is seen along with large platelets in the peripheral smear. What is the likely diagnosis?

- a) Bernard-Soulier syndrome
- b) Glanzmann's thrombasthenia
- c) von Willebrand disease
- d) Evans syndrome

Question 11:

A panicked father brings his adopted daughter to your OPD with complaints of bleeding gums. Hematological investigations reveal a normal platelet count and a prolonged bleeding time. The ristocetin-aggregation test was abnormal. Which of the following would you include in your differential diagnosis?

- a) von Willebrand disease
- b) Bernard-Soulier syndrome
- c) Glanzmann thrombasthenia
- d) Both A and B

Question 12:

An ENT surgeon refers a child who had developed prolonged bleeding post-tonsillectomy to the hematology department for evaluation. Workup revealed a qualitative defect in the von Willebrand factor. Which of the following disease is more likely in this patient?

- a) Type 1 Von Willebrand disease
- b) Type 2 Von Willebrand disease
- c) Type 3 Von Willebrand disease
- d) Type 4 Von Willebrand disease

Question 13:

Which type of von Willebrand disease is associated with thrombocytopenia?

- a) Type 2M
- b) Type 2N
- c) Type 2A
- d) Type 2B

Question 14:

A pediatric resident administers 0.5-1 mg of phytonadione intramuscularly to all newborn infants in the ward. Deficiency of which of the following factors is prevented by this medication?

- a) Factor V
- b) Factor VII

- c) Factor VIII
- d) Factor XI

Question 15:

The most common bleeding manifestation seen in severe hemophilia is _____.

- a) Recurrent hematomas
- b) Recurrent hemarthrosis
- c) Hematuria
- d) Intracranial hemorrhage

Question 16:

Which of the following factor deficiencies are matched to the correct disease?

- a) Factor V deficiency - Stuart-Prower disease
- b) Factor IX deficiency - Parahemophilia
- c) Factor X deficiency - Christmas disease
- d) Factor XI deficiency - Hemophilia C

Question 17:

Dinesh, a 2-year-old boy presented with a painful left ankle swelling and difficulty walking. In the past, he had presented with significant hematomas at immunization sites. On investigation, he was found to have haemophilia A. What is the treatment of choice in this condition?

- a) Cryoprecipitate
- b) Fresh frozen plasma
- c) Prothrombin complex concentrates (PCCs)
- d) Factor VIII concentrate

Question 18:

A grandmother has noticed that her granddaughter's umbilical stump hasn't fallen off even after 4 weeks. On examination, the pediatrician notices mild bruising. Which of the following

clotting factor deficiencies should he suspect?

- a) Factor VII deficiency
- b) Factor IX deficiency
- c) Factor XI deficiency
- d) Factor XIII deficiency

Answer Key

Question No.	Correct Option
1	b
2	b
3	c
4	b
5	b
6	d
7	d
8	c
9	c
10	a
11	d
12	b
13	d
14	b
15	b
16	d
17	d
18	d

Detailed Explanations

Solution to Question 1:

The liver is the predominant site of hematopoiesis in the fetus during mid-pregnancy.

Sites of hematopoiesis:

- 2–6 weeks- yolk sac (mesoblastic phase)
- 8–24 weeks- liver (hepatic phase)
- 28 weeks onwards- bone marrow (myeloid phase)

Note:

There is some overlap between the different phases of hematopoiesis. The ranges given above refer to the predominant site of hematopoiesis.

Solution to Question 2:

Hb Portland is likely to be present as it is embryonic hemoglobin.

Gower-I ($\zeta 2\varepsilon 2$), Gower-II ($\alpha 2\varepsilon 2$), and Hb Portland ($\zeta 2\gamma 2$) are called embryonic hemoglobins and are seen at 8 weeks of intrauterine life.

HbM Iwate (option A) and Hb Auckland (option C) and HbM Boston (option D) result from globin mutations that stabilize iron in the ferric state, leading to congenital methemoglobinemia.

Solution to Question 3:

Hemoglobin synthesis switches from HbF to HbA at 38 weeks of intrauterine life.


Hemoglobin types at various periods:

- Chromosome 16 contains alpha (α) gene cluster (3 genes):
 - zeta (ζ)
 - alpha 1 ($\alpha 1$)
 - alpha 2 ($\alpha 2$)
- Chromosome 11 contains beta (β) gene cluster (5 genes):
 - epsilon (ϵ)
 - 2 gamma (γ)
 - delta (δ)
 - beta (β)

Changes in globin subunits from embryo to early infancy:

Time	Major hemoglobin
8th week of intrauterine life:	Embryonic hemoglobins, Gower-1 ($\zeta 2\varepsilon 2$), Gower-2 ($\alpha 2\varepsilon 2$), and Portland ($\zeta 2\gamma 2$) are formed
9th week to birth	Fetal hemoglobin (HbF) ($\alpha 2\gamma 2$) is the major hemoglobin

Time	Major hemoglobin
24th week	HbF constitutes 90% of the total hemoglobin
38 weeks	The switch to the nearly exclusive synthesis of adult hemoglobin (HbA; $\alpha_2\beta_2$)
At birth	HbF constitutes 70% of the total hemoglobin
6 month	adult pattern HbA > 95%, HbA ₂ < 3.5%, HbF < 2.5%



Solution to Question 4:

HbF is resistant to alkali denaturation.

Elevated levels of HbF are seen in sickle cell disease and β -thalassemia, where there is a mutation of the β globin gene. Increased HbF levels persist in childhood and later.

Properties of fetal hemoglobin (HbF)

- HbF is resistant to alkali denaturation.
- This property forms the basis of the alkali denaturation test (or Apt test) to detect the presence of fetal blood in maternal circulation in suspected fetomaternal hemorrhage.
- The Kleihauer–Betke test (or acid elution test) is used to quantitate the amount of fetomaternal hemorrhage to calculate the dosage of anti-D required in Rh-negative mothers.
- HbF accounts for 70% of the hemoglobin at term.
- Increased HbF can be seen in hereditary persistence of fetal hemoglobin, sickle cell anemia, beta-thalassemia, HbC disease, and HbE disease

- HbF has a higher affinity for oxygen.
- This allows for gas exchange from maternal circulation by binding to oxygen at a relatively low oxygen tension in the maternal blood supplying the chorion.
- HbF has lower affinity for 2,3 bisphosphoglycerate (BPG) than HbA.
- Lack of interaction between HbF and 2,3 BPG increases the affinity of HbF to oxygen.

Solution to Question 5:

The lifespan of neonatal RBCs is 60-90 days.

The short lifespan of neonatal RBCs is the cause for rapid fall in hematocrit immediately after birth. The highest hematocrit during a person's lifetime occurs at birth, and the lowest occurs about 10 weeks later.

The lifespan of RBCs in preterm infants is 35-50 days.

The average lifespan of RBCs in adults is 120 days.

Solution to Question 6:

An otherwise well-baby with acute onset of petechiae and purpura with a preceding history of viral infection 1 to 4 weeks prior to the onset of thrombocytopenia is a classical presentation of (autoimmune) idiopathic thrombocytopenic purpura. Prolonged bleeding time is a feature of ITP.

The most common cause of acute onset of thrombocytopenia in an otherwise well-child is (autoimmune) idiopathic thrombocytopenic purpura (ITP).

ITP is a bleeding disorder and has prolonged bleeding time, low platelet count, and normal coagulation profile (prothrombin time [PT] and activated partial thromboplastin time [aPTT]).

ITP has an acute presentation with a self-limited course in children.

Features of ITP:

- Commonly affects children between 1-4 years
- Preceded by a viral infection
- Mucocutaneous bleeding
- Ecchymoses and petechiae
- Wet purpura (blood blisters in the mouth)
- Retinal hemorrhages
- Bleeding into the CNS, which may be life-threatening
- Hepatomegaly, splenomegaly, and lymphadenopathy, fever, jaundice are absent
- Resolves spontaneously in 80% of the cases in 4-6 weeks.

Laboratory features of ITP:

- Thrombocytopenia (commonly $<20,000/\mu\text{L}$)
- Normal/Increased megakaryocytes in bone marrow
- Prolonged bleeding time
- Normal tests for coagulation (PT/aPTT/INR)
- Eosinophilia and lymphocytosis common
- Peripheral smear is normal
- Bone marrow biopsy shows megakaryocytes

Solution to Question 7:

Other than education, counseling, and watchful expectancy, no therapy is needed for patients with mild ITP.

Classification system for ITP:

- No symptoms.
- Mild symptoms- bruising and petechiae, and occasional minor epistaxis which has very little interference with daily living.
- Moderate symptoms- more severe skin and mucosal lesions, more troublesome epistaxis, and menorrhagia.
- Severe symptoms- bleeding episodes - menorrhagia, epistaxis, and melena - requiring transfusions or hospitalization, symptoms interfering seriously with quality of life.

Treatment of ITP:

- Mild to moderate bleeding symptoms- watchful expectancy, education, and counseling.
- Platelet count $<30,000$ + risk factors for bleeding (antiplatelet drugs/anticoagulants or concomitant bleeding disorders)- oral glucocorticoids/ intravenous immunoglobulin (IVIG) or IV anti-D
- Severe bleeding symptoms- IV methylprednisolone $\pm$ IVIG or IV anti-D
- Life-threatening bleeding (eg. intracranial hemorrhage)- platelet transfusion + IV methylprednisolone + IVIG or IV anti-D

Note:

The role of splenectomy in ITP should be reserved for 2 circumstances:

- The older child (≥ 4 yr) with severe ITP that has lasted >1 year (chronic ITP) and whose symptoms are not easily controlled with therapy
- When life-threatening hemorrhage (ICH) complicates acute ITP and the platelet count cannot be corrected rapidly with transfusion of platelets and administration of IVIG and corticosteroids.

Solution to Question 8:

The given clinical scenario is suggestive of Glanzmann's thrombasthenia which is characterized by defective platelet aggregation.

Decreased CD41 (GPIIb) and CD61 (GPIIIa) expression are diagnostic of Glanzmann's thrombasthenia. Bleeding in Glanzmann thrombasthenia is quite severe and is typically mucocutaneous, including epistaxis, gingival, and GI bleeding.

Bernard-Soulier syndrome and Glanzmann thrombasthenia are severe congenital disorders of platelet function. Patients usually present with petechiae and purpura at birth. The diagnosis is confirmed by flow cytometric analysis of the patient's platelet glycoproteins.

Solution to Question 9:

Von Willebrand disease (VWD) is the most common congenital bleeding disorder.

VWD is thought to affect approximately 0.1- 1% of the population. It is due to abnormal von Willebrand factor (VWF).

VWF is an adhesive protein that acts as a bridge between subendothelial collagen and platelets. It also binds and protects the circulating factor VIII from rapid clearance in circulation.

Von Willebrand disease is mostly inherited as an autosomal dominant trait (rarely autosomal recessive). The types of disease include quantitatively deficient vWF (partial = type 1 or absolute = type 3) or qualitatively abnormal vWF (type 2 = dysproteinemia). The majority (~ 80%) have type 1 disease.

Hemophilia A (factor 8 deficiency) occurs in 1 in 5,000 males. Hemophilia B (factor 9 deficiency) occurs in approximately 1 in 25,000 males.

Solution to Question 10:

Bernard-Soulier syndrome is the likely diagnosis in the given clinical scenario.

The presence of petechiae and purpura at birth, especially after vaginal delivery, along with prolonged bleeding time, normal platelet count, normal aPTT are seen in qualitative platelet defect disorders like Bernard-Soulier syndrome and Glanzmann's thrombasthenia.

The presence of large platelets rules out Glanzmann's thrombasthenia. Evans syndrome is an autoimmune hemolytic disorder with thrombocytopenia. Von Willebrand's disease doesn't usually manifest at birth.

Bernard-Soulier syndrome is an autosomal recessive functional/qualitative platelet disorder. It is characterized by the deficiency of Gp Ib-IX receptor causing defective platelet adhesion. It is due to genetic mutations in the genes forming the GPIb complex of glycoproteins Ib α , Ib β , V, and IX.

Laboratory parameters:

- Normal platelet count (or mild thrombocytopenia)
- Large platelets (megathrombocytes) with dense granules

- Bleeding time abnormally prolonged
- Platelet aggregation tests are normal (with collagen, adenosine diphosphate [ADP], epinephrine)
- Ristocetin aggregation test is abnormal/no response
- von Willebrand factor (vWF) levels are normal.

Solution to Question 11:

The ristocetin-aggregation test is abnormal in both von Willebrand disease and Bernard–Soulier syndrome.

All of the above diseases are inherited bleeding disorders affecting platelet function. Hence, there is prolonged bleeding time but platelet levels are not significantly affected.

Platelet aggregation tests use a panel of platelet agonists (adenosine diphosphate [ADP], epinephrine, collagen, and ristocetin) to measure platelet activation and aggregation in vitro.

Ristocetin induces the binding of VWF to platelets and agglutinates platelets.

Bernard-Soulier syndrome, a severe congenital platelet function disorder, is caused by an absence or severe deficiency of the VWF receptor on the platelet membrane. Hence, in Bernard–Soulier syndrome, no response (or no aggregation) is seen with ristocetin.

The ristocetin-aggregation test is also abnormal in von Willebrand disease. However, it normalizes after the addition of normal plasma (which contains vWF) since there is no defect in binding receptors.

Solution to Question 12:

The given clinical scenario is suggestive of von Willebrand disease (VWD) and a qualitative defect in von Willebrand factor (VWF) is seen in type 2 VWD.

Features of VWD:

- Most common type is Type 1
- Most severe type is Type 3
- Most common inheritance: Autosomal dominant
- Type 1- partial quantitative deficiency of VWF
- Type 2- qualitative deficiency of VWF
- Type 3- complete quantitative deficiency of VWF

There is no type 4 VWD.

Characteristics of the subtypes of VWD are summarized below:

AD: autosomal dominant

AR: autosomal recessive

Type	Inheritance	Mechanism	Important characteristics
1	AD	Partial, quantitative deficiency of VWF	Most common type
3	AR	Complete quantitative deficiency	Most severe VWD
2A	AD	Loss of VWF multimers (increased cleavage by ADAMS13)	Platelet adhesion defect
2B	AD	Increased binding of VWF to platelets causing VWF-platelet complex destruction	Thrombocytopenia
2M	AD	Decreased affinity of VWF to platelets	Platelet adhesion defect, very rare
2N	AR	Decreased affinity of VWF to Factor VIII	Autosomal hemophilia

Solution to Question 13:

von Willebrand disease associated with thrombocytopenia is Type 2B.

Pseudo-VWD: VWF is normal, but platelet has increased affinity to VWF; it presents in a manner similar to VWD 2B with thrombocytopenia.

Type	Inheritance	Mechanism	Important characteristics
1	AD	Partial, quantitative deficiency of VWF	Most common type
3	AR	Complete quantitative deficiency	Most severe VWD
2A	AD	Loss of VWF multimers (increased cleavage by ADAMS13)	Platelet adhesion defect
2B	AD	Increased binding of VWF to platelets causing VWF-platelet complex destruction	Thrombocytopenia
2M	AD	Decreased affinity of VWF to platelets	Platelet adhesion defect, very rare
2N	AR	Decreased affinity of VWF to Factor VIII	Autosomal hemophilia

Solution to Question 14:

Intramuscular injection of 0.5-1 mg of phytonadione is given to all newborn infants because it prevents the deficiency of factor VII that occurs transiently after birth.

Hemorrhagic disease of the newborn is caused by a transient deficiency of vitamin K–dependent factors, factors II, VII, IX, and X.

Causes include:

- Lack of free vitamin K from the mother
- Absence of the bacterial intestinal flora normally responsible for the synthesis of vitamin K.

It is responsive to and prevented by vitamin K therapy (0.5-1 mg of phytonadione).

Solution to Question 15:

Recurrent hemarthrosis is the most common bleeding manifestation of severe hemophilia.

Hemophilia is said to be severe when the residual activity of factor VIII is $\leq 1\%$.

Clinical features of hemophilia:

- Clinically, hemophilia A and B are indistinguishable.
- It presents early in life- bleeding after circumcision or when beginning to walk or crawl.
- Most common bleeding manifestation in severe hemophilia is recurrent hemarthrosis and the most common joint affected is the knee.
- Acute hemarthrosis- local pain, swelling, and erythema; muscle contractures due to fixed positioning.
- Chronic hemarthrosis- synovial thickening and synovitis in response to intraarticular blood.
- Muscle hematomas can lead to compartment syndromes.
- Hematuria is common but self-limited.
- Life-threatening bleeding- bleeding to oropharyngeal spaces, CNS, and retroperitoneum.

Solution to Question 16:

The correct match is factor XI deficiency and hemophilia C.

Clotting factor deficiencies and associated disorders:

Factor V— parahemophilia

Factor VIII— classic hemophilia, hemophilia A

Factor IX— Christmas disease, hemophilia B

Factor X— Stuart-Prower disease

Factor XI— hemophilia C

Solution to Question 17:

The treatment of choice for hemophilia A is factor replacement therapy with factor VIII concentrate.

Cryoprecipitate is used when factor concentrates are not available. Each bag of cryoprecipitate contains 80 IU of factor VIII.

Blood components	Composition	Indications
Factor VIII concentrates	Factor VIII	Hemophilia A
Factor IX concentrates	Factor IX	Hemophilia B
Cryoprecipitate	Fibrinogen Factor XIII Factor VIII von Willebrand factor	Fibrinogen deficiency, Factor XIII deficiency, von Willebrand disease
Fresh Frozen Plasma (FFP)	Coagulation factors Protein C Protein S Antithrombin	Deficiency of prothrombin, Factor V, Factor VII, Factor X, Factor XI
Prothrombin complex concentrates (PCCs)	Prothrombin Factor VII Factor IX Factor X	Deficiency of prothrombin, Factor VII, Factor X

Solution to Question 18:

Delayed separation of the umbilical stump beyond 4 weeks is a characteristic finding seen in factor XIII deficiency. However, these findings can also be seen in septicemia or local infections.

Factor XIII is responsible for the crosslinking of fibrin to stabilize the fibrin clot, symptoms of delayed hemorrhage are secondary to the instability of the clot. Typically, patients have trauma one day and then have a bruise or hematoma the next day.

Results of the usual screening tests for hemostasis are normal in patients with factor XIII deficiency. Screening tests for factor XIII deficiency are based on the observation that there is increased solubility of the clot because of the failure of cross-linking.

Pediatric Anemias

Question 1:

As a part of the Anemia Mukht Bharat program, you screen preschool children for anemia. What Hb values would prompt you to diagnose anemia in this age group?

- a) < 13 g/dL
- b) < 12 g/dL
- c) < 11.5 g/dL
- d) < 11 g/dL

Question 2:

A 7-year-old child is brought to the pediatric OPD by his parents with complaints of excessive sleepiness, and poor scholastic performance. On examination, the child has pallor involving the tongue, conjunctiva, and nail beds. A flow murmur is heard upon auscultation. Which of the following is the most likely cause of the underlying disease?

- a) Congenital hypoplastic anemia
- b) Acquired hypoproliferative anemia
- c) Congenital hemoglobinopathies
- d) Hyperproliferative anemia

Question 3:

Which of the following statements is true regarding iron?

- a) Breast milk has higher iron content than cow's milk.
- b) Cow's milk has higher oral bio-availability of iron than breast milk.
- c) Introduction of cow's milk at 5 months of age decreases the risk of iron deficiency.
- d) Cow's milk protein can cause chronic intestinal blood loss in infants.

Question 4:

An 8-year-old child who has been recently diagnosed with iron deficiency anemia is started on iron therapy. What is the earliest hematological response that you would expect to see following treatment?

- a) Reticulocytosis
- b) Increase in hemoglobin level
- c) Increase in ferritin levels
- d) Bone marrow erythroid hyperplasia

Question 5:

A child who was diagnosed 3 months back with iron deficiency anemia presents with low hemoglobin despite being on oral iron therapy. What is the most likely cause for the non-responsiveness that you should consider in your differential?

- a) Poor compliance
- b) Incorrect dose or formulation
- c) Occult blood loss from parasitic infestations
- d) Thalassemia misdiagnosed as iron-deficiency anemia

Question 6:

Which of the following tests can be used to screen a newborn with a family history of sickle cell disease?

- a) Hemoglobin electrophoresis
- b) High-performance liquid chromatography
- c) Heel-prick Guthrie test
- d) Tandem mass spectrometry

Question 7:

Ravi, an adolescent with sickle cell disease presents to the emergency room with chest pain and respiratory distress. Pulse oximeter shows 85% saturation in room air. Chest X-ray shows new radiodensities. Sputum and blood culture showed no growth. Which of the following are used in the management of this condition?

- a) 1, 2, and 3 only
- b) 1, 3, and 5 only

- c) 2, 4, and 5 only
- d) 1, 2, 3, 4, 5

Question 8:

A 6-year-old boy who is a known case of sickle cell disease presents with acute chest syndrome. Which of the following is the drug of choice to reduce such painful crises in the future?

- a) Hydrocodone
- b) Hydroxyurea
- c) Corticosteroids
- d) Ruxolitinib

Question 9:

Kevin and Anita both have sickle cell traits. They are devastated when they find out both their children have sickle cell anemia. Kirti is an active 9-year-old. Kamal is her 3-month-old brother. Which of the following prophylactic measures would you not recommend to them?

- a) Oral penicillin prophylaxis up to 5 years of age in Kamal
- b) Annual transcranial Doppler ultrasound for Kirti upto age 16
- c) Annual ophthalmology screening from the age of 10 years in both
- d) Hydroxyurea prophylaxis regardless of disease severity or symptoms in both

Question 10:

Which of the following is responsible for the most common cause of congenital hemolytic anemia due to an enzyme deficiency?

- a) Deficiency of an enzyme in glycolysis pathway
- b) Deficiency of an enzyme in Hexose monophosphate pathway
- c) Deficiency of an enzyme in Kreb's cycle
- d) Deficiency of an enzymes in fructose metabolism pathway

Question 11:

Which of the following is not a cause of non-immune hydrops fetalis?

- a) Alpha thalassemia
- b) Parvovirus B19 infection
- c) Rh incompatibility
- d) Twin-twin transfusion syndrome

Question 12:

Which of the following statements about abnormal hemoglobins and their mechanisms is true?

- a) HbS is due to substitution of lysine for glutamine in the 6th position of the beta-globin gene.
- b) HbH is the deletion of three alpha-globin genes.
- c) Hb Bart is the deletion of all four beta-globin genes.
- d) HbC is due to substitution of valine for lysine in the 6th position of the beta-globin gene.

Question 13:

A 5-year-old child presented with pallor, massive hepatosplenomegaly, and hemolytic facies. Which of the following is the most likely iron profile in this patient?

- a) Iron - low, Ferritin - low, TIBC-high, TSAT-low
- b) Iron - high, Ferritin - normal, TIBC-normal, TSAT-normal
- c) Iron - low, Ferritin - high, TIBC-low, TSAT-low
- d) Iron - low, Ferritin - low, TIBC-low, TSAT-low


Question 14:

A toddler presented with progressive pallor and abdominal distention. On investigations, he had low hemoglobin. Which of the following red cell indices can be used to distinguish thalassemia from iron deficiency anemia?

- a) Mean corpuscular volume
- b) Mean corpuscular hemoglobin
- c) Mean corpuscular volume/ Red blood cell count
- d) Mean corpuscular hemoglobin concentration

Question 15:

A 14-month-old child with Hb 6.3 was undergoing repeated blood transfusions. The last transfusion was given in the 9th month. Which of the following investigation is used for a definitive diagnosis?



- a) High performance liquid chromatography
- b) Ferritin levels estimation
- c) Bone marrow aspiration
- d) Globin gene sequencing

Question 16:

You see a child in the hematology unit who has developed anemia which is progressive along with cardiac decompensation requiring transfusion. He is in the latter half of infancy. You suspect transfusion-dependent thalassemia. Which of the following electrophoresis pattern is seen here?

- a) HbA=95%, HbA2=3.5%, HbF =1.5%
- b) HbA=90%, HbA2=5%, HbF=5%
- c) HbA=45%, HbA2=5%, HbF=50%
- d) HbA=0%, HbA2=5%, HbF=95%

Question 17:

You are given the following x-ray in the viva session. Which of the following is the likely reason behind this abnormal appearance?



- a) Effective erythropoiesis
- b) Thinning of trabeculae
- c) Narrowing of the diploic space
- d) Extramedullary hematopoiesis

Question 18:

You are evaluating a 4-year-old child with β -thalassemia major. He has been on chronic transfusion therapy since the age of 1 year. He has developed transfusion-induced iron overload. The child has an increased risk of developing which of the following infections?

- a) *Streptococcus pneumoniae*
- b) *Mycoplasma pneumoniae*
- c) *Yersinia enterocolitica*
- d) *Chlamydia*

Question 19:

Which of the following is true about Diamond Blackfan Anemia?

- a) Short stature, craniofacial abnormalities, and triphalangeal thumb are common features.
- b) Peripheral smear shows microcytic hypochromic anemia.

- c) Bone marrow failure includes erythroid, myeloid and megakaryocytic lineages.
- d) Reticulocyte count is increased.


Question 20:

You see a neonate undergoing phototherapy for hyperbilirubinemia. Blood workup revealed the presence of a hemolytic picture along with spherocytes and polychromatophilic reticulocytes. MCV is low normal along with elevated MCHC. There was a similar history of hemolytic anemia in the mother. The pedigree analysis points towards an autosomal dominant mode of inheritance. The most likely genetic mutation seen in this case is?

- a) Spectrin
- b) Ankyrin
- c) Band 3
- d) Band 4.2

Question 21:

An 11-year-old child, born out of consanguineous marriage developed signs and symptoms of hemolytic anemia and progressive ataxia. He is a known case of retinitis pigmentosa and also had symptoms of fat malabsorption. A peripheral smear of the child is shown. What is the likely diagnosis?



- a) Hemolytic uremic syndrome
- b) Glucose 6-phosphate dehydrogenase deficiency

- c) Hereditary stomatocytosis
- d) Abetalipoproteinemia

Question 22:

A 12-year-old child is receiving iron and folic acid supplements under the Weekly Iron and Folic Acid Supplementation (WIFS) Programme. Which of the following is true about this program?

- a) It was launched by the Ministry of Health and Family Welfare.
- b) It is meant exclusively for adolescent school girls between 10-19 years of age.
- c) It exclusively covers government schools in rural India.
- d) It recommends that administration of 50 mg of elemental iron.

Question 23:

A toddler is brought by his mother with complaints of poor feeding, lethargy, and chronic diarrhea for the past 4 months. Upon clinical examination, the child has pale conjunctiva and is tachycardic. Blood workup reveals low hemoglobin and smear shows macrocytosis, anisopoikilocytosis, and large neutrophils with hyper-segmented nuclei. Serum homocysteine levels are elevated and methylmalonate levels are normal. Bone marrow shows erythroid hyperplasia. Which of the following is true about this condition?

- a) Lactate dehydrogenase levels in serum is low.
- b) Folate-deficient mother's give birth to infants with folate deficiency.
- c) Methylenetetrahydrofolate (MTHFR) deficiency is associated with megaloblastic anemia.
- d) Goat's milk fed infants are more likely to develop this disease

Question 24:

A child presents with megaloblastic anemia that is unresponsive to vitamin C, vitamin B12, and folic acid, but responsive to uridine. What do you suspect?

- a) Thiamine deficiency
- b) Liver disease
- c) Pernicious anemia
- d) Orotic aciduria

Answer Key

Question No.	Correct Option
1	d
2	b
3	d
4	d
5	a
6	b
7	d
8	b
9	d
10	b
11	c
12	b
13	b
14	c
15	d
16	d
17	d
18	c
19	a
20	b
21	d
22	a
23	d
24	d

Detailed Explanations

Solution to Question 1:

Anaemia in a preschool child is defined as a hemoglobin level of <11 g/dL.

Preschool age is defined as 2 to 5 years.

WHO cut-offs for the diagnosis of anaemia:

Age	Hemoglobin (g/dL)
6 months to 5years	<11
5 to 11 years	<11.5
12 to14 years	<12
Non-pregnant women (≥ 15 years)	<12
Pregnant women	<11
Men (≥ 15 years)	<13

Solution to Question 2:

The given clinical signs and symptoms are suggestive of anemia. Acquired hypoproliferative anemia, is the most common cause of anemia in children among the given options.

Iron-deficiency anemia (which is classified as an acquired hypoproliferative anemia) is the most common cause of anemia in children and adults worldwide. Iron deficiency is also the most common nutrient deficiency worldwide.

Solution to Question 3:

Exposure to whole cow's milk protein can cause chronic intestinal blood loss in infants. Therefore, the American Academy of Pediatrics (AAP) recommends delaying the introduction of cow's milk in infancy and limiting it to ≤ 700 mL/day.

Both cow's milk and breast milk have the same iron content (≈ 0.1 mg/dL).

The bioavailability of breast milk iron is 2-3 times higher than cow's milk iron.

Exclusive breastfeeding until 6 months of age and complementary feeding with iron-rich foods thereafter reduces the risk of iron-deficiency anemia in infants.

Solution to Question 4:

Bone marrow erythroid hyperplasia is the earliest hematological response observed following iron therapy in iron-deficiency anemia.

Responses to iron therapy in iron deficiency anemia:

Time after administration	Response
12 to 24 h	Subjective improvements - decreased irritability, increased appetite
36 to 48 h	Bone marrow response- erythroid hyperplasia
48 to 72 h (peaks at 5-7 days)	Reticulocytosis
4 to 30 days	Increase in hemoglobin level
1 to 3 months	Repletion of stores (ferritin)

Solution to Question 5:

Poor compliance is the most likely cause of non-responsiveness to oral iron therapy.

Differential diagnosis of microcytic anemia that fails to respond to oral iron:

- Poor compliance (most common cause)
- Incorrect dose or formulation
- Malabsorption of administered iron
- Ongoing blood loss, including gastrointestinal, menstrual, and pulmonary
- Concurrent infection or inflammatory disorder
- Concurrent vitamin B12 or folate deficiency
- Diagnosis other than iron deficiency
- Thalassemias
- Hemoglobins C and E disorders
- Anemia of chronic disease
- Lead poisoning
- Sickle thalassemias, hemoglobin SC disease
- Iron refractory iron-deficiency anemia (IRIDA)

Solution to Question 6:

High-performance liquid chromatography (HPLC) is used to screen a newborn with a family history of sickle cell disease.

Option A: Hemoglobin electrophoresis is the confirmatory test for sickle cell anemia

Option C: Heel-prick Guthrie test is the neonatal screening test for phenylketonuria.

Option D: Tandem mass spectrometry is being used for neonatal screening of inherited metabolic disorders.

Solution to Question 7:

The given clinical scenario is suggestive of acute chest syndrome and all of the above are used in the management.

Acute chest syndrome (ACS) refers to a life-threatening pulmonary complication of sickle cell disease. It is diagnosed with the presence of a new radiodensity on chest radiography with any 2 of the following:

- Fever
- Respiratory distress
- Hypoxia
- Cough
- Chest pain

In this case, the child with sickle cell anemia satisfies the above criteria. He presented with chest pain and respiratory distress and his chest X-ray image shows new radiodensities.

Most patients with ACS do not have a single identifiable cause. Infection is the most well-known etiology, yet only 30% of ACS episodes will have positive sputum or bronchoalveolar culture. The most common bacterial pathogens implicated are *S. pneumoniae*, *Mycoplasma pneumoniae*, and *Chlamydia* spp.

Oxygen should be supplemented to all patients who demonstrate hypoxia.

Blood transfusion is the only method to abort a rapidly progressing episode of ACS.

Because of the clinical overlap between pneumonia and ACS, all episodes should be treated promptly with antimicrobial therapy, including at least a macrolide and possibly a third-generation cephalosporin.

Regular use of an incentive spirometer at 10-12 breaths every 2 hours can significantly reduce the frequency of subsequent acute chest pain episodes.

Solution to Question 8:

Hydroxyurea is the drug of choice for reducing painful crises in sickle cell anemia.

Hydroxyurea is a myelosuppressive agent that is effective in reducing the frequency of painful episodes like dactylitis and acute chest syndrome, decreasing the rate of hospitalization, and requirement for blood transfusion. It is safe and well-tolerated in children and infants.

Option A: Hydrocodone (opioid) is used in the treatment of pain crisis in sickle cell anemia.

Option C: Corticosteroids is one of the first-line therapies for acute idiopathic thrombocytic purpura (ITP) with risk factors for bleeding.

Option D: Ruxolitinib is the drug of choice for the treatment of polycythemia vera.

Solution to Question 9:

Hydroxyurea prophylaxis regardless of disease severity is recommended in children over 9 months of age. Hydroxyurea can be considered for use in infants between six and nine months with symptomatic disease.

Oral penicillin VK is recommended until at least 5 years of age. It is continued beyond 5 years of age in children with a history of pneumococcal infection. Erythromycin ethyl succinate is used if the patient is allergic to penicillin. Pneumococcal and meningococcal vaccinations are administered because of the risk of functional asplenia.

Transcranial Doppler ultrasound is recommended, starting at age of 2 years. This has led to a decrease in the prevalence of overt stroke among children with sickle cell anemia. Screening should occur annually from age 2-16.

Annual ophthalmology screening should commence from the age of 10 years to screen for retinopathy.

Solution to Question 10:

Deficiency of an enzyme in the hexose monophosphate pathway ie., glucose 6-phosphate dehydrogenase deficiency (G6PD Deficiency), is the most common cause of congenital hemolytic anemia due to an enzyme deficiency

Most common enzymopathies causing hemolytic anemia (in order of most common to least):

- Glucose 6-phosphate dehydrogenase deficiency (G6PD)
- Pyruvate kinase deficiency
- 5' nucleotidase deficiency

Solution to Question 11:

Rh incompatibility causes immune hydrops fetalis, whereas alpha-thalassemia, parvovirus B19 infection, and twin-twin transfusion syndrome cause non-immune hydrops fetalis.

Hydrops fetalis refers to the abnormal fluid collections in fetal soft tissues and serous cavities. Edema, signs of heart failure, and profound intra-uterine anemia are the presenting features in the fetus. Other clinical features include marked hepatosplenomegaly, retardation in brain growth, hydrocephaly, urogenital and cardiovascular deformities, and extramedullary erythropoiesis.

Non-immune hydrops fetalis (NIHF) comprises the subgroup of cases not caused by red cell alloimmunization (e.g., Rh(D), Kell)

Bart's hydrops fetalis (alpha-thalassemia) is caused by the deletion or inactivation of all four alpha-globin alleles, leading to a severe deficiency in alpha-globin chains of Hb and to the

production of gamma-4 tetramers (Hb Bart). Hb Bart tetramers have an extremely high oxygen affinity and are incapable of effective tissue oxygen delivery.

Parvovirus B19 infection can cause fetal anemia, non-immune fetal hydrops, and fetal death.

Solution to Question 12:

HbH is the deletion of three alpha globin chains.

HbS is due to the substitution of valine for glutamic acid in the 6th position of the beta-globin gene (option A).

Hb Bart is the deletion of all four alpha-globin genes. (There are only 2 beta-globin genes) (option C)

HbC is due to the substitution of lysine for glutamic acid in the 6th position of the beta-globin gene (option D).

Solution to Question 13:

The given clinical scenario is suggestive of thalassemia and the most likely iron profile is iron - normal/high, ferritin - normal, total iron - binding capacity (TIBC) - normal, and transferrin saturation (TSAT) - normal.

Iron- low, ferritin-low, TIBC-high, TSAT-low is seen in iron-deficiency anemia.

Iron-low, ferritin-high, TIBC-low, TSAT-low is seen in anemia of chronic disease.

Solution to Question 14:

Mean corpuscular volume/ Red blood cell count (MCV/RBC) also known as Mentzer index can be used to distinguish thalassemia from iron-deficiency anemia (IDA).

MCV (mean corpuscular volume) and MCH (mean corpuscular hemoglobin) are low in both IDA and thalassemia. RBC count may be normal in thalassemia, whereas it is reduced in IDA. A lower RBC count in IDA means the denominator is smaller, yielding a greater MCV/RBC ratio.

Mentzer index is:

- More than 13 in iron-deficiency anemia
- Less than 13 in thalassemia

Solution to Question 15:

The given clinical scenario of repeated transfusion, low hemoglobin level, and target cells in the smear are suggestive of thalassemia. Globin gene sequencing is used for its definitive diagnosis.

High-performance liquid chromatography (HPLC) is used as a screening test for thalassemia (option A).

Thalassemia is a hemoglobinopathy that arises due to deficient synthesis of the globin chains. Depending on the chain affected, it is classified into α -thalassemia and β -thalassemia.

β -thalassemia is caused by point mutations of the β -thalassemia gene on chromosome 11, which diminish the synthesis of β -globin chains. Splicing mutations are the most common cause of (β +) thalassemia.

There is a deficit in the synthesis of HbA with an elevation of HbA₂ and HbF levels in β -thalassemia. Children with severe thalassemia present with distinctive facial features such as maxilla hyperplasia, a flat nasal bridge, and frontal bossing. Other manifestations include pathologic bone fractures, significant hepatosplenomegaly, and cachexia.

Blood smears show marked anisocytosis, poikilocytosis, and microcytic hypochromic anemia. Target cells, basophilic stippling, and fragmented red cells also are common. A radiographic image of the skull shows the classic 'crew cut' appearance.

Hb Electrophoresis and high-performance liquid chromatography (HPLC) are used as screening tests for thalassemia. DNA diagnosis to detect β -thalassemia mutations is used for definitive diagnosis. The Mentzer index is the ratio of MCV to RBC count. It is used to differentiate between iron-deficiency anemia and β -thalassemia trait. The Mentzer index is <13 in thalassemia minor syndrome.

Transfusions are administered every 3-4 weeks to maintain a pretransfusion hemoglobin level of 9.5-10.5 g/dL. Transfusion-induced hemosiderosis is a significant complication in transfusion-dependent thalassemia. Iron deposition occurs in the liver, endocrine organs, and heart, resulting in complications including hypothyroidism, hypogonadotropic hypogonadism, growth hormone deficiency, hypoparathyroidism, diabetes mellitus, heart failure, and arrhythmias.

The image below shows thalassemic facies and crew cut appearance on X-ray:





Solution to Question 16:

HbA=0%, HbA2=5%, HbF=95% is seen on hemoglobin electrophoresis in transfusion-dependent thalassemia.

Given clinical picture where the child becomes symptomatic during the second 6 months of life requiring transfusion is suggestive of beta-thalassemia major or transfusion-dependent thalassemia.

- Normal pattern- HbA=95%, HbA2=3.5%, HbF =1.5%
- Beta-thalassemia trait- HbA=90%, HbA2=5%, HbF=5%
- Beta-thalassemia intermedia- HbA=45%, HbA2=5%, HbF=50%

Alpha- and beta-thalassemia are inherited disorders in which there is reduced production of alpha-globin or beta-globin chains, respectively. This leads to precipitation of the excess (unpaired) chains, interfering with red blood cell (RBC) production and survival. Disease severity is inversely correlated with the number of functioning globin chains. Each individual carries 2 beta-chain genes and 4 alpha-chain genes.

Solution to Question 17:

Crew-cut or hair-on-end appearance on X-ray imaging of the skull is due to extramedullary hematopoiesis, which is seen in thalassemia.

Ineffective erythropoiesis leads to extramedullary hematopoiesis, which causes the widening of the diploic space, thereby causing the thickening of trabeculae. This results in the classical crew-cut or hair-on-end appearance on x-ray imaging of the skull. Extramedullary hematopoiesis also leads to hepatosplenomegaly and enlarged and distorted bony prominences over the face, leading to the characteristic chipmunk facies in thalassemia.

Solution to Question 18:

Thalassemic children with transfusion-induced iron overload are at an increased risk of infections with *Yersinia enterocolitica*.

Iron excess immunologically compromises the host and greatly enhances yersinial growth.

Streptococcus pneumoniae (option A), *Mycoplasma pneumoniae* (option B), and *Chlamydia* species (option D) are the common organisms causing infections that predispose to acute chest syndrome in sickle cell anemia.

Solution to Question 19:

Short stature, craniofacial abnormalities, and triphalangeal thumb are common features is the true statement regarding Diamond–Blackfan anemia (DBA).

Features of DBA (congenital hypoplastic anemia):

- Autosomal dominant inheritance.
- Macrocytosis without hypersegmented neutrophils or other megaloblastic features.
- Reticulocytopenia.
- Absent RBC precursors in the otherwise normally cellular bone marrow. Other lineages are unaffected.
- Craniofacial anomaly is the most common congenital anomaly seen.
- Hypertelorism, snub nose, and high-arched palate seen.
- Thumb abnormalities- flattening of the thenar eminence and triphalangeal thumb
- Short stature

Treatment: Corticosteroids, chronic red cell transfusions, and hematopoietic stem cell transplantation.

Image showing triphalangeal thumb:



Solution to Question 20:

The Ankyrin gene mutation is the most likely genetic mutation seen in the given case of autosomal-dominant hereditary spherocytosis (HS).

Hemolytic manifestation causing indirect hyperbilirubinemia requiring phototherapy or blood transfusion in infancy with positive family history and typical laboratory findings is suggestive of HS.


History of blood transfusions in later life and not infancy is used to classify the severity of the disease.

- Most common mode of inheritance of HS- autosomal dominant
- Most common gene mutation in autosomal-dominant HS- ankyrin > Band 3
- Most common gene mutation in autosomal-recessive HS- spectrin
- Most severe form of HS- spectrin (recessive)

Classification of the severity of HS is based on the requirement of blood transfusion in later life and not during infancy.


RBC Membrane-Cytoskeleton disorders

Image: The peripheral smear below shows regularly spaced, smoothly rounded crenulations, which are characteristic of a burr cell.




Peripheral smear in other options:

Hemolytic uremic syndrome (option A); Numerous schistocytes and nucleated RBCs with extremely low platelets:




Glucose 6-phosphate dehydrogenase deficiency (option B); bite cells or degmacytes or blister cells. In G6PD deficiency uncontrolled oxidative stress causes hemoglobin to denature and form Heinz bodies. Bite cells result from the removal of Heinz bodies by macrophages in the spleen.

The image below shows bite cells represented by blue arrows, black arrows represent burr cells:



Hereditary stomatocytosis (option C); Numerous stomatocytes (Uniconcave cup with mouth-shaped area of central pallor), caused by missense mutation in band 3 or AEP



Solution to Question 22:

Weekly Iron Folic acid Supplementation (WIFS) was launched by the Ministry of Health and Family Welfare is the true statement among the given options.

Salient features of Weekly Iron Folic acid Supplementation (WIFS):

- i. Objective of Weekly Iron Folic acid Supplementation (WIFS): To reduce the prevalence and severity of anemia in the adolescent population (10-19 years).
- ii. Target groups

- School-going adolescent girls and boys in 6th to 12th class enrolled in government/government-aided/municipal schools
- Out-of-school adolescent girls

iii. Intervention

- Administration of supervised weekly iron-folic acid supplements of 60 mg elemental iron and 500 µg folic acid by using a fixed-day approach
- Screening of target groups for moderate/severe anemia and referring these cases to an appropriate health facility
- Biannual de-worming (albendazole 400 mg), six months apart, for the control of helminthic infestation
- Information and counseling for improving dietary intake and for taking actions for prevention of intestinal worm infestation

Solution to Question 23:

The given clinical scenario along with the megaloblastic picture elevated homocysteine levels, and normal methylmalonate levels are suggestive of folate deficiency anemia.

Goat's milk is deficient in folate. So goat milk-fed infants are likely to develop folate deficiency anemia. Human breast milk, infant formulas, and pasteurized cow's milk provide adequate amounts of folic acid.

In folate deficiency anemia, lactate dehydrogenase levels in serum are elevated. Folate-deficient mothers do not give birth to infants with folate deficiency because there is a selective transfer of folate to the fetus via the placental folate receptors.

Methylenetetrahydrofolate (MTHFR) deficiency is the most common inborn error of folate metabolism, and severe cases can produce neurologic and vascular complications; there is no associated megaloblastic anemia.

Malnutrition is the most common cause of folate deficiency in children.

Note: Both homocysteine and methylmalonate levels are elevated in B12 deficiency anemia.

Solution to Question 24:

Orotic aciduria is to be suspected in this child. Megaloblastic anemia that is responsive to uridine is consistent with orotic aciduria.

Orotic aciduria is a rare inborn error of metabolism in children. A defect in the pyrimidine synthesis pathway results in the failure of conversion of orotic acid to uridine.

Supplementation of vitamin B12, folate, and vitamin C rules out the more common nutritional causes of megaloblastic anemia.

Thiamine deficiency (option A) and anemia due to liver disease (option B) do not respond to uridine.

Pernicious anemia (Option C) responds to treatment with vitamin B12.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.