

InBody, February vs May

Two scans, 106 days apart, on different InBody machines. Values from the kg sheet are converted so both tests read in one unit.

● TEST 1 • FEB 12, 2026 • 7:57 AM

UNC Campus Recreation

InBody270S • Age 20 • height entered 5'8" (172.7 cm)

● TEST 2 • MAY 29, 2026 • 9:14 AM

Fitness SF

InBody570 • ID 9199143328 • Age 21 • height entered 170.2 cm (5'7")

WEIGHT

126.3 lb

was 121.0 lb

+5.3 lb

SKELETAL MUSCLE

49.2 lb

was 47.6 lb

▲ +1.6 lb

BODY FAT MASS

35.5 lb

was 32.4 lb

+3.1 lb

PERCENT BODY FAT

28.1%

was 26.8%

+1.3 pt

BMI

19.8

was 18.4

+1.4

BASAL METABOLIC RATE

1,260 kcal

was 1,238 kcal

▲ +22 kcal

Muscle-Fat Analysis

Bars share one scale, so lengths compare across metrics as well as between tests.

● Feb 12 • UNC (270S) ● May 29 • Fitness SF (570)

Weight



Skeletal Muscle Mass



Body Fat Mass



Fat-Free / Lean Body Mass



Segmental Lean Analysis

Percent of the ideal lean mass for each body segment — the dashed line marks 100% (right at ideal).

● Feb 12 · UNC (270S) ● May 29 · Fitness SF (570)

Right Arm



Left Arm



Trunk



Right Leg



Left Leg



100% of ideal

Segmental Fat Analysis

May test only — the InBody270S doesn't report segmental fat. Percent of the ideal fat mass per segment; dashed line marks 100%.

Right Arm



Left Arm



Trunk



Right Leg



Left Leg



100% of ideal

Every Shared Metric

Converted values shown in the selected unit; Δ is Test 2 minus Test 1.

METRIC	FEB 12 (270S)	MAY 29 (570)	Δ
Weight	121.0 lb	126.3 lb	+5.3 lb
Skeletal Muscle Mass	47.6 lb	49.2 lb	+1.6 lb
Body Fat Mass	32.4 lb	35.5 lb	+3.1 lb
Percent Body Fat	26.8%	28.1%	+1.3 pt
Fat-Free / Lean Body Mass	88.6 lb	90.8 lb	+2.2 lb
Total Body Water	64.8 lb	66.4 lb	+1.6 lb
Dry Lean Mass	23.8 lb	24.5 lb	+0.7 lb
BMI	18.4	19.8	+1.4
Basal Metabolic Rate	1,238 kcal	1,260 kcal	+22 kcal

Only on the Feb test (InBody270S)

Whole-body phase angle (50 kHz)	4.5°
Skeletal Muscle Index (SMI)	5.8 kg/m²
Fat-free mass vs normal range	88.6 lb (97.9-119.7 lb)
Recommended change	0.0 lb fat · +20.3 lb lean

Only on the May test (InBody570)

Visceral fat level	7 (under 10 = healthy)
ECW / TBW ratio	0.381
Intracellular / extracellular water	18.6 L / 11.5 L
Recommended change	-4.6 lb fat · +12.6 lb lean

IMPEDANCE Z(Ω) BY SEGMENT

FREQ	RA	LA	TRUNK	RL	LL
5 kHz	490.0	500.1	30.4	381.8	389.9
50 kHz	449.4	460.4	27.0	341.9	349.9
500 kHz	396.0	403.5	20.2	301.3	308.9

Read the comparison loosely. The two scans came from different machines (InBody270S vs the higher-resolution InBody570) with slightly different entered heights (5'8" vs 170.2 cm), which shifts BMI and the "% of ideal" segmental targets — the drop in leg percentages is mostly that recalibration, since the leg mass itself barely moved. Hydration, time of day, and food before the test also nudge every number. For tracking, repeat tests on the same machine under similar conditions.

InBody, February vs May

Two scans, 106 days apart, on different InBody machines. Values from the kg sheet are converted so both tests read in one unit.

● TEST 1 • FEB 12, 2026 • 7:57 AM

UNC Campus Recreation

InBody270S • Age 20 • height entered 5'8" (172.7 cm)

● TEST 2 • MAY 29, 2026 • 9:14 AM

Fitness SF

InBody570 • ID 9199143328 • Age 21 • height entered 170.2 cm (5'7")

WEIGHT

57.3 kg

was 54.9 kg

+2.4 kg

SKELETAL MUSCLE

22.3 kg

was 21.6 kg

▲ +0.7 kg

BODY FAT MASS

16.1 kg

was 14.7 kg

+1.4 kg

PERCENT BODY FAT

28.1%

was 26.8%

+1.3 pt

BMI

19.8

was 18.4

+1.4

BASAL METABOLIC RATE

1,260 kcal

was 1,238 kcal

▲ +22 kcal

Muscle-Fat Analysis

Bars share one scale, so lengths compare across metrics as well as between tests.

● Feb 12 • UNC (270S) ● May 29 • Fitness SF (570)

Weight



Skeletal Muscle Mass



Body Fat Mass



Fat-Free / Lean Body Mass



Segmental Lean Analysis

Percent of the ideal lean mass for each body segment — the dashed line marks 100% (right at ideal).

● Feb 12 · UNC (270S) ● May 29 · Fitness SF (570)

Right Arm



Left Arm



Trunk



Right Leg



Left Leg



100% of ideal

Segmental Fat Analysis

May test only — the InBody270S doesn't report segmental fat. Percent of the ideal fat mass per segment; dashed line marks 100%.

Right Arm



Left Arm



Trunk



Right Leg



Left Leg



100% of ideal

Every Shared Metric

Converted values shown in the selected unit; Δ is Test 2 minus Test 1.

METRIC	FEB 12 (270S)	MAY 29 (570)	Δ
Weight	54.9 kg	57.3 kg	+2.4 kg
Skeletal Muscle Mass	21.6 kg	22.3 kg	+0.7 kg
Body Fat Mass	14.7 kg	16.1 kg	+1.4 kg
Percent Body Fat	26.8%	28.1%	+1.3 pt
Fat-Free / Lean Body Mass	40.2 kg	41.2 kg	+1.0 kg
Total Body Water	29.4 kg	30.1 kg	+0.7 kg
Dry Lean Mass	10.8 kg	11.1 kg	+0.3 kg
BMI	18.4	19.8	+1.4
Basal Metabolic Rate	1,238 kcal	1,260 kcal	+22 kcal

Only on the Feb test (InBody270S)

Whole-body phase angle (50 kHz)	4.5°
Skeletal Muscle Index (SMI)	5.8 kg/m²
Fat-free mass vs normal range	40.2 kg (44.4-54.3 kg)
Recommended change	0.0 kg fat · +9.2 kg lean

Only on the May test (InBody570)

Visceral fat level	7 (under 10 = healthy)
ECW / TBW ratio	0.381
Intracellular / extracellular water	18.6 L / 11.5 L
Recommended change	-2.1 kg fat · +5.7 kg lean

IMPEDANCE Z(Ω) BY SEGMENT

FREQ	RA	LA	TRUNK	RL	LL
5 kHz	490.0	500.1	30.4	381.8	389.9
50 kHz	449.4	460.4	27.0	341.9	349.9
500 kHz	396.0	403.5	20.2	301.3	308.9

Read the comparison loosely. The two scans came from different machines (InBody270S vs the higher-resolution InBody570) with slightly different entered heights (5'8" vs 170.2 cm), which shifts BMI and the "% of ideal" segmental targets — the drop in leg percentages is mostly that recalibration, since the leg mass itself barely moved. Hydration, time of day, and food before the test also nudge every number. For tracking, repeat tests on the same machine under similar conditions.