

WIKIPEDIA

Linear-fractional programming

In mathematical optimization, **linear-fractional programming (LFP)** is a generalization of linear programming (LP). Whereas the objective function in a linear program is a linear function, the objective function in a linear-fractional program is a ratio of two linear functions. A linear program can be regarded as a special case of a linear-fractional program in which the denominator is the constant function one.

Contents

[**Relation to linear programming**](#)[**Definition**](#)[**Transformation to a linear program**](#)[**Duality**](#)[**Properties and algorithms**](#)[**Notes**](#)[**References**](#)[**Further reading**](#)[**Software**](#)

Relation to linear programming

Both linear programming and linear-fractional programming represent optimization problems using linear equations and linear inequalities, which for each problem-instance define a feasible set. Fractional linear programs have a richer set of objective functions. Informally, linear programming computes a policy delivering the best outcome, such as maximum profit or lowest cost. In contrast, a linear-fractional programming is used to achieve the highest *ratio* of outcome to cost, the ratio representing the highest efficiency. For example, in the context of LP we maximize the objective function **profit** = **income** − **cost** and might obtain maximum profit of \$100 (= \$1100 of income − \$1000 of cost). Thus, in LP we have an efficiency of \$100/\$1000 = 0.1. Using LFP we might obtain an efficiency of \$10/\$50 = 0.2 with a profit of only \$10, but only requiring \$50 of investment.

Definition

Formally, a linear-fractional program is defined as the problem of maximizing (or minimizing) a ratio of affine functions over a polyhedron,

$$\begin{array}{ll}\text{maximize} & \frac{\mathbf{c}^T \mathbf{x} + \alpha}{\mathbf{d}^T \mathbf{x} + \beta} \\ \text{subject to} & A\mathbf{x} \leq \mathbf{b},\end{array}$$

where $\mathbf{x} \in \mathbb{R}^n$ represents the vector of variables to be determined, $\mathbf{c}, \mathbf{d} \in \mathbb{R}^n$ and $\mathbf{b} \in \mathbb{R}^m$ are vectors of (known) coefficients, $A \in \mathbb{R}^{m \times n}$ is a (known) matrix of coefficients and $\alpha, \beta \in \mathbb{R}$ are constants. The constraints have to restrict the feasible region to $\{\mathbf{x} | \mathbf{d}^T \mathbf{x} + \beta > 0\}$, i.e. the region on which the denominator is positive.^{[1][2]} Alternatively, the denominator of the objective function has to be strictly negative in the entire feasible region.

Transformation to a linear program

Under the assumption that the feasible region is **non-empty and bounded**, the **Charnes-Cooper transformation**^[1]

$$\mathbf{y} = \frac{1}{\mathbf{d}^T \mathbf{x} + \beta} \cdot \mathbf{x}; \quad t = \frac{1}{\mathbf{d}^T \mathbf{x} + \beta}$$

translates the linear-fractional program above to the equivalent linear program:

$$\begin{array}{ll}\text{maximize} & \mathbf{c}^T \mathbf{y} + \alpha t \\ \text{subject to} & A\mathbf{y} \leq \mathbf{b}t \\ & \mathbf{d}^T \mathbf{y} + \beta t = 1 \\ & t \geq 0.\end{array}$$

Then the solution for \mathbf{y} and t yields the solution of the original problem as

$$\mathbf{x} = \frac{1}{t} \mathbf{y}.$$

Duality

Let the dual variables associated with the constraints $A\mathbf{y} - \mathbf{b}t \leq \mathbf{0}$ and $\mathbf{d}^T \mathbf{y} + \beta t - 1 = 0$ be denoted by \mathbf{u} and λ , respectively. Then the dual of the LFP above is ^{[3][4]}

$$\begin{array}{ll}\text{minimize} & \lambda \\ \text{subject to} & A^T \mathbf{u} + \lambda \mathbf{d} = \mathbf{c} \\ & -\mathbf{b}^T \mathbf{u} + \lambda \beta \geq \alpha \\ & \mathbf{u} \in \mathbb{R}_+^m, \lambda \in \mathbb{R},\end{array}$$

which is an LP and which coincides with the dual of the equivalent linear program resulting from the Charnes-Cooper transformation.

Properties and algorithms

The objective function in a linear-fractional problem is both **quasiconcave** and **quasiconvex** (hence quasilinear) with a **monotone** property, **pseudoconvexity**, which is a stronger property than quasiconvexity. A linear-fractional objective function is both pseudoconvex and pseudoconcave, hence **pseudolinear**. Since an LFP can be transformed to an LP, it can be solved using any LP solution method, such as the **simplex algorithm** (of **George B. Dantzig**),^{[5][6][7][8]} the **criss-cross algorithm**,^[9] or **interior-point methods**.

Notes

1. Charnes, A.; Cooper, W. W. (1962). "Programming with Linear Fractional Functionals". *Naval Research Logistics Quarterly*. **9** (3–4): 181–186. doi:10.1002/nav.3800090303 (https://doi.org/10.1002%2Fnav.3800090303). MR 0152370 (https://www.ams.org/mathscinet-getitem?mr=0152370).
2. Boyd, Stephen P.; Vandenberghe, Lieven (2004). *Convex Optimization* (https://web.stanford.edu/~boyd/cvxbook/bv_cvxbook.pdf) (PDF). Cambridge University Press. p. 151. ISBN 978-0-521-83378-3. Retrieved October 15, 2011.
3. Schaible, Siegfried (1974). "Parameter-free Convex Equivalent and Dual Programs". *Zeitschrift für Operations Research*. **18** (5): 187–196. doi:10.1007/BF02026600 (https://doi.org/10.1007%2FBF02026600). MR 0351464 (https://www.ams.org/mathscinet-getitem?mr=0351464). S2CID 28885670 (https://api.semanticscholar.org/CorpusID:28885670).
4. Schaible, Siegfried (1976). "Fractional programming I: Duality". *Management Science*. **22** (8): 858–867. doi:10.1287/mnsc.22.8.858 (https://doi.org/10.1287%2Fmnsc.22.8.858). JSTOR 2630017 (https://www.jstor.org/stable/2630017). MR 0421679 (https://www.ams.org/mathscinet-getitem?mr=0421679).
5. Chapter five: Craven, B. D. (1988). *Fractional programming*. Sigma Series in Applied Mathematics. Vol. 4. Berlin: Heldermann Verlag. p. 145. ISBN 978-3-88538-404-5. MR 0949209 (https://www.ams.org/mathscinet-getitem?mr=0949209).
6. Kruk, Serge; Wolkowicz, Henry (1999). "Pseudolinear programming". *SIAM Review*. **41** (4): 795–805. CiteSeerX 10.1.1.53.7355 (https://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.53.7355). doi:10.1137/S0036144598335259 (https://doi.org/10.1137%2FS0036144598335259). JSTOR 2653207 (https://www.jstor.org/stable/2653207). MR 1723002 (https://www.ams.org/mathscinet-getitem?mr=1723002).
7. Mathis, Frank H.; Mathis, Lenora Jane (1995). "A nonlinear programming algorithm for hospital management". *SIAM Review*. **37** (2): 230–234. doi:10.1137/1037046 (https://doi.org/10.1137%2F1037046). JSTOR 2132826 (https://www.jstor.org/stable/2132826). MR 1343214 (https://www.ams.org/mathscinet-getitem?mr=1343214).
8. Murty (1983, Chapter 3.20 (pp. 160–164) and pp. 168 and 179)
9. Illés, Tibor; Szirmai, Ákos; Terlaky, Tamás (1999). "The finite criss-cross method for hyperbolic programming". *European Journal of Operational Research*. **114** (1): 198–214. CiteSeerX 10.1.1.36.7090 (https://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.36.7090). doi:10.1016/S0377-2217(98)00049-6 (https://doi.org/10.1016%2FS0377-2217%2898%2900049-6). Zbl 0953.90055 (https://zbmath.org/?format=complete&q=an:0953.90055). Postscript preprint (http://www.cas.mcmaster.ca/~terlaky/files/dut-twi-96-103.ps.gz).

References

- Craven, B. D. (1988). *Fractional programming*. Sigma Series in Applied Mathematics. Vol. 4. Berlin: Heldermann Verlag. p. 145. ISBN 978-3-88538-404-5. MR 0949209 (https://www.ams.org/mathscinet-getitem?mr=0949209).

- Illés, Tibor; Szirmai, Ákos; Terlaky, Tamás (1999). "The finite criss-cross method for hyperbolic programming". *European Journal of Operational Research*. **114** (1): 198–214. CiteSeerX 10.1.1.36.7090 (<https://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.36.7090>). doi:10.1016/S0377-2217(98)00049-6 (<https://doi.org/10.1016%2FS0377-2217%2898%2900049-6>). Zbl 0953.90055 (<https://zbmath.org/?format=complete&q=an:0953.90055>). Postscript preprint (<http://www.cas.mcmaster.ca/~terlaky/files/dut-twi-96-103.ps.gz>).
- Kruk, Serge; Wolkowicz, Henry (1999). "Pseudolinear programming" (<https://doi.org/10.1137%2FS0036144598335259>). *SIAM Review*. **41** (4): 795–805. doi:10.1137/S0036144598335259 (<https://doi.org/10.1137%2FS0036144598335259>). JSTOR 2653207 (<https://www.jstor.org/stable/2653207>). MR 1723002 (<https://www.ams.org/mathscinet-getitem?mr=1723002>).
- Mathis, Frank H.; Mathis, Lenora Jane (1995). "A nonlinear programming algorithm for hospital management". *SIAM Review*. **37** (2): 230–234. doi:10.1137/1037046 (<https://doi.org/10.1137%2F1037046>). JSTOR 2132826 (<https://www.jstor.org/stable/2132826>). MR 1343214 (<https://www.ams.org/mathscinet-getitem?mr=1343214>).
- Murty, Katta G. (1983). "3.10 Fractional programming (pp. 160–164)". *Linear programming*. New York: John Wiley & Sons, Inc. pp. xix+482. ISBN 978-0-471-09725-9. MR 0720547 (<https://www.ams.org/mathscinet-getitem?mr=0720547>).

Further reading

- Bajalinov, E. B. (2003). *Linear-Fractional Programming: Theory, Methods, Applications and Software*. Boston: Kluwer Academic Publishers.
- Barros, Ana Isabel (1998). *Discrete and fractional programming techniques for location models*. Combinatorial Optimization. Vol. 3. Dordrecht: Kluwer Academic Publishers. pp. xviii+178. ISBN 978-0-7923-5002-6. MR 1626973 (<https://www.ams.org/mathscinet-getitem?mr=1626973>).
- Martos, Béla (1975). *Nonlinear programming: Theory and methods*. Amsterdam-Oxford: North-Holland Publishing Co. p. 279. ISBN 978-0-7204-2817-9. MR 0496692 (<https://www.ams.org/mathscinet-getitem?mr=0496692>).
- Schaible, S. (1995). "Fractional programming". In Reiner Horst and Panos M. Pardalos (ed.). *Handbook of global optimization*. Nonconvex optimization and its applications. Vol. 2. Dordrecht: Kluwer Academic Publishers. pp. 495–608. ISBN 978-0-7923-3120-9. MR 1377091 (<https://www.ams.org/mathscinet-getitem?mr=1377091>).
- Stancu-Minasian, I. M. (1997). *Fractional programming: Theory, methods and applications*. Mathematics and its applications. Vol. 409. Translated by Victor Giurgutiu from the 1992 Romanian. Dordrecht: Kluwer Academic Publishers Group. pp. viii+418. ISBN 978-0-7923-4580-0. MR 1472981 (<https://www.ams.org/mathscinet-getitem?mr=1472981>).

Software

- WinGULF (<http://zeus.nyf.hu/~bajalinov/WinGulf/wingulf.html>) – educational interactive linear and linear-fractional programming solver with a lot of special options (pivoting, pricing, branching variables, etc.)
- JOptimizer – Java convex optimization library (open source)

Retrieved from "https://en.wikipedia.org/w/index.php?title=Linear-fractional_programming&oldid=1076327567"

This page was last edited on 10 March 2022, at 15:03 (UTC).

Text is available under the Creative Commons Attribution-ShareAlike License 3.0; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia

Foundation, Inc., a non-profit organization.