

ANNALS OF THE NEW YORK ACADEMY OF SCIENCES

Issue: *Women's and Adolescent Nutrition*
REVIEW

Systematic review on evidence-based adolescent nutrition interventions

Zohra S. Lassi,¹ Anoosh Moin,² Jai K. Das,² Rehana A. Salam,² and Zulfiqar A. Bhutta^{2,3}

¹Robinson Research Institute, University of Adelaide, Adelaide, Australia. ²Centre of Excellence in Women and Child Health, Aga Khan University, Karachi, Pakistan. ³Robert Harding Chair in Global Child Health & Policy, Centre for Global Child Health, the Hospital for Sick Children, Toronto, Canada; and Founding Director, Center of Excellence in Women and Child Health, the Aga Khan University, Karachi, Pakistan

Address for correspondence: Zulfiqar A. Bhutta, Founding Director, Center of Excellence in Women and Child Health, Aga Khan University, Karachi 74800, Pakistan; and Robert Harding Chair in Global Child Health and Policy, Centre for Global Child Health, the Hospital for Sick Children, Toronto, ON, Canada. zulfiqar.bhutta@sickkids.ca

Adolescence is a critical stage in the life cycle, and adequate nutrition is necessary for the proper growth and development of individuals and their offspring. Here, we comprehensively review all published systematic reviews (through October 2016) on adolescents (10–19 years) and women of reproductive age, including pregnant women, which targeted interventions related to nutrition. For interventions where there was no existing systematic review on adolescents, we reviewed primary studies/trials. We included interventions on micronutrient supplementation (iron, folic acid, iron–folic acid (IFA), calcium, vitamin D, vitamin A, zinc, iodine, and multiple micronutrients), food/protein energy supplementation, nutrition education for pregnant adolescents, obesity prevention and management, and management of gestational diabetes. We identified a total of 35 systematic reviews, of which only five were conducted on adolescents, and 107 primary studies on adolescents. Our review suggests that iron alone, IFA, zinc, and multiple micronutrient supplementation in adolescents can significantly improve serum hemoglobin concentration. While zinc supplementation in pregnant adolescents showed improvements in preterm birth and low birth weight, we found a paucity of trials on calcium, vitamin D, vitamin A, and iodine supplementation. We found limited evidence on food/protein energy supplementation in adolescents. Interventions to prevent and manage obesity showed a nonsignificant impact on reducing body mass index. This review underscores the importance of adolescent nutrition interventions. It is imperative that countries design nutritional interventions, particularly for adolescents.

Keywords: adolescent; nutrition; obesity; micronutrient; malnutrition; pregnancy

Introduction

Globally, there are nearly 1.8 billion people 10–24 years of age, making up a quarter of the total population (7.3 billion) of the world, and approximately 89% of these live in low- and middle-income countries (LMICs).¹ The United Nations reports that, with such large numbers of young people, it is imperative that they should be given economic and social power, as well as the right to health and empowerment to handle their own lives and future. Unfortunately, most of the adolescent population grows up in countries with multiple adolescent health problems, including diseases of poverty,

undernutrition, poor sexual and reproductive health, injury and violence, and noncommunicable diseases.² There has been limited focus on adolescents in global initiatives, but more recently there has been an increasing focus on adolescent health, with the launch of a *Lancet* commission on adolescent health and well-being involving a network of academics, policy makers, practitioners, and young health advocates. This Commission outlines the opportunities and challenges for investment in adolescents at both country and global levels.²

Adolescent development is as intricate as children's, as there is a complex interaction between

doi: 10.1111/nyas.13335




Figure 1. Framework for nutrition in adolescents and young women.

puberty, neurocognitive maturity, and social-role transitions.³ In addition to the role of living conditions and environment, especially at home, a key contributor to normal development is nutrition. Consumption of a healthy diet is necessary for proper growth and development during puberty, which necessitates adequate macro- and micronutrition. Recently, there has been an epidemiological shift from undernutrition and growth stunting to increasing rates of obesity, and many LMICs now bear a double burden of nutritional disorders, owing to the emerging issue of overweight and obesity along with the existing high rates of stunting and other micronutrient deficiencies. There has been a growing interest in adolescent girls' nutrition as a means to improve the health of women and children, as maternal preconception nutritional deficiencies have profound consequences for fetal and infant development, with the effects extending to neonatal and early childhood development and mortality.

Adolescent nutrition is relevant for current, future, and intergenerational health, and the causes of adolescent nutritional deficiencies are complex, with contributors at the individual, household, and population levels. This underscores the need for comprehensive research on adolescent nutritional interventions. While an earlier review on adolescent nutrition intervention only included systematic

reviews, it was also only limited to micronutrient supplementation, nutrition interventions for pregnant adolescents, and interventions to prevent obesity.⁴ No efforts were undertaken to uncover data from women of reproductive age in cases where data were not available on adolescents.⁴ Here, we review the current evidence for nutritional and related interventions from studies focused on adolescents only or as part of different population groups. These interventions include micronutrient supplementation, food/protein energy supplementation, and obesity prevention and management, and Figure 1 depicts the framework for pathways for improvement in nutrition in adolescents and young women. Our primary objective here is to analyze the existing literature exploring these interventions and their impact on the health of adolescents (aged 10–19 years), including pregnant teenagers in both high-income countries (HICs) and LMICs.

Methods

We conducted a search to identify existing systematic reviews on adolescents (10–19 years) that targeted the identified nutrition interventions. For interventions where there were no systematic reviews specific to adolescents, we searched for reviews on women of reproductive age including pregnant women. In addition, we also conducted

a new review of trials for interventions where there were no adolescent-specific reviews. The following electronic reference libraries were searched to access the available data: the Cochrane Library, PubMed, CINAHL, and Google Scholar (until October 2016). Studies that included adolescents as a subpopulation along with other population groups were deconstructed to isolate the evidence on the adolescent age group. Detailed examination of cross-references and bibliographies of available data and publications was also performed to identify additional sources of information. The following interventions are reviewed here: (1) micronutrient supplementation (iron, folic acid, iron–folic acid (IFA), calcium, vitamin D, vitamin A, zinc, iodine, and multiple micronutrient (MMN) supplementation); (2) food/protein energy supplementation; (3) nutrition education for pregnant adolescents; (4) obesity prevention and management; and (5) gestational diabetes prevention and management.

For existing systematic reviews

We considered all available published Cochrane and non-Cochrane reviews of randomized or non-randomized controlled trials that fully or partly addressed nutrition-specific interventions for improving the health of adolescents. We included reviews that specifically addressed adolescents or women of reproductive age, including pregnant women. The included systematic reviews were assessed using the AMSTAR (assessment of the methodological quality of systematic reviews) criteria.⁵ We resolved any disagreement by discussion, and the final decision was made by consensus within the team.

We recorded the extracted pooled effect size with 95% confidence intervals (CIs). We contacted the review authors for updated data in case each review had not recently been updated. We also contacted trial authors of studies awaiting classification for updated information. We did not update the individual review, but summary of findings tables were formulated and included for most of the reviews (where applicable). We were very cautious when comparing absolute effects across reviews, as there may be different control group risks.

For primary studies/trials

For the interventions where there was no existing systematic review on adolescent populations, we searched for trials. Pre–post study designs and

observational studies were only included when no higher-quality trials could be found for a specific intervention. The team set up a triage process with standardized criteria for evaluating outputs from the search strategy and primary screening. Following an agreement on the search strategy, the abstracts (and the full sources when abstracts were not available) were screened by two abstractors to identify studies adhering to the inclusion criteria. Any disagreements on selection of studies between these two primary abstractors were resolved by the third reviewer. After full-text retrieval of all the studies that met the inclusion/exclusion criteria, each study was double data abstracted into a standardized form. We extracted the information on characteristics of included studies, including descriptions of methods, participants, interventions, outcomes, extraction of measurement of treatment effects, methodological issues, and quality of included studies.⁶

Search strategies were prepared for each intervention separately to identify the existing reviews (Table S1, online only). The same search strategy was applied to look for primary trials, with some modification, as detailed in Table S1 (Fig. 2). Data analysis of the outcomes was based on an intention-to-treat principle. For dichotomous data, we presented results as summary risk ratio (RR) and as odds ratio (OR) for observational studies with 95% CIs. For continuous data, we used the mean difference (MD), standard mean difference (SMD), or weighted mean difference between trials if outcomes were measured comparably. All the analysis was performed using Review Manager 5.⁷

The outcomes of interest included mortality (death rate, death due to any disease, fatality rate, perinatal/neonatal mortality), pregnancy outcomes (preterm births, preeclampsia, eclampsia, stillbirth, miscarriage, termination of pregnancy), morbidity (obesity, obesity-related complications, hypertension, diabetes, cardiovascular disease, infectious diseases), nutritional deficiencies (anemia or micronutrient deficiency), and anthropometrics (height, weight, body mass index (BMI), stunting).

On the basis of the findings from reviews, the quality of the evidence was assessed following the Grading of Recommendations, Assessment, Development and Evaluation methodology.⁸ The following criteria were taken into account to grade the evidence: study limitations (i.e., risk of bias),




Figure 2. Flow diagram for primary studies on adolescents.

consistency of effect, imprecision, indirectness, and publication bias. The quality of the body of evidence for each key outcome was then rated as “high,” “moderate,” “low,” or “very low.”

Results

We identified 35 systematic reviews, of which five focused on children and adolescents.^{9–13} These reviews examined interventions related to the prevention and management of obesity.^{9–13} On the other hand, 12 reviews targeted women of reproductive age,^{14–25} 12 included pregnant women,^{26–37} seven included children (5–18 years),^{38–44} and one included a mother/child population.⁴⁵ For interventions where evidence on adolescents was scarce, including micronutrient supplementation, food/balanced protein energy supplementation, and prevention of gestational diabetes, we identified and included evidence from primary trials.

Micronutrient supplementation

Iron supplementation. Our search found no systematic reviews of iron supplementation in adolescents; however, we identified reviews on women of reproductive age and pregnant women.^{14,26,27} Our search for primary trials on iron supplementation in adolescents identified 11 studies, of which seven

reported serum hemoglobin as an outcome.^{46–56} Studies were conducted in India, Iran, Pakistan, Sri Lanka, Tanzania, the United States, and Vietnam and provided iron as ferrous sulfate preparations in dosages ranging from 50 to 260 mg. The pooled analysis showed a significant impact of iron supplementation on improving serum hemoglobin levels (SMD 1.83; 95% CI 0.59–3.08; seven studies; *n* = 1257) (Fig. 3A).

Although evidence from studies in adolescents was scarce, reviews focusing on women of reproductive age showed that intermittent iron supplementation (alone or with other nutrients) had a significant impact on improving anemia rates, hemoglobin, and serum ferritin concentration. However, no difference was reported for iron-deficiency anemia, serum iron deficiency, and all-cause morbidity.¹⁴ While intermittent iron supplementation in pregnant women showed no statistically significant effect on anemia, iron-deficiency anemia, low birth weight (LBW), or preterm birth, there was a significant effect on improving serum hemoglobin concentration.²⁶ Daily iron supplementation in pregnant women, on the other hand, showed a significant effect on reducing maternal anemia, maternal iron deficiency, and LBW and improving mean birth weight (Table S2, online only).




Figure 3. Iron supplementation. (A) Iron supplementation versus placebo: hemoglobin concentration (g/L) in adolescents. (B) Weekly/intermittent IFA supplementation versus placebo: hemoglobin (g/L) in adolescents. References: Beasley,⁵³ Bruner,⁴⁶ Eftekhari,⁴⁷ Hettiarachchi,⁴⁸ Mozaffari-Khosravi,⁵⁴ Rezaeian,⁵⁶ Shobha,⁵⁵ Agarwal,⁵⁸ Gilgen,⁶⁰ Hall,⁶¹ Jayatissa,⁶² Roschnik,⁶⁷ Soekarjo.⁷⁰

Folic acid supplementation. We found no systematic review on the supplementation of folic acid in adolescents. However, we found a single study on folic acid supplementation among adolescents from Canada that reported a nonsignificant effect on serum folate concentration (SMD 1.89; 95% CI 1.0–2.79). We found reviews on folic acid supplementation in children,³⁸ women of reproductive age,¹⁵ and pregnant women.²⁸

One review reported that periconceptional folic acid supplementation alone in women of reproductive age showed significant reduction in the number of pregnancies terminated for fetal abnormality and neural tube defects (NTDs). No effects were seen for multiple pregnancy, miscarriages, stillbirths, and congenital anomalies, such as cleft palate, cleft lip, congenital cardiovascular defects, and other birth defects.¹⁵ The same review showed that supplementation with other micronutrients had a nonsignificant effect on pregnancy termination for fetal abnormality, multiple pregnancy, miscarriages, stillbirths, NTDs, cleft palate, cleft lip, congenital cardiovascular defects and other birth defects, and

LBW.¹⁵ On the other hand, folic acid supplementation in pregnant women led to no change in anemia, serum hemoglobin, or folate concentration. However, megaloblastic anemia decreased significantly. No change was seen in stillbirth/neonatal death, preterm birth, LBW, or birth weight (Table S3, online only).²⁸

Iron and folic acid supplementation. We found no systematic review on the supplementation of IFA in adolescents, but we identified two reviews on pregnant women.^{29,30} In our search for primary trials on adolescents, we identified 23 moderate-quality studies from Bangladesh, Kenya, India, Indonesia, Iran, Mali, Malaysia, Mozambique, Nepal, Sri Lanka, and Tanzania.^{57–78} Iron dosages across those studies ranged from 60 to 120 mg/day, and folic acid ranged from 3.5 to 500 mg/day. While there was a lot of variability in study design to detect dose–response results, the overall evidence suggests that daily IFA supplementation reduced the prevalence of anemia (RR 0.52; 95% CI 0.28–0.96; three studies; $n = 1979$) and

improved the serum hemoglobin concentrations (MD 10.07; 95% CI 4.05–16.10; five studies; $n = 822$) among adolescent girls and that weekly IFA supplementation also led to a reduction in anemia (RR 0.73; 95% CI 0.58–0.92; nine studies) and improved the serum hemoglobin concentration (MD 2.24; 95% CI 0.36–4.12; six studies; $n = 2839$) (Fig. 3B).

Multiple micronutrient supplementation. We found no systematic reviews on MMNs among adolescents, but identified reviews on children,^{39,40} general adult populations,¹⁶ and pregnant women.³¹ In our search for primary studies, we identified five moderate- to high-quality studies on adolescents.^{79–83} Of these, four were conducted on nonpregnant adolescents girls from India,⁸⁴ Indonesia,⁸² and Bangladesh,^{80,81} while one was conducted on pregnant adolescents from Brazil.⁸³ MMN supplementation among nonpregnant (SMD 0.55; 95% CI 0.30–0.81) and pregnant girls (SMD 1.30; 95% CI 0.31–2.28) showed a significant improvement in serum hemoglobin concentration.

Data from pregnant women suggest that MMN supplementation, when compared with no supplements, placebo, or supplementation with two micronutrients or fewer, had a significant effect on maternal anemia (third trimester hemoglobin <110 g/L), small for gestational age (SGA), and LBW. However, no change was reported for preeclampsia, miscarriage (loss before 28 weeks), maternal mortality, perinatal mortality, neonatal mortality, and preterm births.³¹ The review also suggests that MMN had similar effects when compared to IFA³¹ (Table S4, online only).

Calcium supplementation. No systematic review was found in adolescent populations, while other reviews included examined children and adolescents,^{41,42} women of reproductive age, and pregnant women.^{32,33}

We included four studies from the primary search: two each on nonpregnant^{85–87} and pregnant adolescents.^{88,89} Calcium supplementation showed a nonsignificant improvement in hip bone mineral density after 1 year of supplementation for nonpregnant adolescents (SMD 1.17; 95% CI –0.45 to 2.79) and serum calcium concentration (SMD 0.17; 95% CI –0.32 to 0.67). Included studies were conducted in Israel, the United States, and China, with calcium dosages ranging from 600 to 1200 mg/day, and are

considered of high quality. Calcium supplementation among pregnant adolescents showed no effect on mean birth weight (SMD 0.47; 95% CI –0.17 to 1.10).

Data from pregnant women showed that low-dose calcium supplementation alone reduces high blood pressure, preeclampsia, and preterm births. No impact was reported in caesarean births or still-birth/death before discharge.⁹⁰ Routine high-dose calcium supplementation (>1 g/day) in women with adequate calcium diet showed significant effects in reducing high blood pressure, while, in a low-calcium intake population, high-dose calcium supplementation showed significant effects on reducing high blood pressure and preeclampsia (Table S5, online only).⁹⁰

Vitamin D supplementation. We found no systematic review of vitamin D supplementation in adolescents. Reviews were mostly conducted on children, adults, and pregnant women.¹⁷ We found three primary studies in our search, two were conducted in Finland^{91,92} and one in India.⁹³ These studies reported nonsignificant change in serum 25(OH)D levels at 1, 2, and 3 years after vitamin D supplementation (Fig. 4). The data from the general pregnant population showed no change in preeclampsia (RR 0.67; 95% CI 0.33–1.35; one study, $n = 400$).¹⁷

Calcium and vitamin D supplementation. We found no systematic reviews on calcium and vitamin D supplementation among adolescents. We did not find any primary studies, either.

Vitamin A supplementation. No systematic reviews were found on vitamin A supplementation in adolescents. Reviews were mostly conducted in children,⁴³ pregnant women,^{34,35} and women of reproductive age.¹⁸ In our search for primary studies, we found three moderate-quality studies on adolescents from Bangladesh,⁵⁷ Kenya,⁹⁴ and Indonesia.⁷⁰ A study from Bangladesh on 138 nonpregnant adolescents who were provided with 2.42 mg retinol as vitamin A supplementation reported a significant reduction in anemia (RR 0.73; 95% CI 0.56–0.93).⁵⁷ The evidence from women of reproductive age showed improvements in maternal serum retinol levels (MD 0.17; 95% CI 0.06–0.28; three studies; $n = 258$).¹⁸ One review on vitamin A supplementation during pregnancy reported




Figure 4. Vitamin D supplementation versus placebo: 25(OH)D (nmol/L) concentration in adolescents. References: Khadilkar,⁹³ Lehtonen-Veromaa.⁹¹

a significant impact on anemia (hemoglobin < 110 g/L) (Table S6, online only).³⁴

Zinc supplementation. No systematic reviews were found on adolescents. Reviews evaluating zinc supplementation were found in children,⁴⁴ adults,¹⁹ and pregnant women.³⁶ We identified seven studies on adolescents from a primary search.^{48,84,95–100} In these studies, which were from Brazil, Chile, India, Sri Lanka, the United Kingdom, and the United States, adolescents were supplemented with zinc ranging from 14 to 20 mg/day. Of these, four studies examined pregnant adolescents.^{96–100} Two studies on 494 nonpregnant adolescents reported significant increases in hemoglobin concentration (SMD 4.81; 95% CI 0.47–8.66; two studies; *n* = 494) (Fig. 5A) and serum zinc concentration (SMD 4.28; 95% CI 0.49–6.06; three studies; *n* = 805) (Fig. 5B). Studies on pregnant adolescent reported a significant reduction in preterm births (RR 0.57; 95% CI 0.46–0.69; two studies) (Fig. 5C) and LBW (RR 0.39; 95% CI 0.15–0.98, one study) (Fig. 5D).

Food fortification with zinc in women of reproductive age led to no change in offspring height growth or weight gain. Neonatal serum zinc levels did not alter significantly, nor did neonatal serum hemoglobin.¹⁹ Zinc supplementation in pregnant women did not significantly alter stillbirth/neonatal mortality risk. Preterm birth decreased significantly. Birth weight, LBW, and SGA did not change (Table S7, online only).³⁶

Iodine supplementation. We found no systematic reviews on iodine supplementation in adoles-

cents. Reviews were mainly conducted on children and pregnant women.⁴⁵ We conducted a search for any trials of iodine supplementation in adolescents but found only one study that fulfilled the criteria. That study, from Iran, provided a single oral dose of 190 mg iodine to high school girls and reported no change in serum thyroid-stimulating hormone (TSH) (MD 0.30; 95% CI –0.06 to 0.66, *n* = 47) (Fig. 6).⁴⁷ Iodine supplementation in pregnant women led to a significant decrease in cretinism (RR 0.27; 95% CI 0.12–0.6; five studies, *n* = 9500).⁴⁵

Food/protein energy supplementation. We found no systematic reviews of food/balanced energy protein (BEP) supplementation in adolescents; the available systematic reviews examined pregnant women and women of reproductive age, providing oral protein supplementation and nutritional education along with balanced supplementation.^{20–25,101} Our search for adolescent-specific trials identified three trials.^{102–105} A study from the United States provided nutritional supplementation to pregnant adolescents (provided 240 kcal of energy, 14.5 g of protein, 33.2 g of carbohydrate with 4 g of lactose and the rest as sucrose, 5.6 g of fat, 4 mg of iron, and 240 mg of calcium in 240 mL). This study reported a non-significant increase in birth weight (SMD 156.80; 95% CI –1.82 to 315.42).¹⁰² Another study from Peru^{103,104} consisted of participatory training with community kitchen leaders, educational materials, and increased access to heme iron (chicken liver




Figure 5. Zinc supplementation versus placebo: (A) hemoglobin (g/L) concentration in adolescents, (B) serum zinc ($\mu\text{mol/L}$) in adolescents, (C) preterm birth in pregnant adolescents, and (D) low birth weight in pregnant adolescents. References: Chiplonkar,⁸⁴ Hettiarachchi,⁴⁸ Clark,⁹⁵ Castillo-Duran,⁹⁹ Cherry.⁹⁶

and blood) for adolescents in the first 5 months, and reported a reduction in anemia rates (RR 0.32; 95% CI 0.26–0.69; $n = 22$). Another study from India¹⁰⁵ targeted university students and provided them with iron along with energy supplementation and reported a nonsignificant effect on hemoglobin concentration (MD -0.10 ; 95% CI -0.46 to 0.26 ; $n = 30$).

The evidence from a review on children and adolescents 5–18 years of age in school-feeding programs found an increase in weight gain of 0.39 kg (95% CI 0.11 – 0.67) over an average of 19 months of supplementation. Similarly, the school-feeding programs improved school attendance and math scores.¹⁰¹

There is a paucity of information on food supplementation or protein energy supplementation

among at-risk adolescents. However, evidence from studies among general pregnant women reported significant effects on reducing stillbirths, neonatal mortality, LBW, and SGA.²⁵ The review also showed improvement in birth weight and birth length.²⁵ However, Bayley mental scores at 1 year remained unchanged²⁵ (Table S8, online only).

Nutrition education for pregnant adolescents

We found two studies on pregnant adolescents from already established reviews,^{106,107} with no studies found during the updated search. These studies involved healthy-eating education programs. Pooled analysis (three studies; $n = 288$) found no significant difference in maternal weight gain (SMD 0.20 ; 95% CI -0.04 to 0.45); and no difference was found in birth weight (SMD 0.25 ; 95% CI -0.02 to 0.52 ; two studies; $n = 233$) (Fig. 7).




Figure 6. Iodine supplementation versus placebo: TSH (μU/dL) concentration in adolescents. References: Hun,¹⁰⁶ Long.¹⁰⁷

Gestational diabetes prevention and management

We found no systematic reviews or trials aimed specifically at preventing or managing gestational diabetes mellitus (GDM) in adolescents. Our systematic review evaluating GDM interventions on women of reproductive age in general included interventions focusing on dietary, lifestyle, and physical activity advice and interventions. Those reviews reported that preconception counseling alone for preconception women had an impact on decreasing HbA1c levels in the first trimester.³⁷ The reviews also reported a decrease in perinatal mortality and congenital malformations.³⁷ No significant change was seen in the rate of miscarriages or in preterm deliveries³⁷ (Table S9, online only).

A comprehensive systematic review on pregnant women found that early detection of GDM in pregnant women led to a significant decrease in GDM (RR 0.24; 95% CI: 0.16–0.37) and abnormal fasting glucose (RR 0.21; 95% CI 0.11–0.39; one study). However, the results from one study showed no significant change in the rate of caesarean sections (RR 0.83; 95% CI 0.60–1.15).³⁷

Borderline diabetic care in pregnant women did not show any significant change in preterm births but did lead to a significant alteration in mean birth weight.³⁷ The same review reported that continuous insulin dosing compared with standard dosing did not lead to any change in caesarean sections or hypoglycemia. No change was seen in stillbirths/intrauterine fetal deaths, preterm deliveries, mean birth weight, SGA babies, or large for gestational age babies.³⁷

The comparison of intensive care against standard care did not show any significant difference in stillbirths. Comparing GDM treatment with standard care showed no change in caesarean sections or in the proportion of children with birth weight

above the 90th percentile.³⁷ Comparing optimal and suboptimal glucose control showed no significant change in stillbirths. A significant decrease was seen in perinatal deaths.³⁷ Exercise with or without dietary changes led to significant postpartum weight loss.³⁷

Obesity prevention and management

Obesity prevention. We found three systematic reviews focusing on adolescents,^{11–13} of which two reported computer-mediated interventions^{11,12} and one reported surgery.¹³ Systematic reviews conducted in the general population of children and women of reproductive age were more comprehensive and included school-based, primary care-based, community-based, behavioral, diet, and supplementary interventions.^{108–111} We identified specific trials (*n* = 31) from those reviews on adolescent populations and updated the search (*n* = 28). The interventions involved diet, physical activity, behavior change, and combinations of these. They were provided via school, community, family, technology, or a combination of those methods.

Interventions, including health promotion and education, counseling on diet, physical activity/lifestyle support, and dietary advice, showed marginal impact on reducing BMI (SMD –0.05; 95% CI –0.11 to 0.01; 15 studies; *n* = 14,912) (Fig. 8A) in adolescents. Moreover, there is an increasing evidence that the types of fluids that we drink can have a long-term impact on health, influencing the development of overweight, obesity, or metabolic diseases. Therefore, plain water is recommended as a healthy option over sweetened beverages for prevention of obesity.¹¹²

Obesity management. We identified two systematic reviews that were conducted on children.^{9,10} We also identified 12 high-quality studies on adolescents^{113–124} from HICs that showed a significant




Figure 7. Interventions for prevention of obesity in pregnant adolescent: birth weight. References: Amaro,¹¹³ Bonsergent,¹¹⁴ Ebbeling,¹¹⁵ Foster,¹¹⁶ Gentile,¹¹⁷ Haerens,¹¹⁸ Kain,¹¹⁹ Lubans,¹²⁰ Neumark-Sztainer,¹²¹ Reed,¹²² Robinson,¹²³ Sichieri,¹²⁴ Simon,¹²⁵ Singh,¹²⁶ Webber.¹²⁷

reduction in BMI postintervention that involved diet, physical activity, behavior change, and combinations of those (SMD -0.24 ; 95% CI -0.36 to -0.13 ; 12 studies; $n = 1541$) (Fig. 8B). Subgroups showed the largest effect for family-based interventions (SMD -0.43 ; 95% CI -0.76 to -0.09 ; two studies; $n = 215$) and the combination of the interventions (SMD -0.23 ; 95% CI -0.38 to -0.08 ; two studies; $n = 720$).

Discussion

To the best of our knowledge, this is the first comprehensive systematic review on adolescent nutrition interventions. Although we aimed to include data from existing systematic reviews, we searched for primary trials when data from systematic reviews on adolescents were not available. We further summarized the data from nutritional interventions from recently published systematic reviews on women of reproductive age, including pregnant women, in order to contextualize the evidence to support nutrition among adolescent girls and young women, especially those in LMICs.

Our review suggests that iron supplementation among adolescents can significantly improve serum

hemoglobin concentration. Data from women of reproductive age, including pregnant women, suggest effects on improving anemia rates, serum hemoglobin and ferritin concentrations, and birth weight. Studies on folic acid supplementation alone were scarce, but IFA supplementation showed effects on improving serum hemoglobin concentration. Data from reviews on periconceptional folic acid supplementation showed significant effects on reducing NTDs. Data from studies on MMN supplementation in adolescent and pregnant adolescent populations showed significant improvement in serum hemoglobin concentration. We found a paucity of trials on calcium and vitamin D supplementation in adolescents, and those that exist did not show improvement in bone mass density. Data from reviews on pregnant women showed that low-dose calcium improves preeclampsia and preterm birth. Limited evidence was found on vitamin A supplementation and iodine supplementation. Pooled analysis on zinc supplementation in adolescents showed improvement in serum zinc and hemoglobin concentration, whereas supplementation in pregnant adolescents showed significant improvement in preterm birth and LBW.




Figure 8. Interventions for management of obesity in adolescents: BMI. References: Mihas,¹²⁸ Melnik,¹²⁹ Doyle,¹³⁰ Jones,¹³¹ Ptomey,¹³² Brennan,¹³³ Toulabi,¹³⁴ Nguyen,¹³⁵ Dâmaso,¹³⁶ Davis,¹³⁷ Kong.¹³⁸

We found a limited number of studies that supplemented food/protein energy in adolescents, and one study from Peru on only a few adolescents showed a reduction in anemia rates. The data from pregnant women showed significant impact of balanced energy/protein on stillbirths, neonatal mortality, and LBW. Similarly, only two studies on very few pregnant adolescents reported the impacts of education interventions on maternal weight gain and birth weight, and those were not significant.

It should be noted that the majority of trials on adolescent micronutrient supplementation were conducted in LMICs, making the findings

relevant since these countries stand the highest burden of undernutrition and micronutrient deficiencies. However, these findings should be interpreted with caution owing to imprecision and high heterogeneity. Moreover, we could not find any study on adolescents related to other micronutrients, such as vitamin B₁₂.

Adolescents are living in increasingly obesogenic environments, and the world is witnessing increasing rates of overweight and obesity. Our review suggests that interventions to promote health and prevent obesity had nonsignificant effects on reducing BMI. On the other hand, interventions

to manage obesity showed significant effects on reducing BMI. The higher rates of overweight and obesity have consistently been reported in high-income settings, particularly among those who are disadvantaged and live with food insecurity. The majority of the studies that we identified were conducted in HICs, limiting the generalizability of findings to HICs only. However, with the current increasing trends of double burden of malnutrition, there is a need for research in LMICs on this issue. It is also imperative to promote health and education for those at risk of obesity and prediabetes in school/community settings. It is also important to arrange for counseling for appropriate dietary intake and dietary advice to prevent obesity and to improve and increase physical activity and lifestyle support.

We aimed to complete a review of adolescent nutritional interventions to evaluate their effectiveness; however, this review has a number of limitations. First, there was a paucity of reviews and clinical trials on adolescent population. Second, the majority of the trials conducted on women of reproductive age (including pregnant women) did not provide disaggregated findings on adolescent populations. Third, certain trials on adolescents have a varied age range and, although a majority of the study population was within our specified range, the findings need to be interpreted with caution. Fourth, a few trials were reported as quasiexperimental, but we found that they used a pre-post method of reporting final data, which limited our ability to pool those studies with other randomized trials. Fifth, some studies did not report the numbers of participants in each group or reported them only in a graph, which again limited our ability to pool the data. Sixth, there was a paucity of trials on nutrition-sensitive interventions. Seventh, most of the studies that we included were ranked as moderate in quality, owing to lack of blinding and/or the lack of reporting of randomization and allocation-concealment methods. Since most of these interventions included educational activities, blinding may have been difficult if not entirely impossible to perform. Therefore, there is a strong need for detailed and comprehensive studies evaluating nutritional interventions conducted on adolescents. Eighth, the majority of the studies that we identified contained very few adolescents, and the results presented high heterogeneity. Last, we did not include ongoing studies on

trial registries; however, we plan to update the evidence on nutritional interventions for adolescents at a subsequent time.

While Figure 1 depicts the nutrition-specific and nutrition-sensitive interventions for improving the health of adolescents, it is important to underscore that our review has only assessed evidence of nutrition-specific interventions and their impact on improved adolescent nutrition and birth outcomes. It is equally important to keep in mind that nutrition-sensitive interventions also have the goals of addressing issues that are indirectly related to malnutrition, such as poverty, environmental contamination, tobacco control, infections, poor education and health literacy, and gender inequalities. Therefore, it is imperative that countries design nutritional interventions, particularly for this population. Implementation through community and school platforms may work. Interventions to reduce the rates of overweight and obesity should also be designed for rapidly urbanizing countries. Similarly, there is a need to implement micronutrient and food supplementation programs for countries with the highest burden of undernutrition and stunting.

Conclusions

This review is limited by the information available on nutrition-specific interventions in adolescents from LMICs. In relation to wasting, while we found limited evidence related to BEP supplementation among adolescents, the general evidence from pregnant women provides evidence of benefits in reducing the risk of stillbirths and neonatal mortality. Similarly, the evidence on interventions to tackle micronutrient deficiencies among adolescents is limited, although most of the studies are from LMICs. The evidence indicates that iron, IFA, and MMN supplements can significantly improve hemoglobin concentration. There is a paucity of trials on calcium and vitamin D supplementation among adolescents; however, the evidence from reviews on pregnant women at risk of low calcium intake shows positive maternal and newborn outcomes and hence could be recommended for this age group as well. Our review found some evidence supporting strategies for obesity prevention and management among adolescents, although the evidence on benefits of interventions to prevent and reduce obesity was mostly from high-income settings with limited generalizability to LMICs. The

evidence summarized in this paper suggests incorporating interventions for prevention and management of obesity as well as those pertaining to food supplementation and micronutrient supplementation in at-risk populations.

Our review shows that there is a general dearth of studies on food and micronutrient supplementation in this population. Given the high proportion of adolescents in studies of nutrition interventions in pregnancy, almost all prenatal supplementation trials have included this age group, but they have not examined or reported disaggregated outcomes among adolescents. In the absence of a specific interaction with age and extra requirements for growth, one can assume that the direction of effect and possible impacts would be comparable between adolescents and young women. We therefore also included available evidence from women of reproductive age, including pregnant women, to assess plausible interventions and strategies.

Supporting Information

Additional supporting information may be found in the online version of this article.

Table S1. Search strategies.

Table S2. Summary estimates of interventions for iron supplementation in pregnant women and women of reproductive age.

Table S3. Summary estimates of interventions for folic acid supplementation in pregnant women and women of reproductive age.

Table S4. Summary estimates of interventions for MMN in pregnant women.

Table S5. Summary estimates of interventions for calcium supplementation in pregnant women.

Table S6. Summary estimates of interventions for vitamin A supplementation in pregnant women.

Table S7. Summary estimates of interventions for zinc supplementation in pregnant and women of reproductive age.

Table S8. Summary estimates of interventions for protein energy supplementation in pregnant women.

Table S9. Summary estimates of interventions for GDM in preconception women and women of reproductive age.

Competing interests

The authors declare no competing interests.

References

1. UNFPA. 2014. The Power of 1.8 billion: Adolescents, youth and the transformation of the future. New York: United Nations Population Fund. ISBN 978-0-89714-972-3.
2. Patton, G.C. *et al.* 2016. Our future: a Lancet commission on adolescent health and wellbeing. *Lancet* **387**: 2423–2478.
3. Sawyer, S.M. *et al.* 2012. Adolescence: a foundation for future health. *Lancet* **379**: 1630–1640.
4. Salam, R.A. *et al.* 2016. Interventions to improve adolescent nutrition: a systematic review and meta-analysis. *J. Adolesc. Health* **59**: S29–S39.
5. Shea, B.J. *et al.* 2007. Development of AMSTAR: a measurement tool to assess the methodological quality of systematic reviews. *BMC Med. Res. Methodol.* **7**: 10.
6. The Cochrane Collaboration. 2011. *Cochrane Handbook for Systematic Reviews of Interventions*. Version 5.1.0. Updated March 2011. www.cochrane-handbook.org.
7. The Cochrane Collaboration. 2008. Review Manager (RevMan) [computer program]. Version 5.0. Copenhagen: The Nordic Cochrane Centre.
8. Guyatt, G.H. *et al.*; GRADE Working Group. 2008. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* **336**: 924–926.
9. Oude Luttikhuis, H. *et al.* 2009. Interventions for treating obesity in children. *Cochrane Database Syst. Rev.* DOI: 10.1002/14651858.CD001872.pub2.
10. Whitlock, E.P. *et al.* 2010. Effectiveness of weight management interventions in children: a targeted systematic review for the USPSTF. *Pediatrics* **125**: e396–e418.
11. Ajie, W.N. & K.M. Chapman-Novakofski. 2014. Impact of computer-mediated, obesity-related nutrition education interventions for adolescents: a systematic review. *J. Adolesc. Health* **54**: 631–645.
12. Chen, J. & M.E. Wilkosz. 2014. Efficacy of technology-based interventions for obesity prevention in adolescents: a systematic review. *Adolesc. Health Med. Ther.* **5**: 159–170.
13. Ells, L.J. *et al.* 2015. Surgery for the treatment of obesity in children and adolescents. *Cochrane Database Syst. Rev.* DOI: 10.1002/14651858.CD011740.
14. Fernandez-Gaxiola, A.C. & L.M. De-Regil. 2011. Inter-mittent iron supplementation for reducing anaemia and its associated impairments in menstruating women. *Cochrane Database Syst. Rev.* DOI: 10.1002/14651858.CD009218.pub2.
15. De-Regil, L.M. *et al.* 2010. Effects and safety of periconceptional folate supplementation for preventing birth defects. *Cochrane Database Syst. Rev.* DOI: 10.1002/14651858.CD007950.pub2.
16. Allen, L.H., J.M. Pearson & D.K. Olney. 2009. Provision of multiple rather than two or fewer micronutrients more effectively improves growth and other outcomes in micronutrient-deficient children and adults. *J. Nutr.* **139**: 1022–1030.

17. De-Regil, L.M. *et al.* 2012. Vitamin D supplementation for women during pregnancy. *Cochrane Database Syst. Rev.* CD008873.
18. Oliveira-Menegozzo, J.M. *et al.* 2010. Vitamin A supplementation for postpartum women. *Cochrane Database Syst. Rev.* DOI: 10.1002/14651858.CD005944.pub2.
19. Das, J.K. *et al.* 2013. Systematic review of zinc fortification trials. *Ann. Nutr. Metab.* **62**(Suppl. 1): 44–56.
20. Potter, J., P. Langhorne & M. Roberts. 1998. Routine protein energy supplementation in adults: systematic review. *BMJ* **317**: 495–501.
21. Dwarkanath, P. *et al.* 2016. Energy and protein supplementation does not affect protein and amino acid kinetics or pregnancy outcomes in underweight indian women. *J. Nutr.* **146**: 218–226.
22. Ota, E. *et al.* 2015. Antenatal dietary education and supplementation to increase energy and protein intake. *Cochrane Database Syst. Rev.* **9**: CD000032.
23. Stevens, B. *et al.* 2015. The effect of balanced protein energy supplementation in undernourished pregnant women and child physical growth in low- and middle-income countries: a systematic review and meta-analysis. *Matern. Child Nutr.* **11**: 415–432.
24. Imdad, A. & Z.A. Bhutta. 2011. Effect of balanced protein energy supplementation during pregnancy on birth outcomes. *BMC Public Health* **11**: S17.
25. Imdad, A. & Z.A. Bhutta. 2012. Maternal nutrition and birth outcomes: effect of balanced protein-energy supplementation. *Paediatr. Perinat. Epidemiol.* **26**(Suppl. 1): 178–190.
26. Peña-Rosas, J.P. *et al.* 2015. Intermittent oral iron supplementation during pregnancy. *Cochrane Database Syst. Rev.* DOI: 10.1002/14651858.CD009997.pub2.
27. Peña-Rosas, J.P. *et al.* 2012. Daily oral iron supplementation during pregnancy. *Cochrane Database Syst. Rev.* **12**: CD004736.
28. Lassi, Z.S. *et al.* 2013. Folic acid supplementation during pregnancy for maternal health and pregnancy outcomes. *Cochrane Database Syst. Rev.* **3**: CD006896.
29. Yakoob, M.Y. & Z.A. Bhutta. 2011. Effect of routine iron supplementation with or without folic acid on anemia during pregnancy. *BMC Public Health* **11**: S21.
30. Imdad, A. & Z.A. Bhutta. 2012. Routine iron/folate supplementation during pregnancy: effect on maternal anaemia and birth outcomes. *Paediatr. Perinat. Epidemiol.* **26**: 168–177.
31. Haider, B.A. & Z.A. Bhutta. 2015. Multiple-micronutrient supplementation for women during pregnancy. *Cochrane Database Syst. Rev.* **11**: CD004905.
32. Imdad, A. & Z.A. Bhutta. 2012. Effects of calcium supplementation during pregnancy on maternal, fetal and birth outcomes. *Paediatr. Perinat. Epidemiol.* **26**(Suppl. 1): 138–152.
33. Hofmeyr, G.J., J.M. Belizan & P. von Dadelzen. 2014. Low-dose calcium supplementation for preventing pre-eclampsia: a systematic review and commentary. *BJOG* **121**: 951–957.
34. Thorne-Lyman, A.L. & W.W. Fawzi. 2012. Vitamin A and carotenoids during pregnancy and maternal, neonatal and infant health outcomes: a systematic review and meta-analysis. *Paediatr. Perinat. Epidemiol.* **26**(Suppl. 1): 36–54.
35. McCauley, M.E. *et al.* 2015. Vitamin A supplementation during pregnancy for maternal and newborn outcomes. *Cochrane Database Syst. Rev.* DOI: 10.1002/14651858.CD008666.pub3.
36. Ota, E. *et al.* 2015. Zinc supplementation for improving pregnancy and infant outcome. *Cochrane Database Syst. Rev.* DOI: 10.1002/14651858.CD000230.pub5.
37. Lassi, Z.S. & Z.A. Bhutta. 2013. Risk factors and interventions related to maternal and pre-pregnancy obesity, pre-diabetes and diabetes for maternal, fetal and neonatal outcomes: a systematic review. *Expert Rev. Obstet. Gynecol.* **8**: 639–660.
38. Lohner, S. *et al.* 2012. Effect of folate supplementation on folate status and health outcomes in infants, children and adolescents: a systematic review. *Int. J. Food Sci. Nutr.* **63**: 1014–1020.
39. Irlam, J.H. *et al.* 2013. Micronutrient supplementation for children with HIV infection. *Cochrane Database Syst. Rev.* **10**: CD010666.
40. Eilander, A. *et al.* 2010. Multiple micronutrient supplementation for improving cognitive performance in children: systematic review of randomized controlled trials. *Am. J. Clin. Nutr.* **91**: 115–130.
41. Vatanparast, H. & S.J. Whiting. 2006. Calcium supplementation trials and bone mass development in children, adolescents, and young adults. *Nutr. Rev.* **64**: 204–209.
42. French, S.A., J.A. Fulkerson & M. Story. 2000. Increasing weight-bearing physical activity and calcium intake for bone mass growth in children and adolescents: a review of intervention trials. *Prev. Med.* **31**: 722–731.
43. Iannotti, L.L., I. Trehan & M.J. Manary. 2013. Review of the safety and efficacy of vitamin A supplementation in the treatment of children with severe acute malnutrition. *Nutr. J.* **12**: 125.
44. Mayo-Wilson, E. *et al.* 2014. Zinc supplementation for preventing mortality, morbidity, and growth failure in children aged 6 months to 12 years of age. *Cochrane Database Syst. Rev.* **5**: CD009384.
45. Zimmermann, M.B. 2012. The effects of iodine deficiency in pregnancy and infancy. *Paediatr. Perinat. Epidemiol.* **26**(Suppl. 1): 108–117.
46. Bruner, A.B. *et al.* 1996. Randomised study of cognitive effects of iron supplementation in non-anaemic iron-deficient adolescent girls. *Lancet* **348**: 992–996.
47. Eftekhari, M.H. *et al.* 2006. Effects of administration of iron, iodine and simultaneous iron-plus-iodine on the thyroid hormone profile in iron-deficient adolescent Iranian girls. *Eur. J. Clin. Nutr.* **60**: 545–552.
48. Hettiarachchi, M. *et al.* 2008. The efficacy of micronutrient supplementation in reducing the prevalence of anaemia and deficiencies of zinc and iron among adolescents in Sri Lanka. *Eur. J. Clin. Nutr.* **62**: 856–865.
49. Rousham, E.K. *et al.* 2013. The effect of a school-based iron intervention on the haemoglobin concentration of school children in north-west Pakistan. *Eur. J. Clin. Nutr.* **67**: 1188–1192.

50. Van Thuy, P. *et al.* 2003. Regular consumption of NaFeEDTA-fortified fish sauce improves iron status and reduces the prevalence of anemia in anemic Vietnamese women. *Am. J. Clin. Nutr.* **78**: 284–290.
51. Van Thuy, P. *et al.* 2005. The use of NaFeEDTA-fortified fish sauce is an effective tool for controlling iron deficiency in women of childbearing age in rural Vietnam. *J. Nutr.* **135**: 2596–2601.
52. Meier, P. *et al.* 2003. Prevention of Iron deficiency anemia in adolescent and adult pregnancies. *Clin. Med. Res.* **1**: 29–36.
53. Beasley, N.M. *et al.* 2000. The impact of weekly iron supplementation on the iron status and growth of adolescent girls in Tanzania. *Trop. Med. Int. Health* **5**: 794–799.
54. Mozaffari-Khosravi, H. *et al.* 2010. Once weekly low-dose iron supplementation effectively improved iron status in adolescent girls. *Biol. Trace Elem. Res.* **135**: 22–30.
55. Shobha, S. & D. Sharada. 2003. Efficacy of twice weekly iron supplementation in anemic adolescent girls. *Indian Pediatr.* **40**: 1186–1190.
56. Rezaeian, A. *et al.* 2014. Effects of iron supplementation twice a week on attention score and haematologic measures in female high school students. *Singapore Med. J.* **55**: 587–592.
57. Ahmed, F., M.R. Khan & A.A. Jackson. 2001. Concomitant supplemental vitamin A enhances the response to weekly supplemental iron and folic acid in anemic teenagers in urban Bangladesh. *Am. J. Clin. Nutr.* **74**: 108–115.
58. Agarwal, K.N. *et al.* 2003. Anemia prophylaxis in adolescent school girls by weekly or daily iron-folate supplementation. *Indian Pediatr.* **40**: 296–302.
59. Februhartanty, J., D. Dillon & H. Khusun. 2002. Will iron supplementation given during menstruation improve iron status better than weekly supplementation? *Asia Pac. J. Clin. Nutr.* **11**: 36–41.
60. Gilgen, D. & C.G.N. Mascie Taylor. 2001. The effect of weekly iron supplementation on anaemia and on iron deficiency among female tea pluckers in Bangladesh. *J. Hum. Nutr. Diet.* **14**: 185–190.
61. Hall, A. *et al.* 2002. A randomised trial in Mali of the effectiveness of weekly iron supplements given by teachers on the haemoglobin concentrations of schoolchildren. *Public Health Nutr.* **5**: 413–418.
62. Jayatissa, R. & C. Piyasena. 1999. Adolescent schoolgirls: daily or weekly iron supplementation? *Food Nutr. Bull.* **20**: 429–434.
63. Kanani, S.J. & R.H. Poojara. 2000. Supplementation with iron and folic acid enhances growth in adolescent Indian girls. *J. Nutr.* **130**: 452S–455S.
64. Kianfar, H., M. Kimiagar & M. Ghaffarpour. 2000. Effect of daily and intermittent iron supplementation on iron status of high school girls. *Int. J. Vitam. Nutr. Res.* **70**: 172–177.
65. Leenstra, T. *et al.* 2009. The effect of weekly iron and vitamin A supplementation on hemoglobin levels and iron status in adolescent schoolgirls in western Kenya. *Eur. J. Clin. Nutr.* **63**: 173–182.
66. Muro, G.S. *et al.* 1999. Increase in compliance with weekly iron supplementation of adolescent girls by an accompanying communication programme in secondary schools in Dar-es-Salaam, Tanzania. *Food Nutr. Bull.* **20**: 435–444.
67. Roschnik, N., V. Phiri & M. Mukaka. 2003. The impact of weekly school-based iron supplementation: Mangochi District, Malawi (report). Save the Children.
68. Saha, K.K. *et al.* 2009. Household food security is associated with growth of infants and young children in rural Bangladesh. *Public Health Nutr.* **12**: 1556–1562.
69. Shah, B.K. & P. Gupta. 2002. Weekly vs daily iron and folic acid supplementation in adolescent Nepalese girls. *Arch. Pediatr. Adolesc. Med.* **156**: 131–135.
70. Soekarjo, D.D. *et al.* 2004. Effectiveness of weekly vitamin A (10 000 IU) and iron (60 mg) supplementation for adolescent boys and girls through schools in rural and urban East Java, Indonesia. *Eur. J. Clin. Nutr.* **58**: 927–937.
71. Tee, E.S. *et al.* 1999. School-administered weekly iron-folate supplements improve hemoglobin and ferritin concentrations in Malaysian adolescent girls. *Am. J. Clin. Nutr.* **69**: 1249–1256.
72. Yusoff, H., W.N. Wan Daud & Z. Ahmad. 2012. Nutrition education and knowledge, attitude and hemoglobin status of Malaysian adolescents. *Southeast Asian J. Trop. Med. Public Health* **43**: 192–200.
73. Horjus, P. *et al.* 2005. School-based iron and folic acid supplementation for adolescent girls: findings from Manica Province, Mozambique. *Food Nutr. Bull.* **26**: 281–286.
74. Vyas, S. *et al.* 2010. Leaf concentrate as an alternative to iron and folic acid supplements for anaemic adolescent girls: a randomised controlled trial in India. *Public Health Nutr.* **13**: 418–423.
75. Deshmukh, P.R., B.S. Garg & M.S. Bharambe. 2008. Effectiveness of weekly supplementation of iron to control anaemia among adolescent girls of Nashik, Maharashtra, India. *J. Health Popul. Nutr.* **26**: 74–78.
76. Dongre, A.R., P.R. Deshmukh & B.S. Garg. 2011. Community-led initiative for control of anemia among children 6 to 35 months of age and unmarried adolescent girls in rural Wardha, India. *Food Nutr. Bull.* **32**: 315–323.
77. Kotecha, P.V., S. Nirupam & P.D. Karkar. 2009. Adolescent girls' anaemia control programme, Gujarat, India. *Indian J. Med. Res.* **130**: 584–589.
78. Shah, S.P. *et al.* 2016. Effectiveness and feasibility of weekly iron and folic acid supplementation to adolescent girls and boys through peer educators at community level in the tribal area of Gujarat. *Indian J. Community Med.* **41**: 158–161.
79. Ahmed, F. *et al.* 2010. Long-term intermittent multiple micronutrient supplementation enhances hemoglobin and micronutrient status more than iron+ folic acid supplementation in Bangladeshi rural adolescent girls with nutritional anemia. *J. Nutr.* **140**: 1879–1886.
80. Ahmed, F. *et al.* 2012. Effect of long-term intermittent supplementation with multiple micronutrients compared with iron-and-folic acid supplementation on Hb and micronutrient status of non-anaemic adolescent school-girls in rural Bangladesh. *Br. J. Nutr.* **108**: 1484–1493.
81. Hyder, S.M. *et al.* 2007. A multiple-micronutrient-fortified beverage affects hemoglobin, iron, and vitamin A status and growth in adolescent girls in rural Bangladesh. *J. Nutr.* **137**: 2147–2153.

82. Angeles-Agdeppa, I. *et al.* 1997. Weekly micronutrient supplementation to build iron stores in female Indonesian adolescents. *Am. J. Clin. Nutr.* **66**: 177–183.
83. Correia-Santos, A.M. *et al.* 2011. Dietary supplements for the lactating adolescent mother: influence on plasma micronutrients. *Nutr. Hosp.* **26**: 392–398.
84. Chipлонkar, S.A. & R. Kawade. 2012. Effect of zinc- and micronutrient-rich food supplements on zinc and vitamin A status of adolescent girls. *Nutrition* **28**: 551–558.
85. Ilich-Ernst, J.Z. *et al.* 1998. Iron status, menarche, and calcium supplementation in adolescent girls. *Am. J. Clin. Nutr.* **68**: 880–887.
86. Ho, S.C. *et al.* 2005. A prospective study of the effects of 1-year calcium-fortified soy milk supplementation on dietary calcium intake and bone health in Chinese adolescent girls aged 14 to 16. *Osteoporos. Int.* **16**: 1907–1916.
87. Dodiuk-Gad, R.P. *et al.* 2005. Sustained effect of short-term calcium supplementation on bone mass in adolescent girls with low calcium intake. *Am. J. Clin. Nutr.* **81**: 168–174.
88. Chan, G.M. *et al.* 2006. Effects of dietary calcium intervention on adolescent mothers and newborns: a randomized controlled trial. *Obstet. Gynecol.* **108**: 565–571.
89. Lopez-Jaramillo, P. *et al.* 1997. Calcium supplementation and the risk of preeclampsia in Ecuadorian pregnant teenagers. *Obstet. Gynecol.* **90**: 162–167.
90. Hofmeyr, G.J. *et al.* 2014. Calcium supplementation during pregnancy for preventing hypertensive disorders and related problems. *Cochrane Database Syst. Rev.* DOI: 10.1002/14651858.CD001059.pub4.
91. Lehtonen-Veromaa, M. *et al.* 2002. The effect of conventional vitamin D [sub 2] supplementation on serum 25 (OH) D concentration is weak among peripubertal Finnish girls: a 3-y prospective study. *Eur. J. Clin. Nutr.* **56**: 431–437.
92. Viljakainen, H.T. *et al.* 2006. A positive dose–response effect of vitamin D supplementation on site specific bone mineral augmentation in adolescent girls: a double-blinded randomized placebo-controlled 1-year intervention. *J. Bone Miner. Res.* **21**: 836–844.
93. Khadilkar, A.V. *et al.* 2010. Vitamin D supplementation and bone mass accrual in underprivileged adolescent Indian girls. *Asia Pacific J. Clin. Nutr.* **19**: 465.
94. Talsma, E.F. *et al.* 2016. Biofortified yellow cassava and vitamin A status of Kenyan children: a randomized controlled trial. *Am. J. Clin. Nutr.* **103**: 258–267.
95. Clark, P.J., R. Eastell & M.E. Barker. 1999. Zinc supplementation and bone growth in pubertal girls. *Lancet* **354**: 485.
96. Cherry, F.F., H.H. Sandstead & A.R. Wickremasinghe. 1993. Adolescent pregnancy. Weight and zinc supplementation effects. *Ann. N.Y. Acad. Sci.* **678**: 334–337.
97. Cherry, F.F. *et al.* 1989. Adolescent pregnancy: associations among body weight, zinc nutriture, and pregnancy outcome. *Am. J. Clin. Nutr.* **50**: 945–954.
98. Hunt, I.F. *et al.* 1985. Zinc supplementation during pregnancy in low-income teenagers of Mexican descent: effects on selected blood constituents and on progress and outcome of pregnancy. *Am. J. Clin. Nutr.* **42**: 815–828.
99. Castillo-Duran, C. *et al.* 2001. Controlled trial of zinc supplementation in Chilean pregnant adolescents. *Nutr. Res.* **21**: 715–724.
100. Nogueira Ndo, N., J.V. Parente & S.M. Cozzolino. 2003. [Changes in plasma zinc and folic acid concentrations in pregnant adolescents submitted to different supplementation regimens]. *Cad. Saude Publica* **19**: 155–160.
101. Kristjansson, B. *et al.* 2007. School feeding for improving the physical and psychosocial health of disadvantaged students. *Cochrane Database Syst. Rev.* **1**: CD004676.
102. Paige, D.M. *et al.* 1981. Nutritional supplementation of pregnant adolescents. *J. Adolesc. Health* **1**: 261–267.
103. Creed-Kanashiro, H.M. *et al.* 2003. Formative research to develop a nutrition education intervention to improve dietary iron intake among women and adolescent girls through community kitchens in Lima, Peru. *J. Nutr.* **133**: 3987S–3991S.
104. Creed-Kanashiro, H.M. *et al.* 2000. Improving dietary intake to prevent anemia in adolescent girls through community kitchens in a periurban population of Lima, Peru. *J. Nutr.* **130**: 459S–461S.
105. Mann, S.K., S. Kaur & K. Bains. 2002. Iron and energy supplementation improves the physical work capacity of female college students. *Food Nutr. Bull.* **23**: 57–64.
106. Hun, D.J. *et al.* 2002. Effects of nutrition education programs on anthropometric measurements and pregnancy outcomes of adolescents. *J. Am. Diet. Assoc.* **102**: S100–S102.
107. Long, V.A., T. Martin & C. Janson-Sand. 2002. The great beginnings program: impact of a nutrition curriculum on nutrition knowledge, diet quality, and birth outcomes in pregnant and parenting teens. *J. Am. Diet. Assoc.* **102**: S86–S89.
108. Waters, E. *et al.* 2011. Interventions for preventing obesity in children. *Cochrane Database Syst. Rev.* DOI: 10.1002/14651858.CD001871.pub3.
109. Cai, L. *et al.* 2014. Effect of childhood obesity prevention programs on blood pressure: a systematic review and meta-analysis. *Circulation* **129**: 1832–1839.
110. Wang, Y. *et al.* 2013. Childhood obesity prevention programs: comparative effectiveness review and meta-analysis. Agency for Health Care Research and Quality, Rockville, MD.
111. National Health and Medical Research Council. 2013. *Clinical practice guidelines for the management of overweight and obesity in adults, adolescents and children in Australia: systematic review*. Melbourne: National Health and Medical Research Council.
112. Armstrong, L.E. *et al.* 2012. Recommendations for healthier hydration: addressing the public health issues of obesity and type 2 diabetes. *Clin. Obes.* **2**: 115–124.
113. Amaro, S. *et al.* 2006. Kaledo a new educational board-game, gives nutritional rudiments and encourages healthy eating in children: a pilot cluster randomized trial. *Eur. J. Pediatr.* **165**: 630–635.
114. Bonsergent, E. *et al.* 2013. Overweight and obesity prevention for adolescents: a cluster randomized controlled trial in a school setting. *Am. J. Prev. Med.* **44**: 30–39.

115. Ebbeling, C.B. *et al.* 2006. Effects of decreasing sugar-sweetened beverage consumption on body weight in adolescents: a randomized, controlled pilot study. *Pediatrics* **117**: 673–680.
116. Foster, G.D. *et al.* 2008. A policy-based school intervention to prevent overweight and obesity. *Pediatrics* **121**: e794–e802.
117. Gentile, D.A. *et al.* 2009. Evaluation of a multiple ecological level child obesity prevention program: Switch[®] what you Do, View, and Chew. *BMC Med.* **7**: 49.
118. Haerens, L. *et al.* 2006. Evaluation of a 2-year physical activity and healthy eating intervention in middle school children. *Health Educ. Res.* **21**: 911–921.
119. Kain, J., R. Uauy, C. Albala, *et al.* 2004. School-based obesity prevention in Chilean primary school children: methodology and evaluation of a controlled study. *Int. J. Obes. Relat. Metab. Discord.* **28**: 483–493.
120. Lubans, D.R. *et al.* 2012. Preventing obesity among adolescent girls: one-year outcomes of the nutrition and enjoyable activity for teen girls (NEAT Girls) cluster randomized controlled trial. *Arch. Pediatr. Adolesc. Med.* **166**: 821–827.
121. Neumark-Sztainer, D. *et al.* 2003. New Moves: a school-based obesity prevention program for adolescent girls. *Prev. Med.* **37**: 41–51.
122. Reed, K.E. *et al.* 2008. Action schools! BC: a school-based physical activity intervention designed to decrease cardiovascular disease risk factors in children. *Prev. Med.* **46**: 525–531.
123. Robinson, T.N. *et al.* 2003. Dance and reducing television viewing to prevent weight gain in African-American girls: the Stanford GEMS pilot study. *Ethn. Dis.* **13**: S1–S65.
124. Sichieri, R. *et al.* 2009. School randomised trial on prevention of excessive weight gain by discouraging students from drinking sodas. *Public Health Nutr.* **12**: 197–202.
125. Simon, C. *et al.* 2008. Successful overweight prevention in adolescents by increasing physical activity: a 4-year randomized controlled intervention. *Int. J. Obes.* **32**: 1489–1498.
126. Singh, A.S. *et al.* 2009. Dutch obesity intervention in teenagers: effectiveness of a school-based program on body composition and behavior. *Arch. Pediatr. Adolesc. Med.* **163**: 309–317.
127. Webber, L.S. *et al.* 2008. Promoting physical activity in middle school girls: trial of activity for adolescent girls. *Am. J. Prev. Med.* **34**: 173–184.
128. Mihas, C. *et al.* 2009. Evaluation of a nutrition intervention in adolescents of an urban area in Greece: short- and long-term effects of the VYRONAS study. *Public Health Nutr.* **13**: 712–719.
129. Melnyk, B.M. *et al.* 2007. The COPE healthy lifestyles TEEN program: feasibility, preliminary efficacy, & lessons learned from an after school group intervention with overweight adolescents. *J. Pediatr. Health Care* **21**: 315–322.
130. Doyle, A.C. *et al.* 2008. Reduction of overweight and eating disorder symptoms via the internet in adolescents: a randomized controlled trial. *J. Adolesc. Health* **43**: 172–179.
131. Jones, M. *et al.* 2008. Randomized, controlled trial of an internet-facilitated intervention for reducing binge eating and overweight in adolescents. *Pediatrics* **121**: 453–462.
132. Ptomey, L.T. *et al.* 2015. The use of technology for delivering a weight loss program for adolescents with intellectual and developmental disabilities. *J. Acad. Nutr. Diet.* **115**: 112–118.
133. Brennan, L. *et al.* 2013. Physiological and behavioural outcomes of a randomised controlled trial of a cognitive behavioural lifestyle intervention for overweight and obese adolescents. *Obes. Res. Clin. Pract.* **7**: e23–e41.
134. Toulabi, T. *et al.* 2012. The influence of a behavior modification interventional program on body mass index in obese adolescents. *J. Formos. Med. Assoc.* **111**: 153–159.
135. Nguyen, B. *et al.* 2013. Two-year outcomes of an adjunctive telephone coaching and electronic contact intervention for adolescent weight-loss maintenance: the Loozit randomized controlled trial. *Int. J. Obes.* **37**: 468–472.
136. Dâmaso, A.R. *et al.* 2014. Aerobic plus resistance training was more effective in improving the visceral adiposity, metabolic profile and inflammatory markers than aerobic training in obese adolescents. *J. Sports Sci.* **32**: 1435–1445.
137. Davis, J.N. *et al.* 2009. Randomized control trial to improve adiposity and insulin resistance in overweight Latino adolescents. *Obesity* **17**: 1542–1548.
138. Kong, A.P. *et al.* 2014. A randomized controlled trial to investigate the impact of a low glycemic index (GI) diet on body mass index in obese adolescents. *BMC Public Health* **14**: 180.