

DNA extraction protocol (Mag-Bind Environmental DNA)

Maria Rebolleda-Gomez

05 February, 2021

Aim

Extract DNA of pellets from isolates using magnetic beads.

Materials

- 96-plate magnetic rack
- Thermomixer (eppendorf)
- Water bath set to 70C
- Deep well plate
- PCR 96 plate
- Mag-Bind Environmental DNA96 kit (Omega Bio-Tek)

Procedure

Preparation steps:

- ☐ Take 10ml ($100\mu l \times 100\text{samples}$) of elution buffer into a falcon tube and warm it to 70C.
- ☐ Put **P2** buffer and samples on ice.
- ☐ Prepare 70% ethanol ($1\text{ml}/\text{sample} \approx 100\text{ml}/\text{plate}$)
- ☐ Make sure VHB buffer has ethanol. Otherwise add 56ml per bottle.
- ☐ Make mastermix of SLX-Mlus Buffer and RNase A.
 - SLX-Mlus Buffer $525\mu l \times 100 = 52.5\text{ml}$
 - RNase A $2\mu l \times 100 = 200\mu l$

Lysis

- ☐ Briefly centrifuge the E-Z disruptor plate (1 min at 2000 rpm)
- ☐ Re-suspend pellet samples in $527\mu l$ of mastermix.
- ☐ Seal the plate with caps
- ☐ Shake a maximum speed in the thermomixer for 5min
- ☐ Centrifuge at **500rcf** for 10 seconds
- ☐ Add $53\mu l$ of **DS Buffer**
- ☐ Shake a maximum speed in the thermomixer for 1min to mix.
- ☐ Incubate at 70C shaking very slowly (300 rpm) for 10min.
- ☐ Increase the temperature at 95C and incubate 2 more minutes.
- ☐ Centrifuge **2500rcf** for **10 min**.
- ☐ Transfer $200\mu l$ supernatant to a new set of 96-well racked tubes.

- ☐ Add 67 μ l of pre-chilled P2 buffer and 100 μ l of cHTR Reagent (shake before using).
- ☐ Seal the plate with new caps.
- ☐ Centrifuge at **2500rcf** for **5 min**.

DNA extraction

- ☐ Transfer supernatant to a 96-well plate that fits in the magnetic rack. *Our normal plates do not work. I am using the ones from the QIAGEN kit*
- ☐ Add 1 volume XP1 buffer and 20 μ l of Mag-bind particles. Mix by pipetting up and down well.
- ☐ Mix at room temperature at 500rpm for 10min.
- ☐ Place plate on magnetic rack for 5 minutes, until particles migrate to the bottom. *If particles do not migrate, pipette up and down a little to move them closer to the magnetic field.*
- ☐ Aspirate and discard cleared supernatant.
- ☐ Remove plate from magnetic rack and add 500 μ l of VHB buffer (diluted with ethanol).
- ☐ Mix for 1min at **2000rpm**.
- ☐ Place plate in magnetic rack and let sit at room temperature until particles are cleared from solution.
- ☐ Aspirate and discard the cleared supernatant without disturbing the magnetic beads.
- ☐ Remove plate from magnetic rack and add 500 μ l of 70% ethanol.
- ☐ Mix for 1min at **2000rpm**.
- ☐ Place plate in magnetic rack and let sit at room temperature until particles are cleared from solution.
- ☐ Aspirate and discard the cleared supernatant without disturbing the magnetic beads.
- ☐ Repeat for a second ethanol wash.
- ☐ Leave plate in magnetic rack for 1 min. Remove any residual liquid with a pipette.
- ☐ Leave the magnetic beads to dry for 10 minutes (in the magnetic rack)
- ☐ Add 100 μ l elution buffer preheated to 70C.
- ☐ Mix for 5min at **2000rpm**.
- ☐ Place the plate into magnetic rack until beads are cleared from liquid.
- ☐ Transfer supernatant to a PCR skirted plate and seal with film. Store at -20C.