

Newdata

Chang Lu

2025-03-27

```
library(readr)
library(dplyr)
```

```
##
## Attaching package: 'dplyr'

## The following objects are masked from 'package:stats':
##
##   filter, lag

## The following objects are masked from 'package:base':
##
##   intersect, setdiff, setequal, union
```

```
library(lubridate)
```

```
##
## Attaching package: 'lubridate'

## The following objects are masked from 'package:base':
##
##   date, intersect, setdiff, union
```

```
library(stringr)
library(purrr)
library(readxl)
```

```
# Computing Total_Quit_Time_Computed
submit_time_cols <- names(combined_df)[str_detect(names(combined_df), "-T_Page Submit")]

combined_df$Total_Quit_Time_Computed <- combined_df %>%
  select(all_of(submit_time_cols)) %>%
  mutate_all(as.numeric) %>%
  rowSums(na.rm = TRUE)

combined_df %>% select(...1, Total_Quit_Time_Computed) %>% head()
```

```
## # A tibble: 6 x 2
##   ...1      Total_Quit_Time_Computed
##   <chr>          <dbl>
## 1 Student 1          38.9
## 2 Student 2        1243.
## 3 Student 3        1209.
## 4 Student 4         172.
## 5 Student 5        1495.
## 6 Student 6        2465.
```

```
length(combined_df$Total_Quit_Time_Computed)
```

```
## [1] 32
```

```
# Rename demographic groups
combined_df <- combined_df %>%
  rename(
    Accommodations = Q21,
    Gender          = Q22,
    Ethnicity       = Q23,
    English_Proficiency = Q24,
    Country_You     = Q25_1,
    Country_Mother  = Q25_2,
    Country_Father  = Q25_3,
    Home_Language   = Q69
  ) %>%
select(-Q68) # Drop Q68 for a few people answered
```

Normality Check

```
# # Accomodations
# group_yes <- combined_df %>% filter(Accommodations == "Yes") %>% pull(Total_Quit_Time_Computed)
# group_no  <- combined_df %>% filter(Accommodations == "No") %>% pull(Total_Quit_Time_Computed)
#
# # Shapiro-Wilk test
# shapiro.test(group_yes)
# shapiro.test(group_no)
```

```
# par(mfrow = c(2, 2)) # Plot layout
#
# # Histogram
# hist(group_yes, main = "Accommodations: YES", xlab = "Total Time", col = "skyblue")
# hist(group_no,  main = "Accommodations: NO", xlab = "Total Time", col = "salmon")
#
# # Q-Q plots
# qqnorm(group_yes); qqline(group_yes, col = "blue")
# qqnorm(group_no); qqline(group_no,  col = "red")
```

Accomodations Error in shapiro.test(group_yes) : sample size must be between 3 and 5000 -> Mann-Whitney U test

```
# Group sizes
table(combined_df$Gender)
```

Gender

```
##
## Female    Male
##      23      8
```

```
# Normality check
by(combined_df$Total_Quit_Time_Computed, combined_df$Gender, shapiro.test)
```

```
## combined_df$Gender: Female
##
##  Shapiro-Wilk normality test
##
## data:  dd[x, ]
## W = 0.75886, p-value = 9.016e-05
##
## -----
## combined_df$Gender: Male
##
##  Shapiro-Wilk normality test
##
## data:  dd[x, ]
## W = 0.95465, p-value = 0.7579
```

We tested whether total quit time differed by gender. The Female group violated the assumption of normality (Shapiro-Wilk $p < 0.001$), so we used a non-parametric Mann-Whitney U test.

```
table(combined_df$Ethnicity)
```

Race

```
##
##              Asian Black or African American      Hispanic or Latino
##              10              1              5
##              Other              White
##              2              13
```

```
# Kruskal-Wallis test (non-parametric ANOVA)
kruskal.test(Total_Quit_Time_Computed ~ Ethnicity, data = combined_df)
```

```
##
##  Kruskal-Wallis rank sum test
##
## data:  Total_Quit_Time_Computed by Ethnicity
## Kruskal-Wallis chi-squared = 1.0582, df = 4, p-value = 0.9008
```

A Kruskal-Wallis test revealed no significant difference in total quit time across ethnic groups ($\chi^2(4) = 1.06$, $p = 0.901$).

```
# Clean and recode English proficiency
combined_df <- combined_df %>%
  mutate(English_Proficiency = case_when(
    str_trim(English_Proficiency) %in% c("Native", "Native Speaker") ~ "Native",
    str_trim(English_Proficiency) == "Fluent" ~ "Fluent",
    TRUE ~ English_Proficiency # keep as-is in case there's other unexpected values
  ))

table(combined_df$English_Proficiency)
```

English Proficiency

```
##
## Fluent Native
##      13      18
```

```
wilcox.test(Total_Quit_Time_Computed ~ English_Proficiency, data = combined_df)
```

```
##
## Wilcoxon rank sum exact test
##
## data: Total_Quit_Time_Computed by English_Proficiency
## W = 113, p-value = 0.8902
## alternative hypothesis: true location shift is not equal to 0
```

A Wilcoxon rank-sum test found no significant difference in total quit time between Fluent and Native English speakers ($W = 113$, $p = 0.89$)

```
# You
combined_df <- combined_df %>%
  mutate(Born_in_US = ifelse(Country_You == "United States", "Yes", "No"))

wilcox.test(Total_Quit_Time_Computed ~ Born_in_US, data = combined_df)
```

Born in US

```
##
## Wilcoxon rank sum exact test
##
## data: Total_Quit_Time_Computed by Born_in_US
## W = 92, p-value = 0.729
## alternative hypothesis: true location shift is not equal to 0
```

A Wilcoxon rank-sum test found no significant difference in total quit time between students born in the U.S. and those born elsewhere ($W = 92$, $p = 0.73$).

```
# Mother
combined_df <- combined_df %>%
  mutate(Born_in_US_M = ifelse(Country_Mother == "United States", "Yes", "No"))

wilcox.test(Total_Quit_Time_Computed ~ Born_in_US_M, data = combined_df)
```

```
##
## Wilcoxon rank sum exact test
##
## data: Total_Quit_Time_Computed by Born_in_US_M
## W = 156, p-value = 0.09266
## alternative hypothesis: true location shift is not equal to 0
```

A Wilcoxon rank-sum test comparing total quit time by mother's country of birth showed a trend toward significance ($W = 156$, $p = 0.093$), but did not reach conventional significance levels. It may need larger sample size to support.

```
# Father
combined_df <- combined_df %>%
  mutate(Born_in_US_F = ifelse(Country_Father == "United States", "Yes", "No"))

wilcox.test(Total_Quit_Time_Computed ~ Born_in_US_F, data = combined_df)
```

```
##
## Wilcoxon rank sum exact test
##
## data: Total_Quit_Time_Computed by Born_in_US_F
## W = 149, p-value = 0.2107
## alternative hypothesis: true location shift is not equal to 0
```

A Wilcoxon rank-sum test showed no significant difference in total quit time based on father's country of birth ($W = 149$, $p = 0.211$).

```
combined_df <- combined_df %>%
  mutate(English_Home = ifelse(Home_Language == "English", "Yes", "No"))

wilcox.test(Total_Quit_Time_Computed ~ English_Home, data = combined_df)
```

Home Language

```
##
## Wilcoxon rank sum exact test
##
## data: Total_Quit_Time_Computed by English_Home
## W = 152, p-value = 0.1297
## alternative hypothesis: true location shift is not equal to 0
```

A Wilcoxon rank-sum test showed no significant difference in total quit time between students who spoke English at home and those who did not ($W = 152$, $p = 0.130$).

To qiuyi

Hi Qiuyi, when comparing total quit time across demographic groups, here's how to decide between using a t-test or the Mann–Whitney U test:

Use t-test if: Both groups have at least 10 observations and the distribution of total quit time is approximately normal in both groups (check via Shapiro-Wilk or Q-Q plot)(I didnt draw plots) \

Use Mann–Whitney U test if: 1.One or both groups violate normality

2.Sample size is small (especially < 10)