

Digital Signal Processing

Discrete Time Signal in Time Domain

Wenhui Xiong
NCL
UESTC



群名称: 格拉斯哥DSP 4005
群 号: 596673294

Outline

- Time Domain Representation
- Typical Sequence & Sequence Representation
- Operation on Sequences
- Operation on Finite Length Sequences
- Sampling Process
- Correlation of Signals

Outline

- Time Domain Representation
- Typical Sequence & Sequence Representation
- Operation on Sequences
- Operation on Finite Length Sequences
- Sampling Process
- Correlation of Signals

Time Domain Representation

Discrete Time Signal: a sequences of numbers

$$x = \{x[n]\}, \quad -\infty < n < \infty$$

Time Domain Representation

Discrete Time Signal: a sequences of numbers

$$x = \{x[n]\}, \quad -\infty < n < \infty$$

- Ways to present:

- Vector: $\mathbf{x} = [x[0], x[1] \cdots x[N-1]]^T$

Time Domain Representation

Discrete Time Signal: a sequences of numbers

$$x = \{x[n]\}, \quad -\infty < n < \infty$$

- Ways to present:

- Vector: $\mathbf{x} = [x[0], x[1] \cdots x[N-1]]^T$

- sequence of num.: $\{x[n]\} = \{\cdots x[-1], \underset{\uparrow}{x[0]}, x[1] \cdots\}$


Time Domain Representation

Discrete Time Signal: a sequences of numbers

$$x = \{x[n]\}, \quad -\infty < n < \infty$$

- Ways to present:

- Vector: $\mathbf{x} = [x[0], x[1] \cdots x[N-1]]^T$
- Sequence of num.: $\{x[n]\} = \{\cdots x[-1], \textcolor{red}{x}[0], x[1] \cdots\}$
- Graph



Time Domain Representation

Length of a discrete time signal

- Finite-length (finite-duration)
 - sequence defined only for interval: $N_1 \leq n \leq N_2$
 - Length: $N = N_2 - N_1 + 1$

$$\mathbf{x} = [x[0], x[1] \cdots x[N-1]]^T$$

Zero-padding: append zero valued samples

Time Domain Representation

Types of a discrete time signal

- Infinite-length (infinite-duration)

Time Domain Representation

Types of a discrete time signal

- Infinite-length (infinite-duration)
 - **Right-sided** sequence : $x[n]$ is zero for $n < N_1$
 - Causal: if $N_1 \geq 0$

Time Domain Representation

Length of a discrete time signal

- Infinite-length (infinite-duration)
 - **Right-sided** sequence : $x[n]$ is zero for $n < N_1$
 - Causal: if $N_1 \geq 0$
 - **Left-sided** sequence: $x[n]$ is zero for $n > N_2$
 - Anti-causal: if $N_2 < 0$

Time Domain Representation

Length of a discrete time signal

- Infinite-length (infinite-duration)
 - **Right-sided** sequence : $x[n]$ is zero for $n < N_1$
 - Causal: if $N_1 \geq 0$
 - **Left-sided** sequence: $x[n]$ is zero for $n > N_2$
 - Anti-causal: if $N_2 < 0$
 - **Two-sided sequence**: n for positive & negative

Outline

- Time Domain Representation
- Typical Sequence & Sequence Representation
- Operation on Sequences
- Operation on Finite Length Sequences
- Sampling Process
- Correlation of Signals

Typical Sequence—Basic Sequence

Unit Sample Sequence(**impulse**)

$$\delta[n] = \begin{cases} 0 & n \neq 0 \\ 1 & n = 0 \end{cases}$$

Typical Sequence—Basic Sequence

Unit Sample Sequence(impulse)

$$\delta[n] = \begin{cases} 0 & n \neq 0 \\ 1 & n = 0 \end{cases}$$

Any sequence can be represented by impulse

$$x[n] = \sum_{k=-\infty}^{\infty} x[k] \delta[n-k]$$

Typical Sequence—Basic Sequence

Unit Sample Sequence(impulse)

$$\delta[n] = \begin{cases} 0 & n \neq 0 \\ 1 & n = 0 \end{cases}$$

Any sequence can be represented by impulse

$$x[n] = \sum_{k=-\infty}^{\infty} x[k] \delta[n - k]$$

Unit Step

$$u[n] = \begin{cases} 1 & n \geq 0 \\ 0 & n < 0 \end{cases}$$

Typical Sequence—Basic Sequence

Unit Sample Sequence(**impulse**)

$$\delta[n] = \begin{cases} 0 & n \neq 0 \\ 1 & n = 0 \end{cases}$$

Any sequence can be represented by impulse

$$x[n] = \sum_{k=-\infty}^{\infty} x[k] \delta[n-k]$$

Unit Step

$$u[n] = \begin{cases} 1 & n \geq 0 \\ 0 & n < 0 \end{cases}$$

Exponential sequence

$$x[n] = A\alpha^n$$

Typical Sequence—Basic Sequence

Sinusoidal

$$x[n] = A \cos(\omega_0 n + \phi) \quad \text{for all } n$$

Typical Sequence—Basic Sequence

Sinusoidal

$$x[n] = A \cos(\omega_0 n + \phi) \quad \text{for all } n$$

How about change ω_0 ?

$$A \cos((\omega_0 + 2k\pi)n + \phi)$$

Typical Sequence—Basic Sequence

Sinusoidal

$$x[n] = A \cos(\omega_0 n + \phi) \quad \text{for all } n$$

How about change ω_0 ?

$$\begin{aligned} & A \cos((\omega_0 + 2k\pi)n + \phi) \\ &= A \cos(\omega_0 n + \phi) \end{aligned}$$

Typical Sequence—Basic Sequence

Sinusoidal

$$x[n] = A \cos(\omega_0 n + \phi) \quad \text{for all } n$$

How about change ω_0 ?

$$\begin{aligned} & A \cos((\omega_0 + 2k\pi)n + \phi) \\ &= A \cos(\omega_0 n + \phi) \end{aligned}$$

$$-\pi < \omega_0 < \pi \text{ or } 0 < \omega_0 < 2\pi$$

Typical Sequence—Basic Sequence


Sinusoidal

$$x[n] = A \cos(\omega_0 n + \phi) \quad \text{for all } n$$

How about change ω_0 ?

$$\begin{aligned} A \cos((\omega_0 + 2k\pi)n + \phi) \\ = A \cos(\omega_0 n + \phi) \end{aligned}$$

$$-\pi < \omega_0 < \pi \text{ or } 0 < \omega_0 < 2\pi$$



Typical Sequence—Basic Sequence



Sinusoidal

$$x[n] = A \cos(\omega_0 n + \phi) \quad \text{for all } n$$

How about change ω_0 ?

$$\begin{aligned} & A \cos((\omega_0 + 2k\pi)n + \phi) \\ &= A \cos(\omega_0 n + \phi) \end{aligned}$$

$$-\pi < \omega_0 < \pi \text{ or } 0 < \omega_0 < 2\pi$$



Typical Sequence—Basic Sequence


Sinusoidal

$$x[n] = A \cos(\omega_0 n + \phi) \quad \text{for all } n$$

How about change ω_0 ?

$$\begin{aligned} & A \cos((\omega_0 + 2k\pi)n + \phi) \\ &= A \cos(\omega_0 n + \phi) \end{aligned}$$

$$-\pi < \omega_0 < \pi \text{ or } 0 < \omega_0 < 2\pi$$



Typical Sequence—Basic Sequence

Sinusoidal

$$x[n] = A \cos(\omega_0 n + \phi) \quad \text{for all } n$$

Are these signal periodic?

$$x_1(t) = A \cos(0.1\pi t)$$

Typical Sequence—Basic Sequence

Sinusoidal

$$x[n] = A \cos(\omega_0 n + \phi) \quad \text{for all } n$$

Are these signal periodic?

$$x_1(t) = A \cos(0.1\pi t)$$

$$x_2[n] = A \cos(0.1\pi n)$$

Typical Sequence—Basic Sequence

Sinusoidal

$$x[n] = A \cos(\omega_0 n + \phi) \quad \text{for all } n$$

Are these signal periodic?

$$x_1(t) = A \cos(0.1\pi t)$$

$$x_2[n] = A \cos(0.1\pi n)$$

$$x_3(t) = A \cos(0.1t)$$

$$x_4[n] = A \cos(0.1n)$$

Typical Sequence—Basic Sequence

Sinusoidal

$$x[n] = A \cos(\omega_0 n + \phi) \quad \text{for all } n$$

Are these signal periodic?

$$x_1(t) = A \cos(0.1\pi t)$$

$$x_2[n] = A \cos(0.1\pi n)$$

$$x_3(t) = A \cos(0.1t)$$

$$x_4[n] = A \cos(0.1n)$$

$$x_5[n] = A \cos(0.1\pi n) + B \cos(0.5\pi n)$$

Typical Sequence—Basic Sequence

Sinusoidal

$$x[n] = A \cos(\omega_0 n + \phi) \quad \text{for all } n$$

Are these signal periodic?

$$x_1(t) = A \cos(0.1\pi t)$$

$$x_2[n] = A \cos(0.1\pi n)$$

$$x_3(t) = A \cos(0.1t)$$

$$x_4[n] = A \cos(0.1n)$$

$$x_5[n] = A \cos(0.1\pi n) + B \cos(0.5\pi n)$$

$$x[n + N] = x[n]$$

Typical Sequence—Basic Sequence

Sinusoidal

$$x[n] = A \cos(\omega_0 n + \phi) \quad \text{for all } n$$

Are these signal periodic?

$$x_1(t) = A \cos(0.1\pi t)$$

$$x_2[n] = A \cos(0.1\pi n)$$

$$x_3(t) = A \cos(0.1t)$$

$$x_4[n] = A \cos(0.1n)$$

$$x_5[n] = A \cos(0.1\pi n) + B \cos(0.5\pi n)$$

$$x[n + N] = x[n] \longrightarrow x[n + N] = A \cos(\omega_0(n + N) + \phi)$$

Typical Sequence—Basic Sequence

Sinusoidal

$$x[n] = A \cos(\omega_0 n + \phi) \quad \text{for all } n$$

Are these signal periodic?

$$x_1(t) = A \cos(0.1\pi t)$$

$$x_2[n] = A \cos(0.1\pi n)$$

$$x_3(t) = A \cos(0.1t)$$

$$x_4[n] = A \cos(0.1n)$$

$$x_5[n] = A \cos(0.1\pi n) + B \cos(0.5\pi n)$$

$$x[n + N] = x[n] \longrightarrow x[n + N] = A \cos(\omega_0(n + N) + \phi)$$

Periodic if $\omega_0 N = 2r\pi$

Typical Sequence—Basic Sequence

Sinusoidal

$$x[n] = A \cos(\omega_0 n + \phi) \quad \text{for all } n$$

Are these signal periodic?

$$x_1(t) = A \cos(0.1\pi t)$$

$$x_2[n] = A \cos(0.1\pi n)$$

$$x_3(t) = A \cos(0.1t)$$

$$x_4[n] = A \cos(0.1n)$$

$$x_5[n] = A \cos(0.1\pi n) + B \cos(0.5\pi n)$$

$$x[n + N] = x[n] \longrightarrow x[n + N] = A \cos(\omega_0(n + N) + \phi)$$

$$\text{Periodic if } \omega_0 N = 2r\pi \longrightarrow N = 2r\pi/\omega_0 \in \mathbb{Z}$$

Typical Sequence—Basic Sequence

Sinusoidal

$$x[n] = A \cos(\omega_0 n + \phi) \quad \text{for all } n$$

Are these signal periodic?

$$x_1(t) = A \cos(0.1\pi t)$$

$$x_2[n] = A \cos(0.1\pi n)$$

$$x_3(t) = A \cos(0.1t)$$

$$x_4[n] = A \cos(0.1n)$$

$$x_5[n] = A \cos(0.1\pi n) + B \cos(0.5\pi n)$$

$$x[n + N] = x[n] \longrightarrow x[n + N] = A \cos(\omega_0(n + N) + \phi)$$

$$\text{Periodic if } \omega_0 N = 2r\pi \longrightarrow N = 2r\pi/\omega_0 \in \mathbb{Z}$$

Not All Sequences are Periodic

Discrete Time Signals—Basic Sequence

Example: Combining Basic Sequence

$$u[n] = \delta[n] + \delta[n-1] + \dots$$

$$= \sum_{k=0}^{\infty} \delta[n-k]$$

Discrete Time Signals—Basic Sequence

Example: Combining Basic Sequence

$$u[n] = \delta[n] + \delta[n-1] + \dots$$

$$= \sum_{k=0}^{\infty} \delta[n-k]$$

$$u[n] = \sum_{k=-\infty}^n \delta[k]$$

Discrete Time Signals—Basic Sequence

Example: Combining Basic Sequence

$$u[n] = \delta[n] + \delta[n-1] + \dots$$

$$= \sum_{k=0}^{\infty} \delta[n-k]$$

$$u[n] = \sum_{k=-\infty}^n \delta[k]$$

$$x[n] = \begin{cases} A\alpha^n & n \geq 0 \\ 0 & n < 0 \end{cases}$$
$$= A\alpha^n u[n]$$

Discrete Time Signals—Basic Sequence

Example: Combining Basic Sequence

$$u[n] = \delta[n] + \delta[n-1] + \dots$$

$$= \sum_{k=0}^{\infty} \delta[n-k]$$

$$x[n] = \begin{cases} A\alpha^n & n \geq 0 \\ 0 & n < 0 \end{cases}$$

$$= A\alpha^n u[n]$$

$$u[n] = \sum_{k=-\infty}^n \delta[k]$$

Any sequence can be represented by impulse

$$x[n] = \sum_{k=-\infty}^{\infty} x[k] \delta[n-k]$$

Outline

- Time Domain Representation
- Typical Sequence & Sequence Representation
- Operation on Sequences
- Operation on Finite Length Sequences
- Sampling Process
- Correlation of Signals

Operation on Sequence

Elementary Operation

Product

$$w[n] = x[n] \cdot y[n]$$

Operation on Sequence

Elementary Operation

Product

$$w[n] = x[n] \cdot y[n]$$

Add

$$w[n] = x[n] + y[n]$$

Operation on Sequence

Elementary Operation

Product

$$w[n] = x[n] \cdot y[n]$$

Add

$$w[n] = x[n] + y[n]$$

Multiplication

$$w[n] = Ax[n]$$

Operation on Sequence

Elementary Operation

Product

$$w[n] = x[n] \cdot y[n]$$

Add

$$w[n] = x[n] + y[n]$$

Multiplication

$$w[n] = Ax[n]$$

Delay/Advance

$$w[n] = x[n - n_d]$$

Operation on Sequence

Elementary Operation

Product

$$w[n] = x[n] \cdot y[n]$$

Add

$$w[n] = x[n] + y[n]$$

Multiplication

$$w[n] = Ax[n]$$

Delay/Advance

$$w[n] = x[n - n_d]$$

Time Reversal

$$w[n] = x[-n]$$

Operation on Sequence

Convolution Sum

$$y[n] = x[n] \circledast h[n]$$

$$\begin{aligned} y(t) &= x(t) \circledast h(t) \\ &= \int x(\tau) y(t - \tau) d\tau \end{aligned}$$

Operation on Sequence

Convolution Sum

$$y[n] = x[n] \circledast h[n]$$

$$y(t) = x(t) \circledast h(t)$$

$$= \int x(\tau) y(t - \tau) d\tau$$

$$y[n] = x[n] \circledast h[n]$$

$$= \sum_{k=-\infty}^{\infty} x[k] h[n - k]$$

Operation on Sequence

Convolution Sum

$$y[n] = x[n] \circledast h[n]$$

$$y(t) = x(t) \circledast h(t)$$

$$= \int x(\tau) h(t - \tau) d\tau$$

$$y[n] = x[n] \circledast h[n]$$

$$= \sum_{k=-\infty}^{\infty} x[k] h[n - k]$$


- The computation is simply the sum of products
- Only delays, additions, and multiplications

Operation on Sequence

Computation of Convolutional Sum

Eg.1 $x[n] = h[n] = \delta[n] + \delta[n-1] + \delta[n-2]$

$$\begin{aligned} y[n] &= \sum_{k=-\infty}^{\infty} x[k] h[n-k] \\ &= \delta[n] + 2\delta[n-1] + 3\delta[n-2] + 2\delta[n-3] + \delta[n-4] \end{aligned}$$



A sequence of length M convolved with a sequence of length N , then resultant sequence is of length $M+N-1$

Operation on Sequence

Computation of Convolutional Sum

Eg.2

$$x[n] = (1/2)^{n-1} u[n-2]$$

$$h[n] = u[n+2]$$

$$\begin{aligned} y[n] &= \sum_{k=-\infty}^{\infty} x[k] h[n-k] \\ &= \sum_{k=-\infty}^{\infty} \left(\frac{1}{2}\right)^{k-2} u[k-2] u[(n-k)+2] \\ &= \sum_{k=2}^{n+2} \left(\frac{1}{2}\right)^{k-2} = \left[2 - \left(\frac{1}{2}\right)^n\right] u[n] \end{aligned}$$

Outline


- Time Domain Representation
- Typical Sequence & Sequence Representation
- Operation on Sequences
- Operation on Finite Length Sequences
- Sampling Process
- Correlation of Signals

Operation on Finite Length Sequence

Circular Shift

$$x_c[n - n_0] = x[\langle n - n_0 \rangle_N]$$

$$= \begin{cases} x[n - n_0] & \text{for } n_0 < n \leq N - 1 \\ x[N + n - n_0] & \text{for } 0 < n \leq n_0 \end{cases}$$




Outline

- Time Domain Representation
- Typical Sequence & Sequence Representation
- Operation on Sequences
- Operation on Finite Length Sequences
- Sampling Process
- Correlation of Signals


Sampling Process

$$x[n] = x_a(nT)$$



Sampling Process

$$x[n] = x_a(nT)$$



Example : $x_a(t) = A \cos(2\pi f_0 t)$


$$x[n] = A \cos(2\pi f_0 nT) = A \cos\left(\frac{2\pi \Omega_0}{\Omega_T} n\right)$$

$$nT = \frac{1}{F_T} n = \frac{2\pi}{\Omega_T} n$$

$$\Omega_0 = 2\pi f_0$$

Sampling Process

$$x[n] = x_a(nT)$$



Example : $x_a(t) = A \cos(2\pi f_0 t)$


$$x[n] = A \cos(2\pi f_0 nT) = A \cos\left(\frac{2\pi \Omega_0}{\Omega_T} n\right) \quad nT = \frac{1}{F_T} n = \frac{2\pi}{\Omega_T} n$$

$F_T = 1/T$ Sampling Frequency (**samples/sec**)

$$\Omega_0 = 2\pi f_0$$

Sampling Process

$$x[n] = x_a(nT)$$



Example : $x_a(t) = A \cos(2\pi f_0 t)$

$$x[n] = A \cos(2\pi f_0 nT) = A \cos\left(\frac{2\pi \Omega_0}{\Omega_T} n\right) \quad nT = \frac{1}{F_T} n = \frac{2\pi}{\Omega_T} n$$


$F_T = 1/T$ Sampling Frequency (**samples/sec**)

$$\Omega_0 = 2\pi f_0$$

$\Omega_T = 2\pi F_T$ Sampling **angular** Frequency(**radians/sample**)

Sampling Process

$$x[n] = x_a(nT)$$



Example : $x_a(t) = A \cos(2\pi f_0 t)$

$$x[n] = A \cos(2\pi f_0 nT) = A \cos\left(\frac{2\pi \Omega_0}{\Omega_T} n\right) \quad nT = \frac{1}{F_T} n = \frac{2\pi}{\Omega_T} n$$

$F_T = 1/T$ Sampling Frequency (**samples/sec**)


$$\Omega_0 = 2\pi f_0$$

$\Omega_T = 2\pi F_T$ Sampling **angular** Frequency(**radians/sample**)

$\omega_0 = \frac{2\pi f_0}{\Omega_T}$ **normalized** angular Frequency(**radians/sample**)

Sampling Process

$$x[n] = x_a(nT)$$



Example : $x_a(t) = A \cos(2\pi f_0 t)$

$$x[n] = A \cos(2\pi f_0 nT) = A \cos\left(\frac{2\pi \Omega_0}{\Omega_T} n\right) \quad nT = \frac{1}{F_T} n = \frac{2\pi}{\Omega_T} n$$

$$F_T = 1/T \text{ Sampling Frequency (samples/sec)} \quad \Omega_0 = 2\pi f_0$$

$$\Omega_T = 2\pi F_T \text{ Sampling angular Frequency (radians/sample)}$$

$$\omega_0 = \frac{2\pi f_0}{\Omega_T} \text{ normalized angular Frequency (radians/sample)}$$

$$-\pi < \omega_0 < \pi \text{ or } 0 < \omega_0 < 2\pi$$

Sampling Process

Example: $x_a(t) = A \cos(2\pi f_i t)$

$$f_0 = 3 \text{ Hz} \quad f_1 = 7 \text{ Hz}$$

$$f_2 = 13 \text{ Hz}$$

$$F_T = 10 \text{ samples/sec}$$

$$x_0[n] = \cos(0.6\pi n)$$

$$x_1[n] = \cos(1.4\pi n) = \cos((-2\pi + 1.4\pi)n) = \cos(-0.6\pi n)$$

$$x_2[n] = \cos(2.6\pi n) = \cos((2\pi + 0.6\pi)n) = \cos(0.6\pi n)$$


Sampling Process

Example: $x_a(t) = A \cos(2\pi f_i t)$

$$f_0 = 3\text{Hz} \quad f_1 = 7\text{Hz}$$

$$f_2 = 13\text{Hz}$$

$$F_T = 10 \text{ samples/sec}$$



$$x_0[n] = \cos(0.6\pi n)$$

$$x_1[n] = \cos(1.4\pi n) = \cos((-2\pi + 1.4\pi)n) = \cos(-0.6\pi n)$$

$$x_2[n] = \cos(2.6\pi n) = \cos((2\pi + 0.6\pi)n) = \cos(0.6\pi n)$$


Sampling Process

Example: $x_a(t) = A \cos(2\pi f_i t)$

$$f_0 = 3\text{Hz} \quad f_1 = 7\text{Hz}$$

$$f_2 = 13\text{Hz}$$

$$F_T = 10 \text{ samples/sec}$$



$$x_0[n] = \cos(0.6\pi n)$$

$$x_1[n] = \cos(1.4\pi n) = \cos((-2\pi + 1.4\pi)n) = \cos(-0.6\pi n)$$

$$x_2[n] = \cos(2.6\pi n) = \cos((2\pi + 0.6\pi)n) = \cos(0.6\pi n)$$

Alias

Outline

- Time Domain Representation
- Typical Sequence & Sequence Representation
- Operation on Sequences
- Operation on Finite Length Sequences
- Sampling Process
- Correlation of Signals

Correlation of Signals

Correlation

$$r_{xy}[l] = \sum_{n=-\infty}^{\infty} x[n]y[n-l]$$

Correlation of Signals

Correlation

$$r_{xy}[l] = \sum_{n=-\infty}^{\infty} x[n]y[n-l]$$

Auto Correlation

$$r_{xx}[l] = \sum_{n=-\infty}^{\infty} x[n]x[n-l]$$

Correlation of Signals

Correlation

$$r_{xy}[l] = \sum_{n=-\infty}^{\infty} x[n]y[n-l]$$

Auto Correlation

$$r_{xx}[l] = \sum_{n=-\infty}^{\infty} x[n]x[n-l]$$

Cross Correlation

$$\begin{aligned} r_{xy}[l] &= \sum_{n=-\infty}^{\infty} x[n]y[n-l] \\ &= r_{xy}[-l] \end{aligned}$$

Correlation of Signals

Correlation

$$r_{xy}[l] = \sum_{n=-\infty}^{\infty} x[n]y[n-l]$$

Auto Correlation

$$r_{xx}[l] = \sum_{n=-\infty}^{\infty} x[n]x[n-l]$$

Cross Correlation

$$\begin{aligned} r_{xy}[l] &= \sum_{n=-\infty}^{\infty} x[n]y[n-l] \\ &= r_{xy}[-l] \end{aligned}$$

$$r_{xy}[l] = \sum_{n=-\infty}^{\infty} x[n]y[n-l] = \sum_{n=-\infty}^{\infty} x[n]y[-(l-n)]$$

Correlation of Signals

Correlation

$$r_{xy}[l] = \sum_{n=-\infty}^{\infty} x[n]y[n-l]$$

Auto Correlation

$$r_{xx}[l] = \sum_{n=-\infty}^{\infty} x[n]x[n-l]$$

Cross Correlation

$$\begin{aligned} r_{xy}[l] &= \sum_{n=-\infty}^{\infty} x[n]y[n-l] \\ &= r_{xy}[-l] \end{aligned}$$

$$\begin{aligned} r_{xy}[l] &= \sum_{n=-\infty}^{\infty} x[n]y[n-l] = \sum_{n=-\infty}^{\infty} x[n]y[-(l-n)] \\ &= x[l] \circledast y[-l] \end{aligned}$$