

Equilibrium Operators, Variational Inequalities and Welfare:

What Is and Is Not a Theorem

Luciano Silva Alarco

Pontificia Universidad Católica del Perú

ORCID 0000-0002-3395-3129

Working draft v0.5 — research line EQO001 — 3 October 2026

Abstract

Research line EQO001 asked whether a formal relationship between equilibrium operators, variational inequalities and welfare criteria could yield results beyond the existing ones; its public record calls the formulation reviewed a combination of known components without an independent theorem. This note records that assessment. With equilibria the solutions of $VI(F, K)$ on a closed convex set K and welfare a C^1 functional on K , we prove the classical facts on which any such relationship rests: potentials, monotonicity, the equilibrium–welfare gap (Pigou, 4/3) and the first-order coincidence whose repair is marginal-cost pricing. The one small addition concerns the cone $\Lambda(x^*)$ of welfare weights for which a Nash equilibrium maximises $\sum_i \lambda_i u_i$: a closed convex cone inside a first-order cone $\Lambda_1(x^*)$ built from $F(x^*)$, the normal cone and the marginal externalities, with equality under joint concavity; (F, K) does not determine it. Exact examples show that $\Lambda(x^*)$ need not be polyhedral (three players; three commodities with affine costs) and that $\Lambda \neq \Lambda_1$ for two commodities. Deciding $\lambda \in \Lambda(x^*)$ is co-NP-complete when the number of players is part of the input, even for strongly monotone quadratic games (a standard reduction from box-constrained quadratic programming). We prove that the same holds for the per-commodity costs of multi-commodity Wardrop routing with affine costs, with explicit path sets or all paths of an acyclic network, when the number of commodities is part of the input, even with a unique equilibrium and a unit weight vector, and we identify two polynomial cases. Numerical checks on 400 random monotone games confirm equality in every tested direction of every concave instance, and brute-force checks on small graphs confirm both reductions. No new equilibrium concept is offered.

1 Introduction

The variational-inequality formulation of equilibrium is one of the unifying ideas of applied mathematics: a Wardrop traffic equilibrium [36], a Nash equilibrium of a concave game [24, 29] and a Walrasian or spatial price equilibrium can all be written as: find x^* in a closed convex set K with $\langle F(x^*), x - x^* \rangle \geq 0$ for all $x \in K$ [6, 10, 12, 14, 17, 23]. Welfare is an optimisation problem over the same K , and the temptation that motivated research line EQO001 is obvious: to look for a “formal relationship” between the inequality and the optimisation problem from which welfare statements about equilibria would follow.

The public record of the line is candid: the formulation reviewed combined known components without establishing an independent theorem, and the line was paused. This note discharges the line’s objective in the only way that seems honest to us: by writing down, with complete proofs where they are short, what is a theorem in this area, what is a definition in disguise, and what remains open. We add one small result on the set of welfare weights that make a given equilibrium optimal; Remark 5.2 states which parts of it are standard and which are, as far as we could check, only repackaged here.

Contributions and scope. Section 2 fixes the setting; Section 3 proves the classical results; Section 4 gives the first-order characterisation of welfare-optimal equilibria and marginal-cost pricing; Section 5 contains the welfare-weight cone (Theorem 5.1), the impossibility statement (Proposition 5.3), five exact examples, three of them negative (Examples 5.6, 5.7, 5.8), and the complexity of deciding membership, for games (Proposition 5.9, a standard reduction) and for multi-commodity routing (Theorem 5.12, with two tractable cases in Proposition 5.15); Sections 6–7 report the numerical checks, what can be claimed, limitations and the open questions. The longer proofs are in Appendix B. Everything is finite-dimensional and deterministic; every number in the text is produced by the scripts of Appendix A with fixed seeds.

2 Setting

Throughout, $K \subseteq \mathbb{R}^n$ is a nonempty closed convex set, $U \supseteq K$ an open set and $F: U \rightarrow \mathbb{R}^n$ a continuous map, the *equilibrium operator*. We write $\langle \cdot, \cdot \rangle$ for the Euclidean inner product.

Definition 2.1 (Variational inequality, normal cone). $\text{VI}(F, K)$ is the problem of finding $x^* \in K$ such that $\langle F(x^*), x - x^* \rangle \geq 0$ for all $x \in K$; its solution set is $\text{SOL}(F, K)$. The *normal cone* of K at $x \in K$ is $N_K(x) = \{v \in \mathbb{R}^n : \langle v, y - x \rangle \leq 0 \ \forall y \in K\}$, a closed convex cone, equal to $\{0\}$ when x is an interior point. By definition,

$$x^* \in \text{SOL}(F, K) \iff -F(x^*) \in N_K(x^*). \quad (1)$$

Definition 2.2 (Monotonicity). F is *monotone* on K if $\langle F(x) - F(y), x - y \rangle \geq 0$ for all $x, y \in K$; *strictly monotone* if the inequality is strict for $x \neq y$; μ -*strongly monotone* if $\langle F(x) - F(y), x - y \rangle \geq \mu \|x - y\|^2$ with $\mu > 0$; and L -*Lipschitz* if $\|F(x) - F(y)\| \leq L \|x - y\|$.

Definition 2.3 (Game). A *game* has N players, strategy sets $K_i \subseteq \mathbb{R}^{n_i}$ closed and convex, $K = K_1 \times \dots \times K_N$, and payoffs $u_i: U \rightarrow \mathbb{R}$ of class C^1 on an open $U \supseteq K$, with $u_i(\cdot, x_{-i})$ concave on K_i for every x_{-i} . A *Nash equilibrium* is $x^* \in K$ with $u_i(x^*) \geq u_i(y_i, x_{-i}^*)$ for all i and $y_i \in K_i$. The *pseudo-gradient* (equilibrium operator) is $F(x) = (-\nabla_{x_1} u_1(x), \dots, -\nabla_{x_N} u_N(x))$, where ∇_{x_i} is the gradient with respect to player i 's own variables. The *marginal externality* of player j on player $i \neq j$ at x is $\nabla_{x_j} u_i(x) \in \mathbb{R}^{n_j}$.

Lemma 2.4 (Nash equilibria are VI solutions). *In the setting of Definition 2.3, x^* is a Nash equilibrium if and only if $x^* \in \text{SOL}(F, K)$. [classical, 29; 10]*

Proof. For a concave differentiable function ϕ on a convex set C , z maximises ϕ over C iff $\langle \nabla \phi(z), y - z \rangle \leq 0$ for all $y \in C$: necessity because $t \mapsto \phi(z + t(y - z))$ is defined on $[0, 1]$ and has right derivative $\langle \nabla \phi(z), y - z \rangle$ at 0, which must be ≤ 0 ; sufficiency because $\phi(y) \leq \phi(z) + \langle \nabla \phi(z), y - z \rangle$. Apply this to $\phi = u_i(\cdot, x_{-i}^*)$ on K_i : x^* is Nash iff $\langle -F_i(x^*), y_i - x_i^* \rangle \leq 0$ for all i and $y_i \in K_i$. Summing over i gives $\langle F(x^*), y - x^* \rangle \geq 0$ for every $y \in K$; conversely, choosing $y = (y_i, x_{-i}^*)$ in the VI recovers the i -th inequality. $\square$

Definition 2.5 (Nonatomic routing). A network has edges E , commodities $k = 1, \dots, m$ with demands $d_k > 0$ and path sets P_k , $P = \bigsqcup_k P_k$. A path flow $f \in \mathbb{R}_+^P$ is feasible if $\sum_{p \in P_k} f_p = d_k$ for each k ; K is the set of feasible flows, a product of scaled simplices. Edge flows are $f_e = \sum_{p \ni e} f_p$, edge costs $c_e: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ are continuous and nondecreasing, path costs are $c_p(f) = \sum_{e \in p} c_e(f_e)$, and the equilibrium operator is $F(f) = (c_p(f))_{p \in P}$. A *Wardrop equilibrium* is a feasible f^* such that, for every k and $p, q \in P_k$ with $f_p^* > 0$, $c_p(f^*) \leq c_q(f^*)$. The *social cost* is $C(f) = \sum_p f_p c_p(f) = \sum_e f_e c_e(f_e)$, and welfare is $W = -C$.

Lemma 2.6 (Normal cone of a face of a simplex). *Let $\Delta = \{x \in \mathbb{R}_+^P : \sum_{p \in P_k} x_p = d_k, \ k = 1, \dots, m\}$ and $x \in \Delta$ with support $S = \{p : x_p > 0\}$. Then*

$$N_\Delta(x) = \{v : \exists \gamma \in \mathbb{R}^m \text{ with } v_p = \gamma_k \text{ for } p \in S \cap P_k, \ v_p \leq \gamma_k \text{ for } p \in P_k \setminus S\}.$$

This is the polar of the tangent cone $\{h : \sum_{p \in P_k} h_p = 0 \ \forall k, \ h_p \geq 0 \ (p \notin S)\}$, computed by Farkas' lemma [28].

Lemma 2.7 (Wardrop equilibria are VI solutions). *f^* is a Wardrop equilibrium iff $f^* \in \text{SOL}(F, K)$ with $F = (c_p)_p$. [classical, 33; 6]*

Proof. By (1) and Lemma 2.6, $f^* \in \text{SOL}(F, K)$ iff for each k there is γ_k with $c_p(f^*) = -\gamma_k$ on used paths of P_k and $c_p(f^*) \geq -\gamma_k$ on unused ones, which is Wardrop's condition. $\square$

Definition 2.8 (Welfare functionals). A *welfare functional* is a C^1 function $W : U \rightarrow \mathbb{R}$ to be maximised over K . For a game, the *weighted welfare* with weights $\lambda \in \mathbb{R}_+^N$ is $W_\lambda = \sum_i \lambda_i u_i$; $\lambda = \mathbf{1}$ is the utilitarian criterion. For routing, $W = -C$.

3 The classical results, with proofs

3.1 Potentials

A C^1 field F with symmetric Jacobian on an open convex U is a gradient (Poincaré's lemma): $P(x) = \int_0^1 \langle F(x_0 + t(x - x_0)), x - x_0 \rangle dt$ satisfies $\partial_i P(x) = \int_0^1 \frac{d}{dt} [t F_i(x_0 + t(x - x_0))] dt = F_i(x)$, by differentiation under the integral and $\partial_j F_i = \partial_i F_j$.

Theorem 3.1 (Equilibria of gradient operators). *Suppose $F = \nabla P$ on $U \supseteq K$ with $P \in C^1(U)$. Then*

$$\text{SOL}(F, K) = \{x \in K : \langle \nabla P(x), y - x \rangle \geq 0 \ \forall y \in K\},$$

the set of stationary points of P over K ; it contains $\arg \min_K P$, and equals it when P is convex on K . If P is strictly convex, $\text{SOL}(F, K)$ has at most one point. [classical: 2 for routing, 20 for potential games; proved here]

Proof. The first identity is Definition 2.1 with $F = \nabla P$. If x minimises P over K then, as in the proof of Lemma 2.4, the right derivative of $t \mapsto P(x + t(y - x))$ at 0 is nonnegative, which is the stationarity inequality. If P is convex and x is stationary, $P(y) \geq P(x) + \langle \nabla P(x), y - x \rangle \geq P(x)$ for all $y \in K$. If P is strictly convex its minimiser over the convex set K is unique. $\square$

Remark 3.2 (The three classical instances). (a) *Routing* [2]. For continuous c_e , $P(f) = \sum_e \int_0^{f_e} c_e(s) ds$ is C^1 with $\partial P / \partial f_p = \sum_{e \in p} c_e(f_e) = c_p(f)$ (fundamental theorem of calculus), convex when the c_e are nondecreasing; Wardrop equilibria minimise P , with unique edge flows when all c_e are strictly increasing. $P \neq C$: the potential integrates the cost, the social cost multiplies it (Section 3.3). (b) *Concave games* [29]. Lemma 2.4 and Theorem 3.4(a) give uniqueness under Rosen's diagonal strict concavity, which for unit weights is strict monotonicity of the pseudo-gradient. (c) *Potential games* [20]. A C^1 game has exact potential Π iff $F = -\nabla \Pi$; its equilibria are the stationary points of $-\Pi$ over K . The potential is bookkeeping, not welfare (Example 5.5(i)).

3.2 Monotonicity: uniqueness, existence and a solver

Lemma 3.3 (Projection). *Let Π_K be the Euclidean projection onto K . Then $z = \Pi_K(y)$ iff $z \in K$ and $\langle y - z, x - z \rangle \leq 0$ for all $x \in K$; Π_K is nonexpansive; and for every $\gamma > 0$, $x \in \text{SOL}(F, K)$ iff $x = \Pi_K(x - \gamma F(x))$. [classical, 17, Chapter I]*

Theorem 3.4 (Monotone operators). (a) *If F is strictly monotone on K , $\text{SOL}(F, K)$ has at most one point.* (b) *If F is μ -strongly monotone and L -Lipschitz on K , $\text{SOL}(F, K)$ is a singleton $\{x^*\}$, and for $0 < \gamma < 2\mu/L^2$ the iteration $x_{t+1} = \Pi_K(x_t - \gamma F(x_t))$ satisfies $\|x_{t+1} - x^*\| \leq q \|x_t - x^*\|$ with $q = \sqrt{1 - 2\gamma\mu + \gamma^2 L^2} < 1$; for $\gamma = \mu/L^2$, $q = \sqrt{1 - \mu^2/L^2}$. [classical, 10, 17; proved here]*

Proof. (a) If $x, y \in \text{SOL}(F, K)$ then $\langle F(x), y - x \rangle \geq 0$ and $\langle F(y), x - y \rangle \geq 0$; adding, $\langle F(x) - F(y), x - y \rangle \leq 0$, so $x = y$ by strict monotonicity. (b) Let $T(x) = \Pi_K(x - \gamma F(x))$. By nonexpansiveness, $\|Tx - Ty\|^2 \leq \|(x - y) - \gamma(F(x) - F(y))\|^2 = \|x - y\|^2 - 2\gamma\langle F(x) - F(y), x - y \rangle + \gamma^2\|F(x) - F(y)\|^2 \leq q^2\|x - y\|^2$, and $q < 1$ iff $\gamma < 2\mu/L^2$. Banach's fixed-point theorem on the complete set K gives a unique fixed point, which is the unique VI solution by Lemma 3.3, and the contraction estimate is the stated rate. $\square$

Remark 3.5. For merely continuous F on compact convex K , existence is the Hartman–Stampacchia theorem (14, Lemma 3.1), obtained from Brouwer's theorem applied to T ; we do not reproduce it. All solvers in Section 6 are the iteration of Theorem 3.4(b) with $\gamma = \mu/L^2$, so each computed equilibrium comes with a uniqueness proof and a residual certificate; the exception is Example 5.6 (E2c), where F is constant, $\gamma = 1$ is used and uniqueness follows from dominance.

3.3 Equilibrium is not welfare: Pigou and the price of anarchy

Example 3.6 (Pigou). Two parallel links, unit demand, $c_1 \equiv 1$ and $c_2(x) = x^d$, $d \geq 1$ [27]. Every user on link 2 pays $1 \leq c_1$, so $f^* = (0, 1)$ is a Wardrop equilibrium with $C(f^*) = 1$, and it is the only one: a feasible flow with $f_1 > 0$ has $f_2 < 1$, so its users on link 1 pay $1 > f_2^d$. The social cost of $(1 - x, x)$ is $1 - x + x^{d+1}$, minimised at $x = (d+1)^{-1/d}$ with value $1 - \frac{d}{d+1}(d+1)^{-1/d}$. For $d = 1$ the optimum is $(\frac{1}{2}, \frac{1}{2})$ with cost $\frac{3}{4}$ and ratio $\frac{4}{3}$; the ratio tends to infinity with d , since $(d+1)^{-1/d} \rightarrow 1$. The computed values are in `results/tables_traffic.md`.

Theorem 3.7 (Price of anarchy for affine costs). *If every $c_e(x) = a_e x + b_e$ with $a_e, b_e \geq 0$, then for every Wardrop equilibrium f and every feasible f' , $C(f) \leq \frac{4}{3}C(f')$. The constant is attained in Example 3.6 with $d = 1$. [classical, 31, see also 30; the proof below follows 5]*

Proof. By Lemma 2.7, $\sum_p c_p(f)(f'_p - f_p) \geq 0$; regrouping by edges, $C(f) = \sum_e c_e(f_e)f_e \leq \sum_e c_e(f_e)f'_e$. For each edge, $c_e(f_e)f'_e = c_e(f'_e)f'_e + a_e(f_e - f'_e)f'_e \leq c_e(f'_e)f'_e + \frac{1}{4}a_e f_e^2 \leq c_e(f'_e)f'_e + \frac{1}{4}c_e(f_e)f_e$, since $y \mapsto a_e(f_e - y)y$ is maximised at $y = f_e/2$ and $a_e f_e^2 \leq (a_e f_e + b_e)f_e$. Summing, $C(f) \leq C(f') + \frac{1}{4}C(f)$. $\square$

4 When an equilibrium is a welfare maximiser

Proposition 4.1 (Necessary and sufficient first-order conditions). *Let $x^* \in \text{SOL}(F, K)$ and let W be differentiable at x^* .*

- (a) *If $x^* \in \arg \max_K W$ then $\nabla W(x^*) \in N_K(x^*)$ and $-F(x^*) \in N_K(x^*)$. If W is concave on K , the converse holds: $x^* \in \arg \max_K W$ iff $\nabla W(x^*) \in N_K(x^*)$.*
- (b) *If x^* is an interior point of K , (a) forces $F(x^*) = 0 = \nabla W(x^*)$. If $N_K(x^*) = \mathbb{R}_+ \nu$ is a ray and $F(x^*) \neq 0$, (a) forces $\nabla W(x^*) = -\theta F(x^*)$ for some $\theta \geq 0$.*
- (c) (Simplices.) *If K is a product of simplices (Lemma 2.6), both $F(x^*)$ and $\nabla W(x^*)$ are constant on the support of x^* within each commodity (hence parallel there when both are non-zero), and each satisfies Wardrop's principle with respect to its own coordinates. On $K = \{x \geq 0 : Ax = b\}$ the same holds with “constant on the support” replaced by “equal on the support to an element of the row space of A ”.*
- (d) *If F is strictly monotone, W is strictly concave and both $\text{SOL}(F, K)$ and $\arg \max_K W$ are nonempty, then $\text{SOL}(F, K) = \arg \max_K W$ iff the unique equilibrium satisfies $\nabla W(x^*) \in N_K(x^*)$; otherwise the two sets are disjoint.*

[proved here; elementary]

Proof. (a) The first-order necessary condition for a maximum of a differentiable function over a convex set (proof of Lemma 2.4) is $\langle \nabla W(x^*), y - x^* \rangle \leq 0$ for all $y \in K$, i.e. $\nabla W(x^*) \in N_K(x^*)$; the second membership is (1); sufficiency under concavity is the same argument as in Theorem 3.1. (b) For interior x^* , $N_K(x^*) = \{0\}$. If $N_K(x^*) = \mathbb{R}_+\nu$, write $-F(x^*) = \alpha\nu$ and $\nabla W(x^*) = \beta\nu$ with $\alpha, \beta \geq 0$; $F(x^*) \neq 0$ gives $\alpha > 0$ and $\theta = \beta/\alpha$. (c) Lemma 2.6 applied to $-F(x^*)$ and to $\nabla W(x^*)$; for $\{x \geq 0 : Ax = b\}$ the polar of the tangent cone is $\{A^\top w - v : v \geq 0, v_S = 0\}$ by Farkas' lemma. (d) Both sets are singletons by Theorem 3.4(a) and strict concavity; apply (a). $\square$

Proposition 4.1 is a statement about first-order conditions and nothing more: it says that an equilibrium can be a welfare optimum only if the two problems happen to share their KKT point, and it gives no mechanism producing that coincidence. The one mechanism that always works is to change the operator.

Corollary 4.2 (Marginal-cost pricing and the toll set). *Let $C \in C^1(U)$ be a cost to be minimised over K and define the toll field $\tau = \nabla C - F$. Then*

$$\text{SOL}(F + \tau, K) = \{x \in K : -\nabla C(x) \in N_K(x)\} \supseteq \arg \min_K C,$$

with equality when C is convex on K , and at most one point when C is strictly convex; the tolled operator is a gradient field with potential C . For separable routing costs with $c_e \in C^1$, $\tau_p(f) = \sum_{e \in p} f_e c'_e(f_e)$, the Pigou–Knight–Beckmann marginal-cost toll. More generally, for a target $\bar{x} \in K$ the set of constant toll vectors $t \in \mathbb{R}^n$ with $\bar{x} \in \text{SOL}(F + t, K)$ is the translated cone $-F(\bar{x}) - N_K(\bar{x})$, polyhedral when K is; choosing in it the toll closest to a reference vector is an inverse optimisation problem in the sense of Ahuja and Orlin [1], and for routing it is the toll-set description of Hearn and Ramana [15]. [classical, 2, 15, 18, 27; proved here]

Proof. $F + \tau = \nabla C$, so the first statement is Theorem 3.1 with $P = C$. For routing, $\partial C / \partial f_p = \sum_{e \in p} (c_e(f_e) + f_e c'_e(f_e))$, whence τ_p . Finally $\bar{x} \in \text{SOL}(F + t, K)$ iff $-F(\bar{x}) - t \in N_K(\bar{x})$ by (1). $\square$

5 The cone of welfare weights

Fix a game as in Definition 2.3 and a Nash equilibrium x^* . Write $u = (u_1, \dots, u_N) : K \rightarrow \mathbb{R}^N$ and $G(x^*) = Du(x^*) \in \mathbb{R}^{N \times n}$ for the Jacobian of the payoff vector, whose i -th row is $\nabla u_i(x^*)^\top$ (all variables, own and others'). Define

$$\Lambda(x^*) = \{\lambda \in \mathbb{R}_+^N : x^* \in \arg \max_K W_\lambda\}, \quad \Lambda_1(x^*) = \{\lambda \in \mathbb{R}_+^N : G(x^*)^\top \lambda \in N_K(x^*)\}.$$

For a set $D \subseteq \mathbb{R}^N$, $D^\circ = \{\lambda : \langle \lambda, d \rangle \leq 0 \ \forall d \in D\}$ is its polar cone.

Theorem 5.1 (Welfare-weight cone). (a) $\Lambda(x^*) = \mathbb{R}_+^N \cap \{u(x) - u(x^*) : x \in K\}^\circ$. In particular $\Lambda(x^*)$ is a closed convex cone.

(b) $\Lambda(x^*) \subseteq \Lambda_1(x^*)$, and $\Lambda_1(x^*)$ is a closed convex cone, polyhedral whenever K is polyhedral.

(c) If every u_i is concave on K (jointly in all variables), then $\Lambda(x^*) = \Lambda_1(x^*)$. More generally, equality holds whenever W_λ is concave on K for every $\lambda \in \Lambda_1(x^*)$.

(d) (Block form.) $\lambda \in \Lambda_1(x^*)$ iff for every player j

$$-\lambda_j F_j(x^*) + \sum_{i \neq j} \lambda_i \nabla_{x_j} u_i(x^*) \in N_{K_j}(x_j^*). \quad (2)$$

Since $-F_j(x^*) \in N_{K_j}(x_j^*)$ already, a sufficient condition is that the λ -weighted externality $\sum_{i \neq j} \lambda_i \nabla_{x_j} u_i(x^*)$ lies in $N_{K_j}(x_j^*)$ for every j ; when x_j^* is interior to K_j , (2) says exactly that this weighted externality vanishes.

(e) $\Lambda(x^*) \neq \{0\}$ implies that x^* is weakly Pareto efficient in K (no $x \in K$ with $u_i(x) > u_i(x^*)$ for all i); under the concavity of (c), the converse holds. Without any concavity, if x^* is locally weakly Pareto efficient then $\Lambda_1(x^*) \neq \{0\}$.

[proved here; (a) and (e) are the standard scalarisation argument of multi-objective optimisation [9, 11], and $\Lambda(x^*)$ is the cell at $u(x^*)$ of the weight-set decomposition studied for linear problems by Benson and Sun [3]]

Proof. (a) $\lambda \in \Lambda(x^*)$ iff $\lambda \geq 0$ and $\langle \lambda, u(x) - u(x^*) \rangle \leq 0$ for all $x \in K$, which is the stated intersection; polar cones and the orthant are closed convex cones. (b) If x^* maximises the differentiable W_λ over K , the first-order condition gives $\nabla W_\lambda(x^*) = \sum_i \lambda_i \nabla u_i(x^*) = G(x^*)^\top \lambda \in N_K(x^*)$. $\Lambda_1(x^*)$ is the preimage of the closed convex cone $N_K(x^*)$ under a linear map, intersected with $\mathbb{R}_+^N$; when K is polyhedral so is $N_K(x^*)$, hence so is $\Lambda_1(x^*)$. (c) For $\lambda \geq 0$, W_λ is concave when the u_i are, and the first-order condition is then sufficient (Proposition 4.1(a)); the general statement is the same argument. (d) For a product set, $v \in N_K(x^*)$ iff $v_j \in N_{K_j}(x_j^*)$ for each j : taking $y \in K$ that differs from x^* in block j only shows necessity, and summing the block inequalities shows sufficiency. Block j of $G(x^*)^\top \lambda$ is $\sum_i \lambda_i \nabla_{x_j} u_i(x^*) = \lambda_j \nabla_{x_j} u_j(x^*) + \sum_{i \neq j} \lambda_i \nabla_{x_j} u_i(x^*)$ and $\nabla_{x_j} u_j(x^*) = -F_j(x^*)$. The sufficient condition follows because $N_{K_j}(x_j^*)$ is a convex cone containing $-F_j(x^*)$ by Lemma 2.4; for interior x_j^* , $N_{K_j}(x_j^*) = \{0\}$ forces $F_j(x^*) = 0$ and then (2) reads $\sum_{i \neq j} \lambda_i \nabla_{x_j} u_i(x^*) = 0$. (e) If $\lambda \in \Lambda(x^*) \setminus \{0\}$ and $u(x) > u(x^*)$ componentwise, then $\langle \lambda, u(x) \rangle > \langle \lambda, u(x^*) \rangle$, a contradiction. Conversely, under concavity the set $A = u(K) - \mathbb{R}_+^N$ is convex; weak efficiency means $u(x^*) + \varepsilon \mathbf{1} \notin A$ for every $\varepsilon > 0$, so $u(x^*)$ is a boundary point of A and a supporting hyperplane gives $\lambda \neq 0$ with $\langle \lambda, a \rangle \leq \langle \lambda, u(x^*) \rangle$ for all $a \in A$; since $u(x^*) - \mathbb{R}_+^N \subseteq A$, $\lambda \geq 0$, and $a = u(x)$ gives $\lambda \in \Lambda(x^*)$. Finally, if $\Lambda_1(x^*) = \{0\}$, the convex cone $\{sG(x^*)(x - x^*) : x \in K, s \geq 0\}$ meets the open orthant: otherwise a separating hyperplane (the Gordan–Motzkin alternative) gives $\lambda \geq 0$, $\lambda \neq 0$, with $\langle \lambda, G(x^*)(x - x^*) \rangle \leq 0$ on K , i.e. $\lambda \in \Lambda_1(x^*)$; and $G(x^*)(x - x^*) > 0$ gives $u(x^* + \tau(x - x^*)) > u(x^*)$ for small $\tau > 0$. $\square$

Remark 5.2 (What is standard and what is not). The scalarisation in (a) and (e) is textbook, and the set of weights under which a given point maximises the weighted sum is a standard object of multi-objective optimisation (the weight-set decomposition, polyhedral in the linear case [3]); Proposition 5.3 repackages the Nikaidô–Isoda gap function and is an instance of best-response equivalence [21], and Proposition 5.9 is a standard reduction (see its status). What we did not find in the literature we consulted, and claim only as packaging, is the reading of (2): the cone is cut out by the equilibrium face (through $N_{K_j}(x_j^*)$ and $F_j(x^*)$, both determined by (F, K)) and by the marginal externalities $\nabla_{x_j} u_i(x^*)$, which are *not*. We did not find Theorem 5.12 in the work we consulted on the complexity of multiclass and multi-commodity equilibria, on toll design [15] and on Stackelberg routing; its reduction has to respect the special form of the per-commodity routing costs, to which Proposition 5.9 does not apply.

Proposition 5.3 ((F, K) does not determine welfare). *Let the u_i be continuous on $K = \prod K_i$ with K_i compact, and let x^* be a Nash equilibrium. Define $v_i(x_{-i}) = \max_{y_i \in K_i} u_i(y_i, x_{-i})$ and $u'_i = u_i - v_i$. Then the game (u'_i) has the same best-response correspondences and the same Nash equilibria as (u_i) ; wherever $\nabla_{x_i} u_i$ exists, $\nabla_{x_i} u'_i = \nabla_{x_i} u_i$, so the two games have the same equilibrium operator; and x^* maximises every $W'_\lambda = \sum_i \lambda_i u'_i$, $\lambda \in \mathbb{R}_+^N$, over K : $\Lambda'(x^*) = \mathbb{R}_+^N$. Consequently welfare optimality of an equilibrium is not a property of (F, K) : every pair (F, K) that arises from a game is compatible with $\Lambda(x^*) = \mathbb{R}_+^N$; if $N \geq 2$ and some K_{-i} is not a singleton, it is also compatible with $e_i \notin \Lambda(x^*)$ (and the pair of Example 5.5(i) with $\Lambda(x^*) = \{0\}$). [proved here; $\sum_i (v_i - u_i)$ is the Nikaidô–Isoda gap function [25], and $u \mapsto u'$ a best-response equivalence [21]]*

Proof. v_i does not depend on x_i , so subtracting it changes neither $\arg \max_{y_i} u_i(y_i, x_{-i})$ nor $\nabla_{x_i} u_i$. By construction $u'_i \leq 0$ on K , and $u'_i(x^*) = 0$ because x_i^* is a best response to x_{-i}^* ; hence $W'_\lambda \leq 0 = W'_\lambda(x^*)$ for every $\lambda \geq 0$. For the second claim, add to u_i the term $c\langle d, x_{-i} - x_{-i}^* \rangle$, with d such that $\delta = \langle d, x_{-i} - x_{-i}^* \rangle > 0$ for some $x_{-i} \in K_{-i}$: neither F nor the equilibria

change, the game stays in the class of Definition 2.3, and $x = (x_i^*, x_{-i})$ beats x^* for W_{e_i} once $c\delta > u_i(x^*) - u_i(x)$. $\square$

The modified game need not be C^1 , so $\Lambda'(x^*)$ is understood in the exact sense of Theorem 5.1(a). When best responses are unique, Danskin's theorem [7] gives $v_i \in C^1$ with $\nabla v_i(x_{-i}) = \nabla_{x_{-i}} u_i(b_i(x_{-i}), x_{-i})$; the modified game is then in the class of Definition 2.3 and has *zero marginal externalities at x^** , so $\Lambda'_1(x^*) = \mathbb{R}_+^N$ by Theorem 5.1(d), the first-order reason for the proposition.

Example 5.4 (Two players, exact cone). Let $K = [0, 1]^2$, $\alpha > 1$, $\beta_1, \beta_2 \geq 0$, and

$$u_1(x) = \alpha x_1 - \frac{1}{2}x_1^2 - \beta_1 x_2, \quad u_2(x) = \alpha x_2 - \frac{1}{2}x_2^2 - \beta_2 x_1.$$

Each player's action harms the other linearly. The equilibrium operator is $F(x) = (x_1 - \alpha, x_2 - \alpha)$, with Jacobian I , so F is 1-strongly monotone and 1-Lipschitz and does not depend on β ; the unique equilibrium is $x^* = (1, 1)$, at which $N_K(x^*) = \mathbb{R}_+^2$ and $F(x^*) = (1 - \alpha)\mathbf{1}$. Here $G(x^*) = \begin{pmatrix} \alpha-1 & -\beta_1 \\ -\beta_2 & \alpha-1 \end{pmatrix}$, so by (2)

$$\Lambda_1(x^*) = \{\lambda \geq 0 : (\alpha - 1)\lambda_1 \geq \beta_2 \lambda_2, (\alpha - 1)\lambda_2 \geq \beta_1 \lambda_1\}.$$

Both payoffs are jointly concave (Hessians $-\text{diag}(1, 0)$ and $-\text{diag}(0, 1)$), so $\Lambda(x^*) = \Lambda_1(x^*)$ by Theorem 5.1(c). Thus the cone is a wedge, non-trivial iff $\beta_1 \beta_2 \leq (\alpha - 1)^2$ and two-dimensional iff the inequality is strict; for $\alpha = 2$, $\beta_1 = \beta_2 = \frac{1}{2}$ it is $\{\frac{1}{2}\lambda_2 \leq \lambda_1 \leq 2\lambda_2\}$: the corner equilibrium is welfare-optimal exactly when neither player is weighted more than twice the other. Games with different β have the same F and different cones: Proposition 5.3 in miniature.

Example 5.5 (Cournot duopoly). Let $u_i(q) = q_i(a - q_1 - q_2)$ on $K = [0, 1]^2$: each u_i is concave in q_i and $F(q) = (2q_1 + q_2 - a, q_1 + 2q_2 - a)$ is strongly monotone. (i) For $a = 1$ the equilibrium $q^* = (\frac{1}{3}, \frac{1}{3})$ is interior and $\partial u_1 / \partial q_2 = -q_1^* \neq 0$, so $\Lambda(q^*) = \{0\}$ by Theorem 5.1(d); with (e), this is the first-order mechanism behind the inefficiency theorem of Dubey [8] for interior equilibria. Collusion at $(\frac{1}{4}, \frac{1}{4})$ gives each firm $\frac{1}{8} > \frac{1}{9}$. The exact potential $\Pi(q) = a(q_1 + q_2) - q_1^2 - q_2^2 - q_1 q_2$ is maximised at the equilibrium: a potential is not a welfare functional. (ii) For $a = 4$ the equilibrium is the corner $(1, 1)$ and (2) gives $\Lambda_1(q^*) = \{\lambda_1 = \lambda_2 \geq 0\}$; the payoffs are not jointly concave, but $W_{(1,1)} = (q_1 + q_2)(4 - q_1 - q_2)$ is, so $\Lambda(q^*) = \Lambda_1(q^*)$: concavity is sufficient, not necessary.

Example 5.6 (Three players: a non-polyhedral cone). Let $K = [0, 1]^3$ and

$$u_1 = x_1, \quad u_2 = x_2, \quad u_3 = x_3 - \frac{1}{2}x_1^2 + \frac{1}{4}x_1 + \frac{1}{2}x_1 x_2 - \frac{1}{2}x_2.$$

Each payoff is linear, hence concave, in the player's own variable, so the game is in the class of Definition 2.3; $F \equiv (-1, -1, -1)$, $x^* = (1, 1, 1)$ is the unique Nash equilibrium and $N_K(x^*) = \mathbb{R}_+^3$. Since $\nabla u_3(x^*) = (-\frac{1}{4}, 0, 1)$, $G(x^*)^\top \lambda = (\lambda_1 - \frac{1}{4}\lambda_3, \lambda_2, \lambda_3)$ and $\Lambda_1(x^*) = \{\lambda \geq 0 : \lambda_1 \geq \frac{1}{4}\lambda_3\}$. We compute $\Lambda(x^*)$ exactly. On $\{\lambda_3 = 0\}$, $W_\lambda = \lambda_1 x_1 + \lambda_2 x_2$ is maximised at x^* , so $\mathbb{R}_+^2 \times \{0\} \subseteq \Lambda(x^*)$; as $\Lambda(x^*)$ is a cone it suffices to fix $\lambda_3 = 1$. Then W_λ is increasing in x_3 , so $x_3 = 1$, and affine in x_2 , so the maximum over x_2 is attained at $x_2 \in \{0, 1\}$. For $x_2 = 1$, $W_\lambda = \frac{1}{2} + \lambda_2 + (\lambda_1 + \frac{3}{4})x_1 - \frac{1}{2}x_1^2$ is maximised on $[0, 1]$ at $x_1 = 1$ iff $\lambda_1 \geq \frac{1}{4}$, with value $W_\lambda(x^*) = \lambda_1 + \lambda_2 + \frac{3}{4}$. For $x_2 = 0$, $W_\lambda = 1 + (\lambda_1 + \frac{1}{4})x_1 - \frac{1}{2}x_1^2$ has maximum $1 + \frac{1}{2}(\lambda_1 + \frac{1}{4})^2$ if $\lambda_1 \leq \frac{3}{4}$ (at $x_1 = \lambda_1 + \frac{1}{4}$) and $\lambda_1 + \frac{3}{4} \leq W_\lambda(x^*)$ otherwise. Hence $\lambda \in \Lambda(x^*)$ iff $\lambda_1 \geq \frac{1}{4}$ and $\lambda_1 + \lambda_2 + \frac{3}{4} \geq 1 + \frac{1}{2}(\lambda_1 + \frac{1}{4})^2$, i.e.

$$\Lambda(x^*) \cap \{\lambda_3 = 1\} = \{\lambda_1 \geq \frac{1}{4}, \lambda_2 \geq \frac{1}{2}(\frac{3}{4} - \lambda_1)^2\}.$$

Its boundary contains the parabolic arc $\lambda_2 = \frac{1}{2}(\frac{3}{4} - \lambda_1)^2$, $\frac{1}{4} \leq \lambda_1 \leq \frac{3}{4}$, which no finite union of segments contains (a line meets a parabola in at most two points); since the slice of a polyhedral

cone by a hyperplane is a polyhedron, $\Lambda(x^*)$ is *not polyhedral*, and $\Lambda(x^*) \subsetneq \Lambda_1(x^*)$ with a curved boundary. For instance $\lambda = (0.3, 0, 1) \in \Lambda_1 \setminus \Lambda$: the point $(0.55, 0, 1)$ improves W_λ by 0.10125. The degeneracy $F \equiv \text{const}$ is inessential. Subtract $\frac{\varepsilon}{2}x_i^2$ from each u_i , $0 < \varepsilon < 1$: $F(x) = \varepsilon x - 1$ is ε -strongly monotone, x^* is unchanged, $\Lambda_1 \cap \{\lambda_3 = 1\} = \{\lambda_1 \geq \frac{1}{4(1-\varepsilon)}\}$, and for $\frac{1}{4(1-\varepsilon)} \leq \lambda_1 \leq \frac{3}{4(1-\varepsilon)}$ a point of the slice $\lambda_3 = 1$ lies in $\Lambda(x^*)$ iff

$$\lambda_2 \geq \varphi(\lambda_1) = \frac{(\lambda_1 + \frac{1}{4})^2 / (2(1 + \varepsilon\lambda_1)) + \frac{1}{4}}{1 - \varepsilon/2} - \lambda_1, \quad \varphi''(\lambda_1) = \frac{(1 - \varepsilon/4)^2}{(1 - \varepsilon/2)(1 + \varepsilon\lambda_1)^3} > 0,$$

which reduces to the parabola for $\varepsilon = 0$; φ decreases to $\varphi(\frac{3}{4(1-\varepsilon)}) = 0$ with zero slope, so the boundary is tangent to the axis as for $\varepsilon = 0$, and $\Lambda(x^*)$ is not polyhedral for a strongly monotone F either (proof in Appendix B). [proved here; verified in E2c]

Example 5.7 (Two commodities: $\Lambda \neq \Lambda_1$ in routing). Theorem 5.1(a),(b) use neither the Nash property nor the product structure: they hold for any finite family of C^1 functions u_k on K and any $x^* \in K$. Apply them to the costs $C_k(f) = \sum_{p \in P_k} f_p c_p(f)$ of the commodities, $u_k = -C_k$, at a Wardrop equilibrium f^* . Two commodities with unit demands: commodity 1 chooses between a private edge of cost $f + 2$ and a shared edge of cost f , commodity 2 between a private edge of cost $f/4$ and the same shared edge. With $y = (f_B, f_D) \in [0, 1]^2$ the flows on the shared edge (path flows are determined by them),

$$C_1(y) = (1 - f_B)(3 - f_B) + f_B(f_B + f_D), \quad C_2(y) = \frac{1}{4}(1 - f_D)^2 + f_D(f_B + f_D),$$

both with indefinite Hessians, so (c) is unavailable. The equilibrium is $f^* = (1, 0)$ (shared cost $1 \leq 2$ for commodity 1, private cost $\frac{1}{4} \leq 1$ for commodity 2), unique because the reduced operator $y \mapsto (c_B - c_A, c_D - c_C)$ has Jacobian $\begin{pmatrix} 2 & 1 \\ 1 & 5/4 \end{pmatrix} \succ 0$. In the coordinates y (the cones are invariant under this affine reparametrisation of K), $N(f^*) = \mathbb{R}_+ \times \mathbb{R}_-$, $\nabla C_1(f^*) = (0, 1)$ and $\nabla C_2(f^*) = (0, \frac{1}{2})$, so $\Lambda_1(f^*) = \mathbb{R}_+^2$. Moreover (Appendix B)

$$\Lambda(f^*) = \{\lambda \geq 0 : \lambda_2 \leq 40\lambda_1\} \subsetneq \Lambda_1(f^*) = \mathbb{R}_+^2.$$

For $\lambda = (0, 1)$, $y = (0, \frac{1}{5})$ lowers C_2 from $\frac{1}{4}$ to $\frac{1}{5}$ (while C_1 rises from 1 to 3). [proved here; verified in E5]

Example 5.8 (Three commodities: a non-polyhedral commodity cone). Three commodities with unit demand; commodity k chooses between a private edge and one shared edge of cost f ; the private costs are $f/4$, $f/4$ and the constant 2. With $y \in [0, 1]^3$ the flows on the shared edge and $S = y_1 + y_2 + y_3$,

$$C_k(y) = \frac{1}{4}(1 - y_k)^2 + y_k S \quad (k = 1, 2), \quad C_3(y) = 2(1 - y_3) + y_3 S.$$

The Wardrop equilibrium is $y^* = (0, 0, 1)$ (shared cost $1 \leq 2$ for commodity 3, private cost $\frac{1}{4} \leq 1$ for commodities 1, 2), unique because the reduced operator $y \mapsto (S - \frac{1}{4}(1 - y_1), S - \frac{1}{4}(1 - y_2), S - 2)$ has Jacobian $\text{diag}(\frac{1}{4}, \frac{1}{4}, 0) + \mathbf{1}\mathbf{1}^\top \succ 0$. At y^* , in reduced coordinates, $N = \mathbb{R}_- \times \mathbb{R}_- \times \mathbb{R}_+$, $\nabla C_1 = (\frac{1}{2}, 0, 0)$, $\nabla C_2 = (0, \frac{1}{2}, 0)$ and $\nabla C_3 = (1, 1, 0)$, so $\Lambda_1(y^*) = \mathbb{R}_+^3$. We claim

$$\Lambda(y^*) = \{\lambda \geq 0 : \lambda_3 + m(\lambda_1, \lambda_2) \geq 0\}, \quad m = \min_{[0, 1]^2} P,$$

$P(u, w) = \frac{\lambda_1}{4}(u^2 - 2u) + \frac{\lambda_2}{4}(w^2 - 2w) + (u + w)(\lambda_1 u + \lambda_2 w)$ (proof in Appendix B). On the slice $\lambda_2 = 1$, for $\frac{2}{3} < \lambda_1 < \frac{3}{2}$, P is strictly convex with minimiser $(3\lambda_1 - 2, \lambda_1(3 - 2\lambda_1)) / ((4 - \lambda_1)(4\lambda_1 - 1))$ in the open square, and $m = -\lambda_1(\lambda_1 + 1) / (4(4 - \lambda_1)(4\lambda_1 - 1))$. So the boundary of $\Lambda(y^*) \cap \{\lambda_2 = 1\}$ over that interval is the graph of the rational, non-affine function $\psi(\lambda_1) = \lambda_1(\lambda_1 + 1) / (4(4 - \lambda_1)(4\lambda_1 - 1))$: $\Lambda(y^*)$ is *not polyhedral*, and $\Lambda \subsetneq \Lambda_1 = \mathbb{R}_+^3$. (Outside $(\frac{2}{3}, \frac{3}{2})$ the minimiser of P lies on an edge and the boundary is linear, $\lambda_3 = \frac{1}{20}$ for $\lambda_1 \leq \frac{2}{3}$ and $\lambda_3 = \lambda_1/20$ for $\lambda_1 \geq \frac{3}{2}$; checked numerically, not used.) For instance $\psi(1) = \frac{1}{18}$, and for $\lambda = (1, 1, \frac{1}{20})$ the flow $y = (\frac{1}{9}, \frac{1}{9}, 0)$ lowers Φ_λ by $\frac{1}{180}$. [proved here; verified in E7]

Proposition 5.9 (Deciding membership is co-NP-complete). *For games whose players have actions in boxes and quadratic payoffs with rational coefficients, given with a rational Nash equilibrium x^* (and, for the first problem, a rational $\lambda \geq 0$), deciding whether $\lambda \in \Lambda(x^*)$ is co-NP-complete, and so is deciding whether $\Lambda(x^*) = \Lambda_1(x^*)$, when the number of variables (players, or the dimension of their actions) is part of the input. Hardness holds with scalar actions in $[0, 1]$, and also with $N = 2$; and with F strongly monotone and independent of the instance, x^* a vertex in strictly dominant strategies, $\Lambda_1(x^*) = \mathbb{R}_+^N$ and, for membership, λ a unit vector. [proved here; in substance the standard MAX-CUT encoding of box-constrained quadratic programming, transported to games by Proposition 5.3 (the equilibrium operator of the reduction does not depend on the instance); proof in Appendix B; checked in E6]*

The hardness lives entirely in the externalities, as Proposition 5.3 predicts. With $N = 2$, $\Lambda(x^*)$ is polyhedral and membership is still hard, while Example 5.6 is non-polyhedral with $N = 3$ fixed and decidable in constant time: hardness and non-polyhedrality are independent phenomena.

Remark 5.10 (Descriptions of $\Lambda(x^*)$). For polynomial payoffs on a polyhedral K , $\Lambda(x^*)$ is semialgebraic (Tarski–Seidenberg, 4), polyhedral for $N = 2$ (as is every closed convex cone of $\mathbb{R}^2$) and not in general for $N \geq 3$ (Example 5.6). By Proposition 5.9, unless $P = NP$ there is no explicit description of $\Lambda(x^*)$ by polynomially many polynomial inequalities of polynomial size computable in polynomial time, so a characterisation “in terms of the active-face pattern and the Hessians” has no efficiently checkable answer in that generality. With scalar actions and N fixed, membership is polynomial (3^N faces); supermodular W_λ (as in Example 5.6) is a candidate tractable subclass that we do not study.

Routing: the commodity cone is hard as well

Throughout this subsection the network is as in Definition 2.5, with affine costs $c_e(x) = a_e x + b_e$, $a_e, b_e \geq 0$ rational, and the path sets P_k listed explicitly as part of the input (each path a set of edges). As in Example 5.7, $C_k(f) = \sum_{p \in P_k} f_p c_p(f) = \sum_e c_e(f_e) f_e^k$, where f_e^k is the flow of commodity k on e ; $\Lambda(f^*)$ and $\Lambda_1(f^*)$ are defined with $u_k = -C_k$, and we write $\Phi_\lambda = \sum_k \lambda_k C_k$, so that $\lambda \in \Lambda(f^*)$ iff $f^* \in \arg \min_K \Phi_\lambda$. With $g_e = \sum_k \lambda_k f_e^k$ the λ -weighted flow on e ,

$$\Phi_\lambda(f) = \sum_e (a_e f_e + b_e) g_e. \quad (3)$$

Lemma 5.11 (Zero-weight commodities enter affinely). *Let $Q = \{k : \lambda_k > 0\}$ and R its complement. For fixed flows of the commodities in Q , Φ_λ is affine in the flows of the commodities in R , jointly; hence $f^R \mapsto \min_{f^Q} \Phi_\lambda(f^Q, f^R)$ is concave and Φ_λ has a minimiser over K in which every commodity of R uses a single path. If moreover $\lambda = \mu \mathbf{1}_Q$ for some $\mu > 0$, then Φ_λ is convex in f^Q for fixed f^R , so $\min_K \Phi_\lambda$ is the minimum, over the pure profiles of the commodities of R that share an edge with a path of Q , of a convex quadratic programme. [proved here, Appendix B; elementary]*

The lemma says where non-convexity can come from: a commodity of weight zero is never penalised for the congestion it causes, and its route choice is a vertex choice. The reduction below exploits exactly this, with one weighted commodity and many weightless ones.

Theorem 5.12 (Membership in the commodity cone is co-NP-complete). *Given a network as above, a rational Wardrop equilibrium f^* in path flows (a property checked in polynomial time, the paths being listed) and a rational $\lambda \geq 0$, deciding whether $\lambda \in \Lambda(f^*)$ is co-NP-complete, and so is deciding whether $\Lambda(f^*) = \Lambda_1(f^*)$, when the number of commodities is part of the input. Both remain co-NP-hard when all of the following hold: $\lambda = e_W$ is a unit vector; f^* is the unique Wardrop equilibrium, it is a vertex of K , and F is strongly monotone on K ; $\Lambda_1(f^*)$ is the whole orthant; every edge lies on paths of at most two commodities; demands are integers; and all commodities but one have exactly two paths. [proved here, Appendix B; checked by an independent internal referee (round 4) and in E8]*

Proof outline. Membership in co-NP is the argument of Proposition 5.9. For hardness, one weighted commodity W chooses between a route O of constant cost and a route T_ℓ per orientation $\ell = (i, j)$ of an edge of a MAX-CUT instance; T_ℓ shares an edge with the route S_i of node i , one with the other route U_j of node j , and one with the route Z_1 of a switch commodity, which meets every T_ℓ . Expensive private edges hold the weightless commodities on U_i and Z_1 (so f^* is the unique equilibrium), but C_W does not charge them for the congestion they impose on W ; minimising over W 's flows leaves a concave function (Lemma 5.11) whose vertex values are $C_W(f^*)$ and $C_W(f^*) + k - \frac{1}{2} - \text{cut}_w(x)$. $\square$

Remark 5.13 (Interior weights). Hardness does not need zero weights. With B, k, n as in the proof, $\Gamma = 2B + 2k + 1$ and $\varepsilon = 1/(4(n+1)\Gamma)$, the weight $\lambda_\varepsilon = e_W + \varepsilon \sum_{h \neq W} e_h$ has all entries positive, and $\lambda_\varepsilon \in \Lambda(f^*)$ iff every cut has weight $< k$: if so, $\Lambda(f^*)$ is the orthant; otherwise the flow built from a heavy cut beats f^* by at least $\frac{1}{4}$ for $\Phi_{\lambda_\varepsilon}$ (Appendix B). [proved here, Appendix B; checked by an independent internal referee (round 4) and in E8]

Corollary 5.14 (Networks with all paths). *When the input is a directed acyclic graph with origin–destination pairs and P_k is the set of all paths of commodity k from its origin to its destination, both problems of Theorem 5.12 are co-NP-complete when the number of commodities is part of the input. They remain co-NP-hard when $\lambda = e_W$, f^* is the unique Wardrop equilibrium and a vertex of K , $\Lambda_1(f^*)$ is the whole orthant, demands are integers, every arc lies on paths of at most three commodities, and all commodities but two (W and a switch, which has more than two paths) have exactly two paths. In this construction F is not strongly monotone (Question 7.1(ii)). [proved here, Appendix B; checked by an independent internal referee (round 4) and in E8 on small graphs]*

Proposition 5.15 (Two tractable cases). (a) *If, on every edge e with $a_e > 0$, λ takes one value μ_e on all commodities having a path through e , then Φ_λ is convex on K and $\lambda \in \Lambda(f^*)$ iff $\lambda \in \Lambda_1(f^*)$, a linear feasibility test. This covers $\lambda = \mathbf{1}$ (the system optimum) and networks in which no edge with $a_e > 0$ is shared.*

(b) *(Explicit path sets.) If $\lambda = \mu \mathbf{1}_Q$ and at most r commodities outside Q use an edge of a path of Q , then $\lambda \in \Lambda(f^*)$ can be decided by solving at most $\prod |P_k| \leq (\max_k |P_k|)^r$ convex quadratic programmes over a simplex product, hence in polynomial time for fixed r [19]. In particular membership of a unit vector is polynomial when the number of commodities is fixed.*

[proved here, Appendix B; elementary]

Theorem 5.12 uses one weighted commodity and unboundedly many weightless ones sharing edges with the weighted paths, which, with explicit path sets, is unavoidable, unless $P = NP$, for weights of the form $\mu \mathbf{1}_Q$ (Proposition 5.15(b)); Example 5.7 ($\lambda = (0, 1)$) and Example 5.8 are instances with few commodities. With all paths of an acyclic network, the case of few commodities is open (Question 7.1(iii)).

6 Numerical checks

All equilibria are computed by the projected-gradient iteration of Theorem 3.4(b) (Remark 3.5), run to a fixed-point step below 10^{-12} , with the natural residual $\|x - \Pi_K(x - F(x))\|$ reported; Λ_1 is tested by its linear system and Λ by exact maximisation (Appendix A). Each script states its protocol and pass/fail criteria in its docstring and writes the flags into its output, from which the phrases “all criteria hold” are generated; there is no external pre-registration record (the criteria were written in the same session as the code, before the reference run). The experiments verify statements proved above; they discover nothing. Full tables are in `results/tables_*.md`.

E1: routing. all 4 criteria hold. Pigou: $\text{PoA} = 1.333333$ for $d = 1$ and 4.7268 for $d = 16$ (closed form and grid search agree to 1.5×10^{-11}). Affine parallel links: maximum PoA 1.2094 over 200 instances, 0 violations of $4/3$, maximum residual 3.4×10^{-11} ; the solver matches exact water-filling to 8.8×10^{-11} , which for affine costs is also the only check of the tolls of Corollary 4.2 we offer ($c + \tau$ is the same function as ∇C).

E2–E4: welfare-weight cones. all 13 criteria hold. Example 5.4: closed form, linear test and exact maximisation agree on 181/181 directions; Cournot (i): 0/61 grid directions in Λ , (ii): agreement 61/61; Nikaidô–Isoda: $\max W'_\lambda \leq 2.8 \times 10^{-17}$. Example 5.6 (E2c): 0 disagreements on 7381 slice points, bisected boundary within 1.0×10^{-8} of the parabola, and within 1.1×10^{-8} of φ for $\varepsilon = 0.2$ (test tolerance 10^{-8}). In the 200 concave random games (E3) every tested point agrees with $\Lambda = \Lambda_1$; in the 200 non-concave ones (E4) $\Lambda \subseteq \Lambda_1$ is never violated and 58 of the 103 games with a non-trivial cone carry a certified witness of strict inclusion (a lower bound: witnesses are sought only among the cone points).

E5: two commodities. all 5 criteria hold. Example 5.7: $f^* = (1, 0)$ (residual 0), boundary $\lambda_1 = 0.025000$ on $\lambda_2 = 1$. In 200 random instances Λ_1 is two-dimensional in 16 and a ray in 75; 21 carry a certified witness of $\Lambda \neq \Lambda_1$ (8 at a direction with both weights positive), none of the 62 with an interior equilibrium and 4 of the 27 with every $b_e = 0$; 0 violations of $\Lambda \subseteq \Lambda_1$.

E6: the reduction of Proposition 5.9. all 5 criteria hold. On all graphs with unit weights and 3–5 nodes and on random graphs with 6 nodes (unit weights, and weights in $\{1, 2, 3\}$), 1696 graphs and 13686 pairs (G, k) with $\varepsilon = 0.2$: $e_{n+2} \in \Lambda(x^*)$ holds iff the maximum cut, found by enumeration, is $< k$ (0 mismatches; 3598 members and 10088 non-members), and $\Lambda = \Lambda_1 = \mathbb{R}_+^{n+2}$ exactly in those cases. Membership was decided by exact maximisation over all faces of the box for the 276 pairs with $n \leq 4$, and over its vertices for the other 13410; the latter relies on the multilinearity used in the proof (36 cross-checks against face enumeration agree).

E7: three commodities. all 5 criteria hold. Example 5.8: $y^* = (0, 0, 1)$ (residual 0); the bisected boundary on $\lambda_2 = 1$ at 16 values $\lambda_1 \in [0.7, 1.45]$ is within 1.0×10^{-8} of ψ ; the description $\{\lambda_3 + m \geq 0\}$ agrees with the exact test at 1878 grid points (0 mismatches); witness gap 0.0055555556 ($= 1/180$).

E8: the reduction of Theorem 5.12. all criteria hold (theory/check_routing_hardness.py; numbers parsed from its saved output, which a re-run reproduced, where the details are). On 381 pairs (G, k) from 130 graphs (the 70 labelled graphs with unit weights on 3 and 4 nodes, and $30 + 30$ random graphs on 5 and 6 nodes, 20 of them with weights in $\{1, 2, 3\}$), $\min_K C_W$, computed without the closed form, gives $e_W \in \Lambda(f^*)$ iff the maximum cut is $< k$ (0 mismatches, margin 0.5000); the equilibrium, Λ_1 , the witness and Remark 5.13 are checked in exact arithmetic; and the acyclic network of Corollary 5.14 (up to 66 paths for W and 20 for the switch) passes the same tests on 16 pairs.

7 What can be claimed, limitations and open question

Every statement carries its status in line; the claim-by-claim table is `results/CLAIMS.md`. In three lines: the classical results are theorems, proved above; the “formal relationship” between equilibrium and welfare is the first-order coincidence of Proposition 4.1, a definition rather than a mechanism; and (F, K) carries no welfare information (Proposition 5.3), so welfare statements need the externalities as separate data, and even with them deciding whether a weighting makes an equilibrium optimal is co-NP-complete when the number of agents is part of the input, for

quadratic games (Proposition 5.9, a standard reduction) and for multi-commodity routing with affine costs (Theorem 5.12, Corollary 5.14).

Limitations. (1) Everything is finite-dimensional, deterministic and C^1 . (2) Theorem 5.1 is a first-order statement; its only general sufficient condition for equality is concavity, which excludes Cournot-type payoffs and multi-commodity routing costs (for routing, Proposition 5.15(a) is a variant). (3) The experiments verify theorems on quadratic games over boxes and on small networks; they are exact only because $N \leq 4$ (E2–E5, E7), $n \leq 6$ (E6) or the networks are small (E8), and the reported frequencies describe one sampling distribution. (4) We have not surveyed the toll-pricing and multi-objective literature exhaustively (Remark 5.2).

Open question 7.1 (Routing with few commodities, and acyclic networks). Theorem 5.12 settles the complexity of membership in multi-commodity routing with affine costs when the number of commodities is part of the input, with explicit path sets or with all paths of an acyclic network (Corollary 5.14); with explicit path sets, a fixed number of commodities and weights of the form $\mu \mathbf{1}_Q$, membership is polynomial (Proposition 5.15(b)). Three questions remain. (i) What is the complexity of deciding $\lambda \in \Lambda(f^*)$ when the number of commodities is fixed (already two) and the weights are positive and distinct, so that two weighted commodities with many paths each share edges? (ii) Does Corollary 5.14 hold with F strongly monotone? Our acyclic construction is not: a path of the switch through q_0 , dz , the arcs $p_\ell, e_\ell^1, e_\ell^2, e_\ell^z$ of some T_ℓ and the tail of Z_1 is not separated from Z_0 by arcs of positive slope that W does not also use. (iii) What is the complexity with all paths of an acyclic network and a fixed number of commodities, already for two commodities and $\lambda = e_W$? Then $\min_K \Phi_\lambda$ is the minimum over W ’s arc flows of a convex quadratic plus d_R times the shortest-path length of the other commodity R for the arc lengths $a_e f_e^W$, a concave function, and Proposition 5.15(b) does not apply because the paths may be exponentially many.

Next steps. Settle Question 7.1; run E4 and E5 with a global solver for larger N and m ; and confirm with the author whether the “formulation reviewed” in the line’s record was an instance of Proposition 4.1 or of Corollary 4.2, which is our reading.

Acknowledgements. Four rounds of internal review by independent referees improved this note. They pointed out the counterexample of Example 5.6, suggested its strongly monotone variant, the three-commodity setting of Example 5.8 and the reduction behind Proposition 5.9, asked that the latter be labelled as standard, and checked Theorem 5.12, Remark 5.13 and Corollary 5.14 independently.

A Reproducibility

The directory contains `experiments/vi_solver.py` (projections and the iteration of Theorem 3.4(b)), `traffic_poa.py` (E1), `welfare_cone.py` (E2–E4), `commodity_cone.py` (E5), `hardness_maxcut.py` (E6) and `three_commodity.py` (E7); E8 is `theory/check_routing_hardness.py` (seed 20261003, 27 s of CPU in the saved run), whose text output is parsed by `make_numbers.py`. Seed 20260930, consumed sequentially by one generator per script; Python 3.11.15, NumPy 2.4.6 [13], SciPy 1.17.1 [35] (linprog/HiGHS for the geometry of Λ_1); E1 took 2 s, E2–E4 114 s, E5 7 s, E6 85 s and E7 2 s on shared cores. The scripts write `results/results_*.json` (with a `criteria` block) and `results/tables_*.md`; `experiments/make_numbers.py` turns them into the macros used here, so every number in the text is generated.

Exact tests and random families. $\lambda \in \Lambda$ is decided by enumerating the 3^N faces of the box and solving the stationarity system of W_λ on each (a maximiser is stationary in the relative interior of its face): $\lambda \in \Lambda$ iff $W_\lambda(x^*) \geq \max_K W_\lambda - 10^{-8}$, and a *certified witness* improves W_λ by more than 10^{-6} . The random quadratic games of E3–E4 ($\frac{1}{2}$ -strongly monotone F ; 64–75% of equilibrium coordinates on the boundary), the cross-check against multistart projected ascent (13 instances, discrepancy 1.4×10^{-17}), the 3 directions of E5 accepted only within tolerance (first-order violation $\leq 5.1 \times 10^{-5}$), the grids and the pass/fail criteria are documented in the script docstrings and in `results/tables_*.md`.

B Deferred proofs

We use the notation of Section 5, in particular (3).

Proof of the strongly monotone variant in Example 5.6. Fix $\lambda_3 = 1$ and $\frac{1}{4(1-\varepsilon)} \leq \lambda_1 \leq \frac{3}{4(1-\varepsilon)}$. As for $\varepsilon = 0$, $x_3 = 1$ at a maximiser; on the edge $x_2 = 0$ the maximum, at $x_1 = \frac{\lambda_1+1/4}{1+\varepsilon\lambda_1} \leq 1$, is $1 - \frac{\varepsilon}{2} + \frac{(\lambda_1+1/4)^2}{2(1+\varepsilon\lambda_1)}$, which exceeds $W_\lambda(x^*) = (1 + \lambda_1 + \lambda_2)(1 - \frac{\varepsilon}{2}) - \frac{1}{4}$ iff $\lambda_2 < \varphi(\lambda_1)$. Conversely let $\lambda_2 \geq \varphi(\lambda_1)$. If the Hessian of W_λ in (x_1, x_2) is negative semidefinite, $\lambda \in \Lambda_1$ suffices; otherwise it is indefinite (its $(1, 1)$ entry is $-(1 + \varepsilon\lambda_1) < 0$), the maximum lies on an edge, and on the edges $x_1 = 1$, $x_2 = 1$ and $x_1 = 0$ the value is at most $W_\lambda(x^*)$ (increasing in x_2 ; increasing in x_1 because $\lambda_1 \geq \frac{1}{4(1-\varepsilon)}$; at most $(1 + \lambda_2)(1 - \frac{\varepsilon}{2}) < W_\lambda(x^*)$). $\square$

Proof of $\Lambda(f^)$ in Example 5.7.* For Λ fix $\lambda_2 = 1$ ($\lambda_2 = 0$ gives $\lambda \in \Lambda$). The Hessian of $\lambda_1 C_1 + C_2$, $\begin{pmatrix} 4\lambda_1 & \lambda_1+1 \\ \lambda_1+1 & 5/2 \end{pmatrix}$, is positive semidefinite iff $10\lambda_1 \geq (\lambda_1+1)^2$, and then $\lambda \in \Lambda$. Otherwise the minimum lies on an edge, where the restriction is a convex quadratic: on $f_B = 1$ and on $f_D = 0$ it is at f^* (derivatives $\lambda_1 + \frac{1}{2} + \frac{5}{2}t > 0$ and $4\lambda_1(f_B - 1) \leq 0$); on $f_D = 1$ the value is $\geq \frac{15}{8}\lambda_1 + 1 > \lambda_1 + \frac{1}{4}$; on $f_B = 0$ it is $3\lambda_1 + \frac{1}{4}(1-t)^2 + t^2$, minimised at $t = \frac{1}{5}$ with value $3\lambda_1 + \frac{1}{5}$. Hence f^* is optimal iff $\lambda_1 + \frac{1}{4} \leq 3\lambda_1 + \frac{1}{5}$, i.e. $\lambda_1 \geq \frac{1}{40}$, which is the claim after scaling. $\square$

Proof of the description of $\Lambda(y^)$ in Example 5.8.* Write $y = (u, w, t)$, $\Phi_\lambda = \sum_k \lambda_k C_k$ and $c = (\lambda_1 + \lambda_3)u + (\lambda_2 + \lambda_3)w \geq 0$; expanding, $\Phi_\lambda(y) - \Phi_\lambda(y^*) = P(u, w) + \lambda_3 + t(c - 2\lambda_3) + \lambda_3 t^2$. For fixed (u, w) this is convex in t . If $c \geq 2\lambda_3$ its minimum over $t \in [0, 1]$ is at $t = 0$ and equals $P + \lambda_3 \geq m + \lambda_3$. Otherwise it is attained at $t = 1 - c/(2\lambda_3)$ and equals $P + c - c^2/(4\lambda_3) \geq P + \frac{c}{2} \geq P + \frac{1}{2}(\lambda_1 u + \lambda_2 w) \geq \frac{1}{4}(\lambda_1 u^2 + \lambda_2 w^2) \geq 0$. Hence y^* is a minimiser iff $\lambda_3 + m \geq 0$, and when $\lambda_3 + m < 0$ the minimiser of P on the face $t = 0$ beats it. $\square$

Proof of Proposition 5.9. Membership in co-NP. $\lambda \notin \Lambda(x^*)$ iff $W_\lambda(x) > W_\lambda(x^*)$ for some $x \in K$. Let $\hat{x}$ maximise W_λ over K and let $\mathcal{F}$ be the face of K in whose relative interior it lies. The restriction of the quadratic W_λ to the affine hull of $\mathcal{F}$ has a local, hence global, maximum at $\hat{x}$; it is therefore concave there, and every point of the nonempty polytope $\{x \in \mathcal{F} : \nabla_{\mathcal{F}} W_\lambda(x) = 0\}$ ($\nabla_{\mathcal{F}}$ the projection of the gradient on the direction space of $\mathcal{F}$) is a maximiser. A vertex of this polytope has encoding size polynomial in the input [32, Section 10.2] and is the certificate, checked in exact arithmetic. The argument uses only that the objective is quadratic with rational coefficients and that K is a rational polyhedron given by linear inequalities; for quadratic programming over general polyhedra it is the theorem of Vavasis [34]. (Deciding that a given point is *not* a local minimiser of an indefinite quadratic programme is NP-complete [22]; see also Pardalos and Schnitger [26].) For $\Lambda \neq \Lambda_1$, a certificate is an extreme-ray generator λ of the pointed rational polyhedral cone $\Lambda_1(x^*)$, of polynomial size by the same theorem, together with such an x for it: if every generator lay in the convex cone Λ , then $\Lambda_1 \subseteq \Lambda$. *Hardness.* We reduce the complement of MAX-CUT, which is co-NP-complete [16], to membership: YES-instances of MAX-CUT go to non-members. Given a graph (V, E) on n nodes with nonnegative integer weights w_{ij} and an integer $k \geq 1$ (is there a cut of weight $\geq k$?), let $\text{cut}(x) = \sum_{\{i,j\} \in E} w_{ij}(x_i + x_j - 2x_i x_j)$,

$\deg_w(i) = \sum_{j:\{i,j\} \in E} w_{ij}$, $L(x) = \sum_i \deg_w(i)x_i$, which is $\geq \text{cut}(x)$ on $[0, 1]^n$ because $w \geq 0$ (with a negative weight the reduction fails), and $q(x, y) = \text{cut}(x) + y(k - \frac{1}{2} - L(x))$. As q is affine in each variable separately, its maximum over $[0, 1]^{n+1}$ is attained at a vertex, where $q = \text{cut}(x)$ if $y = 0$ and $q = \text{cut}(x) - L(x) + k - \frac{1}{2} \leq k - \frac{1}{2} = q(0, 1)$ if $y = 1$; so $(0, 1)$ maximises q iff every cut has weight $< k$. Take players with actions $x_1, \dots, x_n, y, z \in [0, 1]$, fix $0 < \varepsilon < 1$, and let $u_i = -x_i - \frac{\varepsilon}{2}x_i^2$, $u_{n+1} = y - \frac{\varepsilon}{2}y^2$, $u_{n+2} = z - \frac{\varepsilon}{2}z^2 + q(x, y)$. Each payoff is concave in the own action, $F(v) = \varepsilon v + (\mathbf{1}_n, -1, -1)$ is ε -strongly monotone, and $x^* = (0, \dots, 0, 1, 1)$ is the unique equilibrium (strictly dominant strategies). $W_{e_{n+2}} = u_{n+2}$ separates in z , so $e_{n+2} \in \Lambda(x^*)$ iff every cut has weight $< k$. At x^* , $\nabla_x q = 0$ and $\partial_y q = k - \frac{1}{2} > 0$, so $\Lambda_1(x^*) = \mathbb{R}_+^{n+2}$ by (2); the other unit vectors lie in $\Lambda(x^*)$ by dominance, so $\Lambda(x^*) = \Lambda_1(x^*)$ iff $e_{n+2} \in \Lambda(x^*)$. The construction is polynomial. (Merging the first $n+1$ players into one player with action $(x, y) \in [0, 1]^{n+1}$ gives the same conclusions with $N = 2$; membership in co-NP holds for vector actions in boxes by the same proof.) $\square$

Proof of Lemma 5.11. In (3), g_e involves only f^Q , and $f_e = \sum_k f_e^k$ is linear in the path flows; so for fixed f^Q the expression is affine in f^R . A pointwise infimum of affine functions is concave, and a concave function on the polytope $\prod_{k \in R} \{f^k \geq 0, \sum_p f_p^k = d_k\}$ attains its minimum at a vertex, i.e. at a pure profile; a minimiser of Φ_λ over K is obtained by adding a minimising f^Q for that profile (the minimum over the compact K exists). If $\lambda = \mu \mathbf{1}_Q$, then $g_e = \mu f_e^Q$ and, with $r_e = f_e - f_e^Q$ fixed, $\Phi_\lambda = \mu \sum_e (a_e(f_e^Q)^2 + (a_e r_e + b_e)f_e^Q)$ is convex in f^Q ; commodities of R that use no edge of a path of Q do not appear in it. $\square$

Proof of Theorem 5.12 and Remark 5.13. Membership in co-NP. K is a product of simplices given by explicit linear constraints (the paths are listed in the input) and Φ_λ is a quadratic polynomial with rational coefficients, so the certificate argument in the proof of Proposition 5.9 applies [34]. For $\Lambda \neq \Lambda_1$: $\lambda \in \Lambda_1(f^*)$ is a linear feasibility test by Lemma 2.6 (eliminate γ_k , the value on a used path), and $\Lambda_1(f^*)$ is a pointed rational polyhedral cone whose extreme rays have generators of polynomial size [32, Section 10.2], and a certificate is such a generator together with a point of K that beats f^* for it.

Construction. As in Proposition 5.9 we reduce the complement of MAX-CUT [16], sending YES-instances of MAX-CUT to non-members. An instance is a graph (V, E_G) , $V = \{1, \dots, n\}$, with integer weights $w_{ij} \geq 1$ and an integer $k \geq 1$: is there a cut of weight $\geq k$? We may assume that $E_G \neq \emptyset$ (otherwise the answer is no and we output a fixed member) and that no node is isolated (deleting it changes no cut). Let A be the set of the $M = 2|E_G|$ ordered pairs $\ell = (i, j)$ with $\{i, j\} \in E_G$, $w_\ell = w_{ij}$, and set

$$D = M, \quad \alpha_\ell = 16w_\ell, \quad b_o = 16 \max_\ell w_\ell, \quad a_D = \frac{2k-1}{2D}, \quad B = 2 + 2(D+1) \sum_\ell \alpha_\ell.$$

Edges and costs: dz with cost $a_D x$; e^o with constant cost b_o ; for each $\ell \in A$, p_ℓ with constant cost $b_o - \alpha_\ell \geq 0$, e_ℓ^1 and e_ℓ^2 with cost $\alpha_\ell x$, and e_ℓ^z with cost $2\alpha_\ell x$; for each $i \in V$, s_i with cost $x + B$ and r_i with cost x ; and q_1 with cost x , q_0 with cost $x + B$. Commodities:

- W , demand D , paths $O = \{dz, e^o\}$ and $T_\ell = \{dz, p_\ell, e_\ell^1, e_\ell^2, e_\ell^z\}$, $\ell \in A$;
- for each $i \in V$, commodity i , demand 1, paths $S_i = \{s_i\} \cup \{e_\ell^1 : \ell = (i, \cdot)\}$ and $U_i = \{r_i\} \cup \{e_\ell^2 : \ell = (\cdot, i)\}$;
- a switch z , demand 1, paths $Z_1 = \{q_1\} \cup \{e_\ell^z : \ell \in A\}$ and $Z_0 = \{q_0, dz\}$.

Let f^* send W along O , every i along U_i and z along Z_1 , and $\lambda = e_W$. Every edge lies on paths of at most two commodities, and all numbers have polynomial size.

The objective. Write $x_i \in [0, 1]$ for the flow of i on S_i , $z \in [0, 1]$ for the flow of the switch on Z_1 and $t_\ell \geq 0$ for the flow of W on T_ℓ (so $D - \sum_\ell t_\ell$ on O); this affine reparametrisation of K is one-to-one. Edge dz carries $D + 1 - z$, all of W 's demand; e^o carries $D - \sum_\ell t_\ell$ at the

constant cost b_o ; and for $\ell = (i, j)$, e_ℓ^1 , e_ℓ^2 , e_ℓ^z carry $t_\ell + x_i$, $t_\ell + 1 - x_j$, $t_\ell + z$. Hence, since $(b_o - \alpha_\ell) - b_o + \alpha_\ell = 0$,

$$C_W = a_D D(D + 1 - z) + b_o D + \sum_{\ell=(i,j) \in A} \alpha_\ell (4t_\ell^2 + t_\ell(x_i - x_j + 2z)). \quad (4)$$

For fixed (x, z) the minimum over $\{t \geq 0, \sum_\ell t_\ell \leq D\}$ separates: $t_\ell = \frac{1}{8}(x_j - x_i - 2z)_+ \leq \frac{1}{8}$, so $\sum_\ell t_\ell \leq M/8 \leq D$ is slack, and with $\alpha_\ell/16 = w_\ell$,

$$h(x, z) := \min_t C_W = a_D D(D + 1 - z) + b_o D - \sum_{\ell=(i,j) \in A} w_\ell ((x_j - x_i - 2z)_+)^2.$$

By Lemma 5.11 ($\lambda = e_W$; or directly, (4) is affine in (x, z) for fixed t) h is concave, so its minimum over $[0, 1]^{n+1}$ is attained at a vertex. For $z = 1$, $h = C^* := a_D D^2 + b_o D$, since $x_j - x_i - 2 < 0$. For $z = 0$ and $x \in \{0, 1\}^n$, the pair $\ell = (i, j)$ contributes $-w_\ell$ iff $x_i = 0$, $x_j = 1$, and each edge $\{i, j\}$ with $x_i \neq x_j$ does so through exactly one of its two orientations; so $h = C^* + (k - \frac{1}{2}) - \text{cut}_w(x)$. Therefore

$$\min_K C_W = C^* + \min(0, k - \frac{1}{2} - \text{maxcut}_w), \quad C_W(f^*) = C^*,$$

and $e_W \in \Lambda(f^*)$ iff $\text{maxcut}_w \leq k - \frac{1}{2}$, i.e. iff every cut has weight $< k$. When a cut x has weight $\geq k$, the flow y with that x , $z = 0$ and $t_\ell = \frac{1}{8}$ on the pairs $\ell = (i, j)$ with $x_i = 0$, $x_j = 1$ lowers C_W by $\text{cut}_w(x) - k + \frac{1}{2} \geq \frac{1}{2}$. *Interior weights (Remark 5.13)*. At y every path of a node or of the switch costs at most Γ : $c_{S_i} \leq 1 + B + \frac{9}{8} \sum_\ell \alpha_\ell \leq 2B$, $c_{U_i} \leq 1 + \frac{9}{8} \sum_\ell \alpha_\ell$, $c_{Z_1} = \sum_\ell 2\alpha_\ell t_\ell$ and $c_{Z_0} \leq 1 + B + a_D(D + 1) \leq 1 + B + 2k$. As these demands are 1, $0 \leq C_h(y) \leq \Gamma$ and $C_h(f^*) \geq 0$ for $h \neq W$, so $\Phi_{\lambda_\varepsilon}(y) - \Phi_{\lambda_\varepsilon}(f^*) \leq -\frac{1}{2} + \varepsilon(n + 1)\Gamma = -\frac{1}{4}$.

f^* is the unique equilibrium. For every feasible flow, $c_{U_i} \leq 1 + (D + 1) \sum_\ell \alpha_\ell < B \leq c_{S_i}$ and $c_{Z_1} \leq 1 + 2(D + 1) \sum_\ell \alpha_\ell < B \leq c_{Z_0}$, so at every Wardrop equilibrium $x = 0$ and $z = 1$. Then $c_{T_\ell} - c_O = \alpha_\ell t_\ell + \alpha_\ell(t_\ell + 1) + 2\alpha_\ell(t_\ell + 1) - \alpha_\ell = 4\alpha_\ell t_\ell + 2\alpha_\ell > 0$, so W uses only O . Thus f^* , a vertex of K , is the only equilibrium. F is affine with $DF = \Delta^\top \text{diag}(a)\Delta$, Δ the edge-path incidence matrix. If h is tangent to K ($\sum_{p \in P_k} h_p = 0$ for all k) and $h^\top DFh = \sum_e a_e (\Delta h)_e^2 = 0$, then $(\Delta h)_e = 0$ on s_i, r_i, q_1, q_0 (slope 1, each in a single path), so h vanishes on the paths of the nodes and of the switch; then $(\Delta h)_{e_\ell^1} = h_{T_\ell} = 0$ and $h_O = -\sum_\ell h_{T_\ell} = 0$. So DF is positive definite on the tangent space and F is strongly monotone on K .

$\Lambda_1(f^*)$ is the orthant, and $\Lambda = \Lambda_1$ iff $e_W \in \Lambda$. At f^* the derivatives of C_W with respect to path flows are: $2a_D D + b_o$ on O and $2a_D D + b_o + 2\alpha_\ell$ on T_ℓ ; 0 on S_i and on U_i ; 0 on Z_1 and $a_D D$ on Z_0 . In each commodity the used path has the least derivative, so $e_W \in \Lambda_1(f^*)$ by Lemma 2.6. For a node i , $\sigma_i := 1 + \sum_{\ell=(\cdot, i)} \alpha_\ell = C_i(f^*)$ and, for every feasible flow, $C_i \geq \sigma_i(1 - x_i)^2 + Bx_i$, and $\sigma_i(1 - x)^2 + Bx - \sigma_i = x(\sigma_i x - 2\sigma_i + B) \geq 0$ because $B \geq 2\sigma_i$; so $e_i \in \Lambda(f^*)$. For the switch, $\sigma_z := 1 + 2\sum_\ell \alpha_\ell = C_z(f^*)$, $C_z \geq \sigma_z z^2 + B(1 - z)$, and $\sigma_z z^2 + B(1 - z) - \sigma_z = (1 - z)(B - \sigma_z(1 + z)) \geq 0$ because $B \geq 2\sigma_z$ ($D \geq 1$); so $e_z \in \Lambda(f^*)$. As $\Lambda \subseteq \Lambda_1$ are convex cones (Theorem 5.1(a),(b)), $\Lambda_1(f^*)$ contains every unit vector, hence equals $\mathbb{R}_+^{n+2}$ (there are $n + 2$ commodities), and $\Lambda(f^*) = \mathbb{R}_+^{n+2}$ iff $e_W \in \Lambda(f^*)$. Both decision problems therefore have answer “yes” iff every cut has weight $< k$. $\square$

Proof of Corollary 5.14. Network. Number the pairs $\ell = 1, \dots, M$. Turn every edge of the construction into an arc with two endpoints of its own, add an origin o_k and a destination d_k per commodity, and add connector arcs of constant cost: for W , $o_W \rightarrow dz \rightarrow e^o \rightarrow d_W$ and $dz \rightarrow p_\ell \rightarrow e_\ell^1 \rightarrow e_\ell^2 \rightarrow e_\ell^z \rightarrow d_W$; for node i , $o_i \rightarrow s_i \rightarrow e_{\ell_1}^1 \rightarrow \dots \rightarrow e_{\ell_r}^1 \rightarrow d_i$ and $o_i \rightarrow r_i \rightarrow e_{\ell'_1}^2 \rightarrow \dots \rightarrow e_{\ell'_r}^2 \rightarrow d_i$, along the pairs $\ell_1, \dots, \ell_r$ leaving and $\ell'_1, \dots, \ell'_r$ entering i ; for the switch, $o_z \rightarrow q_1 \rightarrow e_1^z \rightarrow \dots \rightarrow e_M^z \rightarrow d_z$ and $o_z \rightarrow q_0 \rightarrow dz \rightarrow d_z$ (“ $x \rightarrow y$ ” is an arc from the head of x to the tail of y). A connector between two consecutive arcs of the same chain ($e^1 \rightarrow e^1$, $e^2 \rightarrow e^2$, $e^z \rightarrow e^z$) costs $H = b_o + 1$; all other connectors cost 0; and the cost of q_0

becomes $x + B'$ with $B' = B + 2(M - 1)H$. Put dz , e^o , the p_ℓ and the private arcs s_i, r_i, q_0, q_1 on level 0 and $e_\ell^1, e_\ell^2, e_\ell^z$ on levels 1, 2, 3: within level 0 arcs go $q_0 \rightarrow dz \rightarrow \{e^o, p_\ell\}$, chain connectors stay on a level and go forward, all other connectors go up, and nothing goes down, so the graph is acyclic.

Paths. (i) A path from $\mathbf{o}_i$ enters the chain S_i (level 1) or U_i (level 2). It can leave a chain only through a connector of W , into the chain U_j of the head $j \neq i$ of the pair, into Z_1 , or to $\mathbf{d}_W$, and it can never come back, while $\mathbf{d}_i$ is entered only from the last arc of S_i or of U_i . So commodity i has exactly the two paths S_i, U_i . (ii) A path of W other than O and the T_ℓ leaves T_ℓ at the head of e_ℓ^1, e_ℓ^2 or e_ℓ^z along a chain connector (the other out-arcs end at $\mathbf{d}_i, \mathbf{d}_j$ or $\mathbf{d}_z$), so it contains an arc of cost H . (iii) A path of the switch other than Z_1 and Z_0 cannot leave Z_1 (its exits lead to $\mathbf{d}_W$), so it starts $\mathbf{o}_z \rightarrow q_0 \rightarrow dz$, then continues along arcs of W and chain connectors, and reaches $\mathbf{d}_z$ through the tail of the chain Z_1 , ending with e_M^z ; for every ℓ , $q_0, dz, p_\ell, e_\ell^1, e_\ell^2, e_\ell^z$ followed by the chain Z_1 up to e_M^z is such a path, so the switch has at least $M + 2$ paths. (iv) By (i), an arc lies on paths of at most one node, hence on paths of at most three commodities.

Minimum. Moving an amount δ of W 's flow from a path p of type (ii) to O changes $C_W = \sum_e c_e(f_e)f_e^W$ by at most $\delta b_o - \delta H < 0$: arcs of p off O lose $c_e(f_e)f_e^W - c_e(f_e - \delta)(f_e^W - \delta) \geq \delta c_e(f_e - \delta)$, which is δH on the chain connector, and O gains δb_o on e^o . Moving the switch's flow from a path of type (iii) to Z_0 removes flow from arcs and adds it only on the arc $dz \rightarrow \mathbf{d}_z$, which W does not use, so C_W does not increase. Hence $\min_K C_W$ is attained on the face where W uses O and the T_ℓ and the switch uses Z_1 and Z_0 ; there, the connectors used by W cost 0 and the chain connectors carry no flow of W , so C_W is (4). The minimum and $C_W(f^*)$ are those of the theorem.

Equilibrium and Λ_1 . An arc e_ℓ^z carries only W and the switch, so $c_{Z_1} \leq 1 + 2(D + 1) \sum_\ell \alpha_\ell + (M - 1)H < B'$ for every feasible flow, while every other path of the switch contains q_0 : at every equilibrium the switch uses only Z_1 . Both paths of node i contain $\deg(i) - 1$ chain connectors. An arc e_ℓ^1 or e_ℓ^2 may now also carry flow of the switch, along a path of type (iii), so $f_e \leq D + 2$ there instead of $D + 1$; still $c_{U_i} \leq 1 + (D + 2) \sum_\ell \alpha_\ell + (\deg(i) - 1)H < B + (\deg(i) - 1)H \leq c_{S_i}$, because $D + 2 \leq 2(D + 1)$ gives $1 + (D + 2) \sum_\ell \alpha_\ell < 2 + 2(D + 1) \sum_\ell \alpha_\ell = B$. Given $x = 0$ and $z = 1$, for every flow of W , $c_{T_\ell} - c_O \geq -\alpha_\ell + \alpha_\ell f_{e_\ell^2} + 2\alpha_\ell f_{e_\ell^z} \geq 2\alpha_\ell > 0$, since $f_{e_\ell^2}, f_{e_\ell^z} \geq 1$, and a path of type (ii) costs at least $c_O + H - b_o > c_O$; so f^* is the unique equilibrium. At f^* the derivative of C_W along a path of type (ii) exceeds that along O by at least $H - b_o > 0$, and along a path of type (iii) it is at least $a_D D > 0$, the value on Z_1 being 0; so $e_W \in \Lambda_1(f^*)$. The bounds for e_i shift both sides by $(\deg(i) - 1)H$, and for the switch $C_z \geq \sigma_z z^2 + z(M - 1)H + B'(1 - z)$, whose excess over $C_z(f^*) = \sigma_z + (M - 1)H$ is $(1 - z)(B' - \sigma_z(1 + z) - (M - 1)H) \geq 0$, as $B' - (M - 1)H = B + (M - 1)H \geq 2\sigma_z$; so $e_i, e_z \in \Lambda(f^*) \subseteq \Lambda_1(f^*)$, and $\Lambda_1(f^*) = \mathbb{R}_+^{n+2}$ as before.

F is not strongly monotone. Let h move a unit of the switch from Z_0 to the path $\mathbf{o}_z \rightarrow q_0 \rightarrow dz \rightarrow p_M \rightarrow e_M^1 \rightarrow e_M^2 \rightarrow e_M^z \rightarrow \mathbf{d}_z$ of type (iii) and a unit of W from T_M to O . Then dz and q_0 keep their flow, the arcs of T_M gain the switch and lose W , and otherwise only e^o and connectors change, so $(\Delta h)_e = 0$ on every arc with $a_e > 0$; hence $DF h = 0$ and $F(f + \tau h) = F(f)$ whenever f and $f + \tau h$ are feasible: F is not even strictly monotone.

Membership. C_k depends only on the arc flows $(f_e^k)_e$ of the commodities; on an acyclic graph every origin–destination flow of value d_k decomposes into path flows, so $\lambda \in \Lambda(f^*)$ iff the arc flows of f^* minimise Φ_λ over the multicommodity flow polytope, which has a polynomial description, and the argument of Proposition 5.9 applies (that f^* is a Wardrop equilibrium is checked by shortest paths). The first-order condition defining $\Lambda_1(f^*)$ is equivalent to the same condition on that polytope (a linear function of path flows is minimised at f^* iff every used path is shortest for it). By linear-programming duality, $\lambda \in \Lambda_1(f^*)$ iff there are node potentials π_k with nonnegative reduced arc costs for the linearised Φ_λ of each commodity k , zero on the arcs used by k ; so $\Lambda_1(f^*)$ is the projection of a rational polyhedral cone in (λ, π) of polynomial

description and has extreme-ray generators of polynomial size [32, Section 10.2]. $\square$

Proof of Proposition 5.15. (a) Edges with $a_e = 0$ contribute $b_e g_e$, which is linear; on the others $g_e = \mu_e f_e$, so (3) is $\sum_e \mu_e (a_e f_e^2 + b_e f_e)$ plus a linear function, convex since $\mu_e \geq 0$; apply Proposition 4.1(a). (b) Lemma 5.11: enumerate the pure profiles of the r commodities, and for each decide whether the convex quadratic programme in f^Q has value $< \Phi_\lambda(f^*)$. $\square$

References

- [1] Ravindra K. Ahuja and James B. Orlin. Inverse optimization. *Operations Research*, 49(5):771–783, 2001.
- [2] Martin Beckmann, C. Bartlett McGuire, and Christopher B. Winsten. *Studies in the Economics of Transportation*. Yale University Press, New Haven, 1956.
- [3] Harold P. Benson and Erjiang Sun. Outcome space partition of the weight set in multiobjective linear programming. *Journal of Optimization Theory and Applications*, 105(1):17–36, 2000.
- [4] Jacek Bochnak, Michel Coste, and Marie-Françoise Roy. *Real Algebraic Geometry*, volume 36 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)*. Springer, Berlin, 1998.
- [5] José R. Correa, Andreas S. Schulz, and Nicolás E. Stier-Moses. Selfish routing in capacitated networks. *Mathematics of Operations Research*, 29(4):961–976, 2004.
- [6] Stella Dafermos. Traffic equilibrium and variational inequalities. *Transportation Science*, 14(1):42–54, 1980.
- [7] John M. Danskin. The theory of max-min, with applications. *SIAM Journal on Applied Mathematics*, 14(4):641–664, 1966. doi: 10.1137/0114053.
- [8] Pradeep Dubey. Inefficiency of Nash equilibria. *Mathematics of Operations Research*, 11(1):1–8, 1986.
- [9] Matthias Ehrgott. *Multicriteria Optimization*. Springer, Berlin, 2nd edition, 2005.
- [10] Francisco Facchinei and Jong-Shi Pang. *Finite-Dimensional Variational Inequalities and Complementarity Problems*. Springer, New York, 2003. Two volumes.
- [11] Arthur M. Geoffrion. Proper efficiency and the theory of vector maximization. *Journal of Mathematical Analysis and Applications*, 22(3):618–630, 1968.
- [12] Patrick T. Harker and Jong-Shi Pang. Finite-dimensional variational inequality and nonlinear complementarity problems: a survey of theory, algorithms and applications. *Mathematical Programming*, 48:161–220, 1990.
- [13] Charles R. Harris, K. Jarrod Millman, Stéfan J. van der Walt, et al. Array programming with NumPy. *Nature*, 585:357–362, 2020.
- [14] Philip Hartman and Guido Stampacchia. On some non-linear elliptic differential-functional equations. *Acta Mathematica*, 115:271–310, 1966.
- [15] Donald W. Hearn and Motakuri V. Ramana. Solving congestion toll pricing models. In Patrice Marcotte and Sang Nguyen, editors, *Equilibrium and Advanced Transportation Modelling*, pages 109–124. Kluwer, Boston, 1998.
- [16] Richard M. Karp. Reducibility among combinatorial problems. In Raymond E. Miller and James W. Thatcher, editors, *Complexity of Computer Computations*, pages 85–103. Plenum Press, New York, 1972.
- [17] David Kinderlehrer and Guido Stampacchia. *An Introduction to Variational Inequalities and Their Applications*. Academic Press, New York, 1980. Reprinted as SIAM Classics in Applied Mathematics 31, 2000.
- [18] Frank H. Knight. Some fallacies in the interpretation of social cost. *Quarterly Journal of Economics*, 38(4):582–606, 1924.
- [19] M. K. Kozlov, S. P. Tarasov, and L. G. Khachiyan. The polynomial solvability of convex quadratic programming. *USSR Computational Mathematics and Mathematical Physics*, 20(5):223–228, 1980. doi: 10.1016/0041-5553(80)90098-1. Translated from Zh. Vychisl. Mat. Mat. Fiz. 20(5), 1319–1323.
- [20] Dov Monderer and Lloyd S. Shapley. Potential games. *Games and Economic Behavior*, 14(1):124–143, 1996.
- [21] Stephen Morris and Takashi Ui. Best response equivalence. *Games and Economic Behavior*, 49:260–287, 2004.
- [22] Katta G. Murty and Santosh N. Kabadi. Some NP-complete problems in quadratic and nonlinear programming. *Mathematical Programming*, 39(2):117–129, 1987.
- [23] Anna Nagurney. *Network Economics: A Variational Inequality Approach*. Kluwer, Dordrecht, 1993.
- [24] John Nash. Non-cooperative games. *Annals of Mathematics*, 54(2):286–295, 1951.
- [25] Hukukane Nikaidô and Kazuo Isoda. Note on non-cooperative convex games. *Pacific Journal of Mathematics*, 5(5):807–815, 1955.
- [26] Panos M. Pardalos and Georg Schnitger. Checking local optimality in constrained quadratic programming is NP-hard. *Operations Research Letters*, 7(1):33–35, 1988. doi: 10.1016/0167-6377(88)90049-1.
- [27] Arthur Cecil Pigou. *The Economics of Welfare*. Macmillan, London, 1920.
- [28] R. Tyrrell Rockafellar. *Convex Analysis*. Princeton University Press, Princeton, 1970.
- [29] J. Ben Rosen. Existence and uniqueness of equilibrium points for concave n -person games. *Econometrica*, 33(3):520–534, 1965.
- [30] Tim Roughgarden. *Selfish Routing and the Price of Anarchy*. MIT Press, Cambridge, MA, 2005.
- [31] Tim Roughgarden and Éva Tardos. How bad is self-

- ish routing? *Journal of the ACM*, 49(2):236–259, 2002.
- [32] Alexander Schrijver. *Theory of Linear and Integer Programming*. Wiley, Chichester, 1986.
 - [33] Michael J. Smith. The existence, uniqueness and stability of traffic equilibria. *Transportation Research Part B*, 13(4):295–304, 1979.
 - [34] Stephen A. Vavasis. Quadratic programming is in NP. *Information Processing Letters*, 36(2):73–77, 1990.
 - [35] Pauli Virtanen, Ralf Gommers, Travis E. Oliphant, et al. SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nature Methods*, 17: 261–272, 2020.
 - [36] John Glen Wardrop. Some theoretical aspects of road traffic research. *Proceedings of the Institution of Civil Engineers, Part II*, 1(3):325–362, 1952. Discussion, pp. 362–378.