

Foundational Geometry:

Invariants Across Formalisms and Structures Preceding a Geometric Description

Luciano Silva Alarco

Pontificia Universidad Católica del Perú

ORCID 0000-0002-3395-3129

Working draft v0.8 — research line MRT001 — 3 October 2026

Abstract

We ask which claims about “structures that precede a geometric description” can be formulated and tested inside explicitly defined mathematical frameworks. We introduce a minimal bookkeeping language: a *formalism* is a class of structures, a *translation* is a map between formalisms, and an *invariant across formalisms* is a quantity that factors through a translation. Geometry then appears as what survives the passage from a richer to a poorer formalism, in the spirit of Klein’s Erlangen programme and of the meaningfulness criterion of measurement theory. The language is worked out in full on one chain where every step can be proved or computed, the finite Euclidean chain $\text{coordinates} \rightarrow \text{distance matrix} \rightarrow \text{ranking of distances} \rightarrow \text{betweenness and congruence}$. Three results are established. (i) On generic finite samples the relational primitives of elementary geometry are empty: exact integer arithmetic over 4000 random configurations produced no non-degenerate instance of betweenness or congruence, while integer grids produce dozens to thousands; the primitives that axiomatise the continuum are therefore not sample-robust. (ii) The ordinal formalism is strictly coarser than similarity geometry for every finite $n \geq 3$, yet a non-metric reconstruction from the ranking alone recovers the configuration up to similarity with a median Procrustes disparity of 2.1×10^{-10} at $n = 256$ in the plane, 7.6 orders of magnitude below its value at $n = 8$; this is the finite-sample face of the asymptotic uniqueness theorem for ordinal embeddings, and the class itself is measured directly: its inradius, in the norm $\max_i \|y_i - x_i\|$, lies between a quarter and a half of the minimal gap between consecutive sorted distances (exactly a quarter when the two extremal pairs are disjoint), a quantity that scales empirically as $n^{-4.0}$. (iii) Standard geometric quantities sort themselves by *formalism depth*: minimum spanning tree, nearest-neighbour graph, relative neighbourhood graph and diameter pair are ordinal invariants, whereas diameter, affine dimension and the Gabriel graph are metric but not ordinal invariants at finite n . The same language is then applied to the causal-order translation $\text{Lorentzian coordinates} \rightarrow \text{causal order}$, followed by a statistical step that adds counting, on uniform sprinklings of the 1+1 Minkowski diamond, where the kernel of the order translation contains the conformal group and coincides with it for some but not all finite configurations, and counting restores proper times and coordinates. We prove that the number of realizers of the causal order of a uniform sample, that is, the number of conformal orbits, up to parity, that make up its kernel class, converges in law to 2^Z with Z Poisson of mean one, at rate $O(1/n)$, and we identify the exceptions to the equality with 2^N , N the number of descending successions of the permutation that sends u -ranks to v -ranks; the proof is self-contained except for Gallai’s theorem on prime comparability graphs, and the combinatorial lemmas it uses are known and reproved. We also identify the first-order term: the total variation distance to 2^Z is $(1 + 1/(2e))/n + O(n^{-2})$, a result checked by an independent internal referee; in this version, and not yet refereed, we make the remainder explicit (at most $422/n^2$ for $n \geq 22$) and show that the number of descending successions is at total variation distance $(3/(4e) + o(1))/n^2$ from $\text{Po}(1 - 1/n)$, while the n^{-2} term of the realizer law is only computed, from the exact law, and its closed form is conjectural. The paper claims no unified theory and reports no experimental physics; its contribution is to turn a recurring foundational phrase into a list of statements with explicit status.

1 Introduction

The idea that geometry is not primitive, that something poorer than a metric or a coordinate system comes first and geometry is recovered from it, is old and recurs in several disciplines under different names. Klein’s Erlangen programme defines a geometry by the group of transformations under which its statements are invariant [20]. Hilbert and Tarski axiomatise Euclidean geometry from the primitive relations of betweenness and congruence [18, 39, 40]. Menger and Blumenthal ask which metric spaces are “Euclidean” and how much of a configuration a distance table determines [4, 28]. Measurement theory calls a statement *meaningful* when it is invariant under the admissible transformations of the scale on which the quantities were measured [22, 31, 38]. In relativity, the causal order of a spacetime determines its topology and its conformal geometry [17, 27], and the causal set programme [7] condenses this, in Sorkin’s later slogan, into “order plus number equals geometry” [37].

Each of these traditions is precise on its own ground, but the phrase “structure preceding geometry” is often used across them without saying which structure, which geometry, and in what sense one precedes the other. The research line to which this draft belongs (MRT001) had, at the time of writing, the modest and explicit objective of *specifying which claims of this kind can be formulated and tested within defined mathematical frameworks*. This paper is an attempt to discharge that objective by three moves.

1. We introduce a small bookkeeping language (Section 2): formalisms, translations between them, the kernel of a translation, invariants along a translation, and the *formalism depth* of a quantity. Nothing in it is new; its purpose is to make “ G precedes F ” a predicate that can be checked.
2. We work the language out completely on one chain where every claim can be proved or computed: finite configurations of points in $\mathbb{R}^d$, read successively as coordinates, as a distance matrix, as a ranking of the pairwise distances, and as Tarski’s betweenness and congruence relations (Sections 3 and 4).
3. We apply the same language to the causal order of a uniform sprinkling of 1+1 Minkowski space (Section 5), where the continuum theorems of Hawking–King–McCarthy and Malament have finite, computable counterparts, and where the finite- n ambiguity of the order obeys an exact limit law (Theorem 5.6).

Scope. We do not propose a unified theory, a new geometry or a physical model; the Lorentzian part is a mathematical model with a fixed background and is used only because it is the sharpest known instance of “order before metric”. Every numerical statement in the paper is produced by four short scripts included with the manuscript, with a fixed seed, and by three more that check the proofs of Section 5 numerically (Appendices A and B). Section 6 lists all claims with their status: proved here, classical, taken from the literature, computationally verified, or conjectural.

2 Formalisms, translations and invariants

Definition 2.1 (Formalism). A *formalism* is a pair $F = (\text{Obj}_F, \cong_F)$ consisting of a class of objects and an equivalence relation on it, read as isomorphism. Two objects that are isomorphic are regarded as the same description.

Definition 2.2 (Translation, kernel). A *translation* $\tau: F \rightarrow G$ is a map $\text{Obj}_F \rightarrow \text{Obj}_G$ such that $X \cong_F Y$ implies $\tau X \cong_G \tau Y$. Its *kernel* is the equivalence relation $X \sim_\tau Y \iff \tau X \cong_G \tau Y$ on Obj_F , which is coarser than (contains) $\cong_F$. Two objects in the same kernel class are *indistinguishable in G* .

Definition 2.3 (Invariant across formalisms). A function Φ on Obj_F is an *invariant along* τ , or a G -invariant of F , if $X \sim_\tau Y$ implies $\Phi(X) = \Phi(Y)$; equivalently, Φ factors through τ . We write $\text{Inv}(\tau)$ for the set of such functions.

Definition 2.4 (Chain, formalism depth). A *chain* is a sequence of translations $F_0 \xrightarrow{\tau_1} F_1 \xrightarrow{\tau_2} \dots \xrightarrow{\tau_k} F_k$. The *formalism depth* of a quantity Φ on Obj_{F_0} is the largest i such that $\Phi \in \text{Inv}(\tau_i \circ \dots \circ \tau_1)$, that is, the poorest formalism in the chain from which Φ can still be computed; the empty composition ($i = 0$) is allowed, so a quantity that is not invariant even along τ_1 has depth 0, the coordinate level.

Lemma 2.5 (Monotonicity). *For composable translations, $\text{Inv}(\tau_2 \circ \tau_1) \subseteq \text{Inv}(\tau_1)$. Hence along a chain the sets of invariants decrease and the kernels increase.*

Proof. If $\tau_1 X \cong \tau_1 Y$ then $\tau_2 \tau_1 X \cong \tau_2 \tau_1 Y$, so $\sim_{\tau_1} \subseteq \sim_{\tau_2 \tau_1}$, and a function constant on the classes of the coarser relation is constant on those of the finer one. $\square$

Definition 2.6 (Faithfulness, precedence). Let $\tau: F \rightarrow G$ and let $\mathcal{C} \subseteq \text{Obj}_F$ be a class of objects of interest. The translation is *faithful on $\mathcal{C}$ up to a group H* acting on Obj_F if, for $X, Y \in \mathcal{C}$, $X \sim_\tau Y$ implies $Y = hX$ for some $h \in H$. For a family of formalisms F_n (below, $\text{Coord}(n, d)$), classes $\mathcal{C}_n \subseteq \text{Obj}_{F_n}$ and a fixed H -invariant distance δ on Obj_{F_n} (below, the Procrustes disparity), translations τ_n are *asymptotically faithful up to H* on $(\mathcal{C}_n)$ if $\sup\{\delta(X, Y) : X \in \mathcal{C}_n, Y \sim_{\tau_n} X\} \rightarrow 0$ as $n \rightarrow \infty$. We say that G *precedes* F on $\mathcal{C}$ when τ has a non-trivial kernel and is nevertheless faithful, or asymptotically faithful, up to a group H that is itself a symmetry group of F .

The definition makes the foundational phrase testable: to claim that a structure G precedes a geometry F one must exhibit the translation, compute or bound its kernel, and prove or measure its faithfulness. Two classical readings are recovered at once.

Remark 2.7 (Erlangen). If Obj_F is a set with an action of a group H and τ is the quotient map to Obj_F/H , then $\text{Inv}(\tau)$ is exactly the set of H -invariants, Klein's geometry of the pair (Obj_F, H) . Euclidean, similarity, affine and projective geometry are the invariants of successively coarser quotients, and Lemma 2.5 is the Erlangen hierarchy.

Remark 2.8 (Meaningfulness). Stevens' scale types are kernels of translations [38]. A ratio scale identifies quantities up to a positive factor, an interval scale up to a positive affine map, an ordinal scale up to a monotone bijection. A statement is *meaningful* in the sense of Krantz et al. [22] and Narens [31] precisely when it is an invariant along the corresponding translation. The chain studied below realises two of these scale types on one object: the distance matrix is a ratio-scale description of a configuration and the ranking of distances is its ordinal-scale description; a configuration read up to similarity is what the ratio scale determines.

3 The finite Euclidean chain

Fix $n \geq 3$ and $d \geq 1$. Write $[n] = \{1, \dots, n\}$ and P_n for the set of unordered pairs of distinct elements of $[n]$, $m = |P_n| = \binom{n}{2}$. All objects are labelled; quotienting by relabelling adds a graph-isomorphism layer that is orthogonal to the questions asked here.

- **Coord**(n, d): injective configurations $X = (x_1, \dots, x_n) \in (\mathbb{R}^d)^n$ (distinct points), with equality as isomorphism.
- **Met**(n): symmetric matrices $D \in \mathbb{R}^{n \times n}$ with zero diagonal and positive off-diagonal entries, with equality as isomorphism.
- **Ord**(n): total preorders $\preceq$ on P_n (generically total orders), with equality as isomorphism.

- $\text{Rel}(n)$: pairs (B, C) of a ternary relation $B \subseteq [n]^3$ (betweenness) and a binary relation $C \subseteq P_n \times P_n$ (congruence), with equality as isomorphism.

The translations are the obvious ones. $\tau_1: \text{Coord} \rightarrow \text{Met}$ sends X to its distance matrix $D_X = (\|x_a - x_b\|)_{a,b}$. $\tau_2: \text{Met} \rightarrow \text{Ord}$ sends D to the preorder $\{a, b\} \preceq \{c, d\} \iff D_{ab} \leq D_{cd}$. $\tau_3: \text{Met} \rightarrow \text{Rel}$ sends D to

$$B_D = \{(a, b, c) \text{ distinct} : D_{ab} + D_{bc} = D_{ac}\}, \quad C_D = \{(p, q) \in P_n^2 : p \neq q, D_p = D_q\}.$$

The congruence part of τ_3 factors through τ_2 (congruent pairs are ties of the preorder) but the betweenness part does not, since it involves an additive equality; the chain therefore branches at Met .

3.1 Coordinates to metric: the classical step

Theorem 3.1 (Schoenberg; Young and Householder). *Two labelled configurations $X, Y \in (\mathbb{R}^d)^n$ have the same distance matrix if and only if $Y = gX$ for a Euclidean motion $g \in E(d)$, possibly improper. A matrix D is the distance matrix of a configuration in $\mathbb{R}^d$ if and only if $-\frac{1}{2}JD^{(2)}J$ is positive semidefinite of rank at most d , where $D^{(2)}$ is the entrywise square and $J = I - \frac{1}{n}\mathbf{1}\mathbf{1}^\top$.*

See Blumenthal [4], Schoenberg [33], Young and Householder [46]. In the language of Section 2: $\ker \tau_1$ is the orbit relation of $E(d)$, τ_1 is faithful up to $E(d)$ on all of Coord , and $\text{Inv}(\tau_1)$ is Euclidean geometry in Klein's sense. The metric formalism precedes the coordinate one in the strongest possible way: nothing is lost except position and orientation.

3.2 Metric to relations: the primitives are empty on samples

Tarski's system axiomatises elementary Euclidean geometry with the single sort "point" and the two primitives betweenness and congruence, and the system is complete and decidable [39, 40]. The restriction of these primitives to a finite sample behaves very differently from their behaviour on the continuum.

Proposition 3.2 (Generic emptiness). *Let $d \geq 2$ and $n \geq 3$, and let $\mathcal{G} \subseteq (\mathbb{R}^d)^n$ be the set of configurations in which no three points are collinear and all m pairwise distances are distinct. Then $\mathcal{G}$ is open, dense and of full Lebesgue measure, and for every $X \in \mathcal{G}$*

$$B_{D_X} = \emptyset \quad \text{and} \quad C_{D_X} = \emptyset.$$

Consequently $\tau_3 \circ \tau_1$ is constant on $\mathcal{G}$: its kernel identifies all generic configurations of a given size, and every relational invariant is constant on $\mathcal{G}$.

Proof. Collinearity of three distinct points x_a, x_b, x_c is the vanishing of all 2×2 minors of the $d \times 2$ matrix $[x_b - x_a \mid x_c - x_a]$, a system of polynomial equations in the coordinates that is not identically satisfied when $d \geq 2$; its solution set is a proper real-algebraic subset, hence closed, nowhere dense and null. Equality $\|x_a - x_b\|^2 = \|x_c - x_d\|^2$ for two distinct pairs is a single non-constant polynomial equation (non-constant because moving x_a alone changes the left side and, when $a \notin \{c, d\}$, not the right; when the pairs share a point the same argument applies to the point that is not shared), with the same consequences. $\mathcal{G}$ is the complement of a finite union of such sets, so it is open, dense and of full measure. For $X \in \mathcal{G}$: if $D_{ab} + D_{bc} = D_{ac}$ with a, b, c distinct then, by the equality case of the triangle inequality for the Euclidean norm (strict convexity), $x_b - x_a$ and $x_c - x_b$ are non-negative multiples of one another, so x_b lies on the segment $[x_a, x_c]$ and the three points are collinear, contradicting $X \in \mathcal{G}$. A congruence (p, q) with $p \neq q$ contradicts distinctness of the distances. The final assertion follows because $\tau_3\tau_1 X = (\emptyset, \emptyset)$ for all $X \in \mathcal{G}$. $\square$

n	d	configurations	betweenness instances	congruence instances
5	2	500	0	0
10	2	500	0	0
20	2	500	0	0
50	2	500	0	0
5	3	500	0	0
10	3	500	0	0
20	3	500	0	0
50	3	500	0	0
grid		n	betweenness instances	congruence instances
3×3	9	8	138	
4×4	16	44	976	
5×5	25	152	4242	

Table 1: E1. Exact counts of non-degenerate betweenness triples and congruent pair-pairs in random integer configurations (upper panel, 500 configurations per row) and in square integer grids (lower panel). A betweenness instance is a pair $(b, \{a, c\})$ with b strictly between a and c (a row of four collinear points contributes $1 + 1 + 2 = 4$); a congruence instance is an unordered pair of distinct point-pairs at equal distance.

Remark 3.3 (The line). For $d = 1$ every triple is collinear and B_{D_X} is the betweenness relation of the linear order of the sample, which is far from empty; the proposition then holds only for C . On the line the relational formalism carries exactly the order type of the sample, which is the structure Section 5 takes as its starting point.

Computational check (E1). Proposition 3.2 is a statement about a full-measure set, and a floating-point test of the equalities defining B and C would be misleading: the betweenness gap $D_{ab} + D_{bc} - D_{ac}$ is quadratic in the distance from x_b to the line through x_a, x_c , so a relative tolerance of 10^{-9} admits every triple whose middle point lies within about $10^{-4.5}$ of a line and, at $n = 50$, produced of the order of one spurious triple per configuration in a preliminary floating-point run. We therefore sample integer coordinates uniformly in $\{0, \dots, 2^{30} - 1\}^d$ and test collinearity and equality of squared distances *exactly* in 64-bit integer arithmetic (the sampling grid is fine enough that the result is the generic one with overwhelming probability, and coarse enough that an exact coincidence, should one occur, would be counted). Table 1 reports the totals: across 8 settings and 4000 configurations, 0 betweenness instances and 0 congruence instances were found. The integer grids in the lower panel, where the relations are rich, show that the test itself is not blind.

Remark 3.4 (Axiomatic versus observational primitives). The primitives that make Tarski’s theory complete on the plane carry no information about a generic finite sample of the plane. Any empirical test of a geometric claim uses finitely many measured points, so such a test must pass through the metric or the ordinal formalism; betweenness and congruence can only be *inferred* from them, with a tolerance whose meaning depends on n and on the chosen tolerance, not on the configuration. This is the first precise sense in which “the structure that precedes geometry” cannot be the axiomatic primitives.

3.3 Metric to ordinal: finite non-uniqueness

Proposition 3.5 (Finite non-uniqueness). *Let $n \geq 3$, $d \geq 1$. For every configuration X of distinct points whose pairwise distances are distinct, the kernel class of X under $\tau_2 \circ \tau_1$ contains a neighbourhood of X in $(\mathbb{R}^d)^n$, and that neighbourhood contains configurations that are not similar to X . In particular $\tau_2\tau_1$ is not faithful up to $\text{Sim}(d)$ on any open set, for any finite n .*

Proof. The distances are continuous functions of X , and the ordinal pattern of X is defined by finitely many strict inequalities between them, which persist in a neighbourhood U of X ; so U lies in the kernel class. Pick three labels a, b, c and move x_b along the ray from x_a through x_b by a small amount: D_{ab} changes while D_{ac} does not, so the ratio D_{ab}/D_{ac} changes. Similarities preserve all distance ratios, hence the moved configuration, which lies in U for a small enough move, is not similar to X . $\square$

Example 3.6 (E3). On the line, $\{0, 1, 3\}$ and $\{0, 1, 2.5\}$ have the same ordinal pattern (distances $1 < 2 < 3$ and $1 < 1.5 < 2.5$ in the same pair order) and are not similar. In the plane, the triangles with sides $(3, 4, 5)$ and $(3, 4, 6)$ have the same pattern and angles $(36.9, 53.1, 90.0)$ and $(26.4, 36.3, 117.3)$ degrees respectively. For $n = 3$ the ordinal formalism records exactly the order of the three sides, equivalently the order of the three angles, and its generic fibres are the six open cells into which these orderings cut the two-dimensional shape space of triangles.

The ordinal formalism nevertheless retains dimensional information. For $n = 4$ there are 720 possible orderings of the six distances. Sampling 300 000 uniform configurations, we observe 120 distinct orderings on the line but all 720 in the plane and all 720 in space (the least frequent planar ordering was seen 53 times, so the count is not an artefact of sample size). The count on the line has a short explanation: for sorted points with gaps u, v, w the six distances are $u, v, w, u + v, v + w, u + v + w$, and only ten orderings of these are consistent with positivity, each labelled pattern arising from two (labelling, gap-order) combinations under reflection, giving $24 \cdot 10/2 = 120$.

3.4 Metric to ordinal: asymptotic faithfulness

Theorem 3.7 (Kleindessner and von Luxburg; informal statement). *Let $\Omega \subset \mathbb{R}^d$ be bounded, connected and open and satisfy the mild regularity condition imposed in Kleindessner and von Luxburg [21] (for instance, Ω a finite union of open balls), and let $x_1, x_2, \dots$ be points of Ω that become dense in Ω . Then the complete ordinal pattern of $(x_1, \dots, x_n)$, that is, the outcome of all comparisons $D_{ab} < D_{cd}$ over quadruples of labels, determines the configuration up to similarity in the limit: any sequence of configurations $y^{(n)} \in (\mathbb{R}^d)^n$, in the same dimension d , with the same ordinal pattern as $(x_1, \dots, x_n)$ converges, after alignment by similarity transformations, to the original points as $n \rightarrow \infty$, in the sense made precise in Kleindessner and von Luxburg [21].*

We state the theorem informally and refer to the source for the exact regularity condition and for the mode of convergence. Terada and von Luxburg [41] obtain a conclusion of the same kind when only the unweighted k -nearest-neighbour graph of the sample is retained, for points drawn from a density on a suitable domain and $k \rightarrow \infty$ with $k/n \rightarrow 0$, at a rate specified there. In our language, take $\mathcal{C}_n = \{(x_1, \dots, x_n)\}$, the initial segments of one fixed dense sequence, and δ the Procrustes disparity; the theorem, read with the alignment uniform over the labels $i \leq n$, which is what a vanishing disparity requires, says that $\tau_2\tau_1$ is asymptotically faithful up to $\text{Sim}(d)$ on $(\mathcal{C}_n)$. The ordinal formalism therefore precedes similarity geometry on dense samples of such domains, although by Proposition 3.5 it never does so exactly at finite n .

Computational check (E2). For $d \in \{2, 3\}$ and n from 8 to 256 (128 for $d = 3$) we sample n points uniformly in the unit cube, compute the rank matrix of the pairwise distances (so that *only* the ordinal pattern enters), reconstruct a configuration by non-metric multidimensional scaling [25, 36] as implemented in scikit-learn [32] (SMACOF with isotonic regression, four random starts, stress tolerance 10^{-13} , at most 50 000 iterations), and measure the Procrustes disparity between the original and the reconstruction after optimal translation, rotation, reflection and scaling [16, 35]: both configurations are centred and scaled to unit Frobenius norm and the disparity is the residual sum of squares, a similarity-invariant number in $[0, 1]$. Table 2 and Figure 1 report 255 runs. In the plane the median disparity falls from 8.2×10^{-3} at $n = 8$

d	n	runs	median	Q_1	Q_3	max	discordance	local exp.
2	8	30	8.2×10^{-3}	5.8×10^{-3}	1.6×10^{-2}	1.6×10^{-1}	2.1×10^{-2}	–
2	16	30	8.1×10^{-5}	5.4×10^{-5}	1.2×10^{-4}	3.0×10^{-4}	2.4×10^{-3}	6.7
2	32	30	1.8×10^{-6}	1.3×10^{-6}	2.2×10^{-6}	3.3×10^{-6}	3.1×10^{-4}	5.5
2	64	30	3.0×10^{-8}	2.7×10^{-8}	3.6×10^{-8}	4.9×10^{-8}	5.1×10^{-5}	5.9
2	128	30	1.6×10^{-9}	1.5×10^{-9}	1.7×10^{-9}	2.0×10^{-9}	1.2×10^{-5}	4.3
2	256	30	2.1×10^{-10}	2.0×10^{-10}	2.2×10^{-10}	2.3×10^{-10}	4.3×10^{-6}	3.0
3	8	15	4.2×10^{-2}	1.6×10^{-2}	7.5×10^{-2}	1.9×10^{-1}	3.7×10^{-2}	–
3	16	15	3.8×10^{-4}	2.8×10^{-4}	4.9×10^{-4}	6.6×10^{-4}	3.9×10^{-3}	6.8
3	32	15	4.2×10^{-6}	3.3×10^{-6}	5.8×10^{-6}	8.3×10^{-6}	4.6×10^{-4}	6.5
3	64	15	8.3×10^{-8}	7.9×10^{-8}	9.0×10^{-8}	1.2×10^{-7}	7.0×10^{-5}	5.7
3	128	15	4.1×10^{-9}	3.8×10^{-9}	4.4×10^{-9}	4.7×10^{-9}	1.7×10^{-5}	4.3

Table 2: E2. Procrustes disparity (similarity-invariant, in $[0, 1]$) between a uniform random configuration and its reconstruction from the ranking of the pairwise distances alone. Discordance: median fraction of ordinal constraints violated by the reconstruction (Kendall). Local exp.: $-\log(\text{median}_n/\text{median}_{n/2})/\log 2$.




Figure 1: E2. Median and interquartile range of the Procrustes disparity of the ordinal reconstruction as a function of n , for $d = 2$ and $d = 3$.

to 2.1×10^{-10} at $n = 256$, 7.6 orders of magnitude; in space from 4.2×10^{-2} to 4.1×10^{-9} at $n = 128$. The local exponents between successive doublings of n (last column of Table 2) range from 6.7 down to 3.0 in the plane, so a single power law is not supported. Two cautions apply. First, the reconstruction is only approximately in the kernel class: the solver stops at a stress tolerance, and the Kendall discordance between the input ranking and the ranking of the output, that is, the fraction of ordinal constraints the output violates, is 4.3×10^{-6} at $n = 256$ (Table 2); with the looser tolerance 10^{-9} of a preliminary run the disparities were between ten and more than a hundred times larger, depending on n , so the absolute values are partly set by the stopping rule. Second, since X itself lies in the class, a solver output near X measures the distance from X to one approximate member of the class, not the class’s diameter. The numbers are consistent with a rapidly shrinking class, because the solver starts from a random configuration and sees only the ranking, but they do not measure the class; they are consistent with Theorem 3.7, not its quantitative content.

Invariance check (E2b). A monotone distortion $D \mapsto D^p$, $p \in [2, 4]$, leaves the rank matrix literally unchanged, so non-metric scaling, which reads only the rank matrix, returns the identical reconstruction by construction (the reported maximum disparity between the two outputs, 1.9×10^{-31} , is round-off of the alignment); this part of E2b checks the pipeline, not

a fact about geometry. The content is in the metric column: metric multidimensional scaling, which uses the values, reconstructs X from D with disparity 7.9×10^{-13} but from D^p with median disparity 4.5×10^{-2} (minimum 1.3×10^{-2} ; $n = 64$, 20 trials). The distortion changes the object of Met and not the object of Ord; this is what “invariant across formalisms” means operationally.

3.5 Measuring the ordinal class directly

The cautions attached to E2 can be met by measuring the kernel class itself rather than a solver’s approximation to it. One measurement is exact.

Proposition 3.8 (Thickness of the ordinal class). *Let $X \in \text{Coord}(n, d)$ have distinct pairwise distances and let $g(X) > 0$ be the minimal difference between consecutive values in the sorted list of its m pairwise distances. (i) If $\|y_i - x_i\| < g(X)/4$ for all i , then Y has the same ordinal pattern as X . (ii) If the two pairs of points realising $g(X)$ are disjoint, there is an injective configuration Y with $\max_i \|y_i - x_i\| = g(X)/4$ in which those two distances coincide (the moved points stay distinct because $g(X) \leq \min_{a \neq b} D_{ab}$); Y therefore lies outside the kernel class (its preorder has a tie), and every neighbourhood of Y contains configurations in which the two distances are strictly reversed. Consequently the inradius of the kernel class at X , in the norm $\max_i \|y_i - x_i\|$, lies in $[g(X)/4, g(X)/2]$, and equals $g(X)/4$ in case (ii).*

Proof. (i) By the triangle inequality, $|\|y_a - y_b\| - \|x_a - x_b\|| \leq \|y_a - x_a\| + \|y_b - x_b\| < g/2$ for every pair, so every distance moves by less than $g/2$ and the difference between any two distances changes by less than g ; two distances that are consecutive in the sorted order differ by at least g and therefore keep their order, hence so do all pairs. (ii) Let $\{a, b\}$ and $\{c, d\}$ be the disjoint pairs with $D_{cd} - D_{ab} = g$. Move x_a and x_b apart along the line through them by $g/4$ each, and x_c and x_d towards each other along their line by $g/4$ each, leaving the other points fixed: the first distance becomes $D_{ab} + g/2$ and the second $D_{cd} - g/2 = D_{ab} + g/2$, a tie, and the four displacements have norm $g/4$. For the upper bound in general, two distinct pairs share at most one point; if $\{a, b\}$ and $\{a, c\}$ realise g , moving x_b away from x_a and x_c towards x_a by $g/2$ each produces the same tie. In this shared-point case the smallest displacement that closes the gap depends on the angle θ at x_a : moving x_a as well, in the direction of $e_{ac} - e_{ab}$ (e_{ab}, e_{ac} the unit vectors from x_a towards x_b and x_c), changes $D_{ab} - D_{ac}$ at first order by $\|e_{ab} - e_{ac}\| = 2 \sin(\theta/2)$ per unit of displacement, so the closing displacement is $g/(2(1 + \sin(\theta/2))) + O(g^2)$; it equals $g/4$ exactly when x_b, x_a, x_c are collinear with x_a between the other two (a configuration the proposition allows), and it approaches $g/2$ as $\theta \rightarrow 0$. Finally, a configuration outside the class is reached along the segment from X only after two distances that are consecutive at X have tied, so the inradius is the minimum over consecutive pairs of their closing displacements; by (i) no tie occurs below $g/4$, and the constructions above tie the pair realising g at a displacement of at most $g/2$, exactly $g/4$ in case (ii). The perturbed configurations remain injective, because $g \leq \min_{a \neq b} D_{ab}$ for $n \geq 3$: if $\{1, 2\}$ is the closest pair and 3 any other point, $|D_{13} - D_{23}| \leq D_{12}$ by the triangle inequality, so some consecutive gap is at most D_{12} . $\square$

Computational check (E2c). For $d = 2$ and n from 8 to 256, 112 uniform configurations were examined (Table 3, Figure 2). Random displacements of every point by a vector of Euclidean norm below $g/4$ preserved the pattern in all 2240 of 2240 trials, and the explicit displacement of part (ii), applied just beyond $g/4$ so that the tie becomes a strict reversal detectable in floating point, changed it in all 91 of 91 configurations whose extremal pairs were disjoint. The median gap scales as $n^{-4.0}$, which is what one expects, heuristically, for the minimal spacing among $m \approx n^2/2$ weakly dependent values with a bounded density, of order $1/m^2 \propto n^{-4}$: the class is thin because n points carry $n^2/2$ distances. A second, similarity-invariant measurement is a random walk constrained to the class (proposals accepted only if the ranking is unchanged),

n	median g	disjoint	kept below $g/4$	changed beyond $g/4$	walk steps	median walk range
8	6.7×10^{-4}	10/20	400/400	10/10	8000	5.5×10^{-3}
16	3.8×10^{-5}	15/20	400/400	15/15	8000	4.4×10^{-5}
32	4.0×10^{-6}	16/20	400/400	16/16	6000	2.7×10^{-7}
64	1.9×10^{-7}	19/20	400/400	19/19	4000	1.0×10^{-9}
128	9.4×10^{-9}	20/20	400/400	20/20	3000	3.8×10^{-12}
256	6.7×10^{-10}	11/12	240/240	11/11	1500	6.4×10^{-15}

Table 3: E2c (20 configurations per n , 12 at $n = 256$). Median minimal gap g between consecutive sorted distances (the inradius of the class is $g/4$ when the extremal pairs are disjoint), the numerical checks of Proposition 3.8 (the displacement of part (ii) is applied with $g/4$ enlarged by $\max(10^{-6}g/4, 10^{-12})$ so that the tie becomes a strict reversal), and the median walk range (largest Procrustes disparity from X visited by a random walk confined to the class; a non-saturated lower bound on the extent of the class). Interquartile ranges are shown in Figure 2 and listed in the result files.




Figure 2: E2c. Median and interquartile range of the minimal gap g and of the walk range against n ($d = 2$); the walk range depends on the step budget and is a lower bound only.

which records the largest Procrustes disparity from X among the configurations visited. This *walk range* is a lower bound on the extent of the class at X in that measure, and a loose one: the recorded value grows roughly linearly with the number of steps within the budgets of Table 3 and has not saturated, so neither its value nor its scaling with n ($n^{-7.9}$ for that schedule) measures the class; the scaling is what one expects from an accepted step size proportional to g and a disparity quadratic in the displacement. The walk nevertheless establishes that the class is far wider than its thickness in generic directions: at $n = 64$ it stays in the class after displacing every point by several hundred times $g/4$, whereas displacing every point by $g/4$ produces a disparity of order $3g^2/8$ (the squared displacement $n(g/4)^2$ divided by the expected squared centred norm $n/6$ of a uniform sample of the unit square; an order-of-magnitude estimate, not a bound valid for every X), that is 1.4×10^{-14} for the median g at $n = 64$, against a walk range of 1.0×10^{-9} . Whether the class extends as far as the residual 3.0×10^{-8} at which the E2 solver stopped (with discordance 5.1×10^{-5}) is not decided by these measurements. What the two measurements do establish is the following. No ball of sup-norm radius larger than $g/2$ centred at X fits inside the class, and g decays empirically like n^{-4} , so the thickness of the class at a configuration of the unit square shrinks at that rate (Proposition 3.8 gives thickness in $[g/4, g/2]$, hence exactly at that rate); this bounds the thickness of the class, not its diameter modulo similarity, which is what Theorem 3.7 controls. An upper bound on that diameter is not established here.

quantity	depth	rigid motion	similarity	monotone distortion
centroid (position)	coordinate	0.00	0.00	–
diameter (value)	metric	1.00	0.00	0.00
affine dimension	metric	1.00	1.00	0.35
Gabriel graph	metric	1.00	1.00	0.03
diameter pair	ordinal	1.00	1.00	1.00
k -nearest-neighbour graph ($k = 3$)	ordinal	1.00	1.00	1.00
minimum spanning tree	ordinal	1.00	1.00	1.00
relative neighbourhood graph	ordinal	1.00	1.00	1.00

Table 4: E4. Formalism depth of standard quantities and the fraction of 300 trials in which each is preserved under rigid motions (which generate $\ker \tau_1$), similarities, and monotone distortions of the distance matrix (which generate $\ker \tau_2$ on **Met**; the distorted matrix is in general not Euclidean, so this column tests the defining formula on **Met**, a sufficient condition for ordinal depth). The centroid is undefined for a bare metric.

4 Formalism depth of geometric quantities

Lemma 2.5 orders the invariants of the chain, and Section 3 locates the boundaries: position is lost at τ_1 , scale at τ_2 , everything at τ_3 on generic samples. Table 4 classifies familiar quantities by the poorest formalism from which they can be computed, together with a direct computational check (E4): each quantity is computed for 300 random configurations of 30 points in the plane, then recomputed after a random rigid motion, after a random similarity, and after a random monotone distortion $D \mapsto D^p$, $p \in [0.3, 3]$, and the fraction of trials in which the value is preserved is reported.

The ordinal rows have one-line proofs. The diameter pair is the argument of a maximum, which depends only on the order. The k -nearest-neighbour graph is defined by comparisons $D_{ij} < D_{ik}$ within each row. The minimum spanning tree is produced by Kruskal’s algorithm from the sorted list of edge weights and is therefore invariant under any strictly monotone reweighting [24]. The relative neighbourhood graph [42] contains ij iff no k satisfies $\max(D_{ik}, D_{jk}) < D_{ij}$, again a comparison. The metric rows fail for equally simple reasons: the diameter changes under scaling; the affine dimension is the number of positive eigenvalues of the doubly centred squared-distance matrix (Theorem 3.1), and a power D^p of a planar metric is not planar: for $p < 1$ it is a Euclidean metric of dimension at most $n - 1$ [34], for $p > 2$ the centred matrix has many positive eigenvalues, and only for $1 < p < 2$ did it have exactly two positive eigenvalues in every trial, so the value is preserved in a fraction 0.35 of trials, essentially the probability that $p \in (1, 2)$; the Gabriel graph [12] contains ij iff no k satisfies $D_{ik}^2 + D_{jk}^2 < D_{ij}^2$, an additive condition, and is preserved in a fraction 0.03 of trials. That the relative neighbourhood graph is ordinal and the Gabriel graph is not, although both are “proximity graphs” of the same family, illustrates that depth is a property of the defining formula, not of the intuitive kind of the quantity. The metric rows are also genuinely non-ordinal on **Coord**, which the distortion column alone does not show, since the distorted matrix leaves the Euclidean locus: a collinear configuration with distinct distances and a small generic perturbation of it share the ordinal pattern (Proposition 3.5) and have affine dimensions 1 and 2; a triangle with a right angle at k , perturbed to acute or to obtuse without changing the order of its sides, keeps the pattern and gains or loses the Gabriel edge ij . Both witnesses are checked in the code (E4b: patterns equal, yes, dimensions 1 and 2; patterns equal, yes, edge present yes and no).

Theorem 3.7 adds an asymptotic column to the table: every *continuous* similarity invariant (distance ratios, angles) is asymptotically determined by the ordinal pattern, because the pattern determines the configuration up to similarity in the limit; for discrete invariants such as the Gabriel graph the statement holds only away from configurations near the defining equality, since a dense sample always contains near-borderline pairs. The finite- n classification and the

asymptotic one therefore differ, and a claim that “order determines geometry” must say which of the two is meant.

5 A second translation: causal order and counting

The Euclidean chain has an analogue in which the poorer structure is a partial order rather than a ranking. Consider the causal diamond of 1+1 Minkowski space in null coordinates $u = t - x$, $v = t + x$, so that $ds^2 = du dv$, the diamond is the unit square $(0, 1)^2$, and the proper time between causally related events is $\tau(p, q) = \sqrt{(u_q - u_p)(v_q - v_p)}$. The causal order is the product order, $p \prec q \iff u_p < u_q$ and $v_p < v_q$; unrelated pairs are written $p \parallel q$. The formalisms are:

- **Coord_{1,1}(n)**: labelled configurations $(u_i, v_i)_{i \leq n}$ in the diamond;
- **Caus(n)**: partial orders on $[n]$ (labelled); for configurations with pairwise distinct u - and pairwise distinct v -coordinates (a full-measure open set, assumed throughout), the image of σ_1 below is the set of labelled orders of Dushnik–Miller dimension at most two [11], the orders realised by two linear orders, here $<_u$ and $<_v$;
- **Caus+N(n)**: the same partial order together with the statistical hypothesis that the configuration is a uniform sample of the diamond, so that counts of points estimate areas; its “invariants” are estimators, consistent as $n \rightarrow \infty$, rather than functions factoring through a translation.

The translation $\sigma_1: \text{Coord}_{1,1} \rightarrow \text{Caus}$ forgets the coordinates and keeps the order (the abstract notion of a causal space goes back to Kronheimer and Penrose [23]); **Caus+N** is not a formalism in the sense of Section 2 but a formalism together with a sampling model, which is what “number” means in the causal set slogan. The sequence is therefore a translation followed by a statistical step, not a chain in the sense of Definition 2.4. All statements below are about uniform samples of a fixed background; none is a statement about physics.

Proposition 5.1 (Kernel of the order translation). *Every map $(u, v) \mapsto (f(u), g(v))$ with f, g strictly increasing bijections of $(0, 1)$, and the swap $(u, v) \mapsto (v, u)$, preserve the causal order of every configuration. Boosts $(u, v) \mapsto (\lambda u, v/\lambda)$, as maps of $\mathbb{R}^{1,1}$, preserve the order and in addition all proper times; a general pair (f, g) does not.*

Proof. $u_p < u_q \iff f(u_p) < f(u_q)$ for increasing f , and likewise for g ; the swap exchanges the two coordinates in the conjunction defining $\prec$. Under a boost $(\lambda \Delta u)(\Delta v/\lambda) = \Delta u \Delta v$. For $f(u) = u^2$, $g = \text{id}$, the proper time between $(0, 0)$ and (u, v) becomes $u\sqrt{v}$. $\square$

The maps (f, g) with f, g increasing homeomorphisms of $(0, 1)$ are conformal maps of the diamond onto itself that preserve time and space orientation, and together with the swap (parity) they form a group $(\text{Homeo}^+(0, 1))^2 \rtimes \mathbb{Z}_2$ of causal automorphisms of the diamond. The group is infinite-dimensional, which is the well-known reason why Zeeman’s theorem, that the causal automorphisms of Minkowski space are generated by the orthochronous Lorentz group, translations and dilations, requires dimension at least three [47]. In the language of Section 2, $\ker \sigma_1$ contains the conformal orbits, the analogue of the monotone distortions that generate $\ker \tau_2$ in the Euclidean chain, so $\text{Inv}(\sigma_1)$ is contained in conformal geometry; the inclusion is strict for some configurations at finite n and an equality for others (Proposition 5.2), and it is always an equality in the continuum theorems: a bijection between past- and future-distinguishing spacetimes that preserves the chronological order in both directions is a smooth conformal isometry [17, 27]. The finite-sample counterparts are tested next.

Proposition 5.2 (Finite non-uniqueness, causal). *Let $n \geq 3$ and consider configurations with pairwise distinct u - and pairwise distinct v -coordinates. The kernel class of such a configuration*

under σ_1 is open in $(0,1)^{2n}$, and it is the union of the conformal orbits, modulo the swap, of the realizers $(\prec_u, \prec_v)$ of its order. The number of realizers modulo the swap is 1 for a chain and otherwise equals half the number of transitive orientations of the incomparability graph of the order (the unique realizer $(\prec, \prec)$ of a chain is fixed by the swap, whereas for any other order the reversal of an orientation is a different orientation); it is $n!/2$ for an antichain, and 1 whenever the incomparability graph has a unique transitive orientation up to reversal (for instance when exactly one pair is unrelated). Hence the kernel class is a single orbit of the group of Proposition 5.1 for some configurations and a union of several such orbits for others; both cases occur for every $n \geq 3$.

Proof. Under the standing assumption the order is defined by strict inequalities between coordinates, so it is locally constant and the class is open. Two configurations with the same pair of linear orders $(\prec_u, \prec_v)$ are related by a pair (f, g) of increasing bijections of $(0, 1)$ (interpolate piecewise linearly between the sorted u -values of one and of the other, and likewise for v), and conversely (f, g) preserves both linear orders while the swap exchanges them; so the conformal orbit modulo the swap of a configuration is exactly the set of configurations with the same realizer modulo the swap, and the kernel class is the union of these sets over the realizers of the order. A realizer is determined by the linear order $\prec_u$ alone, since $\prec_v$ is then the reverse of $\prec_u$ on unrelated pairs and agrees with the order on related ones. Let T be the restriction of $\prec_u$ to the unrelated pairs; T orients every edge of the incomparability graph, and it is a transitive orientation: if $a \parallel b$, $b \parallel c$ and $a \prec_u b \prec_u c$, then $v_a > v_b > v_c$, so $a \parallel c$ and $(a, c) \in T$. Conversely, let T be any transitive orientation of the incomparability graph. Then $\prec \cup T$ is a linear order: it is total and antisymmetric because every pair is either related by $\prec$ or oriented by T , and it is transitive because, besides the two pure cases, the mixed case $a \prec b$, $(b, c) \in T$ forces $(a, c) \in \prec \cup T$: the alternatives are $c \prec a$, which gives $c \prec b$ and contradicts $b \parallel c$, and $(c, a) \in T$, which gives $(b, a) \in T$ by transitivity of T and contradicts $a \prec b$; the mirror case $(a, b) \in T$, $b \prec c$ is handled in the same way. Taking $\prec_u = \prec \cup T$ and $\prec_v = \prec \cup T^{-1}$ therefore gives a realizer $(\prec_u \cap \prec_v = \prec)$ because T is antisymmetric), so realizers correspond bijectively to transitive orientations T , and reversing T gives the swapped realizer. For a chain the incomparability graph is empty, its only orientation is the empty one and the unique realizer $(\prec, \prec)$ is its own swap, so the count is 1; for any other order $T \neq T^{-1}$, the orientations pair off and the count is half their number. For an antichain every linear order of $[n]$ is a transitive orientation, giving $n!/2$; when exactly one pair is unrelated the two orientations are reverses of each other. As an explicit example of distinct orbits, three points with $u_2 < u_1 < u_3$ and $v_3 < v_1 < v_2$, and three with $u_1 < u_2 < u_3$ and $v_3 < v_2 < v_1$, are both antichains and lie in different orbits, since (f, g) preserves the u -order and the swap only reverses it. $\square$

The u -order is thus a conformal invariant that is not in general an order invariant: what the order alone leaves undetermined at finite n is exactly the choice of a realizer, and the coordinate reconstruction of E5d below makes that choice with the help of the counts. How much is left undetermined for uniform samples is measured in E5e below: for a substantial fraction of samples the realizer is unique, so the order determines the configuration exactly up to the group of Proposition 5.1, and for the rest the ambiguity is a handful of binary choices.

Computational check (E5a). For 200 uniform samples of $n = 300$ points, random pairs (f, g) of increasing bijections preserved the order in all 200 cases while changing the proper times of related pairs by 23% (median over samples of the per-sample median relative change); random boosts preserved the order in all 200 cases and the proper times to within a relative 10^{-9} ; a Euclidean rotation of the (t, x) plane by 5° to 30° , which is not a conformal map, changed the relation of a median 19.6% of the pairs (never fewer than 5.9%).

D	n	samples	comparable fraction	expected	dimension estimate
2	400	60	0.4989 ± 0.0185	0.5000	2.00 ± 0.05
3	400	60	0.2298 ± 0.0167	0.2286	3.00 ± 0.09
4	400	60	0.1005 ± 0.0121	0.1000	4.00 ± 0.14

Table 5: E5b. Fraction of comparable pairs (mean $\pm$ sd over samples) in uniform samples of causal diamonds of Minkowski space of dimension D , the expected value, and the Myrheim–Meyer estimate of D .

Order alone: dimension

The fraction of comparable pairs in a uniform sample of a causal diamond is an order invariant with a known expectation in D -dimensional Minkowski space, $\Gamma(D+1)\Gamma(D/2)/(2\Gamma(3D/2))$, twice the Myrheim–Meyer ratio $\langle R \rangle/N^2$ of ordered relations to ordered pairs, which is the basis of the Myrheim–Meyer dimension estimator [29, 30]. Table 5 reports the observed fractions for $n = 400$ points in diamonds of dimension 2, 3 and 4 (60 samples each) next to the expected values, and the dimension estimate obtained by inverting the formula. The three dimensions are separated by many standard deviations. This is the Lorentzian analogue of the count of ordinal patterns in Section 3: a bare order knows the dimension of the space it was sampled from.

Order plus number: proper time

For related points $p \prec q$ let $L(p, q)$ be the number of links of the longest chain from p to q , an order invariant. In the 1+1 diamond a chain is an increasing sequence in both coordinates, so $L(p, q) - 1$ is the length of the longest increasing subsequence of the points inside the rectangle spanned by p and q , whose expected number is $N = (n-2)\tau(p, q)^2$. Uniform points in a rectangle induce a uniformly random permutation, so by the Vershik–Kerov and Logan–Shepp theorem the longest increasing subsequence of N of them has length $2\sqrt{N}(1 + o(1))$ [26, 43]; Brightwell and Gregory [8] and Bollobás and Brightwell [5] give the corresponding statements for sprinklings in Minkowski space of any dimension, and Bollobás and Brightwell [6] the concentration of the chain length. Hence

$$L(p, q) \approx 2\sqrt{n}\tau(p, q),$$

and $\hat{\tau} = L/(2\sqrt{n})$ estimates the proper time from the order and the density alone. Table 6 reports, for n from 250 to 2000 and 39 000 related pairs in total, the correlation between L and τ , the least-squares slope through the origin of L against $\sqrt{n}\tau$, and the median relative error of $\hat{\tau}$. At $n = 2000$ the correlation is 0.990 and the slope 1.85 (it is 1.73 at $n = 250$; the mean length of the longest increasing subsequence is $2\sqrt{N}$ minus a correction of order $N^{1/6}$ with a negative constant [3], which is why all slopes lie below 2); the median relative error of $\hat{\tau}$ over pairs with $\tau > 1/4$ falls from 14% to 8%. The same table shows what “number” contributes: after a random conformal reparametrisation (f, g) the order, and hence L , is unchanged (this is asserted in the code for every pair), but the same $\hat{\tau}$ misses the proper times τ' of the reparametrised configuration by a median of 17% at $n = 2000$ (and their correlation with L drops to 0.91); given E5a this is expected rather than independent evidence, and it is reported as an illustration. The chain length estimates the proper time of the frame in which the sample is uniform: the counting measure selects the conformal factor, and without it the order alone cannot.

Order plus number: coordinates

Under uniform density, $|\downarrow q| \approx (n-1)u_q v_q$ and $|\uparrow q| \approx (n-1)(1-u_q)(1-v_q)$ for the past and future of q , so the order and the count give $s = u_q + v_q$ and $p = u_q v_q$ and hence the unordered pair $\{u_q, v_q\}$ as the roots of $x^2 - sx + p$. Which root is u_q is a sign per point. Both kinds of pairs constrain the relative signs: a comparable pair must satisfy $(u_r - u_q)(v_r - v_q) > 0$ and an

n	pairs	$\text{corr}(L, \tau)$	$\text{corr}(L, \tau')$	slope	rel. err. vs τ	rel. err. vs τ'
250	12000	0.967	0.856	1.732	0.143	0.229
500	12000	0.978	0.866	1.739	0.138	0.217
1000	9000	0.988	0.871	1.792	0.104	0.217
2000	6000	0.990	0.909	1.854	0.082	0.166

Table 6: E5c. Longest-chain length L against proper time τ in uniform samples of the 1+1 diamond; τ' is the proper time after a random conformal reparametrisation, under which L is unchanged. Slope: least-squares coefficient through the origin of L on $\sqrt{n}\tau$ (limit 2). Last two columns: median relative error of $\hat{\tau} = L/(2\sqrt{n})$ against τ and against τ' , over pairs with $\tau > \frac{1}{4}$.

n	samples	median RMSE	max RMSE	median RMSE $\times \sqrt{n}$	order disagreement
250	10	0.0732	0.0971	1.16	7.62%
500	10	0.0494	0.0622	1.10	5.24%
1000	6	0.0415	0.0544	1.31	4.53%
2000	4	0.0312	0.0379	1.39	3.68%

Table 7: E5d. Root-mean-square error of the coordinates (u, v) recovered from the causal order and the past/future counts (after fixing the global swap), and the fraction of pairs whose comparability differs between the true and the reconstructed configuration.

unrelated pair $r \parallel q$ must satisfy $(u_r - u_q)(v_r - v_q) < 0$, and whenever exactly one of the two relative orientations satisfies the constraint the pair votes “same” or “opposite” (an ablation that uses the votes of unrelated pairs only changes the median error at $n = 2000$ by +0.4%, from 0.0312 to 0.0313); the votes are resolved globally by the leading eigenvector of the vote matrix, and the remaining global swap is the parity of Proposition 5.1, which no order can fix. The result is a reconstruction of every coordinate, not merely of the configuration up to a group, because the boundary of the diamond is part of the sampling hypothesis. Table 7 reports the root-mean-square coordinate error over n from 250 to 2000: the median falls from 0.073 to 0.031, roughly as $n^{-0.39}$ over the range, slower than the $n^{-1/2}$ fluctuation of the counts; the loss is expected from the ill-conditioning of the roots near $u = v$, where the discriminant $s^2 - 4p$ vanishes and an $O(n^{-1/2})$ error in the counts becomes an $O(n^{-1/4})$ error in the roots (the rate is not established here). The order recomputed from the reconstructed coordinates disagrees with the true order on 3.68% of the pairs at $n = 2000$. Figure 3 shows one instance.




Figure 3: Left (E5d): a uniform sample of the 1+1 diamond and the coordinates recovered from its causal order and counts. Right (E5c): relative error of the chain-length estimate of proper time and relative deviation of the fitted slope from 2, against n .

Order alone: how many realizers?

Proposition 5.2 reduces the finite- n ambiguity of the order to the count of its realizers modulo the swap. E5e computes this count for uniform samples by enumerating the transitive orientations of the incomparability graph over its implication colour classes (Golumbic’s Γ -relation [14, 15]; every transitive orientation contains each implication class or its reverse, so the 2^k choices over the k classes exhaust the candidates) and checking transitivity, cross-checked against a brute-force enumeration of pairs of linear extensions at $n = 6$ (30 of 30 agree). Table 8 shows that the realizer is unique for a fraction 0.34 of the samples at $n = 300$, and that at most eight realizers occur in a fraction 0.98: for those samples the causal order determines the configuration exactly up to the group of Proposition 5.1, or up to a few binary choices. The extra realizers typically come from pairs of points that are adjacent in both coordinate orders with opposite directions. Such a pair consists of incomparable twins (every other point relates to both in the same way) and is therefore a module of the incomparability graph, which a transitive orientation can orient in two ways independently of the rest; in the typical situation for large n , made precise below, these pairs are the only modules that affect the count, and Lemma 5.5 with Theorem 5.3 gives $2 \cdot 2^N$ transitive orientations, hence 2^N realizers modulo the swap, where N is the number of such pairs. N is the number of descending successions of the permutation that sends u -ranks to v -ranks; it has mean $(n - 1)/n$ and is asymptotically Poisson with mean one, a classical result on runs of consecutive elements in random permutations [19, 45] (Theorem 5.6(b) below gives its factorial moments exactly). These pairs are not the only source of ambiguity (at $n = 10$ the largest count observed is 72, which is not a power of two and comes from larger modules of mutually unrelated points), but the other sources become rare as n grows. Version 0.5 of this draft stated the resulting law as a conjecture; it is proved next.

The realizer law. Sort the points of a configuration by u and let $\pi(i)$ be the v -rank of the i -th point. For independent uniform points u and v are independent and continuous, so π is uniform on the symmetric group S_n , and the i -th point precedes the j -th iff $i < j$ and $\pi(i) < \pi(j)$. The unrelated pairs are the inversions of π , so the incomparability graph of the order is the permutation graph G_π of π (vertices $1, \dots, n$, an edge ij iff $(i - j)(\pi(i) - \pi(j)) < 0$), and by Proposition 5.2 the number of realizers modulo the swap is $R(\pi) = 1$ if π is the identity and $R(\pi) = t(G_\pi)/2$ otherwise, where $t(G)$ is the number of transitive orientations of G . A pair of points adjacent in both coordinate orders with opposite directions is an index i with $\pi(i + 1) = \pi(i) - 1$ (a *descending succession*); $N(\pi)$ denotes their number. An *interval* of π is a set of consecutive positions whose values are consecutive integers; it is trivial if it has 1 or n elements, and π is *simple* if all its intervals are trivial. A *module* of a graph is a set M of vertices such that every vertex outside M is adjacent to all or to none of M ; it is trivial if $|M| \leq 1$ or M is the whole vertex set, and the graph is *prime* if all its modules are trivial. For $\sigma \in S_m$ and permutations $\alpha_1, \dots, \alpha_m$, the inflation $\sigma[\alpha_1, \dots, \alpha_m]$ replaces the j -th entry of σ by a block of consecutive positions with consecutive values in the pattern α_j , the blocks being ordered by value as the entries of σ . We use one result from the theory of comparability graphs that is not proved here.

Theorem 5.3 (Gallai [13]). *A prime comparability graph with at least three vertices has exactly two transitive orientations, one the reverse of the other. [literature]*

The restriction to at least three vertices excludes the two-vertex graph without edges, which is prime in the sense above and has only the empty orientation. Permutation graphs are comparability graphs (G_π is the comparability graph of the order $i < j, \pi(i) > \pi(j)$), so Theorem 5.3 applies to them; it was also checked numerically on all 3318 simple permutations of length 4 to 8 (Appendix B). In a transitive orientation we write $x \rightarrow y$; the basic forcing rule [13, 15] is that if x is adjacent to a and b and a, b are not adjacent, then $x \rightarrow a$ iff $x \rightarrow b$, since $a \rightarrow x \rightarrow b$, like $b \rightarrow x \rightarrow a$, would force the edge ab by transitivity. Not every module of

G_π is an interval of π (for the identity, G_π has no edges and $\{1, 3\}$ is a module), but the two notions agree where it matters.

Lemma 5.4 (Intervals and modules). *Every interval of π is a module of G_π . Conversely, if $n \geq 3$ and G_π has a module M with $2 \leq |M| \leq n-1$, then π has an interval with between 2 and $n-1$ elements. Hence, for $n \geq 3$, π is simple iff G_π is prime, and no permutation of length 3 is simple. [known, e.g. [9]; proof included for completeness]*

Lemma 5.5 (Inflation of a simple permutation). *If $\sigma \in S_m$ is simple, $m \geq 4$, and $\pi = \sigma[\alpha_1, \dots, \alpha_m]$, then $t(G_\pi) = 2 \prod_{j=1}^m t(G_{\alpha_j})$. [classical, the prime-node case of Gallai's decomposition [13]; proof included for completeness]*

Theorem 5.6 (Realizer law). *Let R_n be the number of realizers modulo the swap of the causal order of n independent uniform points of the 1+1 diamond, and N_n the number of pairs of points adjacent in both coordinate orders with opposite directions. For $n \geq 20$:*

$$(a) \Pr(R_n \neq 2^{N_n}) \leq 10/n + 166/n^2;$$

$$(b) \mathbb{E}[N_n(N_n - 1) \cdots (N_n - r + 1)] = 1 - r/n \text{ for } 0 \leq r \leq n-1; \text{ hence } N_n \rightarrow Z \sim \text{Po}(1) \text{ in law.}$$

[proved here, using Theorem 5.3]

Consequently $R_n \rightarrow 2^Z$ in law. Theorem 5.8 and Proposition 5.10 below give the rate $O(1/n)$ with explicit constants; in particular the fractions of samples with one, two and at most eight realizers tend to e^{-1} , e^{-1} and $\frac{8}{3}e^{-1}$, each with error $O(1/n)$.

Proposition 5.7 (Where the exceptions come from). $\Pr(R_n \neq 2^{N_n}) = 3(n-2)/(n(n-1)) + 2/n + O(n^{-2}) = 5/n + O(n^{-2})$. *Up to an event of probability $O(n^{-2})$, an exception is a configuration with a single interval of three elements in one of the patterns 321 (then $R_n = \frac{3}{2} 2^{N_n}$), 231 or 312 (then $R_n = 2 \cdot 2^{N_n}$), or with $\pi(1) = n$ or $\pi(n) = 1$, i.e. a point related to no other one and extreme in both coordinate orders (then $R_n = 2 \cdot 2^{N_n}$). [proved here, using Theorem 5.3]*

Sharp rates. Theorem 5.6 gives no rate for N_n , and its bound (a) is about twice the true rate; the exact first-order terms are as follows. Write $p_k = \Pr(N_n = k)$ and $q_k = e^{-1}/k!$ for $k \geq 0$, $q_k = 0$ for $k < 0$, and $Z \sim \text{Po}(1)$. The three results below were added in version 0.7; they are proved in Appendix B and checked numerically there, and they were checked by an independent internal referee in round 5 (step by step, and by exhaustive enumeration of S_n for $n \leq 11$).

Theorem 5.8 (Sharp Poisson rate for N_n). (i) *For $n \geq 1$ and $0 \leq k \leq n-1$,*

$$p_k = q_k \left(1 - \frac{k-1}{n}\right) + r_{n,k}, \quad |r_{n,k}| \leq \frac{1}{n k! (n-k+1)!}.$$

Equivalently $p_k = q_k + \nu(k)/n + r_{n,k}$ with $\nu(k) = q_k - q_{k-1} = -(k-1)q_k$; for fixed k the remainder is smaller than any power of n .

(ii) *For $n \geq 4$, $d_{\text{TV}}(N_n, \text{Po}(1)) = (p_0 - q_0) + (p_1 - q_1)^+$, and*

$$\left| d_{\text{TV}}(N_n, \text{Po}(1)) - \frac{e^{-1}}{n} \right| < \frac{2}{n \cdot n!}.$$

Hence $n d_{\text{TV}}(N_n, \text{Po}(1)) \rightarrow c_1 = \frac{e^{-1}}{2} \sum_{k \geq 0} |k-1|/k! = e^{-1}$. (The identity and the bound also hold for $n \leq 3$, by direct computation; $n \geq 4$ is used only in the proof.)

[proved here, an elementary consequence of the classical exact law [19]; checked by an independent internal referee (round 5)]

Theorem 5.9 (First-order law of the realizer count). *Let μ be the signed measure on $\{2^k\}_{k \geq 0} \cup \{3 \cdot 2^k\}_{k \geq 1}$ given by*

$$\mu(\{2^k\}) = -3q_k + 3q_{k-1} - q_{k-2} = -\frac{e^{-1}}{k!}(k-1)(k-3), \quad \mu(\{3 \cdot 2^k\}) = q_{k-1} = \frac{e^{-1}}{(k-1)!}.$$

Then μ has total mass 0, and uniformly over sets A of positive reals,

$$\Pr(R_n \in A) = \Pr(2^Z \in A) + \frac{\mu(A)}{n} + O(n^{-2}).$$

Consequently

$$d_{\text{TV}}(R_n, 2^Z) = \frac{c_2}{n} + O(n^{-2}), \quad c_2 = \frac{1}{2}\|\mu\| = 1 + \frac{1}{2e} = 1.1839\dots$$

In particular, with errors $O(n^{-2})$: $\Pr(R_n = 1) = e^{-1}(1 - 3/n)$, $\Pr(R_n = 2) = e^{-1}$, $\Pr(R_n = 4) = \frac{1}{2}e^{-1}(1 + 1/n)$, $\Pr(R_n = 6) = \Pr(R_n = 12) = e^{-1}/n$, $\Pr(R_n \leq 8) = \frac{8}{3}e^{-1} - \frac{3}{2}e^{-1}/n$, and $\Pr(R_n \text{ is not a power of } 2) = 1/n$. [proved here, using Theorem 5.3; checked by an independent internal referee (round 5)]

Proposition 5.10 (Explicit bounds). *For $n \geq 20$,*

$$\Pr(R_n \neq 2^{N_n}) \leq \frac{5}{n} + \frac{220}{n^2}, \quad d_{\text{TV}}(R_n, 2^Z) \leq \frac{5 + e^{-1}}{n} + \frac{221}{n^2}.$$

[proved here, using Theorem 5.3; checked by an independent internal referee (round 5)]

The measure μ is the sum of three contributions: the first-order term ν of the law of N_n (a mean shift, $\mathbb{E}N_n = 1 - 1/n$); the configurations with a three-element interval of pattern 321, which have probability $1/n + O(n^{-2})$ and $R_n = 6 \cdot 2^{N_n-2}$; and the four configurations with $R_n = 2 \cdot 2^{N_n}$ (patterns 231, 312, $\pi(1) = n$, $\pi(n) = 1$), of total probability $4/n + O(n^{-2})$. The latter move mass from 2^j to 2^{j+1} , atoms that 2^Z charges anyway, so they largely cancel in the total variation distance: the coupling bound $\Pr(R_n \neq 2^{N_n}) + d_{\text{TV}}(N_n, Z)$ has first-order constant $5 + e^{-1}$, the true one is $1 + 1/(2e)$. The positive part of μ consists of the atoms $3 \cdot 2^k$ (mass 1) and the atom 4 (mass $e^{-1}/2$).

Second order. The results below were added in version 0.8; they are proved in Appendix B and checked numerically there, but have not yet been checked by an independent referee. For $\lambda > 0$ write $q_k(\lambda) = e^{-\lambda}\lambda^k/k!$, so that $q_k = q_k(1)$, and $\tilde{q}_k = q_k(1 - 1/n)$; for integers $j, k \geq 0$ let $c_j(k) = \sum_{i=0}^j (-1)^i \binom{k}{i} / (j-i)!$ be the coefficient of x^j in $e^x(1-x)^k$, so that $c_0 = 1$, $c_1(k) = 1 - k$, $c_2(k) = \frac{1}{2}(k^2 - 3k + 1)$ and $\tilde{q}_k = q_k \sum_{j \geq 0} c_j(k)n^{-j}$. The remainder $r_{n,k}$ is that of Theorem 5.8(i).

Theorem 5.11 (Second-order law of N_n). *(i) For $n \geq 1$ and $0 \leq r \leq n-1$, $\mathbb{E}[(N_n)_r] - (1 - 1/n)^r = -\binom{r}{2}n^{-2} + \theta\binom{r}{3}n^{-3}$ with $\theta \in [0, 1]$, where $(N)_r = N(N-1)\cdots(N-r+1)$; and for $0 \leq k \leq n-1$,*

$$p_k = \tilde{q}_k - q_k \sum_{j \geq 2} \frac{c_j(k)}{n^j} + r_{n,k} = \tilde{q}_k - q_k \frac{k^2 - 3k + 1}{2n^2} + O_k(n^{-3}).$$

(ii) For $n \geq 3$,

$$d_{\text{TV}}(N_n, \text{Po}(1 - 1/n)) = \Phi(1/n) + \varepsilon_n, \quad |\varepsilon_n| \leq \frac{2^{n+1} - n - 2}{n(n+1)!},$$

where $\Phi(x) = \frac{1}{2}e^{-1}(3-x)(1 - (1-x)e^x)$.

(iii) For $n \geq 3$, $0 < n^2 \Phi(1/n) - \frac{3}{4e} - \frac{1}{4en} \leq \frac{1}{48en^2}$. Hence

$$n^2 d_{\text{TV}}(N_n, \text{Po}(1 - 1/n)) = \frac{3}{4e} + \frac{1}{4en} + O(n^{-2}) \longrightarrow \frac{3}{4e} = 0.27591 \dots$$

[proved here; not yet independently refereed]

The first-order measure $q + \nu/n$ of Theorem 5.8 has factorial moments exactly $1 - r/n$ for every r (since $\sum_k \nu(k)(k)_r = 1 - \mathbb{E}[(Z + 1)_r] = -r$), consistent with its super-exponentially small distance to the law of N_n ; $\text{Po}(1 - 1/n)$ matches the mean but not the second factorial moment, and the mismatch $-\binom{r}{2}/n^2$ is the term $-q_k c_2(k)/n^2$ of (i). Since $\sum_k q_k c_2(k) = 0$ and $c_2(k) < 0$ only for $k = 1, 2$, $\frac{1}{2} \sum_k q_k |c_2(k)| = \frac{3}{4e}$, the constant of (iii); version 0.7 reported this limit only numerically. The mean-matching Poisson law is not the closest one at this order.

Theorem 5.12 (Best Poisson approximation of N_n). For real a let $L(a) = \sum_{k \geq 0} q_k |c_2(k) + a(k - 1)|$.

(i) For every fixed a , $n^2 d_{\text{TV}}(N_n, \text{Po}(1 - 1/n + a/n^2)) \rightarrow L(a)/2$.

(ii) $L(a) \geq e^{-1}$, with equality if and only if $a = \frac{1}{2}$; for $a \geq -\frac{1}{4}$, $L(a) = e^{-1}(1 + (2a - 1)^+ + (\frac{1}{2} - a)^+)$, so $L(0)/2 = 3/(4e)$.

(iii) $n^2 \inf_{\lambda > 0} d_{\text{TV}}(N_n, \text{Po}(\lambda)) \rightarrow 1/(2e) = 0.18394 \dots$, and $\lambda_n = 1 - 1/n + 1/(2n^2)$ attains this limit.

[proved here; not yet independently refereed]

Proposition 5.13 (Explicit constant in Theorem 5.9). For every $n \geq 22$ and every set A of positive reals, $|\Pr(R_n \in A) - \Pr(2^Z \in A) - \mu(A)/n| \leq B(n)$, where $B(n)$ is the explicit function (2) of Appendix B, and $n^2 B(n)$ is non-increasing. Consequently

$$\left| d_{\text{TV}}(R_n, 2^Z) - \frac{c_2}{n} \right| \leq \frac{422}{n^2} \quad (n \geq 22), \quad \leq \frac{286}{n^2} \quad (n \geq 100),$$

and $n^2 B(n) \rightarrow 259 + 10/e = 262.68 \dots$ [proved here, using Theorem 5.3; not yet independently refereed]

The constant is far from sharp: from the exact law of R_n for $n \leq 600$ (Appendix B), $n^2 |d_{\text{TV}}(R_n, 2^Z) - c_2/n|$ is at most 5.72 for $22 \leq n \leq 600$ and at most 8.01 for $4 \leq n \leq 600$. Most of the gap comes from the bad event $\mathcal{E}$ of Lemma B.2 and the occurrences inside it, which the proof charges in full ($90 + 150$ of the limit $259 + 10/e$).

Remark 5.14 (What remains open). The n^{-2} term of the law of R_n is not proved. Numerically (exact laws for $n \leq 600$, polynomial fits in $1/n$ on $200 \leq n \leq 600$), $n^2(\Pr(R_n = x) - \Pr(2^Z = x) - \mu(x)/n)$ converges for each of the 24 atoms $x \in \{2^k, 3 \cdot 2^k\}$ with $k < 12$, $n^2 \Pr(R_n \notin \{2^k, 3 \cdot 2^k\}_{k \geq 0})$ approaches 0.500000, and

$$n^2 \left(d_{\text{TV}}(R_n, 2^Z) - \frac{c_2}{n} \right) \rightarrow \kappa \approx 4.29707,$$

the fits of degree 2, 3 and 4 giving 4.297100, 4.297071 and 4.297072. [open; κ computationally verified (extrapolation of exact values), not proved]

Conjecture 5.15 (Closed form of the second-order term). Let $\mu_2(x) = \lim_n n^2(\Pr(R_n = x) - \Pr(2^Z = x) - \mu(x)/n)$. Then $\mu_2(2^k) = q_k[12\binom{k}{4} - 12\binom{k}{3} + 12\binom{k}{2} - 3k - 5]$, $\mu_2(3 \cdot 2^k) = (k + 1)q_{k-1}$ for $k \geq 1$, and $\mu_2(3) = 0$; with the mass $\frac{1}{2}$ outside these atoms this gives $\kappa = \mu_2(4) + \mu_2(8) + \sum_{k \geq 1} \mu_2(3 \cdot 2^k) + \frac{1}{2} = \frac{7}{2} + \frac{13}{6e} = 4.297072 \dots$ [conjectural: read off the numbers, which agree with these forms within $1.4 \cdot 10^{-6}$ at the 24 atoms of Remark 5.14]

Outline of the proofs. The complete proofs are in Appendix B. Lemma 5.4 is the usual correspondence between intervals and modules, and Lemma 5.5 is the prime-node case of Gallai’s decomposition: the blocks are modules, the quotient is prime, and a transitive orientation is one of the two of the quotient together with independent ones of the blocks. Theorem 5.6 combines them with a first-moment bound on intervals and an exact inclusion–exclusion for successions: if π has no interval with between 3 and $n - 1$ elements, contracting its successions, which are disjoint, gives a simple permutation of length at least 4, and $R_n = 2^{N_n}$; the expected number of intervals with k elements is $(n - k + 1)^2 k! (n - k)! / n!$, that is $6/n + O(n^{-2})$ for $k = 3$, $4/n$ for $k = n - 1$ and $O(n^{-2})$ in total for the other k . This first-moment computation is classical: the number of non-trivial intervals of a uniform permutation converges to a Poisson law of mean 2 [10], and a uniform permutation is simple with probability $e^{-2}(1 - 4/n + O(n^{-2}))$ [2]. What we have not found in the literature, and claim as new, is the assembly of these facts in the causal translation, the explicit constants, and the classification of the exceptions in Proposition 5.7: up to probability $O(n^{-2})$ there is at most one interval with between 3 and $n - 1$ elements, it has 3 or $n - 1$ elements, and its pattern, or the extreme point it leaves out, multiplies the count by $\frac{3}{2}$ or 2. Theorem 5.8 evaluates the alternating tail of the exact inclusion–exclusion of Step 3; Theorem 5.9 conditions on the single exceptional configuration of Proposition 5.7, contracts it, and bounds the event with two or more configurations by second moments. Theorems 5.11 and 5.12 compare the exact law of N_n with Poisson laws through the signs of $e^x(1 - x)^k - 1 - (1 - k)x$, and Proposition 5.13 bounds each error term of the proof of Theorem 5.9.

Sharpness and data. The bound in Theorem 5.6(a) counts every interval and is about twice the true rate $5/n$ of Proposition 5.7; by Theorem 5.8, $d_{TV}(N_n, \text{Po}(1))$ equals e^{-1}/n up to $2/(n \cdot n!)$, which is why the numerical value $n d_{TV}(N_n, \text{Po}(1)) = 0.368$ was found at every n tested (Appendix B). The limits 0.368, 0.368 and 0.981 are consistent with Table 8, where the three fractions at $n = 300$ are 0.34, 0.37 and 0.98, and with the independent samples of E5f (Table 9): the exact count equalled 2^N in a fraction 0.76 of the samples at $n = 20$, in 437 of the 450 samples with $n \geq 100$, and in 49 of 50 at $n = 500$; these samples are too small to resolve the $1/n$ terms, which the larger samples of Table 11 do. The ratios $R/2^N$ among the exceptions are $\frac{3}{2}$, 2, 3, 4 and 6 over all n , but only $\frac{3}{2}$ and 2 for $n \geq 100$, as Proposition 5.7 predicts; the larger ratios come from events of probability $O(n^{-2})$ (Remark B.1). With larger samples of uniform permutations (10000 to 20000 per n , Appendix B), $n \Pr(R_n \neq 2^{N_n})$ lies between 4.83 and 5.40 for n from 20 to 2000; 7 of its 7 confidence intervals at level 95% contain 5, and they widen from $[4.87, 5.11]$ at $n = 20$, where the agreement is partly fortuitous because the event of probability $O(n^{-2})$ is not small (it accounts for 2592 of 4921 exceptions at $n = 20$, Appendix B), to $[3.0, 7.0]$ at $n = 2000$. Thus the realizer of the causal order of a uniform sample is unique with probability tending to e^{-1} , and with probability $1 - O(1/n)$ it is determined up to N_n binary choices, one per pair of points adjacent in both coordinate orders with opposite directions, which can be made independently. This is the finite analogue of the continuum theorems; it differs from the Euclidean ordinal case, where the kernel class is also open (Proposition 3.5) but is never a finite union of similarity orbits, since these preserve all distance ratios and therefore have empty interior.

The dictionary. The two sequences line up term by term. Ranking of distances $\leftrightarrow$ causal order; monotone distortion of the metric $\leftrightarrow$ conformal reparametrisation; similarity class $\leftrightarrow$ conformal class; Theorem 3.7 $\leftrightarrow$ Hawking–King–McCarthy and Malament; the density hypothesis that makes non-metric scaling converge $\leftrightarrow$ the counting measure that fixes the conformal factor. The Euclidean chain is the test bed in which every step is a theorem or an exact computation; the Lorentzian sequence is where the same steps carry their historical weight. What the present draft does *not* do is state or test anything about curved backgrounds, about dimensions above $1+1$ beyond the ordering fraction, or about any physical system.

n	samples	not enumerated	unique	exactly two	at most eight	median colour classes
10	200	0	0.23	0.33	0.91	2
20	200	0	0.32	0.34	0.97	2
50	200	0	0.38	0.35	0.97	2
100	200	0	0.32	0.34	0.98	2
200	100	0	0.35	0.43	0.99	2
300	100	0	0.34	0.37	0.98	2

Table 8: E5e. Distribution of the number of realizers modulo the swap of the causal order of uniform samples of the 1+1 diamond (fractions of samples). Samples whose incomparability graph has more than 14 colour classes are not enumerated.

n	samples	count = 2^N	mean N	unique	exactly two	at most eight
20	200	151/200	0.88	0.32	0.41	0.95
50	200	179/200	0.90	0.36	0.41	0.97
100	200	192/200	0.91	0.40	0.35	0.98
200	100	98/100	0.90	0.38	0.43	0.98
300	100	98/100	1.03	0.32	0.40	0.98
500	50	49/50	0.92	0.36	0.44	0.98

Table 9: E5f. Check of Theorem 5.6 and Proposition 5.7 on independent uniform samples: number of samples whose realizer count modulo the swap equals 2^N , with N the number of pairs adjacent in both coordinate orders with opposite directions; the mean of N (limit 1); and the observed fractions of samples with one, two and at most eight realizers (limits 0.368, 0.368, 0.981). The exception rate is estimated with larger samples in Appendix B.

6 What can be claimed

Table 10 collects the claims of the paper with their status. The list is the deliverable promised by the research line: it separates what is proved from what is measured and from what is only expected.

Limitations. All objects are labelled; quotienting by relabelling turns the kernels into isomorphism problems that are not studied here. The sampling distribution is uniform on a cube or a diamond throughout; the finite- n rates reported (E2, E5c, E5d) are empirical for that distribution and are not proved. The ordinal reconstruction uses one solver with random starts; a local minimum would inflate a disparity, never deflate it. The Lorentzian setting is treated only in 1+1 dimensions, where the conformal group is exceptionally large, and only on a flat background; its second step is statistical, not a translation, so its “invariants” are consistent estimators rather than exact invariants. The realizer law (Theorem 5.6) rests on Gallai’s theorem, which is cited, not reproved. The sharp rates (Theorems 5.8, 5.9, Proposition 5.10) were checked by an internal, not an external, referee; the second-order results (Theorems 5.11, 5.12, Proposition 5.13) only by the author and numerically; the n^{-2} term of the realizer law is numerical.

Next steps for the research line. (a) Prove the n^{-4} law for the minimal gap under uniform sampling (a minimal-spacing statement for weakly dependent distances) and an upper bound on the diameter of the ordinal class in Procrustes disparity, to complement the lower bounds of E2c. (b) Extend E5d to 2+1 dimensions, where the conformal group is finite-dimensional and the count-based reconstruction must be replaced by an embedding of the order. (c) Replace the uniform-sampling hypothesis in Caus+N by a general density and characterise which conformal factors it can and cannot select. None of these is a physical claim.

claim	status
Distance matrix determines the configuration up to $E(d)$ (Thm. 3.1)	classical
Betweenness and congruence are empty on generic finite samples, $d \geq 2$ (Prop. 3.2)	proved here
no instance in 4000 exact random configurations; grids rich (E1)	verified
Ordinal fibres have non-empty interior; ordinal $\neq$ similarity at any finite $n \geq 3$ (Prop. 3.5)	proved here
four points realise 120 of 720 orderings on the line, all in the plane (E3)	verified
Ordinal pattern determines the configuration up to similarity as $n \rightarrow \infty$ (Thm. 3.7)	literature (informal statement)
ordinal reconstruction within 2.1×10^{-10} (Procrustes) of X at $n = 256$ in the plane (E2)	verified
Inradius of the ordinal class is $g/4$ when the two extremal pairs are disjoint (g = minimal gap between sorted distances), in $[g/4, g/2]$ in general (Prop. 3.8)	proved here
median $g \sim n^{-4.0}$ under uniform sampling; walk range shows a class much wider than $g/4$ (E2c)	verified, rate empirical
MST, k -NN graph, RNG, diameter pair are ordinal invariants, for distinct distances (Sec. 4)	proved here
Diameter, affine dimension, Gabriel graph are metric, not ordinal, at finite n (Sec. 4, E4b)	proved here / verified
Kernel of the causal-order translation in 1+1 contains the conformal maps and parity (Prop. 5.1)	proved here
Kernel classes are unions of conformal orbits over realizers; a single orbit iff the order has one realizer up to the swap (Prop. 5.2)	proved here
unique realizer for a fraction 0.34 of uniform samples at $n = 300$ (E5e)	verified
Prime comparability graphs have exactly two transitive orientations (Thm. 5.3)	literature
Intervals of π are modules of G_π , π simple iff G_π prime (Lemma 5.4); inflation of a simple permutation (Lemma 5.5)	known / classical; proofs included
Realizer count = 2^N except with probability $\leq 10/n + 166/n^2$ ($n \geq 20$), $N_n \rightarrow \text{Po}(1)$ (Thm. 5.6); exceptions have rate $5/n + O(n^{-2})$ and factors $\frac{3}{2}$ or 2 (Prop. 5.7)	proved here (uses Gallai and classical interval statistics)
count = 2^N in 437 of 450 samples with $n \geq 100$ (E5f); $n \Pr(R_n \neq 2^{N_n})$ between 4.83 and 5.40 for $20 \leq n \leq 2000$ (App. B)	verified
$d_{\text{TV}}(N_n, \text{Po}(1)) = e^{-1}/n$ up to $2/(n \cdot n!)$ (Thm. 5.8); $d_{\text{TV}}(R_n, 2^Z) = (1 + 1/(2e))/n + O(n^{-2})$ with an explicit first-order law (Thm. 5.9); explicit bounds for $n \geq 20$ (Prop. 5.10)	proved here (5.9, 5.10 use Gallai); checked by an independent internal referee (round 5)
Monte Carlo law of R_n agrees with the first-order law for $100 \leq n \leq 400$; at $n = 50$ the $O(n^{-2})$ term is visible (Table 11)	verified
$n^2 d_{\text{TV}}(N_n, \text{Po}(1 - 1/n)) \rightarrow 3/(4e)$, best Poisson constant $1/(2e)$ (Thms. 5.11, 5.12); $ d_{\text{TV}}(R_n, 2^Z) - c_2/n \leq 422/n^2$, $n \geq 22$ (Prop. 5.13)	proved here (5.13 uses Gallai); not yet independently refereed
exact law of R_n on the atoms 2^k , $3 \cdot 2^k$, $n \leq 600$ (App. B)	verified (exact computation)
$n^2(d_{\text{TV}}(R_n, 2^Z) - c_2/n) \rightarrow \kappa \approx 4.29707$ (Rem. 5.14); $\kappa = 7/2 + 13/(6e)$ (Conj. 5.15)	numerical, not proved; closed form conjectural
Causal order determines topology and conformal structure of the continuum	literature
Ordering fraction estimates the dimension of the diamond (E5b)	literature / verified
Longest chains estimate proper time from order and density, slope $\rightarrow 2$ (E5c)	literature / verified
Coordinates in the diamond recovered from order and counts, error decreasing with n (E5d)	verified; rate not established
Analogous statements for curved backgrounds or physical systems	not addressed

Table 10: Claims and their status. “Verified” means reproduced by the scripts of Appendix A with the fixed seed; it is not a proof.

A Reproducibility

The manuscript directory contains `experiments/formalism_chain.py` (E1–E4), `experiments/ordinal_class.py` (E2c), `experiments/lorentzian_chain.py` (E5a–E5e) and `experiments/realizer_law.py` (E5f). All four take a `-fast` flag for a reduced run. The full runs used seed 20260930 (E5: 20260930+1, E2c: 20260930+2, E5f: 20260930+3), Python 3.11.15, NumPy 2.4.6, SciPy 1.17.1 [44] and scikit-learn 1.9.1; E1–E4 took 1201 seconds, E5 315 seconds, E2c 168 seconds and E5f 128 seconds on four cores. The scripts write `results/results.json`, `results/results_class.json`, `results/results_lorentz.json`, `results/results_realizer_law.json` and Markdown tables; `experiments/make_numbers.py` converts them into the macros and table bodies used by this document, so that every number in the text is generated and none is typed. The numerical checks of the proofs of Section 5, `theory/check_realizer_law.py`, `theory/check_sharp_rate.py` and `theory/check_second_order.py`, are described in Appendix B; their outputs are frozen in `results/check_realizer_law_output.txt`, `results/check_sharp_rate_output.txt` and `results/check_second_order_output.txt` (the last two without timing lines; the last, with `results/second_order_results.json`, written by `experiments/freeze_second_order.py`) together with their SHA-256, which `make_numbers.py` verifies before reading them; the copies in `theory/` are logs with timings, and the files in `results/` are the reference.

E1. Coordinates are drawn uniformly from $\{0, \dots, 2^{30} - 1\}^d$; collinearity is tested by the vanishing of all 2×2 minors and strict betweenness by a negative inner product, congruence by equality of squared distances, all in 64-bit integers.

E2. Points uniform in $[0, 1]^d$; rank matrix of the pairwise distances; `sklearn.manifold.MDS` with `metric_mds=False`, precomputed input, random initialisation, four starts, at most 50 000 iterations, stress tolerance 10^{-13} ; Kendall discordance between the input and output rankings via `scipy.stats.kendalltau`; `scipy.spatial.procrustes` for the disparity.

E2c. For each configuration: g and its extremal pairs from the sorted distances; 20 random displacements of every point by a vector uniform in the Euclidean ball of radius $0.999 g/4$; the explicit displacement of Proposition 3.8(ii) with $g/4$ increased by the larger of a relative 10^{-6} and an absolute 10^{-12} , a margin above double-precision rounding of distances of order one; a Gaussian random walk with step size adapted towards an acceptance rate of about 0.3, accepted only when the ranking is unchanged, recording the maximum Procrustes disparity from X .

E3. Patterns are the permutations sorting the six pairwise distances; ties have probability zero under uniform sampling.

E4. $n = 30$, $d = 2$; rigid motions from a QR factorisation of a Gaussian matrix with random reflection; scale factors in $[0.1, 10]$; exponents $p \in [0.3, 3]$; equality of graphs as sets of edges, of vectors to 10^{-9} , of the numerical rank with relative eigenvalue threshold 10^{-6} .

E5. Uniform samples of the unit square in null coordinates (a fixed- n rather than Poisson sprinkling); diamonds in dimension 3 and 4 by rejection from the bounding box; longest chains by patience sorting on the points inside the rectangle spanned by a related pair; coordinate recovery as described in Section 5, with the sign vector taken from the leading eigenvector of the vote matrix, and an ablation restricted to unrelated pairs; realizer counts (E5e) by implication classes [14] of the incomparability graph computed with connected components, enumeration of the 2^k orientations over the k colour classes when $k \leq 14$, and a transitivity test by boolean matrix product.

E5f. Independent uniform samples of the diamond; for each sample, the exact realizer count of E5e and the number N of indices i with $\pi(i+1) = \pi(i) - 1$, where π sends the u -rank of a point to its v -rank; the exceptions to count $= 2^N$ are listed in `results/tables_realizer_law.md`.

B Proof of the realizer law

This appendix proves Lemmas 5.4 and 5.5, Theorem 5.6 and Proposition 5.7, then the sharp rates of Theorems 5.8 and 5.9 and Proposition 5.10, and finally the second-order results of Theorems 5.11 and 5.12 and Proposition 5.13, with the notation of Section 5. The only result used without proof is Theorem 5.3; the two lemmas are known (see their status labels) and are reproved for completeness.

Proof of Lemma 5.4. A point outside an interval lies on one side of it in position and on one side of it in value, so it forms an inversion with all of its points or with none. For the converse take such an M of minimal size. If $G_\pi[M]$ and its complement are both connected, M is an interval: let $x \notin M$ lie strictly between the smallest and the largest position of M ; if x is adjacent to no point of M , the points of M to its left lie below it and those to its right above it, so there is no edge of $G_\pi[M]$ between the two (non-empty) groups; if x is adjacent to all of M , every pair formed by a left and a right point of M is an edge, so the complement of $G_\pi[M]$ is disconnected; either way a contradiction. The same argument applied to values shows that the values of M are consecutive. If $G_\pi[M]$ is disconnected, each of its components is a module of G_π , so by minimality all are single points, M is independent, every pair in M is a module, and $|M| = 2$; if the complement of $G_\pi[M]$ is disconnected the same argument with co-components gives $|M| = 2$. It remains to treat a module $M = \{a, b\}$, $a < b$. If $\pi(a) < \pi(b)$ (no edge), a point x with $a < x < b$ adjacent to a and b would need $\pi(x) < \pi(a)$ and $\pi(x) > \pi(b)$, so it is adjacent to neither and $\pi(a) < \pi(x) < \pi(b)$; and a point outside $[a, b]$ with value between $\pi(a)$ and $\pi(b)$ would be adjacent to exactly one of a, b . So $[a, b]$ is an interval; if it is all of $[1, n]$ then $\pi(1) = 1$ and $[2, n]$ is an interval with $n - 1 \geq 2$ elements. The case $\pi(a) > \pi(b)$ is symmetric (with $\pi(1) = n$ in the extreme case). Finally, every permutation of length 3 has two consecutive positions with consecutive values. $\square$

Proof of Lemma 5.5. The blocks $B_1, \dots, B_m$ are intervals, hence modules (Lemma 5.4), and the quotient graph is G_σ , which is prime by Lemma 5.4; in particular it has no universal and no isolated vertex, and its only modules with at least two elements are the whole vertex set. Fix a transitive orientation and a block B , and let S be the set of vertices $y \notin B$ adjacent to B for which there are $b_1, b_2 \in B$ with $b_1 \rightarrow y \rightarrow b_2$. If $y \in S$ and $z \notin B$ is adjacent to y but not to B , the forcing rule at y gives $z \rightarrow y$ (from $b_1 \rightarrow y$) and $y \rightarrow z$ (from $y \rightarrow b_2$), which is absurd; so the neighbours of y outside B are adjacent to B . If $y \in S$ and $w \notin B$ is adjacent to B but not to y , the forcing rule at b_1 and at b_2 gives $b_1 \rightarrow w \rightarrow b_2$, so $w \in S$. Hence $B \cup S$ is a module. If $S \neq \emptyset$, the union of the blocks that meet $B \cup S$ is again a module (the union of two modules that intersect is a module) made of at least two blocks, so it corresponds to a module of G_σ with at least two vertices and is everything; then a vertex $z \notin B \cup S$ would make its block a universal or isolated vertex of G_σ , since z sees all of $B \cup S$ in the same way and $B \cup S$ meets every block, and $B \cup S$ equal to everything would make B universal in G_σ . Both are impossible, so $S = \emptyset$: for every $y \notin B$ adjacent to B all edges between y and B point the same way, and therefore all edges between two adjacent blocks point the same way. A transitive orientation of G_π is thus determined by its restriction to one representative per block, a transitive orientation of G_σ , and by its restrictions to the blocks. Conversely, lifting a transitive orientation of G_σ to all edges between blocks and choosing any transitive orientation inside each block gives a transitive orientation: for a path $a \rightarrow b \rightarrow c$, if the three points lie in different blocks or two consecutive ones share a block, transitivity follows from that of the quotient, a and c cannot share a block with b outside it (the quotient would contain $[a] \rightarrow [b] \rightarrow [a]$), and three points in

one block are handled inside the block. With $t(G_\sigma) = 2$ (Theorem 5.3, applicable since $m \geq 4$) the count follows. $\square$

Proof of Theorem 5.6. Step 1 (deterministic). Let $n \geq 5$ and suppose that π has no interval with k elements for $3 \leq k \leq n-1$. Two overlapping successions (descending or ascending) would form an interval with three elements, so the successions are disjoint. Contracting each of them to a point gives $\sigma \in S_m$ with $m \geq \lceil n/2 \rceil \geq 3$ and $\pi = \sigma[\alpha_1, \dots, \alpha_m]$, $\alpha_j \in \{1, 12, 21\}$. An interval of σ with between 2 and $m-1$ elements would expand to an interval of π with between 2 and $n-1$ elements, which is excluded if it has at least three elements and would be an uncontracted succession if it has two; so σ is simple, and $m \geq 4$ by Lemma 5.4. Lemma 5.5 with $t(G_1) = t(G_{12}) = 1$ and $t(G_{21}) = t(K_2) = 2$ gives $t(G_\pi) = 2 \cdot 2^{N(\pi)}$, and since π is not the identity, $R(\pi) = 2^{N(\pi)}$.

Step 2 (first moment). Let I_k be the number of intervals of π with k elements. Choosing the first position, the smallest value and the arrangements inside and outside gives $\mathbb{E}I_k = (n-k+1)^2 k! (n-k)! / n!$. Thus $\mathbb{E}I_3 = 6(n-2)/(n(n-1)) = 6/n - 6/(n(n-1)) \leq 6/n - 6/n^2$, $\mathbb{E}I_{n-1} = 4/n$, $\mathbb{E}I_{n-2} = 18/(n(n-1))$, $\mathbb{E}I_{n-3} = 96/(n(n-1)(n-2))$, $\mathbb{E}I_4 = 24(n-3)/(n(n-1)(n-2))$, $\mathbb{E}I_{n-4} = 600/(n(n-1)(n-2)(n-3))$, and for $5 \leq k \leq n-5$, $\mathbb{E}I_k \leq (n-4)^2 / \binom{n}{5}$, so these $n-9$ terms add up to at most $120(n-4)(n-9)/(n(n-1)(n-2)(n-3)) \leq 120/n^2$ (the last inequality is $7n^2 - 25n - 6 \geq 0$). For $n \geq 20$ the terms $\mathbb{E}I_4$, $\mathbb{E}I_{n-2}$, $\mathbb{E}I_{n-3}$, $\mathbb{E}I_{n-4}$ are at most $24/n^2$, $19/n^2$, $6/n^2$, $3/n^2$, and by Step 1 and the union bound

$$\Pr(R_n \neq 2^{N_n}) \leq \sum_{k=3}^{n-1} \mathbb{E}I_k \leq \frac{10}{n} + \frac{24 + 19 + 6 + 3 + 120 - 6}{n^2} = \frac{10}{n} + \frac{166}{n^2},$$

which is (a); the sum equals $10/n + O(n^{-2})$. This first-moment computation is classical: the number of non-trivial intervals of a uniform permutation converges to a Poisson law of mean 2 [10], and a uniform permutation is simple with probability $e^{-2}(1 - 4/n + O(n^{-2}))$ [2]; only the explicit constants for $n \geq 20$ are needed here.

Step 3 (factorial moments). Let A be a set of r indices in $\{1, \dots, n-1\}$. The event that $\pi(i+1) = \pi(i) - 1$ for all $i \in A$ says that each maximal run of consecutive indices of A , $i, \dots, i+l-1$, occupies positions $i, \dots, i+l$ with consecutive decreasing values; contracting each run to a point is a bijection onto S_{n-r} , so the event has probability $(n-r)!/n!$ whatever A is. Hence $\mathbb{E}[N_n(N_n-1) \cdots (N_n-r+1)] = r! \binom{n-1}{r} (n-r)!/n! = 1 - r/n$, the first part of (b). Since $N_n \leq n-1$, inclusion-exclusion is exact:

$$\Pr(N_n = j) = \frac{1}{j!} \sum_{s=0}^{n-1-j} \frac{(-1)^s}{s!} \left(1 - \frac{j+s}{n}\right),$$

and $N_n \rightarrow \text{Po}(1)$ by the method of moments. This is the classical Poisson limit of successions in a random permutation [19, 45], stated there for ascending successions; reversing the order of the positions exchanges the two kinds and preserves the uniform law. $\square$

Proof of Proposition 5.7. By Step 2 the probability that π has an interval with between 4 and $n-2$ elements is $O(n^{-2})$, and so is the probability of two intervals with 3 or $n-1$ elements: two disjoint three-element intervals have expected number $18(n-4)(n-5)(n-4)!/n! \leq 18n^2(n-4)!/n!$, two overlapping ones form an interval with four or five elements, a three-element interval together with $\pi(1) \in \{1, n\}$ or $\pi(n) \in \{1, n\}$ has probability at most $24/n^2 + 8/(n(n-1)(n-2))$ (given $\pi(1) = n$, an event of probability $1/n$, the rest of π is uniform, the expected number of three-element intervals avoiding position 1 is $6(n-3)/((n-1)(n-2)) \leq 6/n$ and $[1, 3]$ is an interval with probability $2/((n-1)(n-2))$; the four events $\pi(1) \in \{1, n\}$, $\pi(n) \in \{1, n\}$ are alike), and $\pi(1), \pi(n) \in \{1, n\}$ has probability $2/(n(n-1))$. Outside this event, if π has no interval with between 3 and $n-1$ elements, $R_n = 2^{N_n}$ by Step 1. If it has exactly one, J , with three elements,

no succession meets J without lying in it (the union would be an interval with four elements), and contracting J and the successions disjoint from it gives a simple permutation as in Step 1; Lemma 5.5 gives $R_n/2^{N_n} = t(G_\tau)/2^{N(\tau)}$ for the pattern τ of J : this is 1 for 123, 132, 213 ($t = 1, 2, 2$; $N(\tau) = 0, 1, 1$), $6/4 = \frac{3}{2}$ for 321 ($G_\tau = K_3$, $t = 3!$) and 2 for 231 and 312 (G_τ a path on three vertices, $t = 2$, $N(\tau) = 0$). If J has $n - 1$ elements, π is $1 \oplus \pi'$, $\pi' \oplus 1$, $1 \ominus \pi'$ or $\pi' \ominus 1$ (by $\oplus$ and $\ominus$ we mean the direct and skew sums, i.e. $\pi(1) = 1$, $\pi(n) = n$, $\pi(1) = n$, $\pi(n) = 1$), where π' satisfies the hypothesis of Step 1 and is neither a direct nor a skew sum (otherwise π would have a second interval with at least three elements); then G_π is $G_{\pi'}$ plus an isolated vertex in the first two cases and $G_{\pi'}$ plus a vertex adjacent to all others in the last two, and $N(\pi) = N(\pi')$ (a descending succession at the boundary, $\pi(2) = n - 1$ or $\pi(n - 1) = 2$, would create a second interval with $n - 2$ elements). The complement of $G_{\pi'}$, which is the permutation graph of π' with its values reversed, is connected: the components of a permutation graph occupy consecutive positions (if an edge xy of a path jumped over a position j , then j would be adjacent to x or to y), so a disconnected complement would make π' a skew sum. In the last two cases the forcing rule applied to the non-edges of $G_{\pi'}$ therefore makes the extra vertex a source or a sink, so $t(G_\pi) = 2t(G_{\pi'})$ and $R_n = 2 \cdot 2^{N_n}$; in the first two $R_n = 2^{N_n}$. Each three-element pattern has expected number of occurrences $(n - 2)/(n(n - 1))$ and $\Pr(\pi(1) = n) = \Pr(\pi(n) = 1) = 1/n$, so the upper bound follows, and Bonferroni's inequality with the $O(n^{-2})$ bounds on intersections gives the matching lower bound. $\square$

Remark B.1 (Exact count). For a direct sum, $G_{\alpha \oplus \beta}$ is the disjoint union of G_α and G_β ; for a skew sum $\alpha_1 \ominus \dots \ominus \alpha_k$ with skew-indecomposable parts, G_π is the join of the G_{α_i} , whose complements are connected, and the forcing rule orients all edges between two parts the same way. With Lemma 5.5 and the decomposition of every permutation with at least two elements as a direct sum, a skew sum, or an inflation of a simple permutation with at least four elements [1], induction gives, for $\pi \neq \text{id}$, $R(\pi)$ as half the product, over the nodes of the substitution decomposition tree of π , of 2 for a simple node, $k!$ for a skew-sum node with k children and 1 for a direct-sum node: Gallai's product formula [13] for the incomparability graph. Ratios $R/2^N$ other than 1, $\frac{3}{2}$ and 2 require an interval with at least four elements or two of the configurations of Proposition 5.7, events of probability $O(n^{-2})$, and the ratio is not multiplicative over such configurations: an interior block 4231, which contains a 231 and a 312, gives the ratio 6, not 4. The formula is not used in the proofs of Theorem 5.6 and Proposition 5.7; it is used to compute R in the numerical checks. [classical (Gallai; Albert–Atkinson); checked by brute force, see below]

Numerical check. `theory/check_realizer_law.py` (fixed seed, recorded in its output) compares the formula of Remark B.1 with two independent brute-force counts of realizers on all 873 permutations of length at most 6 and with one of them on the 5040 permutations of length 7 (0 mismatches in total) and with the colour-class enumeration of E5e on random permutations of larger length; confirms that each of the 3318 simple permutations of length 4 to 8 has a single realizer modulo the swap (exceptions: 0), as Theorem 5.3 and Lemma 5.4 require; verifies the factorial moments and the law of N_n for $n \leq 8$ in rational arithmetic; evaluates $\sum_{k=3}^{n-1} \mathbb{E}I_k/(10/n + 166/n^2)$ for $20 \leq n \leq 3000$ (largest value 0.9957); computes $d_{\text{TV}}(N_n, \text{Po}(1))$ from the exact law; and estimates $\Pr(R_n \neq 2^{N_n})$ on 10000 to 20000 uniform permutations for each n from 20 to 2000. Its output is frozen as `results/check_realizer_law_output.txt` (SHA-256 beginning 4e011b1e04312e47); the run took 87 seconds on one core.

Proofs of the sharp rates

Proof of Theorem 5.8. By Step 3 of the proof of Theorem 5.6, $p_k = \frac{1}{k!} \sum_{s=0}^{n-1-k} \frac{(-1)^s}{s!} (1 - \frac{k+s}{n})$. Since $\sum_{s \geq 0} (-1)^s/s! = e^{-1}$ and $\sum_{s \geq 0} (-1)^s s/s! = -e^{-1}$, the full series equals $e^{-1}(1 - (k-1)/n)$, so $p_k = q_k(1 - (k-1)/n) - T_k/k!$ with $T_k = \sum_{s \geq n-k} \frac{(-1)^s}{s!} (1 - \frac{k+s}{n})$. The term $s = n - k$ vanishes

and, with $s = n - k + m$, $T_k = -\frac{1}{n} \sum_{m \geq 1} (-1)^{n-k+m} \frac{m}{(n-k+m)!}$; the moduli decrease in m (their ratio is $\frac{m+1}{m(n-k+m+1)} \leq \frac{2}{n-k+2} < 1$), so $|T_k| \leq 1/(n(n-k+1)!)$, which is (i); $kq_k = q_{k-1}$ gives the second form.

(ii) $d_{TV}(N_n, \text{Po}(1)) = \sum_{k \geq 0} (p_k - q_k)^+$. For $k \geq n$, $p_k = 0 < q_k$. For $2 \leq k \leq n-1$, $p_k - q_k = -\frac{1}{n!} (e^{-1}(k-1) + nT_k)$ with $|nT_k| \leq 1/(n-k+1)!$, which is at most $\frac{1}{6} < e^{-1}$ if $k \leq n-2$ and at most $\frac{1}{2} < 2e^{-1} \leq e^{-1}(k-1)$ if $k = n-1 \geq 3$; so $p_k < q_k$. Moreover $p_0 - q_0 = e^{-1}/n - T_0 > 0$ and $p_1 - q_1 = -T_1$, so $d_{TV} = e^{-1}/n - T_0 + (-T_1)^+$, and $|T_0| + |T_1| \leq \frac{1}{n(n+1)!} + \frac{1}{n!} < \frac{2}{n \cdot n!}$. Finally $\sum_k (k-1)/k! = 0$ and the only negative term is $k = 0$, so $\sum_k |k-1|/k! = 2$. $\square$

For the next proofs call an *occurrence* either a pair $\omega = (J, \tau)$ of a window $J = [i, i+2]$ of three positions and a pattern $\tau \in \{321, 231, 312\}$, which *happens* when J is an interval of π with pattern τ (its window of values is then determined by π), with factor $c_\omega = \frac{3}{2}$ for 321 and $c_\omega = 2$ otherwise; or one of the events $\pi(1) = n$, $\pi(n) = 1$, with factor $c_\omega = 2$. A three-element occurrence happens with probability $(n-2)(n-3)!/n! = 1/(n(n-1))$, so each pattern has total weight $(n-2)/(n(n-1))$. Contracting J to the single position $j^* = i$, with the value of J 's lowest point and the values above those of J lowered by two, is a bijection from the permutations in which (J, τ) happens onto S_{n-2} ; so, given that (J, τ) happens, the contracted permutation σ is uniform on S_{n-2} and j^* is a fixed position. (Fixing the window of values as well would fix $\sigma(j^*)$, and σ would no longer be uniform.) Let $\mathcal{E}$ be the event that π has an interval with between 4 and $n-2$ elements or $I_3 + I_{n-1} \geq 2$, where I_k is the number of intervals with k elements. For $n \geq 6$, the proof of Proposition 5.7 shows that outside $\mathcal{E}$ at most one occurrence happens, $R_n = c_\omega 2^{N_n}$ if ω does and $R_n = 2^{N_n}$ if none does.

Lemma B.2 (The bad event). (a) For $n \geq 20$, $\Pr(\mathcal{E}) \leq 223/n^2$. (b) If $\#\omega$ is the number of occurrences that happen, $\mathbb{E}[\#\omega; \mathcal{E}] = O(n^{-2})$. [proved here; checked by an independent internal referee (round 5)]

Proof. (a) By Step 2 of the proof of Theorem 5.6, $\sum_{k=4}^{n-2} \mathbb{E}I_k \leq (24+19+6+3+120)/n^2 = 172/n^2$. Next, $\Pr(I_3 + I_{n-1} \geq 2) \leq \mathbb{E}\binom{I_3}{2} + \mathbb{E}[I_3 I_{n-1}] + \mathbb{E}\binom{I_{n-1}}{2}$. Two distinct three-element intervals are disjoint, overlap in two points (the union then has one of the patterns 1234, 1324, 4231, 4321) or overlap in one point (eight patterns of length five, the common point having the middle value), so $\mathbb{E}\binom{I_3}{2} = \frac{18(n-4)(n-5)}{n(n-1)(n-2)(n-3)} + \frac{4(n-3)}{n(n-1)(n-2)} + \frac{8(n-4)}{n(n-1)(n-2)(n-3)} \leq \frac{18+4+1}{n^2}$. Given $\pi(1) = 1$ the rest of π is uniform, with $6(n-3)/((n-1)(n-2))$ expected three-element intervals, and $[1, 3]$ is an interval with probability $2/((n-1)(n-2))$; with the three symmetric cases, $\mathbb{E}[I_3 I_{n-1}] = 4(6n-16)/(n(n-1)(n-2)) \leq 25/n^2$ (the difference is $(n^2 - 11n + 50)/(n^2(n-1)(n-2)) > 0$). Finally $\mathbb{E}\binom{I_{n-1}}{2} = \Pr(\{\pi(1), \pi(n)\} = \{1, n\}) = 2/(n(n-1)) \leq 3/n^2$. The total is $(172 + 23 + 25 + 3)/n^2$.

(b) Let $\omega = (J, \tau)$ be a three-element occurrence, $J = [i, i+2]$, and $\sigma \in S_m$, $m = n-2$, the contracted permutation, uniform given ω . On $\omega \cap \mathcal{E}$ there is an interval $K \neq J$ with $3 \leq |K| \leq n-1$. Either $K \cap J = \emptyset$, and K is an interval of σ with 3 to $m-1$ elements; or $K \supsetneq J$, and K is an interval of σ with 2 to $m-1$ elements containing j^* ; or K overlaps J partially, and then $U = J \cup K$ is an interval with J at one end and K is U minus one or two points of J , so there are at most two such K for each U ; U is an interval of σ with 2 to $m-1$ elements containing j^* , unless $U = [1, n]$, which requires $i \in \{1, n-2\}$. Let $s_m = \sum_{l=3}^{m-1} \mathbb{E}I_l(m)$, where $\mathbb{E}I_l(m)$ refers to a uniform permutation of length m , and $h_m = \sum_{l=2}^{m-1} \min(l, m-l+1) \mathbb{E}I_l(m)/(m-l+1)$, which bounds the expected number of intervals with 2 to $m-1$ elements containing a fixed position (at most $\min(l, m-l+1)$ windows of l positions contain it, and each is an interval with probability $\mathbb{E}I_l(m)/(m-l+1)$). Hence $\Pr(\omega \cap \mathcal{E}) \leq \Pr(\omega) (s_m + 3h_m + 2 \cdot \mathbf{1}\{i \in \{1, n-2\}\})$. For $\omega = \{\pi(1) = n\}$ the rest π' is uniform on S_{n-1} and another interval is either an interval of π' with 3 to $n-2$ elements or $[1, k]$ with $[2, k]$ an initial interval of π' with 2 to $n-2$ elements, whose expected number is at most h_{n-1} ; the same holds for $\pi(n) = 1$. Summing,

$$\mathbb{E}[\#\omega; \mathcal{E}] \leq \frac{3(n-2)}{n(n-1)} (s_{n-2} + 3h_{n-2}) + \frac{12}{n(n-1)} + \frac{2}{n} (s_{n-1} + h_{n-1}).$$

By Step 2, $s_m = 10/m + O(m^{-2})$, and $h_m \leq 4/m + 3\mathbb{E}I_3(m)/(m-2) + \sum_{l=4}^{m-1} \mathbb{E}I_l(m) = 8/m + O(m^{-2})$ (the term $l=2$ is $4/m$, the factor $\min(l, m-l+1)/(m-l+1)$ is at most 1, and $\mathbb{E}I_{m-1}(m) = 4/m$). So the bound is $O(n^{-2})$. $\square$

Proof of Theorem 5.9. For a set A write $\Pr(R_n \in A) - \Pr(2^Z \in A) = [\Pr(2^{N_n} \in A) - \Pr(2^Z \in A)] + \mathbb{E}[\mathbf{1}\{R_n \in A\} - \mathbf{1}\{2^{N_n} \in A\}]$. By Theorem 5.8(i) the first bracket is $\nu(\{k : 2^k \in A\})/n + \rho_1$ with $|\rho_1| \leq \sum_{k < n} |r_{n,k}| + \sum_{k \geq n} q_k k/n \leq 2^{n+1}/(n(n+1)!) + 1/n!$ (for $k \geq n$, $|p_k - q_k - \nu(k)/n| = q_k|k-1-n|/n \leq q_k k/n$). Outside $\mathcal{E}$, $\mathbf{1}\{R_n \in A\} - \mathbf{1}\{2^{N_n} \in A\} = \sum_{\omega} \mathbf{1}_{\omega} g_{\omega}(N_n)$ with $g_{\omega}(j) = \mathbf{1}\{c_{\omega} 2^j \in A\} - \mathbf{1}\{2^j \in A\}$, the sum running over occurrences; so the second term is $\sum_{\omega} \mathbb{E}[\mathbf{1}_{\omega} g_{\omega}(N_n)] + \rho_2$ with $|\rho_2| \leq \Pr(\mathcal{E}) + \mathbb{E}[\#\omega; \mathcal{E}] = O(n^{-2})$ by Lemma B.2.

Given a three-element occurrence $\omega = (J, \tau)$, contract J as above: then $N_n = N(\tau) + N(\sigma) + \delta$, where $N(\sigma)$ has the law of N_{n-2} and $\delta \neq 0$ only if σ has a descending succession involving j^* (a descending succession of π with one point in J gives one of σ at j^* , and one of σ not involving j^* is one of π , because two consecutive values cannot lie on both sides of the values of J), which has probability at most $2/(n-2)$. Given $\pi(1) = n$, $N_n = N(\pi') + \mathbf{1}\{\pi(2) = n-1\}$ with $N(\pi') \sim N_{n-1}$ and $\Pr(\pi(2) = n-1 \mid \pi(1) = n) = 1/(n-1)$; similarly for $\pi(n) = 1$. Since $|g_{\omega}| \leq 1$, $\mathbb{E}[g_{\omega}(N_n) \mid \omega]$ differs from $\mathbb{E}g_{\omega}(N(\tau) + Z)$ by at most $2\Pr(\delta \neq 0 \mid \omega) + 2d_{\text{TV}}(N_{n-2}, Z)$ (or the analogue with $n-1$), which is $O(1/n)$ by Theorem 5.8(ii). Multiplying by $\Pr(\omega)$ and summing, with total weights $3(n-2)/(n(n-1)) = 3/n + O(n^{-2})$ for the three patterns and $2/n$ for the two boundary events, the second term equals

$$\frac{1}{n} \left([\Pr(6 \cdot 2^Z \in A) - \Pr(4 \cdot 2^Z \in A)] + 4[\Pr(2^{Z+1} \in A) - \Pr(2^Z \in A)] \right) + O(n^{-2}),$$

since the pattern 321 has $N(\tau) = 2$ and $c_{\omega} = \frac{3}{2}$, and the other four kinds have $N(\tau) = 0$ and $c_{\omega} = 2$. All errors are uniform in A , and adding ν/n gives μ/n : at 2^k , $(q_k - q_{k-1}) - q_{k-2} + 4(q_{k-1} - q_k)$, and at $3 \cdot 2^k$ (which is $6 \cdot 2^{k-1}$), q_{k-1} . As μ has total mass 0, $\sup_A |\mu(A)| = \frac{1}{2} \|\mu\|$. For the constant, $3q_{k-1} - 3q_k - q_{k-2} = \frac{e^{-1}}{k!} (3k - 3 - k(k-1)) = -\frac{e^{-1}}{k!} (k-1)(k-3)$; $\sum_k (k-1)(k-3)/k! = 2e - 4e + 3e = e$ and the only negative term is $k=2$, equal to $-\frac{1}{2}$, so $\sum_k |(k-1)(k-3)|/k! = e+1$; the atoms $3 \cdot 2^k$ carry mass $\sum_{k \geq 1} q_{k-1} = 1$. Hence $\|\mu\| = e^{-1}(e+1) + 1 = 2 + e^{-1}$. The particular values follow by evaluating μ at 1, 2, 4, 6, 12, $\{1, 2, 4, 6, 8\}$ and $\{3 \cdot 2^k\}_{k \geq 1}$. $\square$

Proof of Proposition 5.10. Outside $\mathcal{E}$, $R_n \neq 2^{N_n}$ requires an occurrence, whose expected number is $3(n-2)/(n(n-1)) + 2/n \leq 5/n - 3/n^2$; with Lemma B.2(a), $\Pr(R_n \neq 2^{N_n}) \leq 5/n + 220/n^2$. Then $d_{\text{TV}}(R_n, 2^Z) \leq \Pr(R_n \neq 2^{N_n}) + d_{\text{TV}}(2^{N_n}, 2^Z)$, the last term equals $d_{\text{TV}}(N_n, Z)$ because $j \mapsto 2^j$ is injective, and Theorem 5.8(ii) with $2/(n \cdot n!) \leq 1/n^2$ gives the second bound. $\square$

Proofs of the second-order results

Throughout, $x = 1/n$ and $h_k(x) = e^x(1-x)^k - 1 - (1-k)x$, so that $q_k(1-(k-1)/n) - \tilde{q}_k = -q_k h_k(x)$ and, by Theorem 5.8(i),

$$p_k - \tilde{q}_k = -q_k h_k(x) + r_{n,k} \quad (0 \leq k \leq n-1), \quad p_k = 0 \quad (k \geq n). \quad (1)$$

Lemma B.3 (Signs). *For $0 < x < \frac{1}{2}$: $h_0(x) > 0$, $h_1(x) < 0$, $h_2(x) < 0$ and $h_k(x) > 0$ for every integer $k \geq 3$. [proved here]*

Proof. $h_0 = e^x - 1 - x > 0$; $h_1 = e^x(1-x) - 1 < 0$ because $1-x < e^{-x}$; $h_2 = (1-x)h_1 < 0$. For $k=3$, $h_3 > 0$ is equivalent to $\varphi(x) = x + 3\log(1-x) - \log(1-2x) > 0$, and $\varphi(0) = 0$, $\varphi'(x) = 1 - \frac{3}{1-x} + \frac{2}{1-2x} = \frac{x(2x+1)}{(1-x)(1-2x)} > 0$. The second difference $h_{k+1} - 2h_k + h_{k-1} = e^x(1-x)^{k-1}x^2$ is positive, so $k \mapsto h_k(x)$ is convex on the integers; with $h_2 < 0 < h_3$ this gives $h_k \geq h_3 + (k-3)(h_3 - h_2) > 0$ for $k \geq 3$. $\square$

Proof of Theorem 5.11. (i) By Step 3 of the proof of Theorem 5.6, $\mathbb{E}[(N_n)_r] = 1 - r/n$, and $\mathbb{E}[(X)_r] = \lambda^r$ for $X \sim \text{Po}(\lambda)$, so the difference is $-\sum_{i=2}^r \binom{r}{i}(-1/n)^i$; for $r \leq n$ the moduli $\binom{r}{i}n^{-i}$ decrease in i (their ratio is $\frac{r-i}{(i+1)n} < 1$), and the alternating-series bound gives the claim. The expansion of p_k is (1) with $p_k = q_k(1 + c_1(k)x) + r_{n,k}$ and $\tilde{q}_k = q_k \sum_j c_j(k)x^j$, a finite sum times e^x .

(ii) $d_{\text{TV}} = \sum_k (p_k - \tilde{q}_k)^+$, and the terms $k \geq n$ vanish. For $0 \leq k \leq n-1$ put $y_k = -q_k h_k(x)$; since $t \mapsto t^+$ is 1-Lipschitz, $|(p_k - \tilde{q}_k)^+ - y_k^+| \leq |r_{n,k}|$. By Lemma B.3 ($x \leq \frac{1}{3}$), $y_k^+ = q_k |h_k(x)|$ for $k \in \{1, 2\}$, both at most $n-1$, and $y_k^+ = 0$ otherwise. Hence $d_{\text{TV}} = q_1 |h_1| + q_2 |h_2| + \varepsilon_n$ with $|\varepsilon_n| \leq \sum_{k=0}^{n-1} |r_{n,k}| \leq \frac{1}{n(n+1)!} \sum_{k=0}^{n-1} \binom{n+1}{k} = \frac{2^{n+1} - n - 2}{n(n+1)!}$, and $q_1 |h_1| + q_2 |h_2| = e^{-1}(1 - e^x(1 - x))(1 + \frac{1}{2}(1 - x)) = \Phi(x)$.

(iii) $1 - (1 - x)e^x = \sum_{j \geq 2} (j-1)x^j/j!$, so $\Phi(x) = e^{-1} \sum_{j \geq 2} d_j x^j$ with $d_j = \frac{-j^2 + 5j - 3}{2j!}$: $d_2 = \frac{3}{4}$, $d_3 = \frac{1}{4}$, $d_4 = \frac{1}{48}$, $d_j < 0$ for $j \geq 5$, and $\sum_{j \geq 2} d_j = e\Phi(1) = 1$, so $\sum_{j \geq 5} d_j = -\frac{1}{48}$. For $0 < x \leq \frac{1}{3}$ it follows that $\frac{1}{48}(1 - x) \leq \sum_{j \geq 4} d_j x^{j-4} < \frac{1}{48}$, which is the claim; with (ii), $n^2 |\varepsilon_n| \rightarrow 0$. $\square$

Proof of Theorem 5.12. (i) Let $\delta = 1 - \lambda = x - ax^2$, so $0 < \delta \leq 1$ for n large. Since $|c_j(k)|$ is at most the coefficient of x^j in $e^x(1 + x)^k$, $|q_k(\lambda) - q_k(1 + c_1(k)\delta + c_2(k)\delta^2)| \leq q_k e 2^k \delta^3$, and $\sum_k q_k e 2^k = e^2$. For $k \leq n-1$, $p_k = q_k(1 + c_1(k)x) + r_{n,k}$ and $c_1(k)(x - \delta) = (1 - k)ax^2$ give $p_k - q_k(\lambda) = -q_k g_a(k)x^2 + q_k c_2(k)(x^2 - \delta^2) + r_{n,k} + \eta_k$ with $g_a(k) = c_2(k) + a(k-1)$ and $|\eta_k| \leq q_k e 2^k \delta^3$; for $k \geq n$, $|p_k - q_k(\lambda) + q_k g_a(k)x^2| \leq q_k(\lambda) + q_k |g_a(k)|x^2$, whose sum over $k \geq n$ is super-exponentially small. Summing, $|2d_{\text{TV}} - x^2 L(a)| \leq \frac{3}{2e} |x^2 - \delta^2| + e^2 \delta^3 + \sum_k |r_{n,k}| + o(x^3) = O(x^3)$, uniformly for a in compact sets.

(ii) $\sum_k q_k g_a(k) = 0$, since $\mathbb{E}[Z^2 - 3Z + 1] = 0$ and $\mathbb{E}[Z - 1] = 0$. With $f(1) = -1$ and $f(k) = 1$ for $k \neq 1$, $L(a) \geq \sum_k q_k g_a(k) f(k) = -2q_1 g_a(1) = e^{-1}$, because $g_a(1) = -\frac{1}{2}$. Equality requires $g_a(k) \geq 0$ for all $k \neq 1$; $g_a(0) = \frac{1}{2} - a$ and $g_a(2) = a - \frac{1}{2}$ force $a = \frac{1}{2}$, and $g_{1/2}(k) = \frac{1}{2}k(k-2)$ satisfies it. Writing $g_a(k) = \frac{1}{2}[(k-1)(k-2+2a) - 1]$, for $a \geq -\frac{1}{4}$ one has $g_a(3) = \frac{1}{2}(1+4a) \geq 0$ and $g_a(k) \geq \frac{1}{2}(5+6a) > 0$ for $k \geq 4$, so $L(a) = 2 \sum_k q_k g_a(k)^- = 2[\frac{1}{2}q_1 + q_0(a - \frac{1}{2})^+ + q_2(\frac{1}{2} - a)^+]$, which is the formula.

(iii) The upper bound is (i) with $a = \frac{1}{2}$. For the lower bound write $\lambda = 1 - x + ax^2$. If $a \in [-\frac{1}{2}, \frac{3}{2}]$, (i) (uniform in a) and (ii) give $n^2 d_{\text{TV}} \geq \frac{1}{2e} - O(1/n)$. If $a > \frac{3}{2}$, then, as $q_0(\lambda) = e^{-\lambda}$ decreases in λ , $d_{\text{TV}} \geq p_0 - e^{-\lambda} \geq e^{-1}(1 + x - e^{-3x^2/2}) - |r_{n,0}| = e^{-1}x^2 + O(x^3)$; if $a < -\frac{1}{2}$, likewise $d_{\text{TV}} \geq e^{-\lambda} - p_0 \geq e^{-1}(e^{x+x^2/2} - 1 - x) - |r_{n,0}| = e^{-1}x^2 + O(x^3)$. In both cases $n^2 d_{\text{TV}} \geq e^{-1} - o(1) > \frac{1}{2e}$. $\square$

For the explicit constant we sharpen part (a) of Lemma B.2, whose term $120/n^2$ (intervals with 5 to $n-5$ elements) is in fact $O(n^{-3})$. Write $(m)_j = m(m-1)\cdots(m-j+1)$, recall $\mathbb{E}I_k(m) = (m-k+1)^2/\binom{m}{k}$ for a uniform permutation of length m , let s_m, h_m be as in the proof of Lemma B.2(b), and

$$\bar{F}(m) = \frac{42}{(m)_2} + \frac{936}{(m)_3} + \frac{600}{(m)_4} + \frac{4320}{(m)_5}.$$

Lemma B.4 (Intervals, refined). *For $m \geq 12$: (a) $\sum_{k=4}^{m-2} \mathbb{E}I_k(m) \leq \bar{F}(m)$; (b) for $n \geq 12$, $\Pr(\mathcal{E}) \leq \bar{P}(n) = \frac{90}{(n)_2} + \frac{944}{(n)_3} + \frac{600}{(n)_4} + \frac{4320}{(n)_5}$; (c) $s_m \leq \bar{s}_m = \frac{10}{m} + \bar{F}(m)$ and $h_m \leq \bar{h}_m = \frac{8}{m} + \frac{18}{(m)_2} + \bar{F}(m)$. [proved here]*

Proof. (a) Exactly, $\mathbb{E}I_4 = 24(m-3)/(m)_3 \leq 24/(m)_2$, $\mathbb{E}I_{m-2} = 18/(m)_2$, $\mathbb{E}I_{m-3} = 96/(m)_3$, $\mathbb{E}I_{m-4} = 600/(m)_4$, $\mathbb{E}I_5 = 120(m-4)/(m)_4 \leq 120/(m)_3$ and $\mathbb{E}I_{m-5} = 4320/(m)_5$; these six indices are distinct for $m \geq 12$. For the $m-11$ values $6 \leq k \leq m-6$, $\binom{m}{k} \geq \binom{m}{6}$ and $(m-k+1)^2 \leq (m-5)^2$, so they contribute at most $720(m-11)(m-5)/(m)_5 \leq 720/(m)_3$. (b) The proof of Lemma B.2(a) bounds $\Pr(\mathcal{E})$ by $\sum_{k=4}^{n-2} \mathbb{E}I_k$ plus the three pair moments, which it computes exactly; using $(n-4)(n-5) \leq (n-2)(n-3)$, $n-3 \leq n-2$, $n-4 \leq n-3$ and $6n-16 \leq 6(n-2)$

they are at most $\frac{18+4}{(n)_2} + \frac{8}{(n)_3}$, $\frac{24}{(n)_2}$ and $\frac{2}{(n)_2}$. (c) $s_m = \mathbb{E}I_3(m) + \sum_{k=4}^{m-2} \mathbb{E}I_k(m) + \mathbb{E}I_{m-1}(m)$ with $\mathbb{E}I_3(m) = 6(m-2)/(m)_2 \leq 6/m$ and $\mathbb{E}I_{m-1}(m) = 4/m$; the proof of Lemma B.2(b) gives $h_m \leq 4/m + 18/(m)_2 + \sum_{l=4}^{m-1} \mathbb{E}I_l(m)$. $\square$

Proof of Proposition 5.13. Let $\bar{d}_m = e^{-1}/m + 2/(m \cdot m!)$, an upper bound for $d_{\text{TV}}(N_m, \text{Po}(1))$ when $m \geq 4$ (Theorem 5.8(ii)), and for $n \geq 22$

$$\begin{aligned} B(n) = & \frac{2^{n+1}}{n(n+1)!} + \frac{1}{n!} + \bar{P}(n) + \frac{3(n-2)}{(n)_2}(\bar{s}_{n-2} + 3\bar{h}_{n-2}) + \frac{12}{(n)_2} + \frac{2}{n}(\bar{s}_{n-1} + \bar{h}_{n-1}) \\ & + \frac{3(n-2)}{(n)_2} \left(\frac{4}{n-2} + 2\bar{d}_{n-2} \right) + \frac{2}{n} \left(\frac{2}{n-1} + 2\bar{d}_{n-1} \right) + \frac{3}{(n)_2}. \end{aligned} \quad (2)$$

We follow the proof of Theorem 5.9 and bound each error. The first bracket differs from $\nu(\{k : 2^k \in A\})/n$ by $|\rho_1| \leq 2^{n+1}/(n(n+1)!) + 1/n!$. Replacing $\mathbb{E}[\mathbf{1}\{R_n \in A\} - \mathbf{1}\{2^{N_n} \in A\}]$ by $\sum_{\omega} \mathbb{E}[\mathbf{1}_{\omega} g_{\omega}(N_n)]$ costs $|\rho_2| \leq \Pr(\mathcal{E}) + \mathbb{E}[\#\omega; \mathcal{E}]$; the first is at most $\bar{P}(n)$ by Lemma B.4(b), and the second at most the second group of terms of (2), by the bound displayed in the proof of Lemma B.2(b) and Lemma B.4(c) with $m = n-2 \geq 20$ and $m = n-1$. Given a three-element occurrence, g_{ω} takes values in $\{-1, 0, 1\}$, so $|\mathbb{E}[g_{\omega}(N_n) \mid \omega] - \mathbb{E}g_{\omega}(N(\tau) + Z)| \leq 2\Pr(\delta \neq 0 \mid \omega) + 2d_{\text{TV}}(N_{n-2}, Z) \leq \frac{4}{n-2} + 2\bar{d}_{n-2}$, because $\delta \neq 0$ requires a descending succession of the uniform $\sigma \in S_{n-2}$ at one of the two pairs of positions that contain j^* , each of probability $1/(n-2)$; for the two boundary occurrences the bound is $\frac{2}{n-1} + 2\bar{d}_{n-1}$. With the weights $3(n-2)/(n)_2$ and $2/n$ this gives the third group. Finally, replacing the exact weight $(n-2)/(n(n-1))$ of each of the three patterns by $1/n$ changes the main term by at most $1/(n(n-1))$ each, since $|\mathbb{E}g_{\omega}| \leq 1$: the last term. Hence $|\Pr(R_n \in A) - \Pr(2^Z \in A) - \mu(A)/n| \leq B(n)$ for every A ; as $d_{\text{TV}}(R_n, 2^Z) = \sup_A |\Pr(R_n \in A) - \Pr(2^Z \in A)|$ and $c_2/n = \sup_A |\mu(A)|/n$, the same bound holds for $|d_{\text{TV}}(R_n, 2^Z) - c_2/n|$.

Monotonicity: after expanding $\bar{P}$, $\bar{s}$, $\bar{h}$ and $\bar{d}$ and cancelling the factors $n-2$ of the weights, $B(n)$ is a finite sum of terms $c/\prod_{j=1}^d (n-i_j)$ with $c > 0$, $d \geq 2$ and $0 \leq i_j \leq 6$, plus four terms with a factorial $(2^{n+1}/(n(n+1)!))$, $1/n!$ and the two coming from $2/(m \cdot m!)$ in $\bar{d}_m$. For the former, $n^2/((n-i_1)(n-i_2))$ is a product of two positive non-increasing factors and the other factors $1/(n-i_j)$ decrease; for the latter, n^2 times each term decreases (the ratio of consecutive values is below 1 for $n \geq 3$). Hence $n^2 B(n)$ is non-increasing for $n \geq 22$; this was also checked numerically for $22 \leq n \leq 5000$. Evaluating, $22^2 B(22) = 421.49$, $100^2 B(100) = 285.80$, and $n^2 B(n) \rightarrow 90 + 150 + (16 + 10/e) + 3 = 259 + 10/e$. $\square$

Numerical check of the sharp rates. `theory/check_sharp_rate.py` (fixed seed, recorded in its output) computes the law of N_n for $n \leq 60$ in rational arithmetic in three independent ways (inversion of the factorial moments, the closed form $\binom{n-1}{k}(D_{n-k} + D_{n-k-1})/n!$ with D_m the derangement numbers [19], and the recurrence obtained by inserting the largest element), all equal, and equal to brute force for $n \leq 8$. For $4 \leq n \leq 60$, $|r_{n,k}| n k! (n-k+1)! \leq 0.97$ (Theorem 5.8(i): at most 1) and $n! |n d_{\text{TV}}(N_n, \text{Po}(1)) - e^{-1}| \in [0.016, 0.96]$ (at most 2). The pair moments of Lemma B.2(a) agree with exhaustive enumeration for $n \in \{8, 9\}$, and for $20 \leq n \leq 3000$ each of the four pieces of its proof is at most 0.98 times its bound. Table 11 reports the law of R_n estimated on 200000 to 300000 uniform permutations for n from 50 to 400 (method in the frozen output), next to the exact value of the second-order check below. For $n \geq 100$ the estimates of $n d_{\text{TV}}(R_n, 2^Z)$ agree with $c_2 = 1.184$ and with the first-order law. At $n = 50$ the estimate 1.261 exceeds the first-order value 1.201 by 0.06 (2.9 standard errors) and the largest standardised atom-wise deviation is 4.4 (at most 3.3 for the larger n): the n^{-2} term is visible there, since the exact value is 1.280 ($\kappa/n \approx 0.086$, Remark 5.14). Part (F) of the script classifies the exceptions with an exact count of all intervals: the bad event $\mathcal{E}$ of Lemma B.2 has frequency 0.156, 0.046 and 0.008 at $n = 20, 40, 100$ (bounds $223/n^2 = 0.557, 0.139, 0.022$) and accounts for 2592 of the 4921 exceptions at $n = 20$. The

n	samples	$n \Pr(R_n \neq 2^{N_n})$	$n d_{\text{TV}}(R_n, 2^Z)$	exact	first order	max $ z $
50	300000	5.01 ± 0.05	1.261 ± 0.041	1.280	1.201	4.4
100	300000	5.04 ± 0.08	1.231 ± 0.060	1.229	1.192	2.3
200	250000	4.92 ± 0.12	1.165 ± 0.078	1.206	1.188	1.6
400	200000	5.03 ± 0.20	1.206 ± 0.145	1.195	1.186	3.3

Table 11: Monte Carlo check of Theorem 5.9 and Proposition 5.10 (`theory/check_sharp_rate.py`). Intervals are 95% (binomial for the exception rate, jackknife over ten groups for d_{TV} ; the plug-in estimate is biased upwards by sampling noise). Exact: $n d_{\text{TV}}$ from the exact law of R_n (`theory/check_second_order.py`). First order: $n d_{\text{TV}}$ between 2^Z and the first-order law of Theorem 5.9 evaluated with the exact weights $(n-2)/(n(n-1))$ and $1/n$ and the exact laws of N_{n-2} and N_{n-1} . The limit of the fourth to sixth columns is $c_2 = 1 + 1/(2e) = 1.184$; Proposition 5.10 bounds $n d_{\text{TV}}$ by $5 + e^{-1} + 221/n$ and Proposition 5.13 bounds $|n d_{\text{TV}} - c_2|$ by $422/n$. Max $|z|$: largest standardised deviation, over the atoms $1, 2, 4, \dots, 32$ and $6, 12, 24, 48$, between the estimated and the first-order law.

output is frozen as `results/check_sharp_rate_output.txt`, without timing lines (SHA-256 beginning 72d23ffe8f50726c); the run took 116 seconds on one core.

Numerical check of the second-order results. `theory/check_second_order.py` (deterministic) computes the law of N_n in rational arithmetic (closed form, inversion of factorial moments and, for $n \leq 8$, brute force, all equal) and $d_{\text{TV}}(N_n, \text{Po}(1 - 1/n))$ with 150 digits for $3 \leq n \leq 60$ and $n \in \{70, 80, 100, 150, 200, 300, 400, 500, 700, 1000\}$: $|d_{\text{TV}} - \Phi(1/n)|$ is at most 0.194 times the bound of Theorem 5.11(ii) wherever that bound exceeds 10^{-130} , and below 10^{-130} elsewhere; $48en^2(n^2\Phi(1/n) - \frac{3}{4e} - \frac{1}{4en})$ lies in $[0.763, 0.9994]$; and at $n = 1000$, $n^2 d_{\text{TV}} = 0.2760016$ against $\frac{3}{4e} + \frac{1}{4en} = 0.2760016$. Minimising over λ at $n = 1000$ gives $n^2(\lambda - 1 + 1/n) = 0.4997$ and $n^2 d_{\text{TV}} = 0.18388$ (Theorem 5.12: $\frac{1}{2}$ and 0.18394). The script also computes the law of R_n exactly on the atoms 2^k and $3 \cdot 2^k$ from Gallai’s product formula (Remark B.1) and the uniqueness of the substitution decomposition [1]: the generating function of the permutations whose number of transitive orientations is $2^a 3^b$, $b \leq 1$, satisfies a system of equations in which the numbers of simple permutations enter through the compositional inverse of $\sum_{n \geq 1} n! z^n$. The resulting law agrees with the product formula on all permutations of length at most 8, with brute force over pairs of linear extensions for length at most 6, and, in exact integer arithmetic up to $n = 30$, with the floating-point computation (relative error $1.1 \cdot 10^{-15}$) that is used up to $n = 600$. Over $22 \leq n \leq 600$, $|d_{\text{TV}}(R_n, 2^Z) - c_2/n|$ is at most 0.0163 times $B(n)$; atom by atom, $n^2 |\Pr(R_n = x) - \Pr(2^Z = x) - \mu(x)/n| \leq 3.15$ for $100 \leq n \leq 600$, a check of Theorem 5.9 that does not rely on sampling; and the inequalities of Lemma B.4(a), (b) and (c) hold for $12 \leq m \leq 3000$ with ratios at most 0.9942, 0.9970, 0.9998 and 0.9990. The output is frozen as `results/check_second_order_output.txt`, without timing lines (SHA-256 beginning 328efae6b63a2b6c); the run took 51 seconds on one core.

References

- [1] Michael H. Albert and Mike D. Atkinson. Simple permutations and pattern restricted permutations. *Discrete Mathematics*, 300(1–3):1–15, 2005.
- [2] Michael H. Albert, Mike D. Atkinson, and Martin Klazar. The enumeration of simple permutations. *Journal of Integer Sequences*, 6:Article 03.4.4, 2003.
- [3] Jinho Baik, Percy Deift, and Kurt Johansson. On the distribution of the length of the longest increasing subsequence of random permutations. *Journal of the American Mathematical Society*, 12(4):1119–1178, 1999.

- [4] Leonard M. Blumenthal. *Theory and Applications of Distance Geometry*. Clarendon Press, Oxford, 1953.
- [5] Béla Bollobás and Graham Brightwell. Box-spaces and random partial orders. *Transactions of the American Mathematical Society*, 324(1):59–72, 1991.
- [6] Béla Bollobás and Graham Brightwell. The height of a random partial order: concentration of measure. *Annals of Applied Probability*, 2(4):1009–1018, 1992.
- [7] Luca Bombelli, Joohan Lee, David Meyer, and Rafael D. Sorkin. Space-time as a causal set. *Physical Review Letters*, 59(5):521–524, 1987.
- [8] Graham Brightwell and Ruth Gregory. Structure of random discrete spacetime. *Physical Review Letters*, 66(3):260–263, 1991.
- [9] Robert Brignall. A survey of simple permutations. In Steve Linton, Nik Ruškuc, and Vincent Vatter, editors, *Permutation Patterns*, volume 376 of *London Mathematical Society Lecture Note Series*, pages 41–65. Cambridge University Press, Cambridge, 2010. arXiv:0801.0963.
- [10] Sylvie Corteel, Guy Louchard, and Robin Pemantle. Common intervals in permutations. *Discrete Mathematics and Theoretical Computer Science*, 8:189–214, 2006. dmtcs.episciences.org/362.
- [11] Ben Dushnik and Edwin W. Miller. Partially ordered sets. *American Journal of Mathematics*, 63(3):600–610, 1941.
- [12] K. Ruben Gabriel and Robert R. Sokal. A new statistical approach to geographic variation analysis. *Systematic Zoology*, 18(3):259–278, 1969.
- [13] Tibor Gallai. Transitiv orientierbare Graphen. *Acta Mathematica Academiae Scientiarum Hungaricae*, 18:25–66, 1967. doi: 10.1007/BF02020961. English translation by F. Maffray and M. Preissmann in J. L. Ramírez Alfonsín and B. A. Reed (eds.), *Perfect Graphs*, Wiley, 2001.
- [14] Martin Charles Golumbic. Comparability graphs and a new matroid. *Journal of Combinatorial Theory, Series B*, 22(1):68–90, 1977.
- [15] Martin Charles Golumbic. *Algorithmic Graph Theory and Perfect Graphs*. Academic Press, New York, 1980.
- [16] John C. Gower. Generalized Procrustes analysis. *Psychometrika*, 40(1):33–51, 1975.
- [17] Stephen W. Hawking, A. R. King, and P. J. McCarthy. A new topology for curved space-time which incorporates the causal, differential, and conformal structures. *Journal of Mathematical Physics*, 17(2):174–181, 1976.
- [18] David Hilbert. *Grundlagen der Geometrie*. Teubner, Leipzig, 1899.
- [19] Irving Kaplansky. The asymptotic distribution of runs of consecutive elements. *Annals of Mathematical Statistics*, 16(2):200–203, 1945.
- [20] Felix Klein. *Vergleichende Betrachtungen über neuere geometrische Forschungen*. Deichert, Erlangen, 1872. Reprinted in *Mathematische Annalen* 43 (1893), 63–100.
- [21] Matthäus Kleindessner and Ulrike von Luxburg. Uniqueness of ordinal embedding. In *Proceedings of the 27th Conference on Learning Theory (COLT)*, volume 35 of *JMLR Workshop and Conference Proceedings*, pages 40–67, 2014.

- [22] David H. Krantz, R. Duncan Luce, Patrick Suppes, and Amos Tversky. *Foundations of Measurement, Volume I: Additive and Polynomial Representations*. Academic Press, New York, 1971.
- [23] E. H. Kronheimer and Roger Penrose. On the structure of causal spaces. *Proceedings of the Cambridge Philosophical Society*, 63(2):481–501, 1967.
- [24] Joseph B. Kruskal. On the shortest spanning subtree of a graph and the traveling salesman problem. *Proceedings of the American Mathematical Society*, 7(1):48–50, 1956.
- [25] Joseph B. Kruskal. Multidimensional scaling by optimizing goodness of fit to a nonmetric hypothesis. *Psychometrika*, 29(1):1–27, 1964.
- [26] Benjamin F. Logan and Lawrence A. Shepp. A variational problem for random Young tableaux. *Advances in Mathematics*, 26(2):206–222, 1977.
- [27] David B. Malament. The class of continuous timelike curves determines the topology of spacetime. *Journal of Mathematical Physics*, 18(7):1399–1404, 1977.
- [28] Karl Menger. Untersuchungen über allgemeine Metrik. *Mathematische Annalen*, 100:75–163, 1928.
- [29] David A. Meyer. *The Dimension of Causal Sets*. PhD thesis, Massachusetts Institute of Technology, 1989. MIT handle 1721.1/14328; often cited as 1988.
- [30] Jan Myrheim. Statistical geometry. Technical Report CERN-TH-2538, CERN, 1978. CDS record 293594, <https://cds.cern.ch/record/293594>.
- [31] Louis Narens. *Theories of Meaningfulness*. Lawrence Erlbaum, Mahwah, NJ, 2002.
- [32] Fabian Pedregosa, Gaël Varoquaux, Alexandre Gramfort, Vincent Michel, Bertrand Thirion, Olivier Grisel, Mathieu Blondel, Peter Prettenhofer, Ron Weiss, Vincent Dubourg, Jake Vanderplas, Alexandre Passos, David Cournapeau, Matthieu Brucher, Matthieu Perrot, and Édouard Duchesnay. Scikit-learn: machine learning in Python. *Journal of Machine Learning Research*, 12:2825–2830, 2011.
- [33] Isaac J. Schoenberg. Remarks to Maurice Fréchet’s article “sur la définition axiomatique d’une classe d’espaces distanciés vectoriellement applicable sur l’espace de Hilbert”. *Annals of Mathematics*, 36(3):724–732, 1935.
- [34] Isaac J. Schoenberg. On certain metric spaces arising from Euclidean spaces by a change of metric and their imbedding in Hilbert space. *Annals of Mathematics*, 38(4):787–793, 1937.
- [35] Peter H. Schönemann. A generalized solution of the orthogonal Procrustes problem. *Psychometrika*, 31(1):1–10, 1966.
- [36] Roger N. Shepard. The analysis of proximities: multidimensional scaling with an unknown distance function. I. *Psychometrika*, 27(2):125–140, 1962.
- [37] Rafael D. Sorkin. Causal sets: discrete gravity. In Andrés Gomberoff and Donald Marolf, editors, *Lectures on Quantum Gravity*, pages 305–327. Springer, New York, 2005. doi: 10.1007/0-387-24992-3_7.
- [38] Stanley S. Stevens. On the theory of scales of measurement. *Science*, 103(2684):677–680, 1946.

- [39] Alfred Tarski. What is elementary geometry? In Leon Henkin, Patrick Suppes, and Alfred Tarski, editors, *The Axiomatic Method, with Special Reference to Geometry and Physics*, pages 16–29. North-Holland, Amsterdam, 1959.
- [40] Alfred Tarski and Steven Givant. Tarski’s system of geometry. *Bulletin of Symbolic Logic*, 5(2):175–214, 1999.
- [41] Yoshikazu Terada and Ulrike von Luxburg. Local ordinal embedding. In *Proceedings of the 31st International Conference on Machine Learning (ICML)*, volume 32 of *JMLR Workshop and Conference Proceedings*, pages 847–855, 2014.
- [42] Godfried T. Toussaint. The relative neighbourhood graph of a finite planar set. *Pattern Recognition*, 12(4):261–268, 1980.
- [43] Anatoly M. Vershik and Sergei V. Kerov. Asymptotics of the Plancherel measure of the symmetric group and the limiting form of Young tableaux. *Soviet Mathematics Doklady*, 18:527–531, 1977.
- [44] Pauli Virtanen, Ralf Gommers, Travis E. Oliphant, et al. SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nature Methods*, 17:261–272, 2020.
- [45] J. Wolfowitz. Note on runs of consecutive elements. *Annals of Mathematical Statistics*, 15(1):97–98, 1944. doi: 10.1214/aoms/1177731319.
- [46] Gale Young and Alston S. Householder. Discussion of a set of points in terms of their mutual distances. *Psychometrika*, 3(1):19–22, 1938.
- [47] E. Christopher Zeeman. Causality implies the Lorentz group. *Journal of Mathematical Physics*, 5(4):490–493, 1964.