

Deep inelastic scattering on polarized lithium: the physics behind the planned measurements

2026-08-17, revised 2026-08-21 (why lithium, why polarization, fixed target versus collider, beam requirements and uniqueness by observable; fact-checked against primary sources) · companion to the cos 2φ projection report and to plans/01 (physics case) · written for students and colleagues joining the program

Polarized lithium beams would provide the EIC with spin-polarized *nuclei* that are light enough for ab initio computation yet complex enough to exhibit genuine nuclear structure. ⁶Li is a spin-1 nucleus well described as a deuteron bound to an inert α core; ⁷Li is its spin-3/2 partner, an α plus a triton. This note develops, starting from the DIS formalism, the four measurements around which the program is built: the tensor structure function b₁, the double-helicity-flip ("exotic glue") structure function Δ, the polarized EMC effect, and spectator-tagged DIS on the embedded deuteron. It begins with the case itself: why lithium is the right element, why the observables exist only with polarization, and which of them a fixed target could deliver and which require the collider.

1 · Why polarized lithium, and why at a collider

The EIC project scope provides polarized electrons, protons and ³He; polarized deuterons are planned as a later addition, and no heavier polarized ion appears in the conceptual design or in the Yellow Report [29,25]. The EPIOS white paper [1] makes the case for polarized ⁶Li and ⁷Li — "a desirable addition" that it lists with ²H and ³He as critical technology — and the ANL source development [51] specifies the beam: an atomic-beam source with optical pumping at 671 nm feeding EBIS charge breeding, with source targets P_z ≥ 0.90 (vector) and P_{zz} ≥ 0.80 (tensor) at ≥ 10¹³ atoms/s. Four measurements motivate the beam, each addressing a verified gap in the literature: no b₁ prediction or measurement exists for any nucleus beyond the deuteron; no gluon-transversity projection exists for any EIC target; no study of α-tagged DIS on lithium exists; and no lithium-beam luminosity or polarization parameter set has been published. The purpose of the program's simulations is to convert these gaps into figures of merit.

MEASUREMENT	OBSERVABLE	ELECTRON BEAM	ION SPIN	AXIS
Polarized EMC effect (⁷ Li, ⁶ Li)	A → g ₁ ^A	polarized, helicity flipped	vector P _z	longitudinal
Tensor structure function (⁶ Li)	A _{zz} → b ₁	unpolarized	tensor P _{zz}	longitudinal (or along q)
Exotic glue (⁶ Li; ⁷ Li eligible)	cos 2φ → Δ(x,Q ²)	unpolarized	tensor P _{zz}	transverse
Tagged DIS (both isotopes)	tagged versions of all three	as above	as above	as above, correlated with spectator

1.1 · Why lithium as a source: an alkali atom with the right nuclei

Lithium is the lightest alkali metal and its atom is hydrogen-like: a single 2s valence electron whose D lines near 671 nm (670.96 and 670.98 nm in vacuum for ⁷Li) fall within the range of commercial single-frequency diode lasers, and whose ground-state hyperfine splittings, 228.2 MHz for ⁶Li and 803.5 MHz for ⁷Li [35], are small enough for one laser with an electro-optic sideband to address both hyperfine levels. Circularly polarized light pumps the atoms into a single magnetic substate, the hyperfine coupling transfers the electron polarization to the nucleus, and radio-frequency transitions between Zeeman sublevels then set any desired combination of P_z and P_{zz} [51]. The EPIOS paper makes the general point: alkali atoms "are hydrogen-like with a single valence electron and laser cooling of these species has been extensively studied", whereas boron, the other light candidate with a large nuclear spin, has an electronic structure that "presents a challenge for optical pumping" [1]. The low ionization energy of lithium (5.39 eV [35]) eases ionization, and fully stripped Li³⁺ is within reach of the EBIS that is being commissioned to charge-breed the polarized ³He⁺⁺ beam, whose design goal is 2×10¹¹

ions per pulse at $\geq 70\%$ polarization, with the measurement of the polarization transmitted through the EBIS planned for 2026 [34,1].

The nuclei are the other half of the argument. Both isotopes are stable and available (7.6% ${}^6\text{Li}$ and 92.4% ${}^7\text{Li}$ in natural lithium), and they are the only stable isotopes of one element with $J = 1$ and $J = 3/2$: the stable spin-1 nuclei are exactly ${}^2\text{H}$, ${}^6\text{Li}$ and ${}^{14}\text{N}$, and the stable spin-3/2 nuclei below $A = 30$ are ${}^7\text{Li}$, ${}^9\text{Be}$, ${}^{11}\text{B}$, ${}^{21}\text{Ne}$ and ${}^{23}\text{Na}$ [36]. One source, one laser system and one injector therefore serve a spin-1 and a spin-3/2 program. The EPIOS assessment of where an exotic-glue discovery is most likely — "spin-1 ${}^6\text{Li}$ or ${}^{14}\text{N}$, or spin-3/2 ${}^7\text{Li}$ or ${}^{23}\text{Na}$ " [1] — reduces, once optical pumping is required, to lithium and sodium, and lithium is the lighter, cluster-structured and ab initio-computable one (§1.2).

Polarized lithium beams also have a forty-year history at low energy, so the source physics is established rather than speculative. The Heidelberg tandem delivered ${}^6\text{Li}^{3+}$ beams of up to 150 nA with vector polarization 0.57–0.65 in 1977 [30]; by 1987 its atomic-beam source reached $P_z, P_{zz} \geq 0.85$ and accelerated ${}^7\text{Li}$ with an alignment $P_{zz} = 0.69$ on target [31]; the optically pumped Florida State source delivered more than 150 nA of ${}^6\text{Li}^{3+}$ with $P_{yy} = -1.33 \pm 0.02$ and $P_y > 0.74$, with no polarization loss in the linac [32]; and polarized ${}^6\text{Li}$, ${}^7\text{Li}$ and ${}^{23}\text{Na}$ beams served two decades of nuclear-reaction studies [33]. What is new for the EIC is scale and transport: $\geq 10^{13}$ atoms/s, two to three orders of magnitude beyond these sources, and the preservation of the polarization through charge breeding and acceleration to more than 100 GeV/u [51,1].

In the ring, ${}^6\text{Li}$ is the easy case. Its gyromagnetic anomaly, $G \approx -0.18$, is as small as the deuteron's (-0.14), so it crosses only 81 depolarizing resonances on the way to top energy, against 63 for the deuteron and 2457 for ${}^3\text{He}$ [28], and, like the deuteron, it is handled with a partial solenoid snake and tune jumps — full Siberian snakes would be impractically long for so small a G [28,1] — so the deuteron techniques [52] carry over. ${}^7\text{Li}$ ($G \approx +1.52$) crosses 573 resonances and needs partial snakes with jump quadrupoles [1,28]. The rigidity of the hadron ring fixes the top energies at $275 \text{ GeV} \times Z/A$: 137.5 GeV/u for ${}^6\text{Li}$ and 117.9 GeV/u for ${}^7\text{Li}$. Because ${}^6\text{Li}$ has $A/Z = 2$, it runs at the same per-nucleon energy and magnet settings as the deuteron, so $e+d$ and $e+{}^6\text{Li}$ can be compared at identical kinematics.

1.2 · Why lithium as a target: the spin sits on a light cluster

The two stable lithium isotopes are the lightest nuclei beyond $A = 3$ in which a polarized light cluster is bound to an inert core. In ${}^6\text{Li}$ the core is an α particle ($J = 0$, $T = 0$) and the cluster is deuteron-like, bound by 1.47 MeV against $\alpha+d$ breakup; in ${}^7\text{Li}$ the cluster is a triton, bound by 2.47 MeV [36]. Variational Monte Carlo calculations with realistic two- and three-nucleon forces make the picture quantitative: the α -d overlap of the ${}^6\text{Li}$ ground state integrates to $N_{\alpha d} = 0.86$, of which 0.846 is S-wave and 0.017 D-wave, and the α -t overlap of ${}^7\text{Li}$ integrates to $N_{\alpha t} = 1.00$ [17]; the no-core shell model with continuum gives an asymptotic D/S ratio of -0.027 for the $d+\alpha$ configuration [38]. The consequence for polarization is what makes the isotopes useful. In a fully polarized ${}^6\text{Li}$ the α contributes nothing and the two valence nucleons carry the spin: six-body VMC gives spin-up and spin-down valence numbers of 1.924 and 1.076, a polarization of 0.85 [17,19]; three-body $\alpha+n+p$ models, in which the $\alpha+d$ component accounts for 60–70% of the wave function, give a valence-nucleon polarization above 90% [18]; and the SLAC E155 analysis, which used ${}^6\text{LiD}$ as its deuteron target, adopted 87% for the effective deuteron polarization from the same body of calculations [43]. ${}^6\text{Li}$ is therefore a polarized deuteron embedded in the field of an α particle — a spin-1 target whose tensor observables can be set against those of the free deuteron to isolate the medium effect. The only other stable spin-1 ground state is ${}^{14}\text{N}$, whose spin is shared among p-shell nucleons, which has no comparable cluster simplicity, and which lies beyond the reach of exact quantum Monte Carlo ($A \leq 12$) [40].

In ${}^7\text{Li}$ the spin is carried by the unpaired proton of the triton cluster, the neutrons being paired [39]. VMC gives spin-up and spin-down proton numbers of 1.934 and 1.066 and neutron numbers of 1.981 and 2.019 [17], i.e. $P_p = 0.87$ and $P_n = -0.04$ — quoted as 0.866 and -0.037 in the JLab proposal [45] and used in §4.3. The shell model already gives $P_p = 13/15$; the microscopic calculations confirm the proton value and show the neutrons

almost unpolarized [19,14]. Polarized ${}^7\text{Li}$ is thus to the proton what polarized ${}^3\text{He}$ is to the neutron, with the property, decisive for the polarized EMC effect, that the proton sits in a nucleus of normal density. Cloët, Bentz and Thomas state the selection criteria: the nucleus must have $|P_n| \ll |P_p|$, and it must be light, because g_1^A is carried by the valence nucleons and is suppressed relative to F_2^A by roughly $1/A$; their candidates are ${}^{11}\text{B}$, ${}^{15}\text{N}$, ${}^{27}\text{Al}$ and ${}^7\text{Li}$, "where the nuclear polarization is largely dominated by the valence proton" [14], and the gluon-spin study names ${}^7\text{Li}$ "the most promising case for experimental investigation" [16].

Two further properties matter. The cluster structure makes the nuclei computable: Green's-function Monte Carlo with the AV18+IL7 Hamiltonian reproduces the binding energies of both isotopes (-31.93 vs -31.99 MeV for ${}^6\text{Li}$, -39.15 vs -39.25 MeV for ${}^7\text{Li}$), their charge radii to within 3%, and, once two-body currents are included, their magnetic moments (0.835 vs $0.822 \mu_N$; 3.24 vs $3.256 \mu_N$) [40,39], so the nuclear corrections entering every extraction — effective polarizations, momentum distributions, spectral functions — carry controlled uncertainties rather than model spread. The quadrupole moment of ${}^6\text{Li}$ is the one exception: the measured $Q = -0.0806 \text{ fm}^2$, against $+0.286 \text{ fm}^2$ for the free deuteron [36], arises from a delicate cancellation between the deuteron's own quadrupole moment and the D-wave part of the α -d relative motion [19,37], and theory reproduces only its smallness. That smallness is itself an asset. ${}^6\text{Li}$ is nearly spherical, so tensor effects of geometric origin (the deformation term of the coherent channel, §4.2) are bounded and a tensor signal must be dynamical in origin; ${}^7\text{Li}$, whose p-shell triton orbit gives $Q = -4.00 \text{ fm}^2$ [36], is the deformed counterpart. Finally, the two isotopes share one source, one injector and one detector: the α +d versus α +t comparison changes only the cluster, the cleanest controlled variation the program has.

1.3 • Why polarization is indispensable

An unpolarized lithium beam would be one more light ion. Its inclusive cross section is carried by F_1 and F_2 alone — for an unpolarized target the spin-1 cross section has exactly the spin-1/2 form [2] — and it would add an $A = 6, 7$ point to the EMC and shadowing curves that ${}^4\text{He}$ and ${}^{12}\text{C}$ already map. Every item of the program is instead a spin observable, and the two flagship items need the tensor moment specifically. b_1 is *defined* as the difference between the $m = 0$ and $m = \pm 1$ quark distributions; averaging over m erases it, and it can be reached only with tensor polarization — which is why the only existing measurement is HERMES, with the only tensor-polarized target ever used in DIS [5]. Δ is generated by gluon transversity, an interference between gluon helicity states that requires a target spin of at least 1 [3], and its $\cos 2\phi$ signature needs a transverse alignment axis to define the azimuth: the inclusive cross section of an unpolarized target has no azimuthal dependence at all. The polarized EMC effect is by construction a statement about g_1 , and the gluon-spin EMC effect enters only through the Q^2 evolution of g_1 [16]. In tagged DIS the spectator momentum selects S- versus D-wave configurations of the polarized pair, removing depolarization at small p_T and producing tensor asymmetries of order unity near 300 MeV/c [41]: the spin is the signal. Since the EIC project polarizes only protons and ${}^3\text{He}$, with deuterons to follow [29], polarized lithium would be the first polarized ion at a collider with a nuclear core — the first place where the spin structure of *a nucleus*, rather than of a nucleon, is studied at collider kinematics. Tensor polarization is the scarcer resource (source targets $P_{zz} \geq 0.80$ against $P_z \geq 0.90$ [51], with ring transport still to be established), and both b_1 and Δ depend on it.

1.4 • Fixed target or collider?

Polarized lithium has existed for decades as a fixed target, but only in the solid state. Dynamically polarized ${}^6\text{LiD}$ was the deuteron target of SLAC E155 and of COMPASS, ${}^7\text{LiH}$ has been polarized at Saclay (47% for ${}^7\text{Li}$, 56% for the protons) [44], and dynamically polarized ${}^7\text{LiD}$ (with ${}^6\text{LiH}$ or NH_3 as the proton reference) is the material of the approved CLAS12 polarized-EMC experiment [43,42,45]. In these compounds the lithium polarizes along with the hydrogen isotope because all spin species share one spin temperature: COMPASS reached deuteron polarizations of $+54\%/-47\%$, and the equal-spin-temperature relation gives $51 \pm 3\%$ for ${}^6\text{Li}$ and $92 \pm 4\%$ for the ${}^7\text{Li}$ admixture [42]; E155 measured the ${}^6\text{Li}$ polarization at 97% of the deuteron's [43]; in ${}^7\text{LiD}$ at 5 T the CLAS12 proposal expects 65–80% for ${}^7\text{Li}$ [45]. Three limits are structural. (i) Dilution: only a

fraction $f \approx 0.35\text{--}0.45$ of the nucleons in ${}^6\text{LiD}$ are polarizable once the helium coolant is counted — better than 0.22 for ND_3 and 0.15 for NH_3 , but far from the 1.0 of a stored beam [42,43]. (ii) Tensor polarization: in thermal equilibrium the spin-1 tensor moment follows from the vector one, $P_{zz} = 2 - \sqrt{4 - 3P_z^2}$, so $P_z = 0.5$ gives $P_{zz} = 0.20$; radio-frequency "hole burning" has raised it to 29–37% in deuterated butanol at the cost of the vector polarization, and the JLab b1 experiment is conditionally approved on demonstrating 30% [8,46]. An atomic-beam source prepares the tensor moment directly: HERMES ran its gas target at $P_{zz} = +0.89$ and -1.65 in the pure-tensor states, "a combination which is not possible for solid-state polarized targets" [5], and the lithium source targets $P_{zz} \geq 0.80$ [51]. (iii) Spectator loss: a spectator α of 100–300 MeV/c carries only 1.3–12 MeV in the target rest frame and stops within 10–250 μm of lithium deuteride (NIST ASTAR ranges [49]) — inside target cells 2–5 cm long at JLab and 30–60 cm long at COMPASS — and a coherent ${}^6\text{Li}$ recoil at $|t| = 0.05 \text{ GeV}^2$ has only 4.5 MeV. The JLab recoil detectors that do tag slow spectators, BoNuS (protons from 70 MeV/c) and ALERT (α from 230 MeV/c), work with unpolarized gas at a few atmospheres [47,48]; polarized-lithium targets outside dynamically polarized solids are essentially unknown — a 1975 polarized- ${}^6\text{Li}$ target for a low-energy reaction study [50] is the one instance on record — and none has been used in lepton scattering. The collider removes each limit at once. The beam is pure lithium with its polarization prepared atom by atom and measured in the ring; the spectator α or triton carries the beam momentum per nucleon and emerges at a few milliradians into the Roman Pots, off-momentum detectors, BO and zero-degree calorimeter, with $\sim 10\%$ transverse-momentum resolution at 100 MeV/c [22]; the intact coherent recoil can be tagged down to $p_T \approx 0$ with the IR-8 secondary focus [27]; and the centre-of-mass energy — $\sqrt{s_{NN}}$ up to 99 GeV for $e(18 \text{ GeV}) \times {}^6\text{Li}(137.5 \text{ GeV}/u)$ — opens x down to $\sim 10^{-4}$ at $Q^2 \geq 1 \text{ GeV}^2$, where JLab at 11 GeV stops at $x \approx 0.06$ and HERMES stopped at 0.01 [45,5]. That is the region where gluons dominate, where HERMES saw its large b_1 , and where coherence holds. What stays with the fixed target is luminosity: $10^{35} \text{ cm}^{-2} \text{ s}^{-1}$ with dynamically polarized targets (CLAS12 at 10–15 nA, Hall C at 115 nA, with the material annealed every 2–3 days) [45,8] against an electron–nucleon luminosity of $10^{33}\text{--}10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ at the EIC [29,25]. The high- x polarized EMC effect and the high- x b_1 of the deuteron are therefore rightly JLab measurements, and the EIC program concentrates on what the table summarizes: the low- x and high- Q^2 regions, the tensor observables at full polarization, and everything that needs a detected spectator or recoil.

MEASUREMENT	AT A FIXED TARGET	WHAT ONLY THE COLLIDER ADDS
Polarized EMC effect, ${}^7\text{Li}$ (g_A)	Approved: JLab E12-14-001 (CLAS12, polarized ${}^7\text{LiD}$, $0.06 < x < 0.8$, $1 < Q^2 < 15 \text{ GeV}^2$, $P_z \approx 0.8$ assumed, 55 days) [45]	$x < 0.06$; the Q^2 evolution that carries the gluon-spin EMC effect [16]; an undiluted beam; α -tagging to identify the struck triton and control its virtuality (IP6 Roman Pots)
Tensor b_1 , ${}^6\text{Li}$	Deuteron only: E12-13-011 (ND_3 , $0.16 < x < 0.49$, $P_{zz} \geq 0.3$ required) [8]; ${}^6\text{LiD}$ would give $P_{zz} \approx 0.2$ with the deuteron term to be subtracted — no such measurement has been proposed	x down to 10^{-3} , where HERMES found $b_1 \approx 0.1$; pure-tensor fills at $P_{zz} \approx 0.6\text{--}0.8$; spin axis along $\mathbf{q}$ [26]
Exotic glue Δ , ${}^6\text{Li}$ (${}^7\text{Li}$)	Letter of intent only: LOI12-16-006 (${}^{14}\text{N}$ in NH_3 , $x < 0.3$, 2–12% alignment, in-plane spectrometers — $\cos 2\varphi$ "difficult with a fixed target") [13]	Transverse tensor polarization ≥ 0.6 with full 2π azimuthal acceptance; the gluon-dominated region $x \lesssim 0.1$ and a Q^2 lever arm
Spectator-tagged DIS (α , t)	Not feasible: spectators stop within 0.25 mm of a solid target; no polarized lithium gas target exists	Far-forward detection of α and t at beam velocity; pole extrapolation to the free cluster [20,22]
Coherent intact- ${}^6\text{Li}$ channel	Not feasible: needs $x \lesssim 0.01$ and a 2–9 MeV recoil	Roman-Pot p_T tail or the IR-8 secondary focus [27]

1.5 • What each observable requires of the beam, and what lithium alone provides

The requirements below are the program's own, from the fast simulation and the polligen pseudo-experiments (§5), stated for the placeholder in-ring polarizations $P_z = 0.7$ and $P_{zz} = 0.6$, $P_e = 0.7$, and $10 \text{ fb}^{-1}/u$. Every statistical error scales as $1/(P\sqrt{L})$, so a lower polarization is paid for in luminosity squared: halving P_{zz} costs a factor four in running time. Two machine facts shape the list. The stable spin direction in the arcs is vertical, so

transverse polarization at the interaction point is native while longitudinal polarization needs spin rotators, whose lithium settings are not yet defined [29,1]; and tensor fills need $m = 0$ bunches prepared by the source radio-frequency transitions, whose survival through acceleration and bunch-by-bunch patterning are open questions [51]. The observables also demand different spin configurations and therefore share the luminosity; the reach quoted assumes each receives the full $10 \text{ fb}^{-1}/\text{u}$ in its own configuration. (Semi-inclusive channels add the three leading-twist tensor TMDs of a spin-1 target [1]; they need the same tensor polarization and are not projected here.)

OBSERVABLE	BEAM POLARIZATION REQUIRED	ELECTRONS	REACH AT $10 \text{ fb}^{-1}/\text{u}$	BINDING REQUIREMENT
g_1^A , polarized EMC (${}^7\text{Li}$; ${}^6\text{Li}$ for the isoscalar pair)	Vector $P_z \geq 0.7$, longitudinal (spin rotators). A spin-3/2 spin-temperature fill at $P_z = 0.8$ also carries $P_{zz} \approx 0.55$ and $P_{zzz} \approx 0.11$; the octupole term enters the asymmetry numerator and is corrected through the fill model [45]	Polarized, $P_e \approx 0.7$, helicity flipped bunch by bunch	$\delta\Delta R \approx 2.6\text{--}4\%$ per x-bin at $x = 0.3\text{--}0.5$ (three energies combined); CBT-vs-TMT $\approx 1.5\sigma$ per bin, 5σ at $x \approx 0.5\text{--}0.7$ with $100 \text{ fb}^{-1}/\text{u}$	Polarimetry scale $\delta P_z/P_z \leq 3\%$; P_p from VMC [17]; Q^2 lever arm at fixed x (gluon-spin EMC)
b_1 via A_{zz} (${}^6\text{Li}$)	Tensor P_{zz} , longitudinal or along $\mathbf{q}$ [26]; fills of different P_{zz} ("tensor thirds": $m = +1, 0, -1$ -rich), i.e. $m = 0$ bunches from the source	Unpolarized (helicity-averaged)	$\delta A_{zz} \approx 1 \times 10^{-4}$ per x-bin at $x < 0.05$ (7×10^{-5} at $P_{zz} = 0.8$) against scenario $ A_{zz} = (2\text{--}10) \times 10^{-3}$ (HERMES-like b_1 with the 0.29 dilution) or ten times smaller (convolution): both $\geq 10\sigma$ per bin at $x < 0.05$. At $x > 0.3$, $\delta A_{zz} \approx 10^{-3}$ against a signal $\leq 2 \times 10^{-3}$ — JLab's domain	Relative luminosity between fills at the 10^{-4} level (a 10^{-4} offset biases A_{zz} by 1×10^{-4} ; the luminosity-corrected estimator removes it exactly); tensor polarimetry, which needs dedicated development [1]
Δ via $\cos 2\phi$ (${}^6\text{Li}$; ${}^7\text{Li}$ eligible)	Tensor P_{zz} with the alignment axis transverse (the native vertical direction); fills of opposite P_{zz} sign flip the modulation — the decisive systematic check	Unpolarized	$\delta A = (1.5\text{--}4.5) \times 10^{-4}$ per sweet-spot bin ($x \approx 0.02\text{--}0.14$, $Q^2 = 1.1\text{--}14 \text{ GeV}^2$) against moment-constrained amplitudes of $(0.7\text{--}1.2) \times 10^{-2}$, $\geq 21\sigma$; a 3×10^{-3} flat amplitude stays $3\text{--}10\sigma$ per bin even at $P_{zz} = 0.3$	Full azimuthal coverage for the scattered electron (the binned fit tolerates holes); $\delta P_{zz}/P_{zz} \approx 3\%$ scale; tensor radiative corrections not yet calculated
Spectator-tagged DIS	${}^7\text{Li}$ α -tag: any polarization for tagged F_2 ; vector P_z with polarized electrons for tagged g_1 of the bound proton. ${}^6\text{Li}$ α -tag: tensor P_{zz} for the order-unity tagged tensor asymmetries at $p_s \approx 300 \text{ MeV}/c$ [41]	As for the inclusive observable	${}^7\text{Li}$ $\alpha \rightarrow$ Roman Pots ($R = 0.857$), 96–99% acceptance, optics-robust; ${}^6\text{Li}$ α beam-blind at IP6 (3–9% with high-acceptance optics, 1–2% with high-divergence) $\rightarrow$ IR-8 secondary focus	Spectator p_T resolution $\sim 10\%$ at $100 \text{ MeV}/c$ [22]; charge identification of α against $A/Z = 2$ fragments in the Roman Pots (open); pole extrapolation in p_T [20]
Coherent intact- ${}^6\text{Li}$ $\cos 2\phi$	Tensor $P_{zz} \geq 0.6$, transverse	Unpolarized	$x \lesssim 0.01$, $ t < 0.2 \text{ GeV}^2$; 1.1×10^7 tagged events (high-acceptance optics; $f_0 = 0.04$ scenario), best-bin $\delta A = 1.9 \times 10^{-3}$ (5σ floor at a 1.0% modulation); deformation-anchored $\langle a_2 \rangle = 0.036$ (band 0.018–0.059); a flat gluonic $3 \times 10^{-3}\text{--}10^{-2}$ gives $1.6\text{--}5\sigma$, or $5\text{--}16\sigma$ with $100 \text{ fb}^{-1}/\text{u}$	High-acceptance optics (13.5% acceptance against 4×10^{-5} with high divergence) or IR-8; Roman-Pot p_T cut $\leq 0.2 \text{ GeV}$; Z-identification against $\alpha+d$ (both $R = 1$); B0 calorimeter veto of the 3.56 MeV de-excitation γ

What lithium alone provides is clearest species by species. Of the ions the EIC can polarize, only the deuteron and the lithium isotopes have $J \geq 1$, so b_1 and Δ do not exist for protons or ${}^3\text{He}$ at all; among $J \geq 1$ nuclei ${}^6\text{Li}$ is the only $A > 2$ case, and the lattice indication that the deuteron's exotic glue is suppressed [11] is what makes a bound nucleus, rather than the deuteron, the likelier place for a discovery [1]. ${}^7\text{Li}$ is the only bound polarized proton within reach of any collider; ${}^6\text{Li}$ and the deuteron supply the isoscalar companions, so the three together separate the u- and d-quark spin modification — an inference of this program, not yet in the literature. For tagging, ${}^7\text{Li}$ gives the only light-ion spectator that is a spin-0, tightly bound nucleus landing in the Roman-Pot window today, leaving a struck $A = 3$ cluster. In the coherent channel the deuteron's deformation term

dominates ($Q = +0.286 \text{ fm}^2$), ${}^7\text{Li}$ is the deformed reference ($Q = -4.00 \text{ fm}^2$), and ${}^6\text{Li}$ ($Q = -0.081 \text{ fm}^2$) is the deformation null in which a flat-in-t modulation can only be exotic glue.

OBSERVABLE	P	D	${}^3\text{He}$	${}^6\text{Li}$	${}^7\text{Li}$
g_1^A , polarized EMC	free reference	isoscalar pair, weakly bound; D-wave depolarization of a few per cent [20]	neutron ($P_n = 0.86-0.88$) at $A = 3$	isoscalar pair, $P = 0.85$, in the α field	the bound polarized proton , $P_p = 0.87$, at normal density
b_1	—	HERMES; EIC deuterons a later addition	—	the only $A > 2$ case ; ${}^6\text{Li} - d$ isolates the α field	rank-2 analog, no published formalism
Δ (exotic glue)	— ($J = 1/2$)	allowed; lattice suggests suppression [11]	—	allowed, bound ; spin-1 like the ϕ meson of the lattice signal [10]	allowed; $J = 3/2$ as in the bag estimate [9]
Spectator tagging	—	p, n spectators (off-momentum detectors, ZDC)	p spectators into the Roman Pots; double tag [21]	α beam-blind $\rightarrow$ IR-8; tagged tensor asymmetries $O(1)$	α into the Roman Pots now ; struck $A = 3$ cluster
Coherent tensor $\cos 2\varphi$	—	deformation term dominant [23]	—	deformation null ($ Q 3.5\times$ smaller than d, opposite sign)	deformed reference

What an unpolarized lithium beam already gives. Three things, none of them the driver. First, nuclear parton distributions at $A = 6$ and 7 : the high- x EMC ratios of ${}^6\text{Li}$ and ${}^7\text{Li}$ are in the target list of JLab E12-10-008 [54], motivated by the E03-103 finding that the ${}^9\text{Be}$ EMC effect follows the local density of its α clusters rather than its average density [53] — for ${}^6\text{Li}$ and ${}^7\text{Li}$ the local-density picture predicts an EMC effect well below the other models [54] — and the EIC adds the low- x shadowing region. Second, cluster virtuality: the α -tagged F_2 of the triton (${}^7\text{Li}$) or deuteron (${}^6\text{Li}$) cluster versus spectator momentum measures the α -t and α -d momentum distributions that VMC predicts ($N_{\alpha t} = 1.00$ in a relative p-wave, $N_{\alpha d} = 0.86$) [17], the collider counterpart of the JLab tagged-EMC program [48]. Third, coherent diffraction on an α -clustered nucleus, whose t-distribution images the gluon density of the cluster, as studied for ${}^7\text{Li}$ at IR-8 [27]. Polarization turns the same cluster structure from an object of study into a tool: b_1 , Δ , g_1^A and the tagged tensor asymmetries all vanish or are undefined without it (§1.3).

2 • Deep inelastic scattering

An electron with four-momentum k scatters from a nucleus of momentum P by exchanging a virtual photon $q = k - k'$. The kinematic invariants, in the per-nucleon convention used throughout the program (energies and structure functions per nucleon, $x \leq 1$), are

$$Q^2 = -q^2, \quad x = \frac{Q^2}{2P \cdot q}, \quad y = \frac{P \cdot q}{P \cdot k}, \quad Q^2 = xy s, \quad W^2 = M^2 + Q^2 \frac{1-x}{x}$$

where x is the momentum fraction of the struck parton, y the energy-transfer fraction, and W the invariant mass of the hadronic final state. The unpolarized cross section is carried by two structure functions,

$$\frac{d^2\sigma}{dx dQ^2} = \frac{4\pi\alpha^2}{xQ^4} \left[\left(1 - y + \frac{y^2}{2}\right) F_2(x, Q^2) - \frac{y^2}{2} F_L(x, Q^2) \right], \quad F_1 = \frac{F_2}{2x(1+R)}$$

with $R = \sigma_L/\sigma_T$. In the parton model, $F_2 = \sum_q e_q^2 x[q(x) + \bar{q}(x)]$: the cross section measures quark distributions weighted by squared charge. For a nucleus we use F_2^A/A , so rates are quoted against luminosity per nucleon. Polarization introduces additional structure functions beyond F_1 and F_2 ; their number and character depend on the target spin.

3 • Spin-1 targets


A spin- $1/2$ target contributes two spin structure functions, g_1 and g_2 . The principal observable is g_1 , the helicity-weighted quark distribution, measured through the double-spin asymmetry $A_{//} \approx D \cdot g_1/F_1$ (D is a kinematic

depolarization factor) with both beams longitudinally polarized and the electron helicity reversed bunch by bunch.

A spin-1 target carries additional structure. Its spin state along a chosen axis is specified by three populations p_+ , p_0 , p_- of the projections $m = +1, 0, -1$, and two moments characterize the fill:

$$P_z = p_+ - p_- \in [-1, 1], \quad P_{zz} = p_+ + p_- - 2p_0 = 1 - 3p_0 \in [-2, 1]$$

P_z is the familiar vector polarization. P_{zz} , the *tensor* or alignment polarization, is specific to $\text{spin} \geq 1$: it distinguishes spin along $\pm z$ from spin in the transverse plane ($m = 0$), and it is nonzero even when equal numbers point up and down. An unpolarized beam has populations ($1/3, 1/3, 1/3$) and both moments vanish; a pure $m = 0$ beam has $P_z = 0$ but $P_{zz} = -2$.



Hoodbhoy, Jaffe, and Manohar showed [2] that a spin-1 target has, beyond F_1 , F_2 , g_1 , g_2 , four tensor-polarized structure functions b_1 – b_4 , and Jaffe and Manohar identified one more, the double-helicity-flip function Δ [3]. At leading twist the tensor sector reduces to b_1 (with $b_2 = 2x b_1$ as its Callan–Gross partner) and Δ . For unpolarized electrons on a target whose spin axis makes angle θ_m with the beam, the cross section for each pure m state is [2,3]

$$\frac{d\sigma^{(m)}}{dx dy d\phi} = \frac{2y\alpha^2}{Q^2} \left[F_1 + \frac{2}{3} a_m b_1 + \frac{1-y}{xy^2} \left(F_2 + \frac{2}{3} a_m b_2 \right) - \frac{1-y}{y^2} c_m \sin^2 \theta_m \Delta(x, Q^2) \cos 2\phi \right]$$

$$a_m = \frac{1}{4} c_m (3 \cos^2 \theta_m - 1), \quad c_m = 3|\lambda_m| - 2 \rightarrow (1, -2, 1) \text{ for } m = (+1, 0, -1)$$

The weights $(1, -2, 1)$ organize all tensor observables: the $m = \pm 1$ states enter together and against $m = 0$, precisely the combination measured by P_{zz} . Averaging the formula over a fill with populations p_m replaces every c_m by a factor P_{zz} . Two experimental handles follow. With the spin axis *longitudinal*, the rate difference between fills gives $A_{zz} \propto b_1$; with the spin axis *transverse* ($\theta_m = 90^\circ$), the ϕ distribution of the scattered electron acquires a $\cos 2\phi$ modulation proportional to Δ . Both require tensor polarization; neither requires polarized electrons.

^7Li , with spin $3/2$, has four populations and three moments (vector, tensor/quadrupole, and octupole). Its rank-0 and rank-1 sectors are exact analogs of the spin- $1/2$ case, and host the polarized EMC measurement; its rank-2 sector is the formal analog of the spin-1 tensor sector, for which no structure-function basis has been published — an open theory item the program tracks.

4 • The measurement program

4.1 • The tensor structure function b_1

In parton language, b_1 compares quark distributions in an aligned versus a transverse nucleus [2]:

$$b_1(x) = \frac{1}{2} \sum_q e_q^2 \left[q^{(0)}(x) - \frac{q^{(+1)}(x) + q^{(-1)}(x)}{2} \right], \quad \int_0^1 b_1(x) dx = 0 \quad (\text{Close – Kumano, no tensor sea})$$

where $q^{(m)}$ is the quark distribution in a target with spin projection m . For a collection of free spin- $1/2$ nucleons, or for a pure S-wave bound state, all m states are equivalent and $b_1 = 0$. A nonzero b_1 therefore measures directly how binding modifies the partonic content: D-wave admixture, pion exchange, hidden-color components, or a tensor-polarized sea (which would violate the Close–Kumano sum rule [4]).

The only existing measurement is HERMES on the deuteron [5]: A_{zz} up to 2×10^{-2} over $0.01 < x < 0.45$ and $b_1(x = 0.012) = 0.112 \pm 0.055 \pm 0.028$, unexpectedly large at low x , with a zero crossing near $x \approx 0.2$ and a sum-rule integral 2σ from zero. Standard convolution calculations predict $|b_1| < 10^{-3}$ at $x \gtrsim 0.2$, an order of magnitude below the data [6], whereas a model with pions and a hidden-color probability of only 0.15% reproduces it [7]. The tension is unresolved; JLab E12-13-011 will remeasure the region $0.16 < x < 0.49$ with a solid ND₃ target [8], and the EIC adds the low- x region, where HERMES observed the large signal, together with the Q^2 lever arm.

Measurement procedure. The beam is delivered in fills of different tensor polarization ("tensor thirds"). With the estimator $(N_+ + N_- - 2N_0)$ over matched luminosities, all m -independent physics cancels, and the statistical uncertainty on A_{zz} is $\sqrt{(2/N)/P_{zz}}$. ${}^6\text{Li}$ would provide the first b_1 of any $A > 2$ nucleus, for which no prediction exists, and the difference between ${}^6\text{Li}$ and the free deuteron isolates the effect of embedding in the α field. One systematic is tracked from the outset: at collider kinematics a clean b_1 extraction favors orienting the spin axis along the momentum transfer $\mathbf{q}$ rather than along the beam [26].

4.2 · $\Delta(x, Q^2)$: exotic glue

The $\cos 2\phi$ term in the master formula corresponds to the virtual photon flipping its helicity by two units between absorption and emission in the forward Compton amplitude. A spin- $1/2$ particle cannot absorb two units of helicity, so neither quarks nor bound nucleons generate Δ at leading twist; gluons can. Δ is the structure function that the gluon transversity distribution, an interference of gluon helicity states existing only in targets with $J \geq 1$, generates at order α_s through a photon–gluon convolution [3,9]. A nonzero Δ in a nucleus is therefore direct evidence for glue not inherited from individual nucleons, hence the name "exotic glue". It has not been measured for any target; the one prior experimental document is a JLab letter of intent [13].

The measurement uses unpolarized electrons on a tensor-polarized nucleus whose alignment axis lies in the transverse plane, and fits the azimuthal distribution about that axis. Averaged over the fill, the modulation amplitude and its inversion back to Δ are

$$A^{\cos 2\phi} = -\frac{1-y}{y^2} \frac{\Delta}{D_\phi}, \quad D_\phi = F_1 + \frac{1-y}{xy^2} F_2$$

The expected size spans per-mille to percent. The only absolute normalizations available are moment estimates: a bag-model calculation by Sather and Schmidt for the spin- $3/2$ Δ^{++} baryon, $\int x \Delta dx = -0.012 \alpha_s$ [9], and lattice-QCD calculations by Detmold and Shanahan of gluon transversity in the ϕ meson (clearly nonzero, albeit at an unphysical pion mass) [10], with NPLQCD finding the analogous matrix element roughly ten times suppressed in the deuteron [11]. Percent-level asymmetries appear only in proton–deuteron Drell–Yan under the maximal assumption $\Delta_{Tg} = \Delta_g$, described by its authors as optimistic [12]. No calculation exists for ${}^6\text{Li}$ or any $A > 2$ nucleus, and the literature brackets the nuclear Δ in both directions: suppressed, as in the lattice deuteron, or enhanced by binding in analogy with the EMC effect [1]. The projection report transfers the bag moment to ${}^6\text{Li}$ as a labeled scenario, shows that one EIC year measures the amplitude at $\geq 20\sigma$ per bin, and details the caveats.

The coherent channel. The same modulation can be measured in coherent diffraction, $e {}^6\text{Li} \rightarrow e' X {}^6\text{Li}(g.s.)$, in which the nucleus remains in its ground state and recoils intact. There the $\cos 2\phi$ modulation mixes two mechanisms: the nuclear *shape*, in which a deformed gluon density modulates the diffractive slope (the mechanism computed for polarized deuterons in the only existing calculation of this kind [23]), and exotic glue proper. ${}^6\text{Li}$ discriminates between the two because its quadrupole moment is anomalously small ($Q = -0.0806 \text{ fm}^2$, roughly 50 times smaller than that of ${}^7\text{Li}$): the shape term is bounded and linear in t , so a flat-in- t modulation cannot be a shape effect. Detection of the intact recoil is itself nontrivial: the nucleus keeps the beam rigidity, as any coherent recoil does, and emerges within $\sim 2 \text{ mrad}$ of the beam, visible only in the Roman-Pot transverse-momentum tail at IP6 or at the secondary focus of a second interaction region [25,27]; the companion report treats this in detail.

4.3 • The polarized EMC effect: g1 of a bound proton

The EMC effect — the ~10–20% suppression of F2 per nucleon in a nucleus at 0.3 ≲ x ≲ 0.7 — has been measured for forty years without a settled explanation. Its spin analog has never been measured at all: does the *spin* structure of a bound nucleon change more, less, or the same as its momentum structure? The observable is the polarized EMC ratio

ΔRA(x) = g1^A(x) / (Pp g1^p(x) + Pn g1^n(x))

where Pp, Pn are the effective nucleon polarizations from nuclear structure. 7Li is well suited: the variational Monte Carlo spin-projected nucleon numbers of Wiringa et al. give Pp = +0.87 and Pn = -0.04 (0.866 and -0.037 as quoted in the JLab proposal) [17,45], so to a good approximation polarized 7Li is one polarized proton plus six spectators, with computable corrections. Two classes of models make sharply different predictions: Cloët–Bentz–Thomas (mean-field, NJL) obtain a polarized EMC effect about twice the unpolarized one [14], whereas Tronchin–Matevosyan–Thomas obtain equal effects [15]. A measurement at the 5% level discriminates between them, and the two map onto the competing mean-field and short-range-correlation pictures of the EMC effect itself. A handle unique to the EIC is the Q2 evolution of g1, driven by the polarized gluon distribution; the predicted *gluon-spin* EMC effect is larger still, with 7Li identified as the most promising case [16]. JLab E12-14-001 will measure 7Li at x > 0.06 [45]; the EIC extends the coverage to x < 0.06 and adds the Q2 arm, in collider kinematics free of target-fragment absorption.

For this observable both beams are polarized longitudinally and the electron helicity flips bunch-by-bunch; the error scales as 1/(PePz√N), so it uses the vector polarization the source delivers most readily.

4.4 • Spectator-tagged DIS

The cluster description of 6Li is quantitative (§1.2): the α is inert, and the deuteron-like pair carries the nuclear spin with a polarization of 0.85–0.9 in six- and three-body calculations [17,18,19], the 0.87 adopted by E155 [43] being the value the program uses. When the α spectator is detected (it carries beam velocity into the far-forward spectrometers), the struck object is identified as the deuteron cluster, and the 6Li beam acts as a source of polarized deuterons at controlled virtuality. The formal framework is spectator tagging (Cosyn–Weiss [20]): the spectator momentum is measured and extrapolated in its transverse component to the free-nucleon (here free-cluster) pole, recovering free-particle structure with binding effects removed by construction rather than modeled. Tagged *tensor* asymmetries are predicted to reach O(1) at spectator momenta of ~300 MeV/c, where the D-wave dominates [41] — far larger than the inclusive asymmetries above. The benchmark for the method is the e+3He double-tagging study of Frišić et al. [21], in which tagging improved on effective-polarization unfolding by a factor of 2–3, and by a factor of 10 at low x.

Far-forward routing determines where each isotope's tag fragment is detected, and it follows from magnetic rigidity alone: a beam-velocity fragment with mass-to-charge ratio A/Z bends in the hadron optics according to R = (A/Z)frag/(A/Z)beam.

BEAM	TAG FRAGMENT	R	DESTINATION AT IP6
7Li (A/Z = 7/3)	α	0.857	Roman Pots main window (R ∈ 0.60–0.95) — works now
7Li	t	1.286	over-rigid, no coverage
6Li (A/Z = 2)	α or d	1.000	beam rigidity: inside the beam envelope except the pT tail
6Li	p / n	0.50 / -	Off-Momentum Detectors / ZDC (veto handles)


The tagging program therefore begins with 7Li at IP6, while 6Li α-tagging (and the coherent intact-6Li channel, which faces the same rigidity problem) motivates the secondary focus of the second interaction region [27]. This asymmetry between the isotopes was itself a finding of the simulation program.

5 • Rates and statistical precision

All projections use luminosity per nucleon, with $10 \text{ fb}^{-1}/u$ — 30 weeks of operation at $10^{33} \text{ cm}^{-2} \text{ s}^{-1}$ in the Yellow Report's accounting [25], rounded here to "one EIC year" — and $100 \text{ fb}^{-1}/u$ for the ten-year program, and in-ring polarization placeholders $P_z = 0.7$ and $P_{zz} = 0.6$ (ring transport from the source values is an open accelerator question). The statistical uncertainties behind every money plot are


$$\delta A_{\parallel} = \frac{1}{P_e P_z \sqrt{N}}, \quad \delta A_{zz} = \frac{\sqrt{2/N}}{P_{zz}}, \quad \delta A^{\cos 2\phi} = \frac{\sqrt{2/N}}{P_{zz}}$$

The factor $\sqrt{2}$ in the tensor uncertainties follows from $\langle \cos^2 2\phi \rangle = 1/2$ (equivalently, from the three-fill estimator variance); the polarization appears in the denominator because the fill dilutes the physical asymmetry. These formulas, and their behavior under realistic ϕ -acceptance holes (where a binned fit replaces the naive moment), are validated end to end by pseudo-experiments with the polligen event generator. Tensor observables enjoy a structural advantage in systematics: the $\cos 2\phi$ modulation is self-normalized within a single fill, so relative-luminosity errors largely cancel; what remains is the polarimetry scale ($\delta P_{zz}/P_{zz} \approx 3\%$) and radiative corrections, which for tensor observables have not yet been calculated.



Phase-space coverage. Expected events per (x, Q^2) bin in one EIC year at $e(10 \text{ GeV}) \times {}^6\text{Li}(50 \text{ GeV}/u)$. Left: inclusive DIS (4.6×10^9 accepted events) with the bins of the $\cos 2\phi$ analysis overlaid; the measurement is limited by polarization, acceptance, and systematics rather than by statistics. Right: the coherent intact- ${}^6\text{Li}$ channel (1.1×10^7 tagged events), confined to $x \lesssim 0.03$ because coherence requires the photon to resolve the whole nucleus, a condition that fails once $x \gtrsim 0.01$.

Nuclear gluonometry, transversely tensor-polarized ${}^6\text{Li}$, $e(10) \times 6\text{Li}(50/u)$, $P_{zz} = 0.60$
moment_A(variant=mid_x, c=-0.012, A=-0.2941, q2_ref=3.913, dilution=0.3333); ϕ' at 1 yr (10 fb $^{-1}/u$), amplitudes 1 & 10 yr — stat. only; bkg & tensor RC unquantified (plans/0)



Projected extraction. $\cos 2\phi'$ pseudo-data (one EIC year, $P_{zz} = 0.6$) in the four highest-significance (x , Q^2) bins, with the injected moment-constrained Δ model and the binned fit; right, the extracted amplitude versus x against alternative model readings. The projected precision distinguishes between the models rather than only from zero. Assumptions and caveats are detailed in the companion report.

6 • Further reading

Formalism. [2] P. Hoodbhoy, R. L. Jaffe, A. Manohar, Nucl. Phys. B 312 (1989) 571 — the spin-1 DIS structure-function basis. [3] R. L. Jaffe, A. Manohar, Phys. Lett. B 223 (1989) 218 — nuclear gluonometry. [4] F. E. Close, S. Kumano, Phys. Rev. D 42 (1990) 2377 — the b_1 sum rule.

b_1 . [5] A. Airapetian et al. (HERMES), Phys. Rev. Lett. 95 (2005) 242001, hep-ex/0506018 (target polarizations in Table I). [6] W. Cosyn, Y.-B. Dong, S. Kumano, M. Sargsian, Phys. Rev. D 95 (2017) 074036. [7] G. A. Miller, Phys. Rev. C 89 (2014) 045203. [8] K. Slifer et al., JLab PR12-13-011 / E12-13-011, "The deuteron tensor structure function b_1 " (PAC40, 2013; conditional approval requiring $P_{zz} \geq 30\%$); D. Keller, D. Crabb, D. Day, Nucl. Instrum. Meth. A 981 (2020) 164504 (tensor-enhancement results).

Exotic glue. [9] E. Sather, C. Schmidt, Phys. Rev. D 42 (1990) 1424. [10] W. Detmold, P. E. Shanahan, Phys. Rev. D 94 (2016) 014507; W. Detmold, D. Pefkou, P. E. Shanahan, Phys. Rev. D 95 (2017) 114515. [11] F. Winter et al. (NPLQCD), Phys. Rev. D 96 (2017) 094512. [12] S. Kumano, Q.-T. Song, Phys. Rev. D 101 (2020) 054011. [13] J. Maxwell et al., JLab LOI12-16-006, arXiv:1803.11206.

Polarized EMC. [14] I. C. Cloët, W. Bentz, A. W. Thomas, Phys. Lett. B 642 (2006) 210. [15] S. Tronchin, H. H. Matevosyan, A. W. Thomas, Phys. Lett. B 783 (2018) 247. [16] X. G. Wang, W. Bentz, I. C. Cloët, A. W. Thomas, J. Phys. G 49 (2022) 03LT01. [17] R. B. Wiringa, R. Schiavilla, S. C. Pieper, J. Carlson, Phys. Rev. C 89 (2014) 024305, arXiv:1309.3794 — VMC spin-up/spin-down nucleon numbers (Table I) and α -d, α -t overlaps; quoted as effective polarizations in [45]. [53] J. Seely et al., Phys. Rev. Lett. 103 (2009) 202301, arXiv:0904.4448 — EMC effect in ${}^3\text{He}$, ${}^4\text{He}$, ${}^9\text{Be}$, ${}^{12}\text{C}$ (JLab E03-103). [54] J. Arrington, D. Gaskell, A. Daniel et al., JLab PR12-10-008 / E12-10-008, "Detailed studies of the nuclear dependence of F_2 in light nuclei" (Hall C; targets include ${}^6\text{Li}$ and ${}^7\text{Li}$).

Nuclear structure of Li. [18] N. W. Schellingerhout, L. P. Kok, S. A. Coon, R. M. Adam, Phys. Rev. C 48 (1993) 2714; erratum Phys. Rev. C 52 (1995) 439 — three-body $\alpha+n+p$ models of polarized ${}^6\text{Li}$.

[19] B. S. Pudliner et al., Phys. Rev. C 56 (1997) 1720; S. C. Pieper, R. B. Wiringa, Annu. Rev. Nucl. Part. Sci. 51 (2001) 53 — QMC cluster structure of ${}^6\text{Li}$ and ${}^7\text{Li}$, spin-projected densities, $Q({}^6\text{Li})$ cancellation. [36] Nuclear data: N. J. Stone, Table of nuclear electric quadrupole moments, IAEA INDC(NDS)-0833 (2021) and At. Data Nucl. Data Tables 111–112 (2016) 1; Table of recommended nuclear magnetic dipole moments, INDC(NDS)-0794 (2019); IAEA Live Chart ground-state table (spins of stable nuclides); AME2020, M. Wang et al., Chin. Phys. C 45 (2021) 030003. [37] C. Forssén, E. Caurier, P. Navrátil, Phys. Rev. C 79 (2009) 021303(R). [38] G. Hupin, S. Quaglioni, P. Navrátil, Phys. Rev. Lett. 114 (2015) 212502. [39] S. Pastore, S. C. Pieper, R. Schiavilla, R. B. Wiringa, Phys. Rev. C 87 (2013) 035503. [40] J. Carlson et al., Rev. Mod. Phys. 87 (2015) 1067 (Table I: GFMC vs experiment for $A \leq 12$).

Tagging. [20] W. Cosyn, C. Weiss, Phys. Lett. B 799 (2019) 135035; Phys. Rev. C 102 (2020) 065204. [21] I. Friščić et al., Phys. Lett. B 823 (2021) 136726. [22] A. Jentsch, Z. Tu, C. Weiss, Phys. Rev. C 104 (2021) 065205. [26] W. Cosyn et al., Eur. Phys. J. A 61 (2025) 83, arXiv:2410.12764. [41] W. Cosyn, C. Weiss, Phys. Rev. C 114 (2026) 025201 and 025202, arXiv:2603.23699, 2603.23700 — *Semi-inclusive DIS on a polarized spin-1 target*, I. cross section and spin observables; II. deuteron and spectator nucleon tagging ("tensor-polarized spin asymmetries of order unity are achieved for spectator momenta $\gtrsim 300 \text{ MeV}$ "). [47] N. Baillie et al. (CLAS/BoNuS), Phys. Rev. Lett. 108 (2012) 142001. [48] ALERT run group, JLab E12-17-012, arXiv:1708.00888. [49] M. J. Berger et al., NIST Standard Reference Database 124 (ASTAR), doi:10.18434/T4NC7P.

Coherent channel. [23] H. Mäntysaari, F. Salazar, B. Schenke, C. Shen, W. Zhao, Phys. Lett. B 858 (2024) 139053. [24] Y. Hatta, B.-W. Xiao, F. Yuan, Phys. Rev. Lett. 116 (2016) 202301.

Polarized lithium sources. [30] E. Steffens et al., Nucl. Instrum. Meth. 143 (1977) 409 — Heidelberg ${}^6\text{Li}$ source. [31] H. Jänsch et al., Nucl. Instrum. Meth. A 254 (1987) 7 — polarized Li and Na beams. [32] A. J. Mendez et al., Nucl. Instrum. Meth. A 329 (1993) 37; E. G.

Myers et al., Nucl. Instrum. Meth. B 79 (1993) 701 — FSU optically pumped source. [33] D. Fick, G. Grawert, I. M. Turkiewicz, Phys. Rep. 214 (1992) 1 — nuclear physics with polarized heavy ions. [34] A. Zelenski et al., Nucl. Instrum. Meth. A 1055 (2023) 168494 — polarized $^3\text{He}^{++}$ EBIS source. [35] Hyperfine structure: E. Arimondo, M. Inguscio, P. Violino, Rev. Mod. Phys. 49 (1977) 31; M. Allegrini, E. Arimondo, L. A. Orozco, J. Phys. Chem. Ref. Data 51 (2022) 043102; D-line frequencies: R. C. Brown et al., Phys. Rev. A 87 (2013) 032504; ionization energy: NIST Atomic Spectra Database. [51] C. Peng et al. (ANL), "Development of high current and highly polarized $^6,7\text{Li}$ ion sources", proposal to DOE NOFO DE-FOA-0003602 (2026), [docs/ecrp_2026_proposal.pdf](#) .

Fixed targets. [42] J. Ball et al., Nucl. Instrum. Meth. A 498 (2003) 101 (COMPASS ^6LiD); P. Abbon et al. (COMPASS), Nucl. Instrum. Meth. A 577 (2007) 455; J. Koivuniemi et al., "Polarization build up in COMPASS ^6LiD target", SPIN2004 proceedings, p. 796. [43] P. L. Anthony et al. (E155), Phys. Lett. B 463 (1999) 339; S. Bültmann et al., Nucl. Instrum. Meth. A 425 (1999) 23. [44] P. Chaumette et al.,

AIP Conf. Proc. 187 (1989) 1275 — irradiated ^6LiD and ^7LiH . [45] W. K. Brooks, S. E. Kuhn et al., JLab PR12-14-001 / E12-14-001, "The EMC effect in spin structure functions" (PAC42, 2014); target: C. D. Keith, PoS(PSTP2017)008; C. D. Keith et al., PoS(PSTP2022)009. [46] see [8], Keller–Crabb–Day. [50] F. Wittchow, J. Ulbricht, U. Holm, K. D. Stahl, H. Ebinghaus, Phys. Lett. B 59 (1975) 29.

Machine. [1] G. Atoian et al. (EPIOS), Phys. Rev. C 113 (2026) 060501, arXiv:2510.10794. [25] R. Abdul Khalek et al., EIC Yellow Report, Nucl. Phys. A 1026 (2022) 122447. [27] W. Chang et al., Phys. Rev. D 113 (2026) 032018, arXiv:2511.05638. [28] E. Hamwi, G. H. Hoffstaetter, *Polarization transmission in the Electron-Ion Collider's Hadron Storage Ring*, Phys. Rev. Accel. Beams 29 (2026) 073501, arXiv:2509.18558 — resonance counts and snake requirements per species. [29] F. Willeke, J. Beebe-Wang (eds.), EIC Conceptual Design Report, BNL-221006-2021-FOR (2021), doi:10.2172/1765663; S. Nagaitsev, EIC Global Requirements, EIC-ORG-PLN-010 Rev. 03 (2026). [52] H. Huang et al., Phys. Rev. Accel. Beams 23 (2020) 021001 — polarized deuteron acceleration.