A review of the application of data mining techniques for decision making in agriculture

Niketa Gandhi University Dept. of Computer Science University of Mumbai Mumbai, Maharashtra, India niketa@gmail.com

Abstract—This paper provides a review of research on the application of data mining techniques for decision making in agriculture. The paper reports the application of a number of data mining techniques including artificial neural networks, bayesian networks and support vector machines. The review has outlined a number of promising techniques that have been used to understand the relationships of various climate and other factors on crop production. This review proposes that further investigations are needed to understand how these techniques can be used with complex agricultural datasets for crop yield prediction integrating seasonal and spatial factors by using GIS technologies.

Keywords—agriculture; artificial neural network; association rule mining; Bayesian network; data mining; decision making; support vector machines

I. INTRODUCTION

Many computer science techniques such as data mining and machine learning are used to study the influence of various parameters and make predictions of the crop production [1,2,3,4,5,6,7]. Data mining is the process of identifying the hidden patterns from large and complex data [8,9,10,11]. It may provide crucial role in decision making for complex agricultural problems.

A comprehensive review of the application of various techniques such as artificial neural networks, bayesian network, support vector machines and association rule mining will be provided with examples to show how these have been applied to variety of agricultural data sets. The role that these techniques play in the development of decision support tools for agricultural crop production predictions, pest and disease management, will also be explored. This paper reviews the application of data mining techniques applied in the field of agriculture. Recommendations for future research directions for the application of these techniques for decision making in agricultural context will be provided.

II. REVIEW OF LITERATURE

Knowledge discovery from databases and data analysis is an inter-disciplinary concept termed as data mining [12]. A multi-facet approach is used which can include statistical analysis, data visualization, neural networks, knowledge discovery, pattern recognition and data base management [13]. The widely used data mining techniques has been discussed in

Leisa J. Armstrong
University Dept. of Computer Science, University of
Mumbai and School of Science, Edith Cowan University
Perth, Western Australia
l.armstrong@ecu.edu.au

a review by [14]. These techniques have also been applied in the agricultural domain.

For example, a *k*-nearest neighbor technique was applied by [15] to evaluate forest inventories and to estimate forest variables analyzing satellite imagery. The simulation of daily precipitations with other weather variables and forecast of pollution in the atmosphere was carried out using the *k*-means method [16,17]. Classification of soil in combination with GPS-based technologies and plants has also been performed using *k*-means approach [18, 20]. The apples were checked before sending to market by using different classification approaches. Further *k*-means approach is used by [21] to analyse color images of fruits as they run on conveyor belts.

In Targhadia, Gujarat, India the impact of rainfall on crop productivity was studied by [19]. This study considered the dataset of 39 years from 1958 to 1996. Correlation and regression techniques were applied to derive the relationship between rainfall and crop yield. Crop yield prediction model for major crops was developed using the results of this study. In Pakistan, a study was carried out on the excessive usage of pesticides affecting the crop yield. Using data mining techniques interesting patterns of farmer practices were revealed from these clusters along with pesticide usage dynamics [22]. Data mining techniques has also been applied for Western Australian soil data to achieve optimal results compared to the statistical methods [23]. This study showed that the data mining techniques has taken less time compared to the statistical methods and have produced accurate results.

A study by [24] identified Powdery Mildew disease in mango using various data mining techniques. A novel unsupervised algorithm, called CLimate and rEmote sensing Association patteRnsMiner, for mining association patterns on heterogeneous time series from climate and remote sensing data integrated in a remote sensing information system was developed to improve the monitoring of sugar cane fields [25]. Rules generated by this new algorithm showed the association patterns for different periods in each time series. This indicated a time delay between the occurrences of patterns in the series analyzed. This could corroborate the forecasts of specialists without need to analyze numerous data charts.

Using wireless sensor network a simulated experiment was conducted by [26] to interpret the crop-pest-weather relations on groundnut pest thrips. The data was transformed using data mining techniques into effective relations between the dynamic

crop-weather-pest. Using regression models the obtained data was validated. A study by [27] uses data mining techniques on environmental and biotic input variables to predict annual yield of major crops. The study used the findings to make recommendations planting different crops in different districts in Bangladesh.

A number of recent studies have discussed the potential of applying data mining techniques in agriculture [28,29,30,31,32,33,34]. Good crop yield prediction results have been achieved by many researchers after applying data mining techniques under different climatic scenario [35,36,37,38,39, 40, 41].

A. Artificial Neural Network (ANN) in Agriculture

Human brain's biological neural processes form the basis for the ANN structure. A model can be developed using interrelationships of correlated variables. These symbolically represent the interconnected processing neurons or nodes of the human brain that are used. To establish model prediction, large number of input and output example are used to develop a formula to ascertain the relationship [42]. With little a priori knowledge of the functional relationship, nonlinear relationships, which are overlooked by other prediction methods, can be determined [43]. An ANN model requires a minimum of three layers: the input, hidden and output layers. The input and output layers contain nodes that correspond to input and output variables, respectively. Data move between layers across weighted connections.

ANN在农业中的应角gronomic-based models have been developed using artificial neural networks and have been discussed by various researchers. Agronomic ANN applications include crop development modeling [44], pesticide and nutrient loss assessments [45], soil water retention estimations [46], and disease prediction [47]. A study by [48] used ANNs to classify eggs as fertile or infertile. When predicting applied nitrogen leaking below the root zone of turf grass, it was reported that an ANN model performed better than a regression model [49]. Despite the fact that regression model expends more time to be developed, an ANN model can deliver more reliable and accurate crop yield forecasting compared to other methods such as regression models [50]. Aerial pictures have been used for crop yield prediction before harvesting [51]. These photos were able to provide spatial cloud free data of the yields spectral qualities. Another study by [52] used X-ray images of apples to monitor the presence of watercores. A neural network was trained for discriminating between good and bad apples.

For the prediction of rice yield based on weather data, back propagation network was used in a study by [53]. A study by [54] described the use of artificial neural networks to predict maize yield based on rainfall, soil and other parameters and obtained a testing error of 14.8%. Another study by [55] applied principal-component based neural network model and applied it to a water level prediction. A neural network to build a hybrid prediction model and performed a prediction experiment on the precipitation in the northern, central and southern regions of the Guangxi province has also been reported [56]. Data mining techniques were employed for the monitoring of the wine

fermentation process. Taste sensors were used to obtain data from the fermentation process and classified using ANNs [57].

Work by [58] used a back propagation network to predict rice yield using climatic observation data and predicted with a maximum of 45-60kg/ha. A study by [59] used neural networks to predict rice yield based on soil parameters and achieved a testing error of 17.3%. Another study reported the use of ANN in the development of straightforward and precise estimation to predict rice yields [60]. Work by [61] developed a year round air temperature prediction models. These models predicted the horizons of 1 to 12 h using Ward-style ANNs. These models were found to be useful for general decision support by the farmers.

Another study by [62] examined the performance of ANN to predict crop yields in Nepal. The predicted outcome could be utilised for enhancing the paddy yield in Siraha district of Nepal and also in other regions where the topography and vegetation are similar to Siraha district.

Prediction of rice crop production in Phimai district, Thailand showed the effectiveness of a decision support system which used ANN [63]. The application of using ANN using feed forward back propagation for agricultural crop yield prediction is demonstrated by [64]. A study by [65] used ANN for rice crop yield prediction in Maharashtra state, India. This study concluded that ANN is a beneficial tool for crop prediction.

B. Bayesian Networks (BN) in Agriculture

The computer science technique of BN is considered to be ideal for situations enabling diagnostic reasoning on conditional dependencies to assess model structural and as well as parameter uncertainty [66]. BN is a method for representing beliefs and knowledge <u>using probabilities</u>, especially relevant for systems that are highly complex in their structure and functional interactions [67]. BN uses the probabilistic components of a framework, as opposed to deterministic comparisons to describe the connections among variables. [68].

Bayesian Networks are increasingly popular method for modeling uncertain and complex domains. Many studies have reported on the advantages and challenges using BN for complex problems such as environmental modeling [69]. The technique provides a graphical model that encodes probabilistic relationships among variables of interest [70]. This model has several advantages for data modeling when used in conjunction with statistical techniques.

A few studies have reported on the application of BN for the agricultural domain. For example, these BNs have been used for predicting yield response of winter wheat to fungicide programmes [71]. Other studies have reported its use for the estimation of genetic parameters of calving ease in first and second parties of Canadian Holsteins that has also been implemented using BN [72]. The development of a decision support system for growing malting barley without use of pesticides utilized a BN approach [73]. Applications have also been shown where BN has been developed for decisions associated with the selection of irrigation systems for irrigated dairy farms in Northern Victoria [74]. BN are also emerging as a valid approach for modeling and supporting decision making

in other related fields such as water resource management [75]. Bayesian Belief Networks (BBN) have also been used to incorporate characteristics of land managers in the modeling process and to enhance the understanding of land-use change based on the limited and disparate sources of information [76].

Other research has described the use of BN classifiers for modeling of a weed infestation risk inference system that implements a collaborative inference scheme based on rules extracted in a corn crop [77]. Another study by [78] on the best pest control decision making program also based its approach on BN. The study describes the flowchart of a BN and the principles used to calculate the conditional probabilities required in it. It proves that BN is effective tool for crop disease. A study by [79] used BN to model the crop disease coffee rust cause's premature defoliation, weakening the plant and reducing subsequent yield. Improved prediction using BN proved to reduce the use of fungicides, producing healthier quality product and decreasing both economic costs and environmental impact [79].

Research by [80] reported key indicators of rice production and consumption, correlation between them and supply demand prediction. The findings of the study suggested that there were many techniques used for forecasting purpose but FIS and Bayesian technique outperform others. Other recent studies have highlighted the application of BN in an agricultural context [81,82,83,84,85,86,87,88].

C. Support Vector Machines (SVM) in Agriculture

SVMs are one of the newest supervised machine learning techniques. A number of studies have reported on the application of SVMs in an agricultural context. For example, [89] described the use of SVMs to classify crop and [90] highlighted its usefulness in classifying smell of milk. Another study by [91] used SVMs to classify pizza sauce spread, and [92] uses SVMs for detecting weed and nitrogen stress in corn.

A number of other studies have shown the variety of uses for SVMs in agriculture. One study reported on how SVM was applied for reduction of precipitation for climate change scenarios [93]. Data mining techniques were also applied to study sound recognition. Work by [94] demonstrated the use of SVMs to classify the sound of birds and other different sounds. SVM has also been applied for the estimation of crop biophysical parameters using aerial hyper spectral observations [95]. Another application reported for SVM was the modeling of urban land use conversion. This research derived the relationship between rural-urban land use change and various factors [96]. SVM was also applied to provide insights into crop response patterns related to climate conditions by providing the features contribution analysis for agricultural yield prediction [93,97] analysed different possible changes of the weather scenarios using SVMs. Other studies have described the use of discretization based SVM for classification of agricultural datasets [98]. Another study reported how SVM was used to forecast the demand and supply of pulpwood. It was used to minimize the generalization error bound to achieve generalized performance [99]. A study by [100] showed the prediction of rice crop yield for Maharashtra state, India using SVM. This study concluded that SVM could be used for crop prediction.

D. Association Rule Mining in Agriculture

The application of Association Rule Mining have been demonstrated for many cases in agriculture for elucidating hidden patterns and associates between different climate and crop production. For example, one study demonstrated this with remotely sensed and spatial data sets in the agricultural domain [101,102]. Another study by [103] aimed to develop an effective risk management system for tracing frequent occurrences of droughts and floods which both adversely affect the sustainability of Indian agriculture. A data mining algorithm was used to discover association rules between extreme rainfall events and climatic indices. Association rules were generated for the regions of India, which shows strong relationships between the climatic indices chosen, i.e., Darwin sea level pressure, North Atlantic Oscillation, Nino 3.4 and sea surface temperature values, and the extreme rainfall events.

An exhaustive survey on association rule mining in the field of agriculture conducted by [104] found that five major algorithms were most commonly used for the discovery of association rule mining. These included apriori, partition, pincer search, dynamic item set counting and FP-Tree growth algorithm. Another study by [105] also showed the application of association rule mining algorithm for control of agricultural pests. This study found that method significantly improved the quality and efficiency of the data analysis. In another study by [106] on organic cultivated land of the three different municipalities of Thassos, Greece analysed data with a classification algorithm OneR, a k-means clustering algorithm and Apriori association rule mining algorithm. The result of this investigation indicated that organic cultivation could improve the production of olives and olive oil. The climate differences among the three municipalities were also suggested to be a factor involved in production efficacy.

A recent study by [107] presented a theoretical survey on apriori, ID3 and C4.5 algorithms to be used for another agricultural dataset. Another study by [108] provided a comparative study of various association rule mining techniques, which included A Priori Algorithm, Partition Algorithm, Pincer Search Algorithm, Dynamic Item Set (DIC) Algorithm and FP-tree growth algorithm. When these were applied in agricultural domain the best results were generated by FP-Tree growth Algorithm.

Some other recent work has highlighted the use of association rules in agriculture for pattern mining [109,110]. Another study by [111] reported on the use of aprior with WEKA for frequent pattern mining. Other research by [112] conducted another investigation on the performance evaluation of association rule mining on partitioned data. This work found that confusion could exist between the performance of multidimensional database and its association with the rule mining.

Other examples on agricultural datasets include work by [113] that implemented association rule mining for different soil types in agriculture with the aim of establishing meaningful relationships using apriori algorithm. Similar findings in a study by [114] analysed the quality of association rules by computing an interestingness measures. This research

concluded that statistical interestingness measures are really helpful for finding interesting rules among large association rules

A recent study on wheat yield losses caused by stripe rust has also been undertaken by [115]. The work described the development of a computer intelligent system that uses improved apriori algorithm to extract spatial association rules from spatial databases, which helps in a decision support system for wheat crop yellow rust disease management.

III. CONCLUSION

In this paper a review of the application of data mining techniques like artificial neural network, bayesian networks and support vector machines in agricultural domain was discussed. Various studies have reported on the application of data mining techniques. The review has outlined a number of promising techniques that have been used to understand the relations of various climate and other factors on crop production. Further investigations are needed to understand how these techniques can be used with complex agricultural datasets and used for crop yield prediction with an integration of both seasonally and spatially using GIS technologies.

REFERENCES

- W. Guo and H. Xue, "An incorporative statistic and neural approach for crop yield modelling and forecasting", Neural Commuting and Applications, 21, pp. 109-117, 2012.
- [2] W. Guo and H. Xue, "Crop yield forecasting using artificial neural networks: A comparison between spatial and temporal models", Mathematical Problems in Engineering, pp. 1-7, 2014.
- [3] R. Medar and V. Rajpurohit, "A survey on data mining techniques for crop yield prediction", International Journal of Advance Research in Computer Science and Management Studies, vol. 2, no. 9, pp. 59-64, 2014.
- [4] S. Bejo, S. Mustaffha andW. Ismail, "Application of artificial neural network in predicting crop yield: A review", Journal of Food Science and Engineering, vol. 4, pp. 1-9, 2014.
- [5] S. Dahikar and S. Rode, "Agricultural crop yield prediction using artificial neural network approach", International Journal of Innovative Research in Electrical, Electronics, Instrumentation and Control Engineering, vol. 2, no. 1, pp. 683-686, 2014.
- [6] G. Yengoh and J. Ardo, "Crop yield gaps in Cameroon", AMBIO, Springer, vol. 43, pp. 175-190, 2014.
- [7] D. Ramesh and B. Vardhan, "Analysis of crop yield prediction using data mining techniques", International Journal of Research in Engineering and Technology, vol. 4, no. 1, pp. 47-473, 2015.
- [8] J. Abello, PM. Pardalos and M. Resende, "Handbook of massive data sets", Kluwer, New York, 2002.
- [9] W. Klosgen and JM. Zytkow, "Handbook of data mining and knowledge discovery" Oxford University Press, 2002.
- [10] PM. Pardalos, LV. Boginski and A. Vazacopoulos, "Data mining in biomedicine", Springer, New York, 2007.
- [11] PM. Pardalos, P. Hansen, "Data mining and mathematical programming", American Mathematical Society, USA, 2008.
- [12] R. Rupnik, M. Kukar and M. Krisper, "Integrating data mining and decision support through data mining based decision support system", Journal of computer information systems, vol. 47, no. 3, pp.89-104, 2007.
- [13] A. Feelders, H. Daniels and M. Holseimer, "Methodological and practical aspects of data mining", Information and Management, 37, pp. 11, 2000.
- [14] X. Wu, V. Kumar, JR. Quinlan, J. Ghosh, Q. Yang, H. Motoda, GJ. McLachlan, A. Ng, B. Liu, PS. Yu, Z-H. Zhou, M. Steinbach, DJ. Hand and D. Steinberg, Top 10 algorithms in data mining, Knowledge Information Systems, vol. 14, pp. 1–37, 2008.
- [15] P. Holmgren and T. Thuresson, "Satellite remote sensing for forestry planning: a review", Scand J For Res, vol. 13, no. 1, pp. 90–110, 1998.

- [16] B. Rajagopalan and U. Lall, "A k-nearest-neighbor simulator for daily precipitation and other weather variables, Wat Res Res, vol. 35, no. 10, pp. 3089–3101, 1999.
- [17] H. Jorquera, R. Perez, A. Cipriano and G. Acuna, "Short term forecasting of air pollution episodes", In: Zannetti P (eds) Environmental modeling 4. WIT Press, UK, 2001.
- [18] K. Verheyen, D. Adriaens, M. Hermy and S. Deckers, "High-resolution continuous soil classification using morphological soil profile descriptions", Geoderma, vol. 101, pp.31–48, 2001.
- [19] GE. Meyer, JC. Neto, DD. Jones and TW. Hindman, "Intensified fuzzy clusters for classifying plant, soil, and residue regions of interest from color images", Comput Electronics Agric, vol. 42, pp. 161–180, 2004.
- [20] V. Leemans and MF. Destain, "A real time grading method of apples based on features extracted from defects", J Food Eng, vol. 61, pp. 83– 89, 2004.
- [21] D. Mehta, A. Kalola, A. Saradava, A. Yusufzai, "Rainfall variability analysis and its impact on crop productivity – A case study", Indian Journal of Agricultural Research, vol. 36, no. 1, pp. 29-33, 2002.
- [22] A. Abdullah, S. Brobst, I. Pervaiz, M. Umer and A. Nisar, "Agri Data Mining/Warehousing: Innovative Tools for Analysis of Integrated Agricultural & Meteorological Data" The IASTED International Conference on Databases and Applications, Innsbruck, Austria, 2004.
- [23] L.J. Armstrong, D. Diepeveen and R. Maddern, "The application of data mining techniques to characterize agricultural soil profiles", In Proceedings of the sixth Australasian conference on Data mining and analytics, vol. 70, pp. 85-100, 2007.
- [24] R. Jain, M. Sonajharia and V. Ramasubramaniam, "Machine Learning for Forewarning Crop Diseases" Journal of Indian Society of Agricultural Statistics, vol. 63, no. 1, pp. 11, 2009.
- [25] L.A. Romani, A.M.H. de Avila, J. Zullo Jr, R Chbeir, C. Traina Jr, A.J. Traina, "Clearminer: a new algorithm for mining association patterns on heterogeneous time series from climate data", In Proceedings of the 2010 ACM Symposium on Applied Computing, pp. 900-905, 2010.
- [26] M. Divya, T.N. Manjunath, and R.S. Hegadi, "A Study on Developing Analytical Model for Groundnut Pest Management using Data Mining Techniques", IEEE International Conference on Computational Intelligence and Communication Networks (CICN), pp. 691-696, 2014.
- [27] A. Ahamed, N. Mahmood, N. Hossain, M. Kabir, K. Das, F. Rahman, R. Rahman, "Applying data mining techniques to predict annual yield of major crops and recommend planting different crops in different districts in Bangladesh", 16th IEEE/ACIS International Conference on Software Engineering, Artificial Intelligence, Networking and Parallel/Distributed Computing (SNPD), pp. 1-6, 2015.
- [28] A. Raorane and R. Kulkarni, "Review-Role of data mining in Agriculture", International Journal of Computer Science and Information Technologies, vol. 4, no. 2, pp. 270-272, 2013.
- [29] A. Mankar and M. Burange, "Data Mining An evolutionary view of Agriculture", International Journal of Application or Innovation in Engineering and Management, vol. 3, no. 3, pp. 102-105, 2014.
- [30] R. Kalpana, N. Shanthi and S. Arumugam, "A survey on data mining techniques in agriculture", International Journal of Advances in Computer Science and Technology, vol. 3, no. 8, pp. 426-431, 2014.
- [31] G. Fathima and R. Geetha, "Agriculture crop pattern using data mining techniques", International Journal of Advanced Research in Computer Science and Software Engineering, vol. 4, no. 5, pp. 781-786, 2014.
- [32] H. Patel and D. Patel, "A brief survey of data mining techniques applied to agricultural data", International Journal of Computer Applications, vol. 95, no. 9, pp. 6-8, 2014.
- [33] M. Kaur, H. Gulati and H. Kundra, "Data mining in Agriculture on crop price prediction: Techniques and Applications", International Journal of Computer Applications, vol. 99, no. 12, pp. 1-2, 2014.
- [34] M. Geetha, "A survey on data mining techniques in Agriculture", International Journal of Innovative Research in Computer and Communication Engineering, vol. 3, no. 2, pp. 887-892, 2015.
- [35] S. Veenadhari, B. Misra and C. Singh, "Data mining techniques for predicting crop productivity – A review article", International Journal of Computer Science and Technology, vol. 2, no. 1, pp. 98-100, 2011.
- [36] A. Raorane and R. Kulkarni, "Data Mining: An effective tool for yield estimation in the agricultural sector", International Journal of Emerging Trends and Technology in Computer Science, vol. 1, no. 2, pp. 75-79, 2012

- [37] D. Ramesh and B. Vardhan, "Data mining techniques and applications to agricultural yield data", International Journal of Advanced Research in Computer and Communication Engineering, vol. 2, no. 9,pp. 3477-3480, 2013
- [38] D. Ramesh and B. Vardhan, "Analysis of crop yield prediction using data mining techniques", International Journal of Research in Engineering and Technology, vol. 4, no. 1, pp. 47-473, 2015.
- [39] R. Medar and V. Rajpurohit, "A survey on data mining techniques for crop yield prediction", International Journal of Advance Research in Computer Science and Management Studies, vol. 2, no. 9, pp. 59-64, 2014
- [40] N. Gandhi and L. Armstrong, "Rice Crop Yield forecasting of Tropical Wet and Dry climatic zone of India using data mining techniques", IEEE International Conference on Advances in Computer Applications (ICACA), pp. 357-363, 2016.
- [41] N. Gandhi and L. Armstrong, "Applying data mining techniques to predict yield of rice in Humid Subtropical Climatic Zone of India", In Proceedings of the 10th INDIACom-2016, 3rd 2016 IEEE International Conference on Computing for Sustainable Global Development, E-ISBN: 978-9-3805-4421-2, 2016.
- [42] Y. Pachepsky, D. Timlin and G. Várallyay, "Artificial neural networks to estimate soil water retention from easily measurable data", Soil Science Society of America Journal, vol. 60, no. 3, pp. 727-733, 1996.
- [43] D. Elizondo, R. McClendon and G. Hoogenboom, "Neural network models for predicting flowering and physiological maturity of soybean", Transactions of the ASAE, vol. 37, no. 3, pp. 981-988, 1994.
- [44] D. Elizondo, R. McClendon and G. Hoogenboom, "Neural network models for predicting flowering and physiological maturity of soybean", Transactions of the ASAE, vol. 37, no. 3, pp. 981-988, 1994.
- [45] C. Yang, S. Prasher, S. Sreekanth, N. Patni and L. Masse, "An artificial neural network model for simulating pesticide concentrations in soil", Transactions of the ASAE, vol. 40, pp. 1285–1294, 1997.
- [46] M. Schaap and W. Bouten, "Modeling water retention curves of sandy soils using neural networks", Water Resources Research, vol. 32, no. 10, pp. 3033–3040, 1996.
- [47] W. Batchelor, X. Yang and A. Tshanz, "Development of a neural network for soybean rust epidemics", Transactions of the ASAE, vol. 40, pp. 247– 252, 1997.
- [48] K. Das and MD. Evans, "Detecting fertility of hatching eggs using machine vision II: neural network classifiers", Trans. ASABE, vol. 35, no. 6, pp. 2035-2041, 1992.
- [49] S. Starrett and G. Adams, "Using artificial neural networks and regression to predict percentage of applied nitrogen leached under turfgrass", Communications in Soil Science and Plant Analysis, vol. 28, no. 6-8, pp. 497–507, 1997.
- [50] M. Kaul, R. Hill and C. Walthall, "Artificial neural network for corn and soybean prediction", Agricultural System, vol. 85, pp. 1-18, 2005.
- [51] G. Thomas, G. Taylor and J. Wood, "Mapping yield potential with remote sensing", Precision Agriculture, vol.1, pp. 713–720, 1997.
- [52] MA. Shahin, EW. Tollner and RW. McClendon, "Artificial intelligence classifiers for sorting apples based on watercore", Journal of Agric Eng Res, vol. 79, no. 3, pp. 265–274, 2001.
- [53] J. Liu, C. Goering and L. Tian, "A neural network for setting target corn yields", Transaction of the ASAE, vol. 44, no. 3, pp. 705-713, 2001.
- [54] M. O'Neal, B. Engel, D. Ess and J. Frankenberger, "Neural network prediction of maize yield using alternative data coding algorithms", Biosystems Engineering, vol. 83, no. 1, pp. 31-45, 2002.
- [55] K. Nakornphanom, C. Lursinsap and A. Rugchatjaroen, "Fault immunization model for elliptic radial basis function neuron", Proceedings of 9th International Conference on Neural Information Processing, pp.1027-1031, 2002.
- [56] R. Jim and M. Mary, "On the construction and training of reformulated radial basis function neural networks", IEEE Transactions on Neural Networks, 14, 2003.
- [57] Riul A Jr, de Sousa HC, Malmegrim RR, dos Santos DS Jr, Carvalho ACPLF, Fonseca FJ, Oliveira Jr ON, Mattoso LHC, "Wine classification by taste sensors made from ultra-thin films and using neural networks", Sens Actuators, B98, pp. 77–82, 2004.
- [58] S. Putch, M. Rizon, M. Juhari, J. Nor Khairah, S. Siti Kamarudin, B. Aryati, R. Nursalasawati, "Backpropagation algorithm for rice yield prediction", Proceedings of 9th of the Ninth International Symposium on Artificial Life and Robotics, Beppu, Japan, Oita, pp. 586-589, 2004.

- [59] B. Ji, Y. Sun, S. Yang and J. Wan, "Artificial neural networks for riceyield prediction in mountainous regions", Journal of Agricultural Science, vol. 145, pp. 249-261, 2007.
- [60] B. Jietal, "Artificial neural networks for rice yield prediction in mountainous regions", Journal of Agricultural Science, vol. 145, pp. 249– 261, 2007.
- [61] B. Smith, G. Hoogenboom and R. McClendon, "Artificial Neural Networks for Automated Year round Temperature Prediction", Computers and Electronics in Agriculture, vol. 68, pp. 52–6, 2009.
- [62] T. Ranjeet and L. Armstrong, "An artificial neural network for predicting crops yield in Nepal", Proceedings of the 9th Conference of the Asian Federation for Information Technology in Agriculture "ICT's for future Economic and Sustainable Agricultural Systems", Perth, Australia, pp. 376-386, 2014.
- [63] S. Jabjone and S. Wannasang, "Decision Support System Using Artificial Neural Network to Predict Rice Production in Phimai District, Thailand", International Journal of Computer and Electrical Engineering, vol. 6, no. 2, pp. 162-166, 2014.
- [64] S. Dahikar and S. Rode, "Agricultural Crop Yield Prediction Using Artificial Neural Network Approach", International Journal of Innovative Research in Electrical, Electronics, Instrumentation and Control Engineering, vol. 2, no. 1, 2014.
- [65] N. Gandhi, L. Armstrong and O. Petkar, "Rice Crop Yield Prediction using Artificial Neural Networks", In proceedings of International Conference on 2016 IEEE Technological Innovations in ICT for Agriculture and Rural Development (TIAR), 2016.
- [66] N.K. Newlands and L. Townley-Smith, "Predicting Energy Crop Yield Using Bayesian Networks", In Proceedings of the Fifth IASTED International Conference, 711, pp. 014-106, 2010.
- [67] C. Ascough, R. Maier, K. Ravalico and W. Strudley, "Future research challenges for incorporation of uncertainty in environmental and ecological decision-making", Ecological modelling, vol. 219, no. 3, pp. 383-399, 2008.
- [68] N.K. Newlands and L. Townley-Smith, "Predicting Energy Crop Yield Using Bayesian Networks", In Proceedings of the Fifth IASTED International Conference, 711, pp. 014-106, 2010.
- [69] L. Uusitalo, "Advantages and challenges of Bayesian networks in environmental modelling", Ecological modelling, vol. 203, no. 3, pp. 312-318, 2007.
- [70] D. Heckerman, "Bayesian networks for data mining", Data mining and knowledge discovery, vol. 1, no. 1, pp. 79-119, 1997.
- [71] F. Tari, "A Bayesian Network for predicting yield response of winter wheat to fungicide programmes", Computers and electronics in agriculture, vol. 15, no. 2, pp. 111-121, 1996.
- [72] M.F. Luo, P.J. Boettcher, L.R. Schaeffer and J.C.M. Dekkers, "Estimation of genetic parameters of calving ease in first and second parities of Canadian Holsteins using Bayesian methods", Livestock Production Science, vol. 74, no. 2, pp. 175-184, 2002.
- [73] K. Kristensen and I.A. Rasmussen, "The use of a Bayesian network in the design of a decision support system for growing malting barley without use of pesticides", Computers and Electronics in Agriculture, vol. 33, no. 3, pp. 197-217, 2002.
- [74] D. Robertson and Q.J. Wang, "Bayesian networks for decision analyses-An application to irrigation system selection", Animal Production Science, vol. 44, no. 2, pp. 145-150, 2004.
- [75] A. Castelletti and R. Soncini-Sessa, "Bayesian Networks and participatory modelling in water resource management", Environmental Modelling & Software, vol. 22, no. 8, pp. 1075-1088, 2007.
- [76] I. Aalders, "Modeling land-use decision behavior with Bayesian belief networks", Ecology and Society, vol.13, no. 1, pp. 16, 2008.
- [77] G.M. Bressan, V.A. Oliveira, E.R. Hruschka and M.C. Nicoletti, "Using Bayesian networks with rule extraction to infer the risk of weed infestation in a corn-crop", Engineering Applications of Artificial Intelligence, vol. 22, no. 4, pp. 579-592, 2009.
- [78] C. Bi andG. Chen, "Bayesian networks modeling for Crop Diseases", In Computer and Computing Technologies in Agriculture IV, Springer Berlin Heidelberg, pp. 312-320, 2010.
- [79] C. Perez-Ariza, A. Nicholson and M. Flores, "Prediction of coffee rust disease using bayesian networks", In Proceedings of the Sixth European Workshop on Probabilistic Graphical Models, pp. 259-266, 2012.
- [80] A. Sharma and M. Goyal, "Bayesian network model for monthly rainfall forecast", 2015 IEEE International Conference on Research in

- Computational Intelligence and Communication Networks (ICRCICN), pp. 241-246, 2015.
- [81] J. Quinn, R. Monaghan, V. Bidwell and S. Harris, "A Bayesian Belief Network approach to evaluating complex effects of irrigation-driven agricultural intensification scenarios on future aquatic environmental and economic values in a New Zealand catchment", Marine and Freshwater Research, vol. 64, no. 5, pp. 460-474, 2013.
- [82] G. Nearing, H. Gupta and W. Crow, "Information loss in approximately Bayesian estimation techniques: A comparison of generative and discriminative approaches to estimating agricultural productivity", Journal of hydrology, vol. 507, pp. 163-173, 2013.
- [83] S. Maiti and R. Tiwari, "A comparative study of artificial neural networks, Bayesian neural networks and adaptive neuro-fuzzy inference system in groundwater level prediction", Environmental earth sciences, vol. 71, no. 7, pp. 3147-3160, 2014.
- [84] S. Madadgar and H. Moradkhani, "Spatio-temporal drought forecasting within Bayesian networks", Journal of Hydrology, vol. 512, pp. 134-146, 2014
- [85] A. Sharma and M. Goyal, "Bayesian network model for monthly rainfall forecast", 2015 IEEE International Conference on Research in Computational Intelligence and Communication Networks (ICRCICN), pp. 241-246, 2015.
- [86] E. Forio, D. Landuyt, E. Bennetsen, K. Lock, T. Nguyen, D. Ambarita, S. Musonge, P. Boets, G. Everaert, L. Dominguez-Granda, L. Goethals, "Bayesian belief network models to analyse and predict ecological water quality in rivers", Ecological Modelling, vol. 312, pp. 222-238, 2015.
- [87] Y. Shin, M. Ajmal, J. Yoo, T.W. Kim, "A Bayesian Network-Based Probabilistic Framework for Drought Forecasting and Outlook", Advances in Meteorology, 2016.
- [88] N. Gandhi, L. Armstrong and O. Petkar, "Predicting Rice Crop Yield using Bayesian Networks", In proceedings of IEEE 5th International Conference on Advances in Computing, Communications and Informatics, E-ISBN: 978-1-5090-2029-4, 2016.
- [89] G. Camps-Valls, L. Gomez-Chova, J. Calpe-Maravilla, E. Soria-Olivas, JD. Martin-Guerrero and J. Moreno, "Support vector machines for crop classification using hyperspectral data", Lect Notes CompSci, 2652, pp. 134–141, 2003.
- [90] K. Brudzewski, S. Osowski and T. Markiewicz, "Classification of milk by means of an electronic nose and SVM neural network", Sens Actuators, B98, pp. 291–298, 2004.
- [91] C-J, Du and D-W, Sun, "Pizza sauce spread classification using colour vision and support vector machines", J Food Eng, vol. 66, pp. 137–145, 2005.
- [92] Y. Karimi, SO. Prasher, RM. Patel, SH. Kim, "Application of support vector machine technology for Weed and nitrogen stress detection in corn", Comput Electronics Agricult, 51, pp. 99–109, 2006.
- [93] S. Tripathi, V. Srinivas and R. Nanjundiah, "Downscaling of precipitation for climate change scenarios: a support vector machine approach", Journal of Hydrology, vol. 330, no. 3, pp. 621-640, 2006.
- [94] S. Fagerlund, "Bird species recognition using support vector machines", EURASIP J Adv Signal Processing, Article ID 38637, pp. 8, 2007.
- [95] Y. Karimi, SO. Prasher, RM. Patel and SH. Kim, "Application of support vector machine technology for Weed and nitrogen stress detection in corn", Comput Electronics Agricult, vol. 51, pp. 99–109, 2006.
- [96] B. Huang, C. Xie and R. Tay, "Support vector machines for urban growth modeling" Geoinformatica, vol. 14, no. 1, pp. 83-99, 2010.
- [97] S. Brdar, D. Culibrk, B. Marinkovic, J. Crnobarac and V. Cmojevic, "Support Vector Machines with Features Contribution Analysis for Agricultural Yield Prediction", 2011.
- [98] A. Bharadwaj, S. Dahiya and R. Jain, "Discretization based Support Vector Machine (D-SVM) for Classification of Agricultural Datasets", International Journal of Computer Applications, vol. 40, no. 1, pp. 8-12, 2012.
- [99] V. Anandhi and R. Chezian, "Support Vector Regression in forecasting", International Journal of Advanced Research in Computer and Communication Engineering, vol. 2, no. 10, pp. 4148-4151, 2013.
- [100] N. Gandhi, L. Armstrong, O. Petkar and A. Tripathy, "Rice Crop Yield Prediction in India using Support Vector Machines", In proceedings of IEEE The 13th International Joint Conference on Computer Science and Software Engineering (JCSSE), 2016.

- [101] J. Dong, W. Perrizo, Q. Ding and J. Zhou, "The application of association rule mining to remotely sensed data", In Proceedings of the 2000 ACM symposium on Applied computing, vol. 1, pp. 340-345, 2000.
- [102] Q. Ding, Q. Ding and W. Perrizo, "PARM—An efficient algorithm to mine association rules from spatial data" Systems, Man, and Cybernetics, Part B: Cybernetics, IEEE Transactions on, vol. 38, no. 6, pp.1513-1524, 2008
- [103]C.T. Dhanya and D. Nagesh Kumar, "Data mining for evolution of association rules for droughts and floods in India using climate inputs", Journal of Geophysical Research: Atmospheres, vol. 114(D2), 2009
- [104]F. Khan and D. Singh, "Association Rule Mining in the field of Agriculture: A Survey" International Journal of Scientific and Research Publications, p.329, 2014.
- [105]Bao. Xiao'an, Zhang. Na, S. Junmin, Z. Lingling and G. Jun, "Application of association rule mining algorithm based on interestingness to agricultural pest control", vol. 20, no. 4, pp. 52-57, 2011
- [106] T. Theodosiou, S. Valsamidis, G. Hatziliadis, M. Nikolaidis, "Measuring, archetyping and mining Olea europaea production data", Journal of Systems and Information Technology, vol. 14, no. 4, pp. 318-335, 2012.
- [107]R. Thakkar, M. Kayasth and H. Desai, "Rule Based and Association Rule Mining on Agriculture dataset" International Journal of Innovative Research in Computer and Communication Engineering, vol. 2, no. 11, pp. 6381-6384, 2014.
- [108]J. Bhatia and A.Gupta, "Mining of Quantitative Association Rules in Agricultural Data Warehouse: A Road Map", International Journal of Information Science and Intelligent System, vol. 3, no.1, pp. 187-98, 2014
- [109]N. Gandhi and L. Armstrong, "Assessing impact of seasonal rainfall on rice crop yield of Rajasthan, India using Association Rule Mining", In proceedings of IEEE 5th International Conference on Advances in Computing, Communications and Informatics, E-ISBN: 978-1-5090-2029-4, 2016.
- [110]N. Gandhi and L. Armstrong, "Assessing the Impact of Seasonal Rainfall Patterns on Indian Rice Production using Association Rule Mining", World Congress on Computers in Agriculture, Asian Federation for Information Technology in Agriculture (AFITA), pp.50, ISBN: 979-11-958140-0-8, 2016.
- [111]P. Tanna and Y. Ghodasara, "Using Apriori with WEKA for Frequent Pattern Mining", International Journal of Engineering Trends and Technology (IJETT), vol. 12, no. 3, pp. 127-131, 2014.
- [112]J. Ashri and P. Kaur, "Performance evaluation of association rule mining on partitioned data", International Journal of Enterprise Computing and Business Systems" ISSN (Online): 2230-8849,vol. 4, no. 2, pp. 1-7, 2014.
- [113] M. Geetha, "Implementation of Association Rule Mining for different soil types in Agriculture", International Journal of Advanced Research in Computer and Communication Engineering, vol. 4, no. 4, pp. 520-522, 2015.
- [114]J. Manimaran and T. Velmurugan, "Analysing the Quality of Association Rules by Computing an Interestingness Measures", Indian Journal of Science and Technology, vol. 8, no. 15, 2015.
- [115]S. Bisht and S.D. Samantaray, "Extracting Spatial Association Rules in Remotely Sensed Data of Yellow Rust Disease in Wheat Crop at Udham Singh Nagar", 1st International Conference on Next Generation Computing Technologies, Dehradun, India, pp. 847-851, 2015.