Brain Report

Patient Information

Name: Kyle Rosebrook DOB: 01/16/1986 Gender: Male Assessment Date: 07/07/2022

Mental Health Screening Scores

Standardized tools to identify risk of mental health conditions

Depression (PHQ-9)	10 (Moderate)	At Risk: No
Anxiety (GAD-7)	5 (Mild)	At Risk: No
Substance Abuse (CAGE-AID)		At Risk: Yes
PTSD (PC-PTSD-5)		At Risk: No
ADHD		At Risk: No
Social Anxiety		At Risk: No
Sleep Apnea		At Risk: No

Brain Assessment Results

Assessment scores are shown as a percentile from 1-100. A higher percentile reflects a stronger performance.

LOW 1-30 | MEDIUM 30-70 | HIGH 70-100

Emotion - Automatic and nonconscious processes that help us minimize danger and maximize reward.

Capacity	Scale	Score	
Emotion Awareness	Medium	54th	1 — 100
Non-conscious Negativity Bias	Medium	33rd	1 — 100
Emotional Flexibility	N/A	25th	1 — 100

Feeling - Your conscious experience of emotions that relies on feedback from your body reactions.

Stress Control	Low	27th	1 — 100
Anxiety Control	Medium	46th	1 — 100
Depressive Mood Control	Medium	34th	1 — 100


Cognition - Selective awareness of information processing so we can know and remember.

Memory	Medium	39th	1 — 100
Focus	Medium	59th	1 — 100
Planning	Medium	46th	1 — 100
Processing Speed	High	99th	1 — 100
Flexibility	Low	17th	1 — 100
Recall Memory	Low	12th	1 — 100


Self Control - Shaping and planning of our thinking and emotion over time to maximize our well being.			
Conscious Negativity Bias	Low	24th	1 — 100
Resilience	Low	2nd	1 — 100
Social Connectivity	Low	27th	1 — 100

Percentile values from 1-100


Where you score compared to others: Normalized scores based on gender, age, and years of education.



Emotion:



Thinking:




Percentile values from 1-100
Where you score compared to others: Normalized scores based on gender, age, and years of education.
Low
1-30
Medium
30-70
High
70-100
Focus (Controlled Attention)
Trials Completed in 20 seconds (congruent)
Reaction Time (congruent)
Trials Completed in 20 seconds (incongruent)
Reaction Time (interference)
Flexibility
Accuracy
Completion Time
Connection Time
Working Memory
Maximum Recall Span
Correct Trials
Recall Memory
Immediate Recall Accuracy
Delayed Recall Accuracy
Planning
Completion Time
Overall Accuracy
Overrun Errors
Paths Completed
Path Learning Time

DETAILED RESULTS

Readiness to Change (0-10 scale)

On a scale of 0 to 10, how ready are you to make changes to address the problems bothering you?	0 1 2 3 4 5 6 7 8 9 (10)

PHQ-9 + SelfHarm

In the last 2 weeks, how often have you been bothered by the following problems?	Not at all	Several days	More than half the days	Nearly every day
Little interest or pleasure in doing things		✓		
Feeling down, depressed, or hopeless		✓		
Trouble falling or staying asleep, or sleeping too much		✓		
Feeling tired or having little energy		✓		
Poor appetite or overeating		✓		
Feeling bad about yourself, or that you are a failure or have let yourself or your family down			✓	
Trouble concentrating on things, such as reading the newspaper or watching television			✓	
Moving or speaking so slowly that other people have noticed. Or the opposite - being so fidgety or restless that you have been moving around a lot more than usual		✓		
Thoughts that you would be better off dead, or of hurting yourself	✓			

GAD-7

In the last 2 weeks, how often have you been bothered by the following problems?	Not at all	Several days	More than half the days	Nearly every day
Feeling nervous, anxious, or on edge		✓		
Not being able to stop or control worrying		✓		
Worrying too much about different things		✓		
Trouble relaxing		✓		
Being so restless that it is hard to sit still		✓		
Becoming easily annoyed	✓			
Feeling afraid, as if something awful might happen	✓			

CAGE-AID

	Yes	No
Have you felt you ought to cut down on your drinking or drug use?	✓	
Have people annoyed you by criticizing your drinking or drug use?	✓	
Have you ever felt bad or guilty about your drinking or drug use?	✓	
Have you ever had a drink or used drugs first thing in the morning to steady your nerves or to get rid of a hangover (eye-opener)?	✓	

PC-PTSD-5

Have you ever experienced any event(s) that was/were so frightening or upsetting that, in the past month you have:	Yes	No
Had nightmares about the event(s) or thought about the event(s) when you did not want to?		✓
Tried hard not to think about the event(s) or went out of your way to avoid situations that reminded you of the event(s)	✓	
Been constantly on guard, watchful, or easily startled?		✓
Felt numb or detached from people, activities, or your surroundings?		✓
Felt guilty or unable to stop blaming yourself or others for the event(s) or any problems the event(s) may have caused?	✓	

Sleep apnea

This score is determined by the answer to this question plus additional risk factors of age, gender, and BMI	Never	Rarely	1-2 times a week	3-4 times a week	5-7 times a week	Don't know
In the last month, have you experienced or have you been told that while sleeping you have breathing difficulties, snorting, gasping or loud snoring?		✓				

Social Anxiety

	Yes	No
In the past few months, did you feel afraid or uncomfortable doing things in front of other people? For example, speaking among a group of people, speaking to strangers or shop staff, speaking to authority figures, or eating or writing in front of others?		✓

Stress Control

Please answer the following questions:	Never or Almost Never	Occasionally	Often	Very Often	Always or Almost Always
During the past week, I found it hard to wind down		✓			
During the past week, I was jumpy and agitated	✓				
During the past week, I found it difficult to relax				✓	
During the past week, I was intolerant of anything that kept me from getting on with things		✓			
During the past week, I was rather touchy		✓			
During the past week, I felt that I was using a lot of nervous energy		✓			
During the past week, I tended to over-react to situations		✓			

Anxiety Control

	Never or Almost Never	Occasionally	Often	Very Often	Always or Almost Always
During the past week, I was worried about situations in which I might panic and make a fool of myself		✓			
During the past week, I was aware of dryness of my mouth	✓				
During the past week, I experienced breathing difficulty (e.g. excessive rapid breathing, breathlessness in the absence of physical exertion)	✓				
During the past week, I experienced trembling (e.g. in the hands)	✓				
During the past week, I was aware of the action of my heart in the absence of physical exertion	✓				
During the past week, I felt scared without any good reason	✓				
During the past week, I felt I was close to panic		✓			

Conscious Negativity Bias

Please indicate how much each of the following statements applied to you during the past week:	Never or Almost Never	Occasionally	Often	Very Often	Always or Almost Always
During the past week, I was stressed, with my nerves on edge		✓			
During the past week, I lost hope and wanted to give up when something went wrong			✓		
During the past week, I tended to over-react to situations		✓			
During the past week, I felt down-hearted and blue		✓			
During the past week, I felt I wasnâ€™t worth anything		✓			

Depressive Mood Control	Never or Almost Never	Occasionally	Often	Very Often	Always or Almost Always
During the past week, I couldn't seem to experience any positive feeling at all		✓			
During the past week, I found it difficult to work up the initiative to do things		✓			
During the past week, I felt that I had nothing to look forward to	✓				
During the past week, I was unable to become enthusiastic about anything	✓				
During the past week, I felt that life was meaningless		✓			

Resilience Please indicate how much each of the following statements applied to you during the past week:	Never or Almost Never	Occasionally	Often	Very Often	Always or Almost Always
During the past week, there were times when people couldnâ€™t rely on me as much as they should have been able to		✓			
During the past week, I was always successful at completing my tasks even if I had more tasks than others		✓			
During the past week, I felt very satisfied with the way I look and act	✓				
During the past week, I responded best to positive feedback about myself		✓			
During the past week, when receiving negative comments about myself, I looked for positive things to counter balance those comments		✓			

Social Connectivity	Never or Almost Never	Occasionally	Often	Very Often	Always or Almost Always
I love socializing and chatting to other people			✓		
I can sense the mood of a group and how other people feel without them having to tell me		✓			
People tell me that I am a sensitive and understanding person		✓			
I take the initiative and introduce myself to strangers				✓	
I try to have close relationships with people	✓				

VALIDATION OF TOOLS AND RESOURCES

PHQ-9, Range 1-27:
The PHQ-9 is a commonly used brief self-report tool to assess for the presence of depression symptoms over the past 2 weeks. It is largely based on DSM-IV depression criteria and has 88% sensitivity and 88% specificity for diagnosis of Major Depressive Disorder. High reliability and validity have been established (Kroenke et al., 2001).
Total Score Depression Severity: 1-4 Minimal, 5-9 Mild, 10-14 Moderate, 15-19 Moderately severe, 20-27 Severe.

GAD-7, Range 1-21:
The GAD-7 is a commonly used brief self-report tool for assessing the presence of anxiety symptoms associated with Generalized Anxiety Disorder over the past 2 weeks. It has 89% sensitivity and 82% specificity for diagnosing Generalized Anxiety Disorder when compared to independent clinician diagnosis. It will also detect anxiety symptoms that are common across different types of anxiety disorders.
Total Score Severity: 1-4 Minimal, 5-9 Mild, 10-14 Moderate, 15-21 Severe.

CAGE-AID:
The CAGE-AID is a brief 4-item self-report tool to assess for substance abuse (covering both alcohol and drug abuse). It has a 79% sensitivity and 77% specificity when compared to independent clinician diagnosis. A score of 2 or greater is considered clinically significant.

PC-PTSD-5:
The PC-PTSD-5 is a brief 5-item self-report tool to screen for the presence of probable PTSD and is based on DSM-5 criteria. This is intended as a screening tool for PTSD, with a positive result requiring further clinical assessment before a diagnosis can be made. A score of 3 or greater constitutes a positive screen that requires further assessment.

ADHD:
Derived from the method for ADHD diagnosis using cognitive assessment outlined in Williams et al., (2010), the current screening method maintains an acceptable level of sensitivity and specificity, while removing the cognitive test measures that were in the original Williams et al. method but are not included within the Total Brain cognitive assessment. It has a sensitivity of 67% and specificity of 74%.

Sleep Apnea:
Sleep apnea is screened by asking patients three questions about sleep disordered breathing from the Maislin Sleep Questionnaire with body mass index, age, and gender to predict the presence of sleep apnea. This method has a sensitivity of 81% and specificity of 73%.

Social Anxiety:
Social anxiety is screened by asking patients two questions derived from the Dalrymple & Zimmerman (2008) approach, which combines an initial screening question asking about feeling afraid or uncomfortable doing things in front of other people, with a list of specific social fears that improves the accuracy of screen. This method has a sensitivity of 100% and specificity of 63%.

Stress Control:

Emotional Flexibility:
Your primary nonconscious emotional bias, one of four: adaptable, driven, positive, cautious.
Usefulness: Allows for a focused exploration of one's state of being, self-reflection.

Adaptable
Emotional brain networks are structured to have a non-conscious bias towards fear-inducing events, so as to respond rapidly at the time but then to not remain in fear-states the rest of the time. The Adaptable category reflects people who have a well-functioning and flexible non-conscious emotional brain network that is good at both responding to fear and then quickly relaxing back to normal. People who are adaptable have an overall profile of good cognition and emotional functioning compared to others.

Driven
Emotional brain networks are structured to have a non-conscious bias towards fear-inducing events, so as to respond rapidly at the time but then to not remain in fear-states the rest of the time. The Driven category reflects people who have a nonconscious bias to respond to anger in other people as much as they do to fear. They tend to be high performing and highly motivated, but are more impulsive and less socially connected to others.

Positive
Emotional brain networks are structured to have a non-conscious bias towards fear-inducing events, so as to respond rapidly at the time but then to not remain in fear-states the rest of the time. The Positive category reflects people who have a strong bias towards prioritizing positive cues from others as much as they do to fear cues, and they may be strongly motivated to receive only positive feedback from others. They tend to function well emotionally as they are highly positive and resilient, but at the cost of cognitive performance, where they are slower than others across most tasks and may favor getting the task right over trying to do it quickly.

Cautious
Emotional brain networks are structured to have a non-conscious bias towards fear-inducing events, so as to respond rapidly at the time but then to not remain in fear-states the rest of the time. The Cautious category reflects people who also have a strong nonconscious bias towards sadness as well as fear. These people tend to have higher levels of anxiety and depressive mood, and may also have some subsequent impact on cognitive performance.

Emotion Awareness:
The capacity to accurately identify emotional cues in others and yourself (such as fear and happiness).
Usefulness: Associated with effective decision making, contextually appropriate feelings, and social connections.

Non-Conscious Negativity Bias:
The relative capacity to properly read and react to primary emotions, particularly negative ones, such as fear, anger, and sadness.
Usefulness: Triggering the “safety first” response and risk aversion.

REFERENCES

An extreme magnification of negativity, a low mood, a loss of joy, and a sense of hopelessness.
Usefulness: A warning sign to resolve the underlying frustration.

Anxiety Control:
A difficulty relaxing, nervousness, becoming easily upset, irritable, impatient and over reactive.
Usefulness: It can help highlight real or perceived threats.

Depressive Mood Control:
The activation of the fight-flight reflex in response to physical, emotional, or social threats.
Usefulness: Can highlight a negativity bias about oneself, relationships, work and the world. It also highlights a context distortion that should be corrected.

Memory:
The capacity to use information to plan and organize behavior to meet a goal.
Usefulness: Associated with high-functioning individuals who can multi-task, utilization of working memory, and the capacity to recall previous outcomes for optimal decision making.

Focus:
The capacity to sustain attention without being distracted.
Usefulness: Associated with the completion of any task.

Planning:
The capacity to use information to plan and organize behavior to meet a goal.
Usefulness: Associated with functions where on-the-fly decision making is critical. Decisiveness.

Processing Speed:
The capacity for speed of sensori-motor function and information processing.
Usefulness: measures variability of response times

Flexibility
The capacity for information processing efficiency.
Usefulness: measures completion time

Recall Memory:
Short-term working memory capacity.
Usefulness: measures total number of digits recalled

Conscious Negativity Bias:
The capacity for enhancing positivity and not magnifying negative threats.
Usefulness: Associated with choosing the best response to negative threats.

Resilience:
The capacity for forming and sustaining authentic social networks.
Usefulness: Building successful stress-recovery strategies.

Social Connectivity:
The capacity for forming and sustaining authentic social networks.
Usefulness: Associated with appropriate, warm, and open bonding with people and social circles.

Brown RL & Rounds LA (1995). Conjoint screening questionnaires for alcohol and other drug abuse: Criterion validity in a primary care practice. Wisconsin Medical Journal, 94(3): 135-140.

Kroenke K, Spitzer R L, Williams J B (2001). The PHQ-9: validity of a brief depression severity measure. Journal of General Internal Medicine, 16(9): 606-613.

Kroenke, K. & Spitzer, R.L. (2002). The PHQ-9: A new depression and diagnostic severity measure. Psychiatric Annals, 32: 509-521.

Spitzer RL, Kroenke K, Williams JB, Lowe B (2006). A brief measure for assessing generalized anxiety disorder: the GAD-7. Archives of internal medicine, 166(10):1092-1097.

Williams, L. M., Hermens, D. F., Thein, T., Clark, C. R., Cooper, N. J., Clarke, S. D., et al. (2010). Using brain-based cognitive measures to support clinical decisions in ADHD. Pediatric Neurology, 42(2), 118-126.

Dalrymple KL & Zimmerman M (2008). Screening for Social Anxiety Disorder in psychiatric outpatients. Comprehensive Psychiatry, 49(4): 399-406.

Maislin G, Pack AI, Kribbs NB, Smith PL, Schwartz AR, Kline LR, Schwab RJ, & Dinges DF (1995). A survey screen for prediction of apnea. Sleep, 18(3): 158-166.

Williams LM, Hermens DF, Thein T, Clark CR, Cooper NJ, Clarke SD, Lamb C, & Gordon E (2010). Using brain-based cognitive measures to support clinical decisions in ADHD. Pediatric Neurology, 42 (2): 118-26.